

New and little-known stonefly species of the *Rhopalopsole vietnamica* ‘western assemblage’ group (Plecoptera, Leuctridae) from China

Mengyu Li¹, Bingli Wang¹, Ding Yang², Dávid Murányi³, Weihai Li¹, Hongliang Wang¹

¹ Henan International Joint Laboratory of Taxonomy and Systematic Evolution of Insecta, Henan Institute of Science and Technology, Xinxiang, Henan 453003, China

² Department of Entomology, China Agricultural University, 2 Yuanmingyuan West Road, Beijing 100193, China

³ Department of Zoology, Eszterházy Károly Catholic University, Leányka u. 6, Eger H-3300, Hungary

Corresponding author: Weihai Li (lwh7969@163.com)

Abstract

We examined Chinese stonefly specimens of the *Rhopalopsole vietnamica* ‘western assemblage’ group. A new species from Hainan Province, *R. bawanglinga* Li, Li & Yang, **sp. nov.** is described and illustrated from male and female adults, and it is compared to closely related taxa. The hitherto unknown female of *R. hainana* Li & Yang, 2010 is described. Morphological evidence is presented for the identity of *R. dentiloba* Wu, 1973, on the basis of topotypes from Yunnan Province, southwestern China.

Key words: female description, new species, redescription, *Rhopalopsole bawanglinga*, *Rhopalopsole dentiloba*, *Rhopalopsole hainana*, taxonomy



Academic editor: Sven Bradle

Received: 11 March 2023

Accepted: 23 September 2023

Published: 6 November 2023

ZooBank: <https://zoobank.org/E6FB3FB1-F61C-4FC6-B249-508F02FCB0F3>

Citation: Li M, Wang B, Yang D, Murányi D, Li W, Wang H (2023) New and little-known stonefly species of the *Rhopalopsole vietnamica* ‘western assemblage’ group (Plecoptera, Leuctridae) from China. ZooKeys 1183: 125–137. <https://doi.org/10.3897/zookeys.1183.103288>

Copyright: © Mengyu Li et al.
This is an open access article distributed under the terms of the CC0 Public Domain Dedication.

Introduction

The stonefly genus *Rhopalopsole* Klapálek, 1912 is one of the largest leuctrid genera, with about one hundred valid species, distributed in the Oriental and East Palaearctic regions (DeWalt et al. 2023). Over sixty species have been recorded from China by Chen et al. (2022), Chen and Du (2017), Klapálek (1912), Li et al. (2010a, 2010b, 2011, 2017, 2022), Li and Yang (2010, 2011, 2012), Mo et al. (2018), Qian et al. (2014), Qian and Du (2011, 2012a, 2012b, 2013, 2017), Sivec et al. (2008), Wu (1949, 1973), Yang and Du (2021a, 2021b, 2022), Yang et al. (2021, 2022), Yang et al. (2004, 2009), Yang and Li (2006) and Yang and Yang (1991a, 1991b, 1993, 1994, 1995a, 1995b). The ‘western assemblage’ of the *R. vietnamica* group was recognized by Sivec et al. (2008), and currently, thirteen species belong to this group (Wu 1973; Yang and Yang 1993; Sivec et al. 2008; Li and Yang 2010; Li et al. 2017, 2010b, 2022; Qian and Du 2017; Yang et al. 2022; Yang and Du 2022). In this contribution, two species of this species group were described or supplemented, including a new species, *R. bawanglinga* Li, Li & Yang, sp. nov. from Hainan Province in southern China, and the hitherto unknown female of *R. hainana* Li & Yang, 2010. Additionally, we provide a redescription and new images for *R. dentiloba* Wu, 1973, a newly included species of this group, based on topotypes.

Table 1. World list of species belonging to the *Rhopalopsale vietnamica* ‘western assemblage’ group. Abbreviation: M, male; F, female.

Species	Known life stages	Distribution
<i>R. amamiensis</i> Kawai, 1967	M, F	Japan: Kagoshima Prefecture, Okinawa
<i>R. assamensis</i> Sivec & Harper, 2008	M, F	India: Assam
<i>R. bawanglinga</i> sp. nov.	M, F	China: Hainan
<i>R. brevidigitata</i> Qian & Du, 2017	M	China: Yunnan
<i>R. dentiloba</i> (Wu, 1973)	M, F	China: Yunnan
<i>R. fengyangshanensis</i> Yang, Shi & Li, 2009	M	China: Fujian, Zhejiang
<i>R. furcospina</i> (Wu, 1973)	M, F	China: Guangxi, Shaanxi, Sichuan, Zhejiang
<i>R. hainana</i> Li & Yang, 2010	M	China: Hainan
<i>R. nanlinga</i> Yang & Du, 2022	M	China: Guangdong
<i>R. sinensis</i> Yang & Yang, 1993	M	China: Chongqing, Fujian, Gansu, Guangxi, Guizhou, Hubei, Ningxia, Shaanxi, Zhejiang; Vietnam: Laocai
<i>R. singiplatta</i> Yang & Du, 2022	M, F	China: Sichuan
<i>R. sipirokana</i> Sivec & Harper, 2008	M, F	Sipirok: Sumatra
<i>R. vietnamica</i> Sivec & Harper, 2008	M, F	Vietnam: Thinhua
<i>R. yajunae</i> Li & Yang, 2010	M	China: Zhejiang
<i>R. yunnana</i> Sivec & Harper, 2008	M, F	China: Yunnan

Material and methods

The specimens were collected by hand and preserved in 75% ethanol. Most of the studied materials are deposited in the Entomological Museum of China Agricultural University, Beijing, China (**CAU**). Additional specimens are deposited in the Insect Collection of Henan Institute of Science and Technology, Xinxiang, Henan Province, China (**HIST**), as indicated in the text. The color illustrations were made using an Imaging Source CCD attached to a Leica M205FA dissecting microscope. The morphological terminology follows that of Sivec et al. (2008). The association of sexes of *Rhopalopsale hainana* is based on the type locality and barcoding data.

The maps were downloaded from Standard Map Service (<http://bzdt.ch.mnr.gov.cn/>) and edited by Photoshop 2020, map number is GS (2019)1823.

Total genomic DNA was extracted from thorax muscle using the TIANamp Genomic DNA kit (Tiangen Biotechnology, Beijing, China) following the manufacturer’s protocol. The universal primers LCO1490 and HCO2198 (Folmer et al. 1994) were used to amplify the target COI gene region. Polymerase chain reaction (PCR) program was as follows: initial denaturation at 95 °C for 60 s, followed by 40 cycles of 30 s at 95 °C, 50 s at 40–60 °C, and 60 s at 65 °C, and a final extension phase of 65 °C for 10 min. The PCR products were confirmed by 1% agarose gel electrophoresis and then sent to Sangon Biotechnology Co. Ltd. (Shanghai, China) for DNA sequencing. The primary sequences were assembled using Contigexpress software (Vector NTI Informax). Genetic distances between sequences were calculated in MEGA v.5.2 and bootstrap analysis was conducted using 1000 replicates. The GenBank accession numbers for the three specimens are in Table 2.

Table 2. China, Hainan Province specimens used for COI based identification, genetic differentiation, and circumscription analyses of three *Rhopalopsole* species.

Species	Sex	Catalog number	GenBank accessions	Geographic coordinates
<i>R. hainana</i>	1♀	J61	OR435219	19°4'52.32"N, 109°31'10.92"E
<i>R. hainana</i>	1♀	J64	OR435220	19°4'52.32"N, 109°31'10.92"E
<i>R. hainana</i>	1♂	J66	OR435221	19°4'52.32"N, 109°31'10.92"E

Results and discussion

Family Nemouridae Billberg, 1820

Genus *Rhopalopsole* Klapálek, 1912

Rhopalopsole bawanglinga Li, Li & Yang, sp. nov.

<https://zoobank.org/F26FAD81-D6AA-44BA-99B7-88067CD05AB1>

Figs 1, 2

Type material. Holotype: male (CAU), CHINA: Hainan Province, Changjiang County, Bawangling National Forest Park, Dong'er Station, 22.X.2007, 19°15'1.44"N, 109°2'2.4"E, 1000 m, leg. D. Yang. **Paratypes:** 1 female (CAU): same data as holotype; 2 females (HIST), CHINA: Hainan Province, Ledong County, Jianfengling, Tianchi–Mingfenggu, 3.VIII.2016, 18°44'49.2"N, 108°50'57.12"E, 890 m, leg. Weihai Li, Rongfeng Wang.

Diagnosis. Male adult of this species is characterized by the lateral projections of the tergum 10 being bifurcate with the upper spine longer than the lower one in lateral view. The cercus bears a stout dorsoapical spine, and the epiproct is subtrapezoid with deep, circularly incised anterior margin in dorsal view. Females are diagnosed by the posterior margin of sternum 7 forming a narrowly produced pregenital plate, posteromedial portion slightly bilobed.

Description. Adult habitus. Forewing length is 6.8 mm in the male, 7.0–7.4 mm in the females. Head brown to dark brown, slightly wider than pronotum; compound eyes black; antennae and mouthparts brownish to brown. Pronotum brown with dark rugosities; wings subhyaline with darker veins; legs brown. Abdomen brown.

Male (Figs 1, 3). Tergum 9 (Figs 1A, 3A) weakly sclerotized except anterior margin and lateral portions, medial 1/3 with a large trapezoidal membranous area surrounded by lateral sclerotized portion and ornamented posteromedial margin which terminates in a thin, produced sclerite with tiny granules. Sternum 9 (Figs 1B, 3B) longer than wide, distal portion with distinct trapezoidal subgenital plate, the plate about as long as wide, with a subapical constriction; vesicle dark brown, ovum-shaped in ventral view, tongue-like in lateral view, covered with dense hairs. Tergum 10 (Figs 1A, C, 3A, C) with well sclerotized, bifurcate lateral projections, terminating in two acute, parallel-sided points in dorsal aspect, the upper spine is longer than lower spine in lateral view. Central plate (Figs 1A, D, 3A, D) sclerotized and trilobed, lateral lobes subtriangular, together with a circular anteromedial incision of the medial lobe resemble flying wings; medial lobe darkly pigmented, with a small knob-like projection. Transverse bars slightly elevated medially, triangular, posterior margin and inner portion sclerotized. Cercus barely 3× longer than wide, curved dorsally, with a stout subapical spine. Epiproct (Figs 1A, E, 3A, E) strongly sclerotized and darkly




Figure 1. *Rhopalopsole bawanglinga* Li, Li & Yang, sp. nov. (male) **A** terminalia, dorsal view **B** terminalia, ventral view **C** terminalia, lateral view **D** terminalia, dorsolateral view **E** epiproct dorsal view. **ep**: epiproct; **ls**: lower spine; **us**: upper spine.




Figure 2. *Rhopalopsale bawanglinga* Li, Li & Yang, sp. n. (female) **A** terminalia, ventral view (apex of pregenital plate enlarged) **B** terminalia, lateral view.

sclerotized laterally, subtrapezoid with deep, circularly incised anterior margin in dorsal view, longer than wide. Subanal lobe (Figs 1B, 3B) distinctly sclerotized basally, lateral margins and apex membranous, ventral furrows vague.

Female (Fig. 2A, B). Posterior margin of sternum 7 forming a triangularly produced pregenital plate; plate with sclerotized bilobed distal tip. Sternum 8 membranous except a slender, transverse posterior strip.

Etymology. The specific name refers to the Bawangling National Forest Park.

Distribution. China (Hainan).

Remarks. The new species is a typical member of the *R. vietnamica* group, western assemblage sensu Sivec et al. (2008). The male of the new species seems closely related to *R. hainana* from Hainan and *R. cestroidea* Li & Yang, 2017 (in: Li et al. 2017) from Guangxi, as they share a similar wide epiproct. However, it can be differentiated from both by bifurcate lateral projections on tergum 10, with the upper spine being longer than the lower one in lateral view, and the presence of a concave anterior margin of the epiproct. In *R. hainana* and *R. cestroidea*, the terminal bifurcation of lateral projections on tergum 10 is of equal length in lateral view, and the epiproct has a truncate or convex anterior margin. Additionally, the cercus of *R. cestroidea* lacks a spine, and that of *R. hainana* has a sharp subapical spine, whereas the cercal spine in the new species is stout. The female is distinctive due to the bilobed distal tip of the pregenital plate.

***Rhopalopsale hainana* Li & Yang, 2010**

Fig. 4

Rhopalopsale hainana Li & Yang, 2010: 59 (original description).

Material examined. 11 males, 5 females (CAU), CHINA: Hainan Province, Baisha, Yinggeling, Hongxin Village, 23–24.V.2007, 19°12'22.32"N, 109°33'14.4"E, 250 m, leg. J.X. Liu; 1 female (CAU), CHINA: Hainan Province, Lingshui County, Diaoluo Mountain, 1.VII.2011, 18°43'31.08"N, 109°52'8.4"E, 294 m, leg. W.H.




Figure 3. *Rhopalopsole bawanglinga* Li, Li & Yang, sp. n. (male) **A** terminalia, dorsal view **B** terminalia, ventral view **C** terminalia, lateral view **D** terminalia, dorsolateral view **E** epiproct dorsal view. **ep**: epiproct; **ls**: lower spine; **us**: upper spine.




Figure 4. *Rhopalopsole hainana* Li & Yang, 2010 (female) **A** terminalia, ventral view (apex of pregenital plate enlarged) **B** terminalia, lateral view.

Li; 2 females (CAU), CHINA: Hainan Province, Baisha, Yinggeling, 29.X.2011, 19°4'52.32"N, 109°31'10.92"E, 1191 m, leg. W.H. Li.

Description. Female. (Fig. 4A, B) Pregenital plate on sternum 7 sclerotized, distally produced into strongly sclerotized triangular lobe with sinuous lateral margins, covered with long hairs; dorsoventrally flattened in lateral view. Sternum 8 with a trapezoid, transverse band across the posterior half of the segment.

Remarks. Genetic distance refers to the genetic divergence between species and can be used to compare the genetic similarity between difference species (Wang et al. 2013); generally, more than 98% of congeneric species or sister species pairs have greater than 0.02 sequence divergence (Johns and Avise 1998; Hebert et al. 2003; Park et al. 2011). In our study, pairwise distances of the *Rhopalopsole* species were 0–0.9%; lower than the 2% threshold considered for a rough differentiation between intraspecific and interspecific distances (Zhou et al. 2009). The genetic distance of COI between J66 and J61 is 0 and between J66 and J64 is 0.9% (Table 3). The combination of morphology and molecular data suggest the female is *R. hainana*.

Table 3. Genetic distances among sequenced specimens. Upper (right) triangular matrix: standard deviation; lower (left) triangular matrix: genetic distances.

	J61 F <i>R. hainana</i>	J64 F <i>R. hainana</i>	J66 M <i>R. hainana</i>
J61 F <i>R. hainana</i>		0.006	0.0
J64 F <i>R. hainana</i>	0.009		0.006
J66 M <i>R. hainana</i>	0.0	0.009	

We compared the holotype with the specimens from Baisha County, which originated from near the type locality. The hitherto unknown female resembles the female of *R. bawanglinga*, but the tip of the pregenital plate is not bilobed, and the transverse band on sternum 8 is more pronounced.

***Rhopalopsole dentiloba* Wu, 1973**

Figs 5, 6

Rhopalopsole dentiloba Wu, 1973: 105 (original description).

Material examined. 4 males, 1 female (CAU), CHINA: Yunnan Province, Xishuangbanna, Meng shimron botanical garden, 22.IV.2007, 21°55'9.12"N, 101°16'6.96"E, 550 m, leg. D. Hui. 1 male (HIST), CHINA: Yunnan Province,




Figure 5. *Rhopalopsole dentiloba* Wu, 1973 (male) **A** terminalia, dorsal view **B** terminalia, ventral view **C** terminalia, lateral view **D** terminalia, dorsolateral view. **us**: upper spine.




Figure 6. *Rhopalopsole dentiloba* Wu, 1973 (female) **A** terminalia, ventral view **B** terminalia, lateral view.

Xishuangbanna, Mengla, Wangtianshu scenic spot, 9.V.2009, 22°1'53.76"N, 100°52'32.88"E, 840 m, leg. X.S. Yang.

Remarks. We examined several specimens from Xishuangbanna, Yunnan Province, which is the same location where *R. dentiloba* was described. The original description and illustrations did not provide sufficient details for identification. Therefore, we discuss herein the distinctive morphological characters of our specimens. This species is a typical member of the *R. vietnamica* group, western assemblage sensu Sivec et al. (2008). The male tergum 9 (Fig. 5A, C, D) has a small triangular, upraised sclerotized process on hind margin. Tergum 10 (Fig. 5A, B) bears a small bifurcate spine at each posterolateral corner in dorsal and lateral aspects, but the inner/lower spine is not easily observed. In lateral view (Fig. 5C, D), unbranched, sclerotized lateral projections are seen. The epiproct is small, hook-like. The minute teeth of the subanal lobe described for the holotype are possibly the distinct furrows seen in the males observed in this study. The identity of these specimens is primarily based on a similar female and their proximity to the type locality (Mengsong town or Mengsong village in this town). The female (Fig. 6A, B) is identical to the original illustration. However, doubts may arise about association of the original female and the male holotype, as tropical Yunnan is well known for its high biodiversity, where more *Rhopalopsole* species may co-occur.

Conclusion

Thirteen species were previously recorded in the *R. vietnamica* ‘western assemblage’ group, as specified in Table 1. *Rhopalopsole dentiloba* was included in the *R. shaanxiensis* species group but is now transferred to the *R. vietnamica* ‘western assemblage’ group due to its great similarities to *R. sinensis* Yang & Yang, 1993. Considering the geographical (Fig. 7) differences between populations of *R. sinensis*, molecular methods should be applied to confirm the status of *R. dentiloba* in the future. With the description of the new taxon and the inclusion of *R. dentiloba*, the number of species in this group in China is now up to eleven, and more new species may await discovery.




Figure 7. Distribution of species of the *Rhopalopsole vietnamica* ‘western assemblage’ group of China. City records shaded in grey.

Acknowledgments

The specimens used in this study were partially provided by National Animal Collection Resource Center, Beijing, China.

Additional information

Conflict of interest

The authors have declared that no competing interests exist.

Ethical statement

No ethical statement was reported.

Funding

We acknowledge the support of the National Natural Science Foundation of China (No. 31000977) and the Program for Science and Technology Innovation Talents in Universities of Henan (No. 21HASTIT042).

Author contributions

Ding Yang, Dávid Murányi and Weihai Li: Review, commentary or revision; Bingli Wang and Hongliang Wang: Color illustrations acquisition; Mengyu Li: Manuscript writing.

Author ORCIDs

Mengyu Li  <https://orcid.org/0000-0002-7950-4142>

Bingli Wang  <https://orcid.org/0000-0002-1080-549X>

Ding Yang  <https://orcid.org/0000-0002-7685-3478>

Dávid Murányi  <https://orcid.org/0000-0002-3907-5590>

Hongliang Wang  <https://orcid.org/0000-0002-9758-1328>

Data availability

All of the data that support the findings of this study are available in the main text.

References

- Chen ZT, Du YZ (2017) Description of the first new species of Leuctridae (Plecoptera) from Jiangsu Province of China. *Zootaxa* 4319(1): 185–193. <https://doi.org/10.11646/zootaxa.4319.1.11>
- Chen J, Wang HL, Li WH (2022) Two new species of the genus *Rhopalopsole* (Plecoptera: Leuctridae) from China. *Zootaxa* 5129(4): 569–578. <https://doi.org/10.11646/zootaxa.5129.4.6>
- DeWalt RE, Maehr MD, Hopkins H, Neu-Becker U, Stueber G (2023) Plecoptera Species File Online. Version 5.0/5.0. <http://Plecoptera.Species.File.org> [Accessed 14 February 2023]
- Folmer O, Black M, Hoeh W, Lutz R, Vrijenhoek R (1994) DNA primers for amplification of mitochondrial cytochrome c oxidase subunit I from diverse metazoan invertebrates. *Molecular Marine Biology and Biotechnology* 3: 294–299. <https://www.ncbi.nlm.nih.gov/pubmed/7881515>
- Hebert PDN, Ratnasingham S, DeWaard JR (2003) Barcoding animal life: Cytochrome c oxidase subunit 1 divergences among closely related species. *Proceedings of the Royal Society B, Biological Sciences* 270(Supplement): S96–S99. <https://doi.org/10.1098/rsbl.2003.0025>
- Johns GC, Avise JC (1998) A comparative summary of genetic distances in the vertebrates from the mitochondrial cytochrome b gene. *Molecular Biology and Evolution* 15(11): 1481–1490. <https://doi.org/10.1093/oxfordjournals.molbev.a025875>
- Klapálek F (1912) H. Sauter's Formosa-Ausbeute. *Plecoptera. Entomologische Mitteilungen* 1: 342–351. <https://doi.org/10.5962/bhl.part.25903>
- Li WH, Yang D (2010) Two new species of *Rhopalopsole vietnamica* group (Plecoptera: Leuctridae: *Rhopalopsole*) from China. *Zootaxa* 2614(1): 59–64. <https://doi.org/10.11646/zootaxa.2614.1.6>

- Li WH, Yang D (2011) A new species of *Rhopalopsole dentata* group from China (Plecoptera: Leuctridae). Transactions of the American Entomological Society 137(1 & 2): 199–201. <https://doi.org/10.3157/061.137.0112>
- Li WH, Yang D (2012) A review of *Rhopalopsole magnicerca* group (Plecoptera: Leuctridae) from China. Zootaxa 3582(1): 17–32. <https://doi.org/10.11646/zootaxa.3582.1.2>
- Li WH, Li GL, Yang D (2010a) A new species of the genus *Rhopalopsole* (Plecoptera: Leuctridae) from Sichuan, China. Dong Wu Fen Lei Xue Bao 35: 310–312.
- Li WH, Lu WY, Yang D (2010b) A new species of *Rhopalopsole vietnamica* group (Plecoptera: Leuctridae) from China. Entomological News 121: 163–164. <https://doi.org/10.3157/021.121.0207>
- Li WH, Kong FB, Yang D (2011) A new species of *Rhopalopsole shaanxiensis* species group (Plecoptera: Leuctridae) from China. Dong Wu Fen Lei Xue Bao 36: 246–248.
- Li WH, Murányi D, Gamboa M, Yang D, Wantanabe K (2017) New species and records of Leuctridae (Plecoptera) from Guangxi, China, on the basis of morphological and molecular data, with emphasis on *Rhopalopsole*. Zootaxa 4243(1): 165–176. <https://doi.org/10.11646/zootaxa.4243.1.8>
- Li WL, Zhang Y, Yang D, Li WH (2022) *Rhopalopsole magnispina* sp. nov., a new species of Leuctridae (Plecoptera) from China, with notes on *R. horvati* Sivec & Harper, 2008. Zootaxa 5141(1): 094–100. <https://doi.org/10.11646/zootaxa.5141.1.9>
- Mo RR, Wang GQ, Yang D, Li WH (2018) Two new species of the *Rhopalopsole magnicerca* group (Plecoptera: Leuctridae) from China. Zootaxa 4388(3): 444–450. <https://doi.org/10.11646/zootaxa.4388.3.11>
- Park DS, Footitt R, Maw E, Hebert PD (2011) Barcoding bugs: DNA-based identification of the true bugs (Insecta: Hemiptera: Heteroptera). PLoS ONE 6(4): e29419. <https://doi.org/10.1371/journal.pone.0018749>
- Qian YH, Du YZ (2011) Two new species of genus *Rhopalopsole* (Insecta, Plecoptera, Leuctridae) from China. ZooKeys 154: 1–7. <https://doi.org/10.3897/zookeys.154.2234>
- Qian YH, Du YZ (2012a) A new stonefly species, *Rhopalopsole tricuspis* (Leuctridae: Plecoptera), and three new records of stoneflies from the Qinling Mountains of Shaanxi, China. Journal of Insect Science 12(47): 1–9. <https://doi.org/10.1673/031.012.4701>
- Qian YH, Du YZ (2012b) Two new species of the genus *Rhopalopsole* (Plecoptera: Leuctridae) from China. Entomotaxonomia 34: 17–21. <https://doi.org/10.11646/zootaxa.3582.1.2>
- Qian YH, Du YZ (2013) Two new species of *Rhopalopsole* (Plecoptera, Leuctridae) from China. Zootaxa 3599(6): 588–592. <https://doi.org/10.11646/zootaxa.3599.6.7>
- Qian YH, Du YZ (2017) Two new species of *Rhopalopsole* (Plecoptera: Leuctridae) from China. Zootaxa 4273(2): 296–300. <https://doi.org/10.11646/zootaxa.4273.2.10>
- Qian YH, Li HL, Du YZ (2014) A study of Leuctridae (Insecta: Plecoptera) from Shennongjia, Hubei Province, China. The Florida Entomologist 97(2): 605–610. <https://doi.org/10.1653/024.097.0236>
- Sivec I, Harper PP, Shimizu T (2008) Contribution to the study of the Oriental genus *Rhopalopsole* (Plecoptera: Leuctridae). Scopolia 64: 1–122.
- Wang YY, Liu XY, Winterton SL, Yan Y, Chang WC, Yang D (2013) Comparative mitogenomic analysis reveals sexual dimorphism in a rare montane lacewing (Insecta: Neuroptera: Ithonidae). PLoS ONE 8(12): e83986–e83986. <https://doi.org/10.1371/journal.pone.0083986>
- Wu CF (1949) Sixth supplement to the stoneflies of China (Order Plecoptera). Peking Natural History Bulletin 17: 251–256.

- Wu CF (1973) New species of Chinese stoneflies (Order Plecoptera). *Acta Entomologica Sinica* 16: 97–118.
- Yang YB, Du YZ (2021a) Three new synonyms of *Rhopalopsole sinensis* Yang & Yang, 1993 (Plecoptera: Leuctridae). *Zootaxa* 5040(4): 528–538. <https://doi.org/10.11646/zootaxa.5040.4.4>
- Yang YB, Du YZ (2021b) A new synonym with a new species of *Rhopalopsole* (Plecoptera: Leuctridae) from China. *Zootaxa* 5072(5): 485–492. <https://doi.org/10.11646/zootaxa.5072.5.5>
- Yang YB, Du YZ (2022) A new synonym species with description of a new species of *Rhopalopsole* from China (Plecoptera: Leuctridae). *The European Zoological Journal* 89 (1): 183–189. <https://doi.org/10.1080/24750263.2022.2034998>
- Yang D, Li Z (2006) A new species of *Rhopalopsole* (Plecoptera: Leuctridae) from China. *Entomological News* 117(4): 433–435. [https://doi.org/10.3157/0013-872X\(2006\)117\[433:ANSORP\]2.0.CO;2](https://doi.org/10.3157/0013-872X(2006)117[433:ANSORP]2.0.CO;2)
- Yang D, Yang CK (1991a) New species of Plecoptera from Hubei. *Journal of Hubei University* 13: 369–372.
- Yang D, Yang CK (1991b) One new species of *Rhopalopsole* from Zhejiang. *Journal of Zhejiang Forestry College* 8: 78–79.
- Yang D, Yang CK (1993) New and little-known species of Plecoptera from Guizhou Province (III). *Entomotaxonomia* 15: 235–238.
- Yang D, Yang CK (1994) Two new species of *Rhopalopsole* (Plecoptera: Leuctridae) from Shaanxi. *Entomotaxonomia* 16: 189–191.
- Yang D, Yang CK (1995a) Plecoptera: Leuctridae. In: *Insects and Macrofungi of Gutianshan, Zhejiang*, Zhejiang Science and Technology Press, Hangzhou, 20–24.
- Yang D, Yang CK (1995b) Plecoptera: Leuctridae. In: *Insects of Baishanzu Mountain, Eastern China*, China Forestry Publishing House, Beijing, 61–62.
- Yang D, Li WH, Zhu F (2004) Two new species of *Rhopalopsole* (Plecoptera: Leuctridae) from China. *Entomological News* 115: 279–282.
- Yang D, Shi L, Li WH (2009) A new species of *Rhopalopsole* from Oriental China (Plecoptera: Leuctridae). *Transactions of the American Entomological Society* 135(1 & 2): 193–195. <https://doi.org/10.3157/061.135.0207>
- Yang YB, Du YZ, Zhu BQ (2021) Two new species of *Rhopalopsole* (Plecoptera: Leuctridae) from Wuyi Mountains of China. *Zootaxa* 5071(1): 143–150. <https://doi.org/10.11646/zootaxa.5071.1.8>
- Yang YB, Du YZ, Zhu BQ (2022) A new species of *Rhopalopsole* Klapálek, 1912 with preliminary female identifications in this genus. *Zootaxa* 5133(2): 270–278. <https://doi.org/10.11646/zootaxa.5133.2.7>
- Zhou X, Adamowicz SJ, Jacobus LM, DeWalt RE, Hebert PD (2009) Towards a comprehensive barcode library for Arctic life-Ephemeroptera, Plecoptera, and Trichoptera of Churchill, Manitoba, Canada. *Frontiers in Zoology* 6(1): 30. <https://doi.org/10.1186/1742-9994-6-30>