

Leucosolenia qingdaoensis sp. nov. (Porifera, Calcarea, Calcaronea, Leucosolenida, Leucosoleniidae), a new species from China

Yan-Ling Chu^{1,2,3,4*}, Lin Gong^{1,2,3,4*}, Xin-Zheng Li^{1,2,3,4}

1 Institute of Oceanology, Chinese Academy of Sciences, Qingdao 266071, China **2** University of Chinese Academy of Sciences, Beijing 100049, China **3** Center for Ocean Mega-Science, Chinese Academy of Sciences, Qingdao, 266071, China **4** Laboratory for Marine Biology and Biotechnology, Pilot National Laboratory for Marine Science and Technology (Qingdao), Qingdao 266237, China

Corresponding author: *Xin-Zheng Li* (lixzh@qdio.ac.cn)

Academic editor: *P. Stoev* | Received 9 October 2019 | Accepted 4 December 2019 | Published 22 January 2020

<http://zoobank.org/DE8D65F1-1739-4377-8BC4-E42F84E77B08>

Citation: Chu Y-L, Gong L, Li X-Z (2020) *Leucosolenia qingdaoensis* sp. nov. (Porifera, Calcarea, Calcaronea, Leucosolenida, Leucosoleniidae), a new species from China. ZooKeys 906: 1–11. <https://doi.org/10.3897/zookeys.906.47164>

Abstract

A new species of Leucosoleniidae, *Leucosolenia qingdaoensis* sp. nov., is described. This new species was collected in a scallop-breeding pond from the Yellow Sea and preserved in 75% ethanol. This sponge consists of a dense reticulation of ascon tubes, with the surface minutely hispid and the consistency soft and fragile.

Spiculation of the new species consists of diactines, which are smooth, straight or sometimes slightly curved, triactines of two types, and tetractines with short and curved apical actines; spiculation also slightly overlaps and is somewhat irregularly assembled. Together these form a thin layer of skeleton, with a small number of cells, which results in a transparent, white sponge. As a typical asconoid feature, all internal cavities of the sponge are lined with choanocytes, and there is no fully developed inhalant system. Comparisons with other *Leucosolenia* reported from the Pacific Ocean are also made.

Keywords

Sponge, taxonomy, Yellow Sea

* Contributed equally as first authors.

Introduction

The family Leucosoleniidae is characterised by a branched and rarely anastomosed cormus and asconoid aquiferous system; there is neither a common cortex nor a delimited inhalant or exhalant aquiferous system (Minchin 1900). The family includes three genera (Borojevic et al. 2002): *Ascyssa* Haeckel, 1872, *Ascute* Dendy & Row, 1913, and *Leucosolenia* Bowerbank, 1864. They can be easily distinguished by their skeletons: the skeleton of *Ascyssa* contains only diactines; the skeleton of *Ascute* exhibits giant longitudinal diactines forming a continuous layer on the external surface, and includes triactines and tetractines; and the skeleton of *Leucosolenia* lacks any of these obvious characteristics in the above two genera. Instead, the skeleton of *Leucosolenia* is characterised by being composed of diactines, triactines and/or tetractines, without a reinforced external layer on the tubes.

The genus *Leucosolenia* comprises 40 living species worldwide (Van Soest et al. 2019), of which only three species, *L. microspinata* Longo, 2009, *L. salpinx* Van Soest, 2017, and *L. parthenopea* Sarà, 1953, were named after 1950; 11 species were described by Haeckel between 1870 and 1872. The literature of this genus is relatively old, and the descriptions contained therein of the species of *Leucosolenia* were simple, almost without details and illustrations of the body shapes and spicules. Thus, a taxonomic revision of this genus is very difficult, and to date, no worldwide revision of the genus has been made.

The localities of the 15 known species of *Leucosolenia* recorded from the Pacific Ocean are shown in Figure 1. Seven species (*L. eleanor* Urban, 1906, *L. minuta* Tanita, 1943, *L. mollis* Tanita, 1941, *L. pyriformis* Tanita, 1943, *L. serica* Tanita, 1942, *L. tenera* Tanita, 1940, and *L. ventosa* Hôzawa, 1940) were reported from the Japanese waters (Sagimi Sea, Wakayama Prefecture, Onagawa Bay, Mie Prefecture, Matsushima Bay, Izushima, Wagu Miye Prefecture, respectively). *Leucosolenia macquariensis* Dendy, 1918 was reported from the west coast of Macquarie Island; *L. australis* Brøndsted, 1931 was reported from Comau Fjord; *L. albatrossi* Hôzawa, 1918 was reported from Copper Island and the Komandorski Islands; *L. echinata* Kirk, 1893 and *L. rosea* Kirk, 1896 were reported from New Zealand; *L. lucasi* Dendy, 1891 was reported from Port Phillip Heads, Australia; *L. nautilia* Laubenfels, 1930 was reported from California, USA; and *L. feuerlandica* Tanita, 1942 was reported from Tierra del Fuego, South America. The *Leucosolenia* species reported from the coasts of Japan account for most species. The type specimens of new species were found in the Yellow Sea, very close to Japan.

Materials and methods

The specimens were collected in a scallop-breeding pond from the Yellow Sea and were preserved in 75% ethanol. Two specimens were deposited in the Marine Biological Museum of the Institute of Oceanology in the Chinese Academy of Sciences (IOCAS), Qingdao, China.




Figure 1. Distribution of *Leucosolenia* **A** location in the Pacific Ocean **B** detail of the type locality in the Japanese coast: (1) Komandorski Islands (*L. albatrossi* Hôzawa, 1918); (2) Comau Fjord (*L. australis* Brøndsted, 1931); (3) Cook Strait, Poverty Bay, Kawakawa (*L. echinata* Kirk, 1893); (4) Francisco Bay, California; Sukumo ôsima, Kôti Prefecture, Sagimi Sea (*L. eleanor* Urban, 1906); (5) Tierra del Fuego (*L. feuerlandica* Tanita, 1942); (6) Port Phillip Heads, Australia, and New Zealand (*L. lucasi* Dendy, 1891); (7) Macquarie Island (*L. macquariensis* Dendy, 1918); (8) Wakayama Prefecture (*L. minuta* Tanita, 1943); (9) Onagawa Bay (*L. mollis* Tanita, 1941); (10) Monterey Bay, California (*L. nautilia* Laubenfels, 1930); (11) Mie Prefecture (*L. pyriformis* Tanita, 1943); (12) New Zealand (*L. rosea* Kirk, 1896); (13) Yodomi, Sagami Sea (*L. serica* Tanita, 1942); (14) Matsushima Bay, Onagawa Bay, Izushima (*L. tenera* Tanita, 1940); (15) Wagu Miye Prefecture (*L. ventosa* Hôzawa, 1940); (*) Qingdao (*L. qingdaoensis* sp. nov.).

For examination of the spicules, a small piece of specimen was cut and placed in a 1.5 mL microcentrifuge tube to which 1000 µL of sodium hypochlorite solution was added (Kersken et al. 2016). The mixture was then vortexed, placed at environmental temperature, and vortexed occasionally during incubation until it was completely lysed. Next, the sample was centrifuged at 8000 rpm for 2 min, the supernatant was poured off, 1000 µL of distilled water was added, and the sample was again centrifuged at 8000 rpm for 2 min. This procedure was repeated four times, then the spicules were washed three times with 96% ethanol and then the spicules were preserved in the third ethanol solution.

Scanning Electron Microscopy (SEM) was performed with a Hitachi S3400N. Preserved spicules for SEM were adhered to stubs with double-sided carbon conductive tape and coverslip. After dehydration, the spicules were coated with gold in a Hitachi MC1000 (LOPES 2018).

Measurements of at least 20 spicules of each type were performed using an optical microscope (Nikon Eclipse Ni) with a micrometric eyepiece. The length from the tip to the base and the thickness at the base of each actine were measured. The reported numbers refer to the range of measurements for each spicule type. Photographs were taken with a stereomicroscope (Zeiss Stemi 2000-c) and an optical microscope (Nikon Eclipse Ni-U) equipped with a digital camera to evaluate difference between the length of the unpaired and paired actines of each type of triactine. For comparison with the new species, we only selected those species of *Leucosolenia* reported from the Pacific Ocean.

Results

Systematics

Class Calcarea Bowerbank, 1862

Subclass Calcaronea Bidder, 1898

Order Leucosolenida Hartman, 1958

Family Leucosoleniidae Minchin, 1900

Genus *Leucosolenia* Bowerbank, 1864

Leucosolenia qingdaoensis sp. nov.

<http://zoobank.org/F0C1D83E-3940-4D4C-B0BB-60A379ED507D>

Figs 1–4; Tables 1, 2

Type material. *Holotype*: MBM181606, scallop-breeding pond on southeastern Shandong Peninsula, China, June 1988, 0–0.3 m depth, collected by Shue Li, 35°58'N, 120°11'E. *Paratype*: MBM181476, Zhonggang, Qingdao, China, 7 June 1984, 0–0.6 m depth, 36°06'N, 120°21'E.

Type locality. Qingdao, Yellow Sea.

Etymology. The name is derived from the type locality, Qingdao, China.

Description. The sponge is arborescent, consisting of many thin-walled tubes, which are copiously ramified but never anastomosed. The sponge occurs as growth form. The oscula are terminal on erect tubes. The color of the sponge is white after being preserved in alcohol and in vivo. The external walls of the tubes are hairy, with diactines protruding at right or oblique angles from the body; the surface is minutely hispid, and the consistency is soft and fragile. The holotype measures 21.32 × 3.38 mm (height × width). The wall of the sponge body is very thin, and there is no fully developed inhalant system, the gap between the skeleton and the cell on the wall arrange evenly (Fig. 2F); only a small amount of cells is distributed on the thin sponge skeleton (Fig. 2C–F), which is a typical asconoid feature. All internal cavities of the sponge are lined by choanocytes.

Skeletal arrangement. The skeleton consists of multifarious diactines, sagittal triactines of two types, sagittal tetractines with bent apical actines and triactine-like basal actines; together these form the wall of the ascon-type sponge body.

In the apical osculum (Fig. 2C, E), there are paired actines of triactines and tetractines, some additional tangential diactines, together forming a clear line dividing the apical oscula, and some radial diactines projecting beyond the apical osculum with different length.

In the sponge body (Fig. 2C, E), the triactines and tetractines are regularly arranged, their paired actines are parallel to the apical oscula, and the unpaired actines point downward, with slight folding allowed, but never overlapping; in contrast to the triactines and tetractines, the diactines are arranged more irregularly but generally point downward.

In the root-like structures (Fig. 2D, F), the arrangement of triactines and tetractines is the same as that in the body, but the arrangement of diactines is different; most of them tangentially project beyond the surface, which results in the surface having a slightly hispid appearance.




Figure 2. *Leucosolenia qingdaoensis* sp. nov. **A** holotype **B** paratype **C** detail of oscula (stereo microscope) **D** detail of root-like structures (stereo microscope) **E** detail of oscula (optical microscope) **F** detail of root-like structures (optical microscope); arrowhead pointing at the ostium.

By observing the sponge tissue taken from different parts, it is clear that as the diameter of the tubes decreases, the contents of small diactines and small triactines increase. This observation can suggest that in the growth zone spiculogenesis is more intense.

Spicules. Diactines. There is only one type of diactine (Fig. 3A1–3), though the diactines vary in size and shape, their width varies from 24 μm to 61 μm , the length of diactines vary from 43 μm to 421 μm but half of the diactines present a length of 200–300 μm (Fig. 4). The shapes of the diactines are straight or slightly curved in different directions. The variation in *Leucosolenia* is very common and considerable.

Triactines. Two types of triactines are present, with actines straight or undulated. Their ends are generally sharp or asymmetrical (Fig. 3B1–2). The paired actines are slightly curved. Some deformations are present.

Type 1: triactines with paired actines longer than unpaired actines (Fig. 3B1): unpaired actines 42–105 \times 3–5 μm ; paired actines 63–105 \times 3–5 μm .

Type 2: triactines with unpaired actines longer than paired ones (Fig. 3B2): unpaired actines 76–129 \times 3–4 μm ; paired actines 60–104 \times 3–4 μm .

Tetractines. A relatively small number of tetractines are observed, approximately 10 per 100 spicules, with straight and fusiform actines (Fig. 3C1–2). The tetractines are similar to triactines but with the addition of apical actines, the apical actines are fairly stout and short, sharply pointed and curved: unpaired actines 93–119 \times 2–5 μm ; paired actines 50–93 \times 2–5 μm ; apical actines 11–29 \times 2–5 μm .

Remarks. Three species described by Tanita (*L. minuta*, *L. pyriformis*, and *L. serica*) exhibit only regular (equiangular and equiradiate) spicules. This characteristic does not fit the description of *Leucosolenia*, *L. qingdaoensis* sp. nov. can be easily differentiated from the 12 species of *Leucosolenia* reported from the Pacific Ocean. The skeletal compositions of these species are shown in Table 1.




Figure 3. Spicules of *Leucosolenia qingdaoensis* sp. nov. (holotype) A1–3 = diactines; B1 = triactines of type 1; B2 = triactines of type 2; C1–2 = tetractines.

Table 1. Spicules dimensions of *Leucosolenia* Bowerbank, 1864 in the Pacific Ocean. Measurements are reported in μm .

	Triactines		Tetractines			Diactines	References
	Unpaired	Paired	Unpaired	Paired	Apical	Length/Width	
	Length/Width	Length/Width	Length/Width	Length/Width	Length/Width		
<i>L. albatrossi</i>	70–90/8	80–100/8	70–90/8	80–100/8	40–60/6	70–90/8	Hózwawa 1918
	60–90/8	130–240/8	60–90/8	130–240/8	40–60/6–8	–	
<i>L. australis</i>	69–122/6	66–106/6	66–119/6	69–99/7	27–41/4	41–49/1	Azevedo et al. 2009
	–	–	–	–	–	63–347/7	
<i>L. echinata</i>	100/10	130/10	130/15	150/15	70/15	240–730/10–5	Kirk 1893
<i>L. eleanor</i>	80/7	80/7	140/9	140/9	140/9	105/4	Laubenfels 1932
	140/7	140/7	–	–	–	434/9	
<i>L. feuerlandica</i>	50–70/12–18	70–95/12–18	60–70/8–10	75–90/8–10	40–50/6–8	70–90/4–6	Tanita 1942
	60–70/8–10	75–90/8–10	–	–	–	–	
<i>L. lucasi</i>	100/5	70/5	100/5	70/5	<70/5	160/5	Dendy 1891
<i>L. macquariensis</i>	980/9	980/9	980/9	980/9	–	140/6	Dendy 1918
	–	–	–	–	–	90/5	
<i>L. minuta</i>	130–175/14–18	130–175/14–18	60–75/8–10	60–75/8–10	50–60/7–10	–	Tanita 1943
	60–75/8–10	60–75/8–10	–	–	–	–	
<i>L. mollis</i>	70–130/6–8	90–140/6–8	70–130/6–8	90–140/6–8	35–55/6	230–400/7–10	Tanita 1941
<i>L. nautilia</i>	140/9	140/9	140/9	140/9	30/8	400/10	Laubenfels 1932
	–	–	–	–	–	140/4	
	–	–	–	–	–	1000/20	
<i>L. pyriformis</i>	180–190/12–18	180–190/12–18	180–190/12–18	180–190/12–18	150–260/8–15	630–800/40–55	Tanita 1943
<i>L. rosea</i>	300/70	300/70	140/10	140/10	110/8	–	Kirk 1896
	200/18	200/18	–	–	–	–	
<i>L. serica</i>	140–210/7–8	140–210/7–8	140–210/7–8	140–210/7–8	90–135/8–10	–	Tanita 1942
<i>L. tenera</i>	80–180/7–10	90–210/7–10	80–180/7–10	90–210/7–10	30–10/6–8	200–530/8–12	Tanita 1940
<i>L. ventosa</i>	100–120/10	85–100/10	–	–	–	–	Hózwawa 1940
	150–180/20–25	140–150/20–25	–	–	–	–	
	100–120/10–14	70–90/10–14	–	–	–	–	
<i>L. qingdaoensis</i>	42–104/3–5	63–105/3–5	93–119/2–5	50–93/2–5	11–29/2–5	43–422/4–7	Present paper
sp. nov.	76–129/3–4	60–104/3–4	–	–	–	–	

Table 2. Spicules measurements of *Leucosolenia qingdaoensis* sp. nov. (holotype).

	length(μm)				width(μm)				n
	min	mean	max	sd	min	mean	max	sd	
Diactines	43	219	422	93	1	4	7	1.7	50
Triactines 1									
paired	63	83	105	9	3	4	5	0.8	50
unpaired	42	66	105	13	–	–	–	–	–
Triactines 2									
paired	60	79	104	11	3	3	4	0.4	50
unpaired	76	102	129	15	–	–	–	–	–
Tetractines									
paired	50	77	93	12	2	4	5	0.8	20
unpaired	93	104	119	11	–	–	–	–	–
apical	11	21	29	6	–	–	–	–	–

The new species exhibits one type of diactine. In *L. ventosa* and *L. rosea*, there is no record of diactines, and in *L. mollis* and *L. nautilia*, there are two types of diactines. The triactines of *L. ventosa* are 2–8 times thicker than those in the new species; the triactines of *L. rosea* are 10–35 times thicker than in the new species; and *L. mollis* only has one type of triactine and all rays being nearly equally thick. The diactines




Figure 4. *Leucosolenia qingdaoensis* sp. nov. Size-class distribution of diactines (holotype).

of *L. nautilia* are extremely large, with a length of 1 mm and a thickness of 20 μm (Laubenfels 1932), while in the new species the diactines are less than 8 μm thick. Laubenfels (1932) gave few details on the actines, but *L. nautilia* differs from the new species by having only one type of triactine.

The difference between *L. albatrossi* and the new species is obvious. The diactines of *L. albatrossi* are club-shaped, while the diactines of the new species are spindle-shaped.

The sagittal triactines of the new species distinguish it from *L. macquariensis*, *L. tenera*, and *L. eleanor*. The new species have two types of sagittal triactines, while *L. macquariensis* and *L. tenera* only have one type of sagittal triactine, with rays of approximately equal length. *Leucosolenia eleanor* have both sagittal and regular triactines.

The new species, with slender and long diactines, the longest diactines 5 times longer than those of *L. feuerlandica*, is distinct from that species. Additionally, the triactines of the new species are sagittal, and the actines straight or undulated. However, the triactines of *L. feuerlandica* are pseudoderm sagittal and are tripod-shaped.

Leucosolenia echinata, *L. lucasi*, and *L. qingdaoensis* sp. nov. have many features in common, including their body shape, colour in alcohol, general arrangement, shape of diactines, and apical ray, but they show important differences in the shape of their triactines. The new species has two types of triactines; *L. lucasi* and *L. echinata* only have one type of triactine. The triactines of *L. lucasi* are sagittal, but the three angles are roughly equal; the triactines of *L. echinata* are generally regular, and frequently slightly sagittal, with the oral angle largest and the basal ray longest.

Key to the species of *Leucosolenia* in the Pacific Ocean

- 1 Skeleton contains only regular spicules.....2
 1a Skeleton contains sagittal spicules4
 2 Skeleton including diactines *L. pyriformis*
 2a Skeleton without diactines3
 3 Rays are stout.....*L. minuta*
 3a Rays are relatively thin *L. serica*
 4 Skeleton contains diactines, triactines, and tetractines.....5
 4a Skeleton contains triactines and tetractines *L. rosea*
 4b Skeleton contains only triactines*L. ventosa*
 5 Skeleton contains one type of diactine8
 5a Skeleton contains two types of diactines6
 6 Diactines are club-shaped *L. macquariensis*
 6a Diactines are spindle-shaped.....7
 7 Skeleton without large diactines.....*L. mollis*
 7a Skeleton including large diactines *L. nautilia*
 8 One tip of diactines has spines.....*L. australis*
 8a Diactines have no spines9
 9 Skeleton contains one type of triactine.....10
 9a Skeleton contains two types of triactines.....11
 10 Sagittal triactines with rays are of approximately equal in length.... *L. tenera*
 10a Sagittal triactines with rays are of different lengths..... *L. lucasi*
 10b Triactines are generally regular, slightly sagittal.....*L. echinata*
 11 Skeleton including tripod type of triactines..... *L. feuerlandica*
 11a Skeleton without tripod type of triactines12
 12 Diactines have one 'lance head' type ends*L. albatrossi*
 12a Diactines have two smooth and sharply pointed ends13
 13 Skeleton contains both sagittal and regular triactines*L. eleanor*
 13a Skeleton contains only sagittal triactines*L. qingdaoensis* sp.nov.

Acknowledgements

This work was supported by the National Natural Science Foundation of China (no. 41706188), the Science and Technology Basic Work Program (2014FY110500), Biological Resources Programme, Chinese Academy of Sciences KFJ-BRP-017-37, and the Scientific and Technological Innovation Project Financially Supported by Qingdao National Laboratory for Marine Science and Technology (no. 2015ASKJ01). We thank Dr Yuanyuan Sun for assistance with SEM operations, and we appreciate Dr Qinghe Liu for the help with photography. We also thank Mr Hongfa Wang, Dr Lin Ma, Dr Jixing Sui, and Mr Youwei Tzeng for providing important references needed for this study.

References

- Azevedo F, Hajdu E, Willenz P, Klautau M (2009) New records of Calcareous sponges (Porifera, Calcarea) from the Chilean coast. *Zootaxa* 2072 (1): 1–30. <https://doi.org/10.11646/zootaxa.2072.1.1>
- Borojevic R, Boury-Esnault N, Manuel M, Vacelet J (2002) Order Leucosolenida Hartman, 1958. In: Hooper JNA, Van Soest RWM (Eds) *Systema Porifera: A Guide to the Classification of Sponges*. Kluwer Academic/Plenum Publishers, New York, 1157–1184. https://doi.org/10.1007/978-1-4615-0747-5_120
- Brøndsted HV (1931) Die Kalkschwämme der Deutschen Südpolar Expedition 1901–1903. *Deutsche Südpolar-Expedition, 1901–03* 20: 1–47.
- Dendy A (1891) A monograph of the Victorian sponges, I. The organisation and classification of the Calcarea Homocoela, with descriptions of the Victorian Species. *Transactions of the Royal Society of Victoria* 3: 1–81.
- Dendy A (1918) Calcareous sponges. *Scientific Reports of the Australasian Antarctic Expedition, 1911–1914, Series C, Zoology and Botany* 6(1): 1–17.
- Hôzawa S (1918) Reports on the calcareous sponges collected during 1906 by the United States Fisheries Steamer Albatross in the Northwestern Pacific. *Proceedings of the United States National Museum* 54: 525–556. <https://doi.org/10.5479/si.00963801.54-2247.525>
- Hôzawa S (1940) On some calcareous sponges from Japan. *Science Reports of the Tohoku Imperial University* 15: 29–58.
- Kersken D, Feldmeyer B, Janussen D (2016) Sponge communities of the Antarctic Peninsula: influence of environmental variables on species composition and richness. *Polar Biology* 39(5): 851–862. <https://doi.org/10.1007/s00300-015-1875-9>
- Kirk HB (1893) Contribution to the knowledge of the New Zealand sponges. *Transactions of the New Zealand Institute* 26: 175–179. <https://biodiversitylibrary.org/page/9780834>
- Laubenfels MWde (1930) The sponges of California (abstracts of dissertations for the degree of doctor of philosophy). *Stanford University Bulletin* 5(98): 24–29.
- Laubenfels MWde (1932) The marine and fresh-water sponges of California. *Proceedings of the United States National Museum* 81(2927): 1–140. <https://doi.org/10.5479/si.00963801.81-2927.1>
- Lopes MV, Padua A, Condor-Lujan B, Klautau M (2018) Calcareous sponges (Porifera, Calcarea) from Florida: new species, new records and biogeographical affinities. *Zootaxa* 4526(2): 127–150. <https://doi.org/10.11646/zootaxa.4526.2.2>
- Minchin EA (1900) *Sponges. A treatise on zoology* 2: 1–178.
- Tanita S (1940) Calcareous sponges of Matsushima Bay. *Science Reports of the Tōhoku Imperial University* 4(15): 165–177.
- Tanita S (1941) Calcareous sponges obtained from Onagawa Bay and its vicinity. *Science Reports of the Tōhoku Imperial University* 16(3): 263–282.
- Tanita S (1942a) Report on the calcareous sponges obtained by the Zoological Institute and Museum of Hamburg. Part II. *Science Reports of the Tōhoku University* 4(17): 105–135.
- Tanita S (1942b) Calcareous sponges collected in the Kanto District, Japan. *Science Reports of the Tōhoku Imperial University* 17(1): 17–69.

- Tanita S (1943) Studies on the Calcareous of Japan. Science Reports of the Tôhoku Imperial University 17: 353–490.
- Urban F (1906) Kalifornische Kalkschwämme. Archiv für Naturgeschichte 72 (I): 33–76. [pls VI–IX: 36–55]
- Van Soest RWM, Boury-Esnault N, Hooper JNA, Rützler K, de Voogd NJ, Alvarez B, Hajdu E, Pisera AB, Manconi R, Schönberg C, Klautau M, Picton B, Kelly M, Vacelet J, Dohrmann M, Díaz M-C, Cárdenas P, Carballo JL, Ríos P, Downey R (2019) World Porifera Database. *Leucosolenia* Bowerbank, 1864. World Register of Marine Species. <http://www.marinespecies.org/> [Accessed on: 2019-05-07]