

Diversity of flesh flies (Sarcophagidae, Sarcophaginae) of pond habitats in rural areas in the Croatian part of Baranja

Stjepan Krčmar¹

¹ Department of Biology, Josip Juraj Strossmayer University of Osijek, Cara Hadrijana 8/A, HR-31000 Osijek, Croatia

Corresponding author: Stjepan Krčmar (stjepan@biologija.unios.hr)

Academic editor: Marc De Meyer | Received 21 January 2023 | Accepted 7 April 2023 | Published 24 April 2023

<https://zoobank.org/CB922F29-F5C6-4AB0-928D-5C69B15F3B0F>

Citation: Krčmar S (2023) Diversity of flesh flies (Sarcophagidae, Sarcophaginae) of pond habitats in rural areas in the Croatian part of Baranja. ZooKeys 1159: 17–36. <https://doi.org/10.3897/zookeys.1159.100878>

Abstract

The diversity of grey flesh flies (Sarcophagidae: Sarcophaginae) from the Croatian part of Baranja was studied during 2019 to 2021, resulting in 37 species, of which the following are new for the area: *Ravinia pernix* (Harris, 1780); *Sarcophaga* (*Het.*) *depressifrons* Zetterstedt, 1845; *S. (Het.) filia* Rondani, 1860; *S. (Het.) haemorrhooides* Böttcher, 1913; *S. (Het.) pumila* Meigen, 1826; *S. (Het.) vagans* Meigen, 1826; *S. (Lis.) dux* Thomson, 1869; *S. (Lis.) tuberosa* Pandellé, 1896; *S. (Meh.) sexpunctata* (Fabricius, 1805); *S. (Pan.) protuberans* Pandellé, 1896; *S. (Sar.) carnaria* (Linnaeus, 1758); *S. (Sar.) variegata* (Scopoli, 1763), and *S. (Pse.) spinosa* Villeneuve, 1912. New locality records are provided for 25 species. *Sarcophaga* (*Sar.*) *croatica* Baranov, 1941 was the most abundant with 37%, followed by *S. (Sar.) lehmanni* Müller, 1922 (21%), and *S. (Pas.) albiceps* Meigen, 1826 (5%), making up 63% of all collected specimens. Most species (35) were collected in locality of Zmajevac, while the fewest (3) were collected in Bilje locality. During this study, *S. (Pse.) spinosa* was recorded in Croatia for the first time. Combined with previous records, 42 species of flesh flies have been recorded from Croatian Baranja, which comprise 27% of the flesh flies known to occur in Croatia. The total number of species of the family Sarcophagidae currently known in Croatia has increased to 156.

Keywords

Baranja, Croatia, Diptera, flesh flies, Sarcophagidae, Sarcophaginae

Introduction

The European fauna of flesh flies (Sarcophagidae) comprises 310 species (Pape et al. 2015; Whitmore et al. 2020), of which approximately 150 species occur in Central Europe (Povolný and Verves 1997) and 155 species have been recorded from Croatia (Krčmar et al. 2019; Verves and Barták 2021). Verves and Barták (2021) erroneously listed *Paragusia multipunctata* (Rondani, 1859) as new for Croatia, but this species was already listed by Krčmar et al. (2019) under the name *Taxigramma multipunctata* (Rondani, 1859). Also, under a broad concept of *Sarcophaga*, the species name *Heteronychia* (s.str.) *rohdendorfi* (Povolný & Slamečkova, 1959) represents a junior secondary homonym of *Sarcophaga rohdendorfi* Salem, 1936, *Parasarcophaga rohdendorfi* Baranov, 1938, and *Sarcophaga rohdendorfi* Baranov, 1941 (Whitmore 2011), and the valid name for this species is *Sarcophaga (Heteronychia) lederbergi* (Lehrer, 1995). Of the Croatian Sarcophagidae, most species (108) belong to subfamily Sarcophaginae, followed by Miltogramminae (38) and Paramacronychiinae (9) (Krčmar et al. 2019; Verves and Barták 2021). By reviewing the published data of Langhoffer (1920), Baranov (1928, 1929, 1938, 1941, 1942, 1943), Strukan (1964, 1967, 1970), Sisojević et al. (1989), Rucner (1994), as well as the recently published data of Whitmore (2010, 2011), Whitmore et al. (2013), and Krčmar et al. (2019), 90 species of flesh flies have been recorded in the Pannonian–Peripannonian biogeographic region of Croatia (Sarcophaginae: 68; Miltogramminae: 19; Paramacronychiinae: 3). Most of the studies were done and most species were recorded in the northern part of this region. In the recent study of Krčmar et al. (2019), 29 species of flesh flies were collected from only two localities in the Croatian part of Baranja, which is far from the total number of species considering the great landscape diversity of this area. The principal aims of this study are to give the first overview of the flesh fly fauna from subfamily Sarcophaginae in rural settlements in the Croatian part of Baranja and to emphasize the importance of these habitats for flesh fly diversity.

Materials and methods

Study area

Baranja is a Pannonian Plain region of Hungary (its northern portion) and Croatia (its southern portion). It is situated in the eastern part of Croatia and forms part of Osijek-Baranja County. Triangular in shape, it covers an area of 1147 km² between the Drava, the Danube, and the state border with Hungary (Bognar et al. 1975). The Croatian part of Baranja is a predominantly lowland area (elevation ≤ 259 m). Bansko brdo (Bansko Hill) is the most prominent part of Baranja in terms of relief and extends NE-SW for 21 km, whereas its width is much smaller (Bognar et al. 1975). The steppe, the natural vegetation that covers Bansko Hill, has completely disappeared. The belt along the Danube and Drava is a flooded area (~ 63% of the territory) with many secondary tributaries

and wetlands (Kopački rit) (Bognar et al. 1975). The Kopački rit Nature Park is one of the largest fluvial-marshy plains in Europe (Schneider-Jacoby 1994), and the basic ecological features relate to the river dynamics (Schneider-Jacoby 1994; Mihaljević et al. 1999). Forests cover ~ 20% of the Croatian part of Baranja. The climate is moderately continental with significant temperature fluctuations. The average January temperature is ~ -1.3 °C, and the July temperature is ~ 22 °C; the average annual rainfall is ~ 650 mm (Bognar et al. 1975). The Croatian part of Baranja contains 54 settlements, some of which contain pond habitats, which increasingly serve as places for recreation of people and their pets. All seven sampling sites are situated at the periphery of the settlements (Fig. 1). Geographical coordinates of these seven sampling sites are given in Table 1. Pond habitats in settlements of Kotlina, Suza, and Zmajevac are surrounded by species of willow (*Salix*) and poplar (*Populus*). Sampling sites in the settlements of Petlovac and Popovac are overgrown mainly with reeds (*Phragmites australis*), sedges (*Carex* spp.), and rush (*Typha* spp.) without forest vegetation at its edges. A similar type of vegetation is present at the pond in Darda, with the addition of water lilies (*Nymphaea alba*) and nenuphar (*Nuphar luteum*), while pond habitats in Bilje settlement are overgrown with different species of low grasses exposed to the open sun throughout the day.




Figure 1. Sampling sites for flesh flies (Sarcophagidae: Sarcophaginae) in the Croatian part of Baranja. Legend: 1. Bilje; 2. Darda; 3. Kotlina; 4. Petlovac; 5. Popovac; 6. Suza; 7. Zmajevac.

Table 1. List of sampling sites around pond habitats in Croatian part of Baranja.

No. of locality	Locality	Geographical coordinates
1	Bilje	45°36'16"N, 18°44'39"E
2	Darda	45°37'30"N, 18°41'21"E
3	Kotlina	45°47'17"N, 18°44'16"E
4	Petlovac	45°45'31"N, 18°31'41"E
5	Popovac	45°48'22"N, 18°39'34"E
6	Suza	45°46'55"N, 18°46'32"E
7	Zmajevac	45°48'03"N, 18°48'29"E

Sampling and identification

Collections of sarcophagids from pond habitats were made frequently over a period of seven months (April–October) from 2019 to 2021. During 2019 and 2020, from April to October, sampling at the pond habitat in Zmajevac was done 1–5 times a month. Samplings in 2021 were carried out once per month from May to August. Flesh flies were sampled from 1 pm to 5 pm using a standard insect sweep net to selectively collect flies resting on soil and vegetation, or attracted to animal faeces and the remains of discarded food. The collected specimens were preserved in 96% ethanol. Male terminalia were prepared for species identification following the method of Richet et al (2011). After two days in ethanol, male abdomens were dissected and soaked in a 10% KOH solution for 72 h. They were then immersed in 10% acetic acid for 1 min and rinsed with water for 1 min. They were then dehydrated in beech-wood creosote for 4 h. The phallus, pregonites and postgonites, sternite 5, cerci, and surstyli were separated from the rest of abdomen and placed into a plastic vial (volume of 2 ml) with 96% of ethanol solution. Identifications were carried out using keys for Sarcophagidae (Pape 1987; Povolný and Verves 1997; Richet et al. 2011) and descriptions and illustrations in Whitmore (2009, 2010, 2011) and Whitmore et al. (2013). Nomenclature and classification follow the Fauna Europaea database (Pape 2004). For all samples, the following information is provided: locality and date of collection, collector(s), number and sex of specimens, and depository. Also, 33 unidentified specimens from May 2018 are now identified and their data have been included in this study. One species newly recorded for the Croatian fauna is marked with a black triangle (▲). Specimens examined for this study are deposited in the collections of the Department of Biology, Josip Juraj Strossmayer University of Osijek, Osijek, Croatia (DBUO) and the Staatliches Museum für Naturkunde, Stuttgart, Germany (SMNS).

Terminology

Subgeneric names are abbreviated as follows:

Bel.	<i>Bellieriomima</i>
Ber.	<i>Bercaea</i>
Hel.	<i>Helicophagella</i>
Het.	<i>Heteronychia</i>

Kra.	<i>Krameromyia</i>
Lip.	<i>Liopygia</i>
Lis.	<i>Liosarcophaga</i>
Meh.	<i>Mehria</i>

<i>Myo.</i>	<i>Myorhina</i>	<i>Ros.</i>	<i>Rosellea</i>
<i>Pan.</i>	<i>Pandelleana</i>	<i>Sar.</i>	<i>Sarcophaga</i>
<i>Pas.</i>	<i>Parasarcophaga</i>	<i>Sac.</i>	<i>Sarcotachinella</i>
<i>Pad.</i>	<i>Pandelleisca</i>	<i>Ser.</i>	<i>Servaisia</i>
<i>Pse.</i>	<i>Pseudothyrsocnema</i>	<i>Thy.</i>	<i>Thyrsocnema</i>
<i>Rob.</i>	<i>Robineauella</i>		

Results

A total of 1293 flesh flies belonging to 37 species was collected (Table 2). *Sarcophaga (Sar.) croatica* Baranov, 1941 was the most abundant with 37%, followed by *S. (Sar.) lehmanni* Müller, 1922 (21%), *S. (Pas.) albiceps* Meigen, 1826 (5%), *S. (Sar.) baranoffi* Rohdendorf, 1937 (5%), *S. (Thy.) incisilobata* Pandellé, 1896 (5%), *S. (Lis.) emdeni* (Rohdendorf, 1969) (3%), *S. (Lis.) aegyptica* Salem, 1935 (3%), *S. (Hel.) melanura* Meigen, 1826 (3%), *S. (Het.) filia* Rondani, 1860 (3%), *S. (Ros.) aratrix* Pandellé, 1896 (2%), *S. (Hel.) noverca* Rondani, 1860 (2%), *S. (Bel.) subulata* Pandellé, 1896 (1%), *S. (Pad.) similis* Meade, 1876 (1%), and *Ravinia pernix* (Harris, 1780) (1%). These fourteen species comprised 92% of the samples; the remaining 23 species represented 8% (Table 2). Most species (35) and specimens (75%) were collected in Zmajevac (Table 2). The lowest number of species was collected in Bilje (3), whereas in other localities the number of collected species was between 8 and 11. The largest number of specimens and species was collected during 2019 (Table 3). New records for Baranja are provided for 25 species, with the record of *S. (Pse.) spinosa* representing a first record for Croatia.

List of recorded species

Subfamily Sarcophaginae Macquart, 1834

1. *Ravinia pernix* (Harris, 1780)

New records for Croatian Baranja. Zmajevac, 5.IX.2019, S. Krčmar leg. (1♂) (SMNS); same locality, 28.VI.2020, S. Krčmar leg. (2♂) (DBUO); same locality, 13.IX.2020, S. Krčmar leg. (2♂) (DBUO); same locality, 20.VI.2021, S. Krčmar leg. (1♂) (DBUO); Kotlina, 18.V.2021, S. Krčmar leg. (1♂) (DBUO); same locality, 18.VI.2021, S. Krčmar leg. (1♂) (DBUO); Popovac, 11.VI.2021, S. Krčmar leg. (1♂) (DBUO); same locality, 10.VII.2021, S. Krčmar leg. (5♂) (DBUO); Darda, 17.VI.2021, S. Krčmar leg. (1♂) (DBUO).

2. *Sarcophaga (Bellieriomima) subulata* Pandellé, 1896

Records. Zmajevac, 9.V.2018, S. Krčmar leg. (4♂) (DBUO); same locality, 22.IV.2019, S. Krčmar leg. (1♂) (DBUO); same locality, 29.VII.2019, S. Krčmar leg. (8♂) (DBUO); same locality, 8.V.2020, S. Krčmar leg. (1♂) (DBUO); same locality,

12.VI.2020, S. Krčmar leg. (1♂) (DBUO); same locality, 12.VIII.2020, S. Krčmar leg. (2♂) (DBUO); same locality, 11.V.2021, S. Krčmar leg. (1♂) (DBUO).

3. *Sarcophaga (Berceaia) africa* (Wiedemann, 1824)

Records. Zmajevac, 16.V.2019, S. Krčmar leg. (1♂) (DBUO); same locality, 24.VIII.2019, S. Krčmar leg. (1♂) (DBUO).

4. *Sarcophaga (Helicophagella) crassimargo* Pandellé, 1896

Records. Zmajevac, 17.VI.2019, S. Krčmar leg. (1♂) (SMNS); same locality, 24.VIII.2019, S. Krčmar leg. (1♂) (SMNS); same locality, 28.VI.2020, S. Krčmar leg. (3♂) (DBUO); same locality, 28.VII.2020, S. Krčmar leg. (1♂) (DBUO).

5. *Sarcophaga (Helicophagella) hirticrus* Pandellé, 1896

Records. Zmajevac, 5.IX.2019, S. Krčmar leg. (1♂) (SMNS); same locality, 8.V.2020, S. Krčmar leg. (1♂) (DBUO).

6. *Sarcophaga (Helicophagella) melanura* Meigen, 1826

Records. Zmajevac, 9.V.2018, S. Krčmar leg. (2♂) (DBUO); same locality, 19.VIII.2019, S. Krčmar leg. (1♂) (DBUO); same locality, 22.VIII.2019, S. Krčmar leg. (3♂) (DBUO); same locality, 24.VIII.2019, S. Krčmar leg. (1♂) (DBUO); same locality, 5.IX.2019, S. Krčmar leg. (3♂) (DBUO); same locality, 6.IX.2019, S. Krčmar leg. (9♂) (DBUO, SMNS); same locality, 17.IX.2019, S. Krčmar leg. (1♂) (DBUO); same locality, 30.IX.2019, S. Krčmar leg. (2♂) (DBUO); same locality, 28.VI.2020, S. Krčmar leg. (1♂) (DBUO); same locality, 28.VII.2020, S. Krčmar leg. (1♂) (DBUO); same locality, 20.VI.2021, S. Krčmar leg. (9♂) (DBUO).

New records for Croatian Baranja. Popovac, 11.VI.2021, S. Krčmar leg. (1♂) (DBUO); same locality, 10.VII.2021, S. Krčmar leg. (2♂) (DBUO); Petlovac, 18.VI.2021, S. Krčmar leg. (1♂) (DBUO); Darda, 19.VIII.2021, S. Krčmar leg. (1♂) (DBUO).

7. *Sarcophaga (Helicophagella) noverca* Rondani, 1860

Records. Zmajevac, 9.V.2018, S. Krčmar leg. (6♂) (DBUO); same locality, 16.V.2019, S. Krčmar leg. (2♂) (DBUO); same locality, 22.V.2019, S. Krčmar leg. (12♂) (DBUO); same locality, 22.VIII.2019, S. Krčmar leg. (2♂) (DBUO); same locality, 5.IX.2019, S. Krčmar leg. (1♂) (DBUO); same locality, 6.IX.2019, S. Krčmar leg. (1♂) (DBUO); same locality, 13.IX.2019, S. Krčmar leg. (1♂) (SMNS); same locality, 8.V.2020, S. Krčmar leg. (1♂) (DBUO); same locality, 12.VIII.2020, S. Krčmar leg. (2♂) (DBUO); same locality, 13.IX.2020, S. Krčmar leg. (1♂) (DBUO).

8. *Sarcophaga (Heteronychia) depressifrons* Zetterstedt, 1845

New records for Croatian Baranja. Zmajevac, 12.VIII.2020, S. Krčmar leg. (1♂) (DBUO).

9. *Sarcophaga (Heteronychia) filia* Rondani, 1860

New records for Croatian Baranja. Zmajevac, 6.IX.2019, S. Krčmar leg. (1♂) (SMNS); same locality, 12.VI.2020, S. Krčmar leg. (6♂) (DBUO); same locality, 19.VI.2020, S. Krčmar leg. (1♂) (DBUO); same locality, 28.VI.2020, S. Krčmar leg. (17♂) (DBUO); same locality, 28.VII.2020, S. Krčmar leg. (6♂) (DBUO); same locality, 12.VIII.2020, S. Krčmar leg. (1♂) (DBUO); Petlovac, 18.V.2021, S. Krčmar leg. (1♂) (DBUO); same locality, 18.VI.2021, S. Krčmar leg. (2♂) (DBUO); Suza, 18.V.2021, S. Krčmar leg. (2♂) (DBUO).

10. *Sarcophaga (Heteronychia) haemorrhoa* Meigen, 1826

Records. Zmajevac, 9.V.2018, S. Krčmar leg. (1♂) (SMNS); same locality, 16.V.2019, S. Krčmar leg. (1♂) (SMNS); same locality, 29.VI.2019, S. Krčmar leg. (1♂) (DBUO), same locality, 5.IX.2019, S. Krčmar leg. (1♂) (SMNS); same locality, 12.VI.2020, S. Krčmar leg. (1♂) (DBUO); same locality, 28.VI.2020, S. Krčmar leg. (3♂) (DBUO); same locality, 28.VII.2020, S. Krčmar leg. (1♂) (DBUO); same locality, 12.VIII.2020, S. Krčmar leg. (1♂) (DBUO).

11. *Sarcophaga (Heteronychia) haemorrhoides* Böttcher, 1913

New records for Croatian Baranja. Zmajevac, 28.VI.2020, S. Krčmar leg. (1♂) (DBUO); same locality, 12.VIII.2020, S. Krčmar leg. (1♂) (DBUO).

12. *Sarcophaga (Heteronychia) proxima* Rondani, 1860

Records. Zmajevac, 22.V.2019, S. Krčmar leg. (1♂) (SMNS); same locality, 24.VIII.2019, S. Krčmar leg. (1♂) (SMNS); same locality, 30.VIII.2019, S. Krčmar leg. (1♂) (SMNS).

13. *Sarcophaga (Heteronychia) pseudobenaci* (Baranov, 1942)

Records. Zmajevac, 29.VI.2019, S. Krčmar leg. (1♂) (DBUO); same locality, 8.V.2020, S. Krčmar leg. (1♂) (DBUO); same locality, 13.IX.2020, S. Krčmar leg. (2♂) (DBUO).

New records for Croatian Baranja. Petlovac, 29.VIII.2021, S. Krčmar leg. (2♂) (DBUO).

14. *Sarcophaga (Heteronychia) pumila* Meigen, 1826

New records for Croatian Baranja. Popovac, 11.VI.2021, S. Krčmar leg. (1♂) (DBUO); Zmajevac, 30.VII.2021, S. Krčmar leg. (1♂) (DBUO).

15. *Sarcophaga (Heteronychia) schineri* Bezzi, 1891

Records. Zmajevac, 9.V. 2018, S. Krčmar leg. (1♂) (SMNS); same locality, 22.V.2019, S. Krčmar leg. (1♂) (SMNS); same locality, 12.VI.2020, S. Krčmar leg. (1♂) (DBUO); same locality, 28.VI.2020, S. Krčmar leg. (1♂) (DBUO); same locality, 12.VIII.2020, S. Krčmar leg. (1♂) (DBUO).

16. *Sarcophaga (Heteronychia) vagans* Meigen, 1826

New records for Croatian Baranja. Zmajevac, 19.VIII.2019, S. Krčmar leg. (1♂) (SMNS); same locality, 19.VI.2020, S. Krčmar leg. (1♂) (DBUO); same locality, 28.VI.2020, S. Krčmar leg. (1♂) (DBUO).

17. *Sarcophaga (Liopygia) argyrostoma* (Robineau-Desvoidy, 1830)

Records. Zmajevac, 29.VII.2019, S. Krčmar leg. (1♂) (DBUO); same locality, 22.VIII.2019, S. Krčmar leg. (1♂) (SMNS); same locality, 6.IX.2019, S. Krčmar leg. (1♂) (SMNS).

New records for Croatian Baranja. Popovac, 13.VIII.2021, S. Krčmar leg. (1♂) (DBUO).

18. *Sarcophaga (Liopygia) crassipalpis* Macquart, 1839

New records for Croatian Baranja. Popovac, 10.VII.2021, S. Krčmar leg. (1♂) (DBUO).

19. *Sarcophaga (Liosarcophaga) aegyptica* Salem, 1935

Records. Zmajevac, 22.VIII.2019, S. Krčmar leg. (1♂) (DBUO); same locality, 24.VIII.2019, S. Krčmar leg. (2♂) (DBUO); same locality, 30.VIII.2019, S. Krčmar leg. (2♂) (SMNS); same locality, 5.IX.2019, S. Krčmar leg. (5♂) (DBUO); same locality, 6.IX.2019, S. Krčmar leg. (2♂) (SMNS); same locality, 13.IX.2019, S. Krčmar leg. (1♂) (DBUO); same locality, 17.IX.2019, S. Krčmar leg. (1♂) (DBUO); same locality, 30.IX.2019, S. Krčmar leg. (9♂) (DBUO); same locality, 12.VI.2020, S. Krčmar leg. (1♂) (DBUO); same locality, 28.VI.2020, S. Krčmar leg. (1♂) (DBUO); same locality, 13.IX.2020, S. Krčmar leg. (2♂) (DBUO); same locality, 20.VI.2021, S. Krčmar leg. (2♂) (DBUO).

New records for Croatian Baranja. Kotlina, 18.V.2021, S. Krčmar leg. (1♂) (DBUO); Darda, 17.VI.2021, S. Krčmar leg. (1♂) (DBUO); Petlovac, 18.VI.2021, S.

Krčmar leg. (5♂) (DBUO); same locality, 20.VII.2021, S. Krčmar leg. (1♂) (DBUO); Suza, 25.VI.2021, S. Krčmar leg. (1♂) (DBUO); Popovac, 10.VII.2021, S. Krčmar leg. (2♂) (DBUO); same locality, 13.VIII.2021, S. Krčmar leg. (1♂) (DBUO).

20. *Sarcophaga (Liosarcophaga) dux* Thomson, 1869

New records for Croatian Baranja. Darda, 17.VI. 2021, S. Krčmar leg. (1♂) (DBUO).

21. *Sarcophaga (Liosarcophaga) emdeni* (Rohdendorf, 1969)

Records. Zmajevac, 9.V.2018, S. Krčmar leg. (1♂) (DBUO); same locality, 16.V.2019, S. Krčmar leg. (4♂) (DBUO); same locality, 22.V.2019, S. Krčmar leg. (3♂) (DBUO); same locality, 17.VI.2019, S. Krčmar leg. (2♂) (DBUO); same locality, 6.VII.2019, S. Krčmar leg. (4♂) (DBUO); same locality, 29.VII.2019, S. Krčmar leg. (2♂) (DBUO); same locality, 22.VIII.2019, S. Krčmar leg. (1♂) (DBUO); same locality, 24.VIII.2019, S. Krčmar leg. (2♂) (DBUO); same locality, 30.VIII.2019, S. Krčmar leg. (2♂) (DBUO); same locality, 5.IX.2019, S. Krčmar leg. (1♂) (DBUO); same locality, 13.IX.2019, S. Krčmar leg. (1♂) (DBUO); same locality, 8.V.2020, S. Krčmar leg. (1♂) (DBUO); same locality, 12.VI.2020, S. Krčmar leg. (1♂) (DBUO); same locality, 19.VI.2020, S. Krčmar leg. (1♂) (DBUO); same locality, 28.VI.2020, S. Krčmar leg. (4♂) (DBUO); same locality, 13.IX.2020, S. Krčmar leg. (1♂) (DBUO); same locality, 11.V.2021, S. Krčmar leg. (2♂) (DBUO).

New records for Croatian Baranja. Suza, 18.V.2021, S. Krčmar leg. (1♂) (DBUO); same locality, 11.VII.2021, S. Krčmar leg. (7♂) (DBUO); same locality, 12.VIII.2021, S. Krčmar leg. (2♂) (DBUO).

22. *Sarcophaga (Liosarcophaga) tuberosa* Pandellé, 1896

New records for Croatian Baranja. Zmajevac, 17.VI.2019, S. Krčmar leg. (1♂) (DBUO); Suza, 18.V.2021, S. Krčmar leg. (1♂) (DBUO).

23. *Sarcophaga (Mehria) sexpunctata* (Fabricius, 1805)

New records for Croatian Baranja. Zmajevac, 12.VI.2020, S. Krčmar leg. (1♂) (DBUO).

24. *Sarcophaga (Myorhina) nigriventris* Meigen, 1826

Records. Zmajevac, 16.V.2019, S. Krčmar leg. (1♂) (SMNS); same locality, 22.V.2019, S. Krčmar leg. (1♂) (DBUO); same locality, 30.IX.2019, S. Krčmar leg. (1♂) (SMNS); same locality, 28.VI.2020, S. Krčmar leg. (2♂) (DBUO); same locality, 28.VII.2020, S. Krčmar leg. (1♂) (DBUO); same locality, 20.X.2020, S. Krčmar leg. (1♂) (DBUO).

25. *Sarcophaga (Myorbina) soror* Rondani, 1860

Records. Zmajevac, 9.V.2018, S. Krčmar leg. (1♂) (DBUO); same locality, 16.V.2019, S. Krčmar leg. (3♂) (SMNS); same locality, 22.V.2019, S. Krčmar leg. (3♂) (DBUO); same locality, 29.VII.2019, S. Krčmar leg. (2♂) (DBUO); same locality, 19.VI.2020, S. Krčmar leg. (3♂) (DBUO).

26. *Sarcophaga (Pandelleana) protuberans* Pandellé, 1896

New records for Croatian Baranja. Zmajevac, 28.VI.2020, S. Krčmar leg. (1♂) (DBUO).

27. *Sarcophaga (Pandelleisca) similis* Meade, 1876

Records. Zmajevac, 16.V.2019, S. Krčmar leg. (1♂) (DBUO); same locality, 22.V.2019, S. Krčmar leg. (3♂) (DBUO); same locality, 24.VIII.2019, S. Krčmar leg. (1♂) (DBUO); same locality, 30.VIII.2019, S. Krčmar leg. (2♂) (SMNS); same locality, 5.IX.2019, S. Krčmar leg. (1♂) (DBUO); same locality, 6.IX.2019, S. Krčmar leg. (2♂) (DBUO); same locality, 13.IX.2019, S. Krčmar leg. (1♂) (DBUO); same locality, 30.IX.2019, S. Krčmar leg. (1♂) (DBUO); same locality, 28.VI.2020, S. Krčmar leg. (1♂) (DBUO).

New records for Croatian Baranja. Popovac, 11.VI.2021, S. Krčmar leg. (1♂) (DBUO); Petlovac, 18.VI.2021, S. Krčmar leg. (1♂) (DBUO); same locality, 20.VII.2021, S. Krčmar leg. (1♂) (DBUO); Darda, 19.VIII.2021, S. Krčmar leg. (1♂) (DBUO).

28. *Sarcophaga (Parasarcophaga) albiceps* Meigen, 1826

Records. Zmajevac, 16.V.2019, S. Krčmar leg. (1♂) (DBUO); same locality, 22.V.2019, S. Krčmar leg. (3♂) (DBUO); same locality, 29.VII.2019, S. Krčmar leg. (5♂) (DBUO); same locality, 19.VIII.2019, S. Krčmar leg. (1♂) (DBUO); same locality, 22.VIII.2019, S. Krčmar leg. (5♂) (DBUO); same locality, 24.VIII.2019, S. Krčmar leg. (4♂) (DBUO); same locality, 5.IX.2019, S. Krčmar leg. (5♂) (DBUO); same locality, 6.IX.2019, S. Krčmar leg. (16♂) (DBUO, SMNS); same locality, 8.V.2020, S. Krčmar leg. (1♂) (DBUO); same locality, 19.VI.2020, S. Krčmar leg. (2♂) (DBUO); same locality, 28.VI.2020, S. Krčmar leg. (7♂) (DBUO); same locality, 28.VII.2020, S. Krčmar leg. (2♂) (DBUO); same locality, 20.VI.2021, S. Krčmar leg. (7♂) (DBUO).

New records for Croatian Baranja. Bilje, 24.VII.2021, S. Krčmar leg. (6♂) (DBUO); Kotlina, 12.VIII.2021, S. Krčmar leg. (1♂) (DBUO).

29. *Sarcophaga (Robineauella) caerulescens* Zetterstedt, 1838

Records. Zmajevac, 22.V.2019, S. Krčmar leg. (1♂) (SMNS); same locality, 17.VI.2019, S. Krčmar leg. (1♂) (DBUO); same locality, 8.V.2020, S. Krčmar leg. (1♂) (DBUO); same locality, 28.VII.2020, S. Krčmar leg. (1♂) (DBUO).

30. *Sarcophaga (Rosellea) aratrix* Pandellé, 1896

Records. Zmajevac, 9.V.2018, S. Krčmar leg. (1♂) (DBUO); same locality, 22.IV.2019, S. Krčmar leg. (1♂) (SMNS); same locality, 16.V.2019, S. Krčmar leg. (2♂) (DBUO); same locality, 22.V.2019, S. Krčmar leg. (2♂) (SMNS); same locality, 17.VI.2019, S. Krčmar leg. (1♂) (DBUO); same locality, 24.VIII.2019, S. Krčmar leg. (1♂) (DBUO); same locality, 30.VIII.2019, S. Krčmar leg. (5♂) (DBUO); same locality, 5.IX.2019, S. Krčmar leg. (2♂) (DBUO); same locality, 8.V.2020, S. Krčmar leg. (3♂) (DBUO); same locality, 12.VI.2020, S. Krčmar leg. (1♂) (DBUO); same locality, 13.IX.2020, S. Krčmar leg. (2♂) (DBUO); same locality, 20.VI.2021, S. Krčmar leg. (2♂) (DBUO); same locality, 30.VII.2021, S. Krčmar leg. (1♂) (DBUO); same locality, 12.VIII.2021, S. Krčmar leg. (1♂) (DBUO).

New records for Croatian Baranja. Suza, 18.V.2021, S. Krčmar leg. (3♂) (DBUO); same locality, 11.VII.2021, S. Krčmar leg. (1♂) (DBUO); Petlovac, 20.VII.2021, S. Krčmar leg. (1♂) (DBUO); Darda, 24.VII.2021, S. Krčmar leg. (1♂) (DBUO); Popovac, 13.VIII.2021, S. Krčmar leg. (1♂) (DBUO).

31. *Sarcophaga (Sarcophaga) baranoffi* Rohdendorf, 1937

Records. Zmajevac, 9.V.2018, S. Krčmar leg. (2♂) (DBUO); same locality, 16.V.2019, S. Krčmar leg. (8♂) (DBUO, SMNS); same locality, 22.V.2019, S. Krčmar leg. (4♂) (DBUO); same locality, 29.VII.2019, S. Krčmar leg. (7♂) (DBUO); same locality, 22.VIII.2019, S. Krčmar leg. (1♂) (DBUO); same locality, 24.VIII.2019, S. Krčmar leg. (2♂) (DBUO); same locality, 30.VIII.2019, S. Krčmar leg. (4♂) (DBUO); same locality, 5.IX.2019, S. Krčmar leg. (1♂) (DBUO); same locality, 6.IX.2019, S. Krčmar leg. (6♂) (DBUO); same locality, 30.IX.2019, S. Krčmar leg. (1♂) (DBUO); same locality, 19.VI.2020, S. Krčmar leg. (1♂) (DBUO); same locality, 28.VI.2020, S. Krčmar leg. (2♂) (DBUO); same locality, 12.VIII.2020, S. Krčmar leg. (1♂) (DBUO); same locality, 13.IX.2020, S. Krčmar leg. (1♂) (DBUO); same locality, 11.V.2021, S. Krčmar leg. (1♂) (DBUO); same locality, 20.VI.2021, S. Krčmar leg. (1♂) (DBUO); same locality, 30.VII.2021, S. Krčmar leg. (6♂) (DBUO); same locality, 12.VIII.2021, S. Krčmar leg. (3♂) (DBUO).

New records for Croatian Baranja. Kotlina, 18.V.2021, S. Krčmar leg. (2♂) (DBUO); Suza, 25.VI.2021, S. Krčmar leg. (2♂) (DBUO); same locality, 12.VIII.2021, S. Krčmar leg. (2♂) (DBUO); Darda, 24.VII.2021, S. Krčmar leg. (2♂) (DBUO); Petlovac, 29.VIII.2021, S. Krčmar leg. (2♂) (DBUO).

32. *Sarcophaga (Sarcophaga) carnaria* (Linnaeus, 1758)

New records for Croatian Baranja. Zmajevac, 13.IX.2019, S. Krčmar leg. (1♂) (DBUO); same locality, 20.VI.2021, S. Krčmar leg. (1♂) (DBUO); Kotlina, 18.V.2021, S. Krčmar leg. (3♂) (DBUO).

33. *Sarcophaga (Sarcophaga) croatica* Baranov, 1941

Records. Zmajevac, 9.V.2018, S. Krčmar leg. (14♂) (DBUO); same locality, 22.IV.2019, S. Krčmar leg. (17♂) (DBUO, SMNS); same locality, 16.V.2019, S. Krčmar leg. (20♂) (DBUO); same locality, 22.V.2019, S. Krčmar leg. (15♂) (DBUO); same locality, 17.VI.2019, S. Krčmar leg. (10♂) (DBUO); same locality, 6.VII.2019, S. Krčmar leg. (1♂) (DBUO); same locality, 29.VII.2019, S. Krčmar leg. (5♂) (DBUO); same locality, 19.VIII.2019, S. Krčmar leg. (19♂) (DBUO); same locality, 22.VIII.2019, S. Krčmar leg. (19♂) (DBUO); same locality, 24.VIII.2019, S. Krčmar leg. (29♂) (DBUO); same locality, 30.VIII.2019, S. Krčmar leg. (34♂) (DBUO); same locality, 5.IX.2019, S. Krčmar leg. (9♂) (DBUO); same locality, 13.IX.2019, S. Krčmar leg. (8♂) (DBUO); same locality, 17.IX.2019, S. Krčmar leg. (1♂) (DBUO); same locality, 30.IX.2019, S. Krčmar leg. (3♂) (DBUO); same locality, 8.V.2020, S. Krčmar leg. (12♂) (DBUO); same locality, 12.VI.2020, S. Krčmar leg. (8♂) (DBUO); same locality, 19.VI.2020, S. Krčmar leg. (29♂) (DBUO); same locality, 28.VI.2020, S. Krčmar leg. (8♂) (DBUO); same locality, 28.VII.2020, S. Krčmar leg. (2♂) (DBUO); same locality, 12.VIII.2020, S. Krčmar leg. (3♂) (DBUO); same locality, 13.IX.2020, S. Krčmar leg. (22♂) (DBUO); same locality, 11.X.2020, S. Krčmar leg. (1♂) (DBUO); same locality, 20.X.2020, S. Krčmar leg. (5♂) (DBUO); same locality, 11.V.2021, S. Krčmar leg. (8♂) (DBUO); same locality, 20.VI.2021, S. Krčmar leg. (17♂) (DBUO); same locality, 30.VII.2021, S. Krčmar leg. (4♂) (DBUO); same locality, 12.VIII.2021, S. Krčmar leg. (5♂) (DBUO).

New records for Croatian Baranja. Kotlina, 18.V.2021, S. Krčmar leg. (8♂) (DBUO); same locality, 18.VI.2021, S. Krčmar leg. (9♂) (DBUO); Petlovac, 18.V.2021, S. Krčmar leg. (9♂) (DBUO); same locality, 18.VI.2021, S. Krčmar leg. (22♂) (DBUO); same locality, 20.VII.2021, S. Krčmar leg. (10♂) (DBUO); Suza, 18.V.2021, S. Krčmar leg. (6♂) (DBUO); same locality, 25.VI.2021, S. Krčmar leg. (17♂) (DBUO); same locality, 11.VII.2021, S. Krčmar leg. (5♂) (DBUO); same locality, 12.VIII.2021, S. Krčmar leg. (5♂) (DBUO); Popovac, 11.VI.2021, S. Krčmar leg. (1♂) (DBUO); Bilje, 17.VI.2021, S. Krčmar leg. (17♂) (DBUO); same locality, 19.VIII.2021, S. Krčmar leg. (3♂) (DBUO); Darda, 17.VI.2021, S. Krčmar leg. (2♂) (DBUO); same locality, 24.VII.2021, S. Krčmar leg. (16♂) (DBUO); same locality, 19.VIII.2021, S. Krčmar leg. (14♂) (DBUO).

34. *Sarcophaga (Sarcophaga) lehmanni* Müller, 1922

Records. Zmajevac, 22.IV.2019, S. Krčmar leg. (4♂) (DBUO, SMNS); same locality, 16.V.2019, S. Krčmar leg. (10♂) (DBUO); same locality, 22.V.2019, S. Krčmar leg. (19♂) (DBUO); same locality, 17.VI.2019, S. Krčmar leg. (6♂) (DBUO); same locality, 6.VII.2019, S. Krčmar leg. (5♂) (DBUO); same locality, 29.VII.2019, S. Krčmar leg. (3♂) (DBUO); same locality, 22.VIII.2019, S. Krčmar leg. (1♂) (DBUO); same locality, 24.VIII.2019, S. Krčmar leg. (3♂) (DBUO); same locality, 30.VIII.2019, S. Krčmar leg. (4♂) (DBUO); same locality, 5.IX.2019, S. Krčmar leg. (3♂) (DBUO); same locality, 6.IX.2019, S. Krčmar leg. (10♂) (DBUO); same locality, 13.IX.2019, S. Krčmar leg. (7♂) (DBUO); same locality, 17.IX.2019, S. Krčmar leg. (2♂) (DBUO);

same locality, 30.IX.2019, S. Krčmar leg. (13♂) (DBUO); same locality, 29.VI.2019, S. Krčmar leg. (1♂) (DBUO); same locality, 8.V.2020, S. Krčmar leg. (17♂) (DBUO); same locality, 12.VI.2020, S. Krčmar leg. (18♂) (DBUO); same locality, 19.VI.2020, S. Krčmar leg. (12♂) (DBUO); same locality, 28.VI.2020, S. Krčmar leg. (16♂) (DBUO); same locality, 28.VII.2020, S. Krčmar leg. (6♂) (DBUO); same locality, 13.IX.2020, S. Krčmar leg. (6♂) (DBUO); same locality, 11.X.2020, S. Krčmar leg. (2♂) (DBUO); same locality, 20.X.2020, S. Krčmar leg. (4♂) (DBUO); same locality, 11.V.2021, S. Krčmar leg. (13♂) (DBUO); same locality, 20.VI.2021, S. Krčmar leg. (4♂) (DBUO); same locality, 12.VIII.2021, S. Krčmar leg. (2♂) (DBUO).

New records for Croatian Baranja. Kotlina, 18.V.2021, S. Krčmar leg. (9♂) (DBUO); same locality, 18.VI.2021, S. Krčmar leg. (10♂) (DBUO); same locality, 12.VIII.2021, S. Krčmar leg. (1♂) (DBUO); Petlovac, 18.V.2021, S. Krčmar leg. (3♂) (DBUO); same locality, 18.VI.2021, S. Krčmar leg. (3♂) (DBUO); same locality, 20.VII.2021, S. Krčmar leg. (2♂) (DBUO); Suza, 18.V.2021, S. Krčmar leg. (4♂) (DBUO); same locality, 25.VI.2021, S. Krčmar leg. (2♂) (DBUO); same locality, 11.VII.2021, S. Krčmar leg. (29♂) (DBUO); same locality, 12.VIII.2021, S. Krčmar leg. (7♂) (DBUO); Popovac, 11.VI.2021, S. Krčmar leg. (9♂) (DBUO); same locality, 13.VIII.2021, S. Krčmar leg. (2♂) (DBUO); Bilje, 17.VI.2021, S. Krčmar leg. (4♂) (DBUO).

35. *Sarcophaga (Sarcophaga) variegata* (Scopoli, 1763)

New records for Croatian Baranja. Zmajevac, 30.VIII.2019, S. Krčmar leg. (2♂) (SMNS); same locality, 11.V.2021, S. Krčmar leg. (1♂) (DBUO); Petlovac, 18.VI.2021, S. Krčmar leg. (4♂) (DBUO).

36. *Sarcophaga (Thyrsocnema) incisilobata* Pandellé, 1896

Records. Zmajevac, 16.V.2019, S. Krčmar leg. (2♂) (DBUO, SMNS); same locality, 17.VI.2019, S. Krčmar leg. (3♂) (DBUO); 22.VIII.2019, S. Krčmar leg. (1♂) (DBUO); same locality, 24.VIII.2019, S. Krčmar leg. (3♂) (DBUO); same locality, 5.IX.2019, S. Krčmar leg. (4♂) (DBUO); same locality, 6.IX.2019, S. Krčmar leg. (10♂) (DBUO); same locality, 8.V.2020, S. Krčmar leg. (2♂) (DBUO); same locality, 12.VI.2020, S. Krčmar leg. (1♂) (DBUO); same locality, 19.VI.2020, S. Krčmar leg. (2♂) (DBUO); same locality, 28.VI.2020, S. Krčmar leg. (3♂) (DBUO); same locality, 12.VIII.2020, S. Krčmar leg. (3♂) (DBUO); same locality, 11.X.2020, S. Krčmar leg. (1♂) (DBUO); same locality, 20.X.2020, S. Krčmar leg. (4♂) (DBUO); same locality, 20.VI.2021, S. Krčmar leg. (4♂) (DBUO); same locality, 12.VIII.2021, S. Krčmar leg. (1♂) (DBUO).

New records for Croatian Baranja. Petlovac, 18.V.2021, S. Krčmar leg. (3♂) (DBUO); same locality, 18.VI.2021, S. Krčmar leg. (1♂) (DBUO); same locality, 20.VII.2021, S. Krčmar leg. (1♂) (DBUO); Suza, 18.V.2021, S. Krčmar leg. (1♂) (DBUO); same locality, 25.VI.2021, S. Krčmar leg. (1♂) (DBUO); same locality, 11.VII.2021, S. Krčmar leg. (1♂) (DBUO); Darda, 17.VI.2021, S. Krčmar leg.

(1♂) (DBUO); Kotlina, 18.VI.2021, S. Krčmar leg. (1♂) (DBUO); same locality, 20.VII.2021, S. Krčmar leg. (1♂) (DBUO); same locality, 12.VIII.2021, S. Krčmar leg. (3♂) (DBUO); Popovac, 13.VIII.2021, S. Krčmar leg. (1♂) (DBUO).

37. *Sarcophaga (Pseudothrysocnema) spinosa* Villeneuve, 1912 ▲

New records for Croatian Baranja. Zmajevac, 12.VIII.2020, S. Krčmar leg. (1♂) (SMNS). New for Croatia.

Table 2. Number of species and specimens of flesh flies from subfamily Sarcophaginae collected at seven localities in Croatian part of Baranja from 2018 to 2021.

Species / Locality	1 Bilje	2 Darda	3 Kotlina	4 Petlovac	5 Popovac	6 Suza	7 Zmajevac	Σ
<i>Sarcophaga (Sar.) croatica</i>	20	32	17	41	1	33	328	472
<i>S. (Sar.) lehmanni</i>	4	-	20	8	11	42	191	276
<i>S. (Pas.) albiceps</i>	6	-	1	-	-	-	59	66
<i>S. (Sar.) baranoffi</i>	-	2	2	2	-	4	52	62
<i>S. (Thy.) incisilobata</i>	-	1	5	5	1	3	44	59
<i>S. (Lis.) emdeni</i>	-	-	-	-	-	10	33	43
<i>S. (Lis.) aegyptica</i>	-	1	1	6	3	1	29	41
<i>S. (Hel.) melanura</i>	-	1	-	1	3	-	33	38
<i>S. (Het.) filia</i>	-	-	-	3	-	2	32	37
<i>S. (Ros.) aratrix</i>	-	1	-	-	1	5	25	32
<i>S. (Hel.) noverca</i>	-	-	-	-	-	-	29	29
<i>S. (Bel.) subulata</i>	-	-	-	-	-	-	18	18
<i>S. (Pad.) similis</i>	-	1	-	2	1	-	13	17
<i>Ravinia pernix</i>	-	1	2	-	6	-	6	15
<i>S. (Myo.) soror</i>	-	-	-	-	-	-	12	12
<i>S. (Het.) haemorrhoa</i>	-	-	-	-	-	-	10	10
<i>S. (Myo.) nigriventris</i>	-	-	-	-	-	-	7	7
<i>S. (Sar.) variegata</i>	-	-	-	4	-	-	3	7
<i>S. (Hel.) crassimargo</i>	-	-	-	-	-	-	6	6
<i>S. (Het.) pseudobenaci</i>	-	-	-	2	-	-	4	6
<i>S. (Het.) schineri</i>	-	-	-	-	-	-	5	5
<i>S. (Sar.) carnaria</i>	-	-	3	-	-	-	2	5
<i>S. (Lip.) argyrostoma</i>	-	-	-	-	1	-	3	4
<i>S. (Rob.) caerulescens</i>	-	-	-	-	-	-	4	4
<i>S. (Het.) proxima</i>	-	-	-	-	-	-	3	3
<i>S. (Het.) vagans</i>	-	-	-	-	-	-	3	3
<i>S. (Ber.) africa</i>	-	-	-	-	-	-	2	2
<i>S. (Hel.) hirticrus</i>	-	-	-	-	-	-	2	2
<i>S. (Het.) haemorrhoides</i>	-	-	-	-	-	-	2	2
<i>S. (Het.) pumila</i>	-	-	-	-	1	-	1	2
<i>S. (Lis.) tuberosa</i>	-	-	-	-	-	1	1	2
<i>S. (Het.) depressifrons</i>	-	-	-	-	-	-	1	1
<i>S. (Lip.) crassipalpis</i>	-	-	-	-	1	-	-	1
<i>S. (Lis.) dux</i>	-	1	-	-	-	-	-	1
<i>S. (Meh.) sexpunctata</i>	-	-	-	-	-	-	1	1
<i>S. (Pan.) protuberans</i>	-	-	-	-	-	-	1	1
<i>S. (Pse.) spinosa</i>	-	-	-	-	-	-	1	1
Σ = 37	30	41	51	74	30	101	966	1293

Table 3. Number of collected flesh flies from subfamily Sarcophaginae according to sampling years in the Croatian part of Baranja.

Species / Year	2018	2019	2020	2021	Σ
<i>Sarcophaga</i> (Sar.) <i>croatica</i>	14	190	90	178	472
<i>S. (Sar.) lehmanni</i>	-	91	81	104	276
<i>S. (Pas.) albiceps</i>	-	40	12	14	66
<i>S. (Sar.) baranoffi</i>	2	34	5	21	62
<i>S. (Thy.) incisilobata</i>	-	23	16	20	59
<i>S. (Lis.) emdeni</i>	1	22	8	12	43
<i>S. (Lis.) aegyptica</i>	-	23	4	14	41
<i>S. (Hel.) melanura</i>	2	20	2	14	38
<i>S. (Het.) filia</i>	-	1	31	5	37
<i>S. (Ros.) aratrix</i>	1	14	6	11	32
<i>S. (Hel.) noverca</i>	6	19	4	-	29
<i>S. (Bel.) subulata</i>	4	9	4	1	18
<i>S. (Pad.) similis</i>	-	12	1	4	17
<i>Ravinia pernix</i>	-	1	4	10	15
<i>S. (Myo.) soror</i>	1	8	3	-	12
<i>S. (Het.) haemorrhoa</i>	1	3	6	-	10
<i>S. (Myo.) nigriventris</i>	-	3	4	-	7
<i>S. (Sar.) variegata</i>	-	2	-	5	7
<i>S. (Hel.) crassimargo</i>	-	2	4	-	6
<i>S. (Het.) pseudobenaci</i>	-	1	3	2	6
<i>S. (Het.) schineri</i>	1	1	3	-	5
<i>S. (Sar.) carnaria</i>	-	1	-	4	5
<i>S. (Lip.) argyrostoma</i>	-	3	-	1	4
<i>S. (Rob.) caerulescens</i>	-	2	2	-	4
<i>S. (Het.) proxima</i>	-	3	-	-	3
<i>S. (Het.) vagans</i>	-	1	2	-	3
<i>S. (Ber.) africa</i>	-	2	-	-	2
<i>S. (Hel.) hirticrus</i>	-	1	1	-	2
<i>S. (Het.) haemorrhoides</i>	-	-	2	-	2
<i>S. (Het.) pumila</i>	-	-	-	2	2
<i>S. (Lis.) tuberosa</i>	-	1	-	1	2
<i>S. (Het.) depressifrons</i>	-	-	1	-	1
<i>S. (Lip.) crassipalpis</i>	-	-	-	1	1
<i>S. (Lis.) dux</i>	-	-	-	1	1
<i>S. (Meh.) sexpunctata</i>	-	-	1	-	1
<i>S. (Pan.) protuberans</i>	-	-	1	-	1
<i>S. (Pse.) spinosa</i>	-	-	1	-	1
Σ = 37	33	533	302	425	1293

Discussion

The first samples of flesh flies from Croatian Baranja were collected from 2014 to 2017 in the localities Zmajevac and Kamenac, at which time 29 species were recorded (Krčmar et al. 2019). Of those 29, the following five species were not confirmed in the present study: *Blaesoxiphia* (Ser.) *rossica* Villeneuve, 1912; *S. (Het.) benaci* Böttcher, 1913;

S. (Kra.) anaces Walker, 1849; *S. (Lis.) portschinskyi* (Rohdendorf, 1937), and *S. (Sac.) sinuata* Meigen, 1826. Meanwhile, 13 species were newly recorded for the area in this study. Of all the 42 species recorded from Croatian Baranja, 41 are widely distributed in Central Europe (Povolný and Verves 1997), whereas one, *S. (Het.) pseudobenaci*, is restricted to southeastern Europe (Pape 2004). *Sarcophaga (Pse.) spinosa* represents a new record for Croatia, which is not surprising as this species is recorded from neighbouring Hungary and Serbia, and has also been recorded from Albania, French mainland, Italian mainland, North Macedonia, Romania, and Ukraine (Pape 2004). Most species were found in natural and semi-natural habitats, although some of the species recorded in this study, i.e., *S. (Ber.) africa*, *S. (Hel.) hirticrus*, *S. (Het.) depressifrons*, *S. (Lip.) argyrostoma*, *S. (Ros.) aratrix*, and *S. (Sar.) carnaria* are known to be found also in urban habitats (Hwang and Turner 2005). Some species of significance for forensic entomology were collected on several localities around pond habitats in rural settlements, in particular *S. (Lip.) argyrostoma* and *S. (Lis.) dux*, both of which colonize decomposing human remains (Draber-Moňko et al. 2009; Sukontason et al. 2014), whereas *S. (Lis.) dux* has a medical importance as an agent of myiasis (Sukontason et al. 2014). Also, an important role of *S. (Lip.) argyrostoma* in accidental intestinal myiasis was recently confirmed (Najjari et al. 2020). Moreover, *S. (Ber.) africa*, *S. (Lip.) argyrostoma*, *S. (Pad.) similis*, and *S. (Rob.) caerulescens* have been found on corpses in indoor cases in Switzerland and Finland (Cherix et al. 2012; Ren et al. 2018), while *S. (Lip.) crassipalpis* was found on corpses at the earliest stage of decomposition in Australia (Ren et al. 2018). All these five species were recorded in this study (Table 2). *Sarcophaga (Pad.) similis* was recorded in four localities (Darda, Petlovac, Popovac, Zmajevac). *Sarcophaga (Lip.) argyrostoma* was recorded on two localities (Popovac, Zmajevac), *S. (Ber.) africa*, and *S. (Rob.) caerulescens* were only recorded in Zmajevac, while *S. (Lip.) crassipalpis* was recorded only in Popovac (Table 2). In this study, a large number of flesh fly specimens were collected on or nearby pet animal faeces and on discarded leftover food. This is not surprising since it is known that *S. (Pas.) albiceps*, *S. (Thy.) incisilobata*, and *Ravinia pernix* visit excrements from humans and (other) animals (Papp 1992a, 1992b). Several species of flesh flies are also known to visit different animal carcasses (Szpila et al. 2015). Among them, the following species were recorded during this study: *R. pernix*, *S. (Hel.) melanura*, *S. (Lip.) argyrostoma*, *S. (Pad.) similis*, *S. (Pas.) albiceps*, *S. (Rob.) caerulescens*, *S. (Ros.) aratrix*, *S. (Sar.) carnaria*, *S. (Sar.) lehmanni*, *S. (Sar.) variegata*, and *S. (Thy.) incisilobata*. During this study not a single specimen of flesh flies was collected from carcasses. In Yozgat province of Turkey, 21 flesh flies species were recorded visiting small carrion (Pekbey 2019) from which *S. (Ber.) africa*, *S. (Hel.) hirticrus*, *S. (Hel.) melanura*, *S. (Lip.) argyrostoma*, *S. (Lip.) crassipalpis*, *S. (Lis.) aegyptica*, *S. (Lis.) emdeni*, *S. (Lis.) tuberosa*, *S. (Myo.) nigriventris*, *S. (Sar.) lehmanni* and *S. (Thy.) incisilobata* were also collected in this study. Eighty years ago, Baranov (1940) confirmed the presence of five flesh fly species in a laystall in the village of Metajna on the Island of Pag, four of which were also recorded in this study. In a similar study from the Polish Baltic coast, a number of species were recorded from a marshy habitat (15) and a sandy habitat (24) (Kaczorowska 2009), while in this study 37 species were recorded from pond habitats. Regardless of the distance and different climatic conditions, 11 species are present in all three types of habitats: *S. (Ber.) africa*,

S. (Hel.) crassimargo, *S. (Hel.) melanura*, *S. (Het.) haemorrhoa*, *S. (Het.) vagans*, *S. (Myo.) nigriventris*, *S. (Rob.) caerulescens*, *S. (Ros.) aratrix*, *S. (Sar.) carnaria*, *S. (Sac.) sinuata*, and *S. (Thy.) incisilobata*. In marshy and sandy habitats the most numerous species was *S. (Sar.) carnaria* (Kaczorowska 2009), while *S. (Sar.) croatica* was the most abundant around pond habitats in rural settlements. Martínez-Sánchez et al. (2000) recorded 12 species from a holm-oak pasture agroecosystem in Salamanca province (Spain), from which four species *S. (Ber.) africa*, *S. (Hel.) melanura*, *S. (Lip.) crassipalpis*, and *S. (Sar.) lehmanni* were also collected in this study. The large differences in the number of recorded species between Zmajevac and other localities may be explained by the much higher number of samples. This is clearly shown by the fact that in 2021 only 15 species were collected at the Zmajevac locality compared to the total number of 35 species that were sampled during all three years (2019–2021) at this locality. The lower number of recorded species at the Bilje locality was influenced by environmental factors such as open sun throughout the day (with afternoon temperatures ≥ 32 °C) and a lack of animal faeces and remains for food, which reduced the number of recorded species. The seven localities around pond habitats are polluted by different organic contaminations caused by various human activities, which can attract certain species for feeding and breeding.

Conclusions

The species *S. (Sar.) croatica* and *S. (Sar.) lehmanni* were recorded from almost all sampling sites around pond habitats in the Croatian part of Baranja. They belong to the most widespread and abundant flesh flies in Croatia and have been recorded in all three biogeographic regions: Alpine, Mediterranean, and Pannonian–Peripannnonian. By merging data from earlier studies (2014–2017) and this study, altogether 42 species of flesh flies were recorded in the Croatian part of Baranja. The species *S. (Pse.) spinosa* is recorded as new for the Croatian fauna, thus increasing the total number of recorded species of flesh flies in Croatia to 156.

Acknowledgements

I am very grateful to Daniel Whitmore (Naturkunde Museum, Stuttgart, Germany) for the identification of newly recorded species of flesh flies for the Croatian fauna, as well as to Thomas Pape (Natural History Museum of Denmark) for help with literature. I am also grateful to reviewers for all improvements in this paper and to Stephen M. Smith, University of Waterloo, Waterloo, Canada for his linguistic corrections and improvements. This study was funded by the Department of Biology, Josip Juraj Strossmayer University of Osijek under the project OZB-ZP2021 (Sarcophagidae- 31053) “Diversity of flesh flies fauna from subfamily Sarcophaginae (Diptera: Sarcophagidae) in pond habitats in rural settlements in Croatian part of Baranja“ and the Foundation of the Croatian Academy of Sciences and Arts from the project “Comparison of repellent activity of different essential oils and plant extracts against hard ticks (Acari: Ixodidae)“.

References

- Baranov N (1928) Tachinidensammlung des zoologischen Museums in Zagreb. Glasnik Hrvatskoga Prirodoslovnoga Društva 39/40: 196–200.
- Baranov N (1929) Beitrag zur Kenntnis der Gattung *Sarcophaga* (Mg.) Böttcher (Dipt., Tach.). Neue Beiträge zur Systematische Insektenkunde 4: 142–153.
- Baranov N (1938) Raupenfliegen (Tachinidae s.l.) welche auf der Adria-Insel Pag bei trinken von Meerwasser gefangen wurden. Encyclopédie entomologique Série B II (Diptera) 9: 103–107.
- Baranov N (1940) Die Düngerstätten im Dorfe Metajna als Brutplätze der Hausfliegen. Veterinarski Arhiv 10: 193–213. [In Croatian with German abstract]
- Baranov N (1941) Zweiter Beitrag zur Kenntnis der Gattung *Sarcophaga* s.l. Veterinarski Arhiv 11: 361–404. [In Croatian with German abstract]
- Baranov N (1942) Sarcophagen im Unabhängigen Staate Kroatien. Veterinarski Arhiv 12: 497–659. [In Croatian with German abstract]
- Baranov N (1943) *Wohlfahrtia magnifica* Schin., als Erreger der Schweine-Myiasis in Kroatien. Veterinarski Arhiv 13: 447–456. [In Croatian with German abstract]
- Bognar A, Crkvenčić I, Peponik Z, Riđanović J, Roglić J, Sić M, Šegota T, Vresk M (1975) Geography of SR Croatia. Eastern Croatia. Školska knjiga, Zagreb, 256 pp. [In Croatian with English abstract]
- Cherix D, Wyss C, Pape T (2012) Occurrences of flesh flies (Diptera: Sarcophagidae) on human cadavers in Switzerland, and their importance as forensic indicators. Forensic Science International 220(1–3): 158–163. <https://doi.org/10.1016/j.forsciint.2012.02.016>
- Draber-Mońko A, Malewski T, Pomorski J, Łoś M, Ślipiński P (2009) On the morphology and mitochondrial DNA barcoding of the flesh fly *Sarcophaga (Liopygia) argyrostoma* (Robineau-Desvoidy, 1830) (Diptera: Sarcophagidae) – an important species in forensic entomology. Annales Zoologici 59(4): 465–493. <https://doi.org/10.3161/000345409X484865>
- Hwang C, Turner BD (2005) Spatial and temporal variability of necrophagous Diptera from urban to rural areas. Medical and Veterinary Entomology 19(4): 379–391. <https://doi.org/10.1111/j.1365-2915.2005.00583.x>
- Kaczorowska E (2009) Flesh flies (Diptera: Sarcophagidae) of sandy and marshy habitats of the Polish Baltic coast. Entomologica Fennica 20(1): 61–64. <https://doi.org/10.33338/ef.84461>
- Krčmar S, Whitmore D, Pape T, Buenaventura E (2019) Checklist of the Sarcophagidae (Diptera) of Croatia, with new records from Croatia and other Mediterranean countries. ZooKeys 831: 95–155. <https://doi.org/10.3897/zookeys.831.30795>
- Langhoffer A (1920) Beiträge zur Dipterenfauna Kroatiens. Glasnik hrvatskoga prirodoslovnoga društva 32: 57–63. <https://www.biodiversitylibrary.org/page/11380253>
- Martínez-Sánchez A, Rojo S, Marcos-García MA (2000) Necrophagous and coprophagous flesh-flies associated with a holm-oak pasture agroecosystem (Diptera: Sarcophagidae). Boletín de la Asociación Espanola de Entomología 24: 171–185. [In Spanish with English abstract]
- Mihaljević M, Getz D, Tadić Z, Živanović B, Gucunski D, Topić J, Kalinović I, Mikuska J (1999) Kopački rit research survey and bibliography. Hrvatska akademija znanosti i

- umjetnosti, Zavod za znanstveni rad Osijek, Zagreb-Osijek, 188 pp. [In Croatian with English, German and Hungarian abstract]
- Najjari M, Dik B, Pekbey G (2020) Gastrointestinal Myiasis Due to *Sarcophaga argyrostoma* (Diptera: Sarcophagidae) in Mashhad, Iran: a Case Report. Journal of Arthropod-Borne Diseases 14: 317–324. <https://doi.org/10.18502/jad.v14i3.4565>
- Pape T (1987) The Sarcophagidae (Diptera) of Fennoscandia and Denmark. Fauna Entomologica Scandinavica, 19. E. J. Brill/Scandinavian Science Press Ltd., Leiden-Copenhagen, 203 pp. <https://doi.org/10.1163/9789004273436>
- Pape T (2004) Fauna Europaea: Diptera, Sarcophagidae. Fauna Europaea version 2.4. <http://www.faunaeur.org> [Accessed 28 December 2017]
- Pape T, Beuk P, Pont A, Shatalkin A, Ozerov A, Woźnica A, Merz B, Bystrowski C, Raper C, Bergström C, Kehlmaier C, Clements D, Greathead D, Kameneva E, Nartshuk E, Petersen F, Weber G, Bächli G, Geller-Grimm F, Van de Weyer G, Tschorsnig H, de Jong H, van Zuijlen J, Vaňhara J, Roháček J, Ziegler J, Majer J, Hurka K, Holston K, Rognes K, Greve-Jensen L, Munari L, de Meyer M, Pollet M, Speight M, Ebejer M, Martinez M, Carles-Tolrá M, Földvári M, Chvála M, Barták M, Evenhuis N, Chandler P, Cerretti P, Meier R, Rozkosny R, Prescher S, Gaimari S, Zatwarnicki T, Zeegers T, Dikow T, Korneyev V, Richter V, Michelsen V, Tanasijtshuk V, Mathis W, Hubenov Z, de Jong Y (2015) Fauna Europaea: Diptera-Brachycera. Biodiversity Data Journal 3: e4187. <https://doi.org/10.3897/BDJ.3.e4187>
- Papp L (1992a) Flies (Diptera) visiting human faeces in mountain creek valleys in Hungary. Parasitologia Hungarica 25: 85–96. <https://www.publication.nhmus.hu/pdf/parasitologia/bannales.php>
- Papp L (1992b) Fly communities in pasture dung: some results and problems (Diptera). Acta Zoologica Hungarica 38: 75–88. <http://real-j.mtak.hu/id/eprint/4317>
- Pekbey G (2019) A preliminary study on determination of small carrion visitor Sarcophagidae (Diptera) species from Yozgat (Turkey), with two new records. Turkish Journal of Agricultural and Natural Sciences 6: 354–362. <https://doi.org/10.30910/turkjans.595085>
- Povolný D, Verves YuG (1997) The Flesh-Flies of Central Europe (Insecta, Diptera, Sarcophagidae). Spixiana (Suppl. 24): 1–264. <https://www.zobodat.at/pdf/SpixSupp0240001–0260.pdf>
- Ren L, Shang Y, Chen W, Meng F, Cai J, Zhu G, Chen L, Wang Y, Deng J, Guo Y (2018) A brief review of forensically important flesh flies (Diptera: Sarcophagidae). Forensic Sciences Research 3(1): 16–26. <https://doi.org/10.1080/20961790.2018.1432099>
- Richet R, Blackith RM, Pape T (2011) *Sarcophaga* of France (Diptera: Sarcophagidae). Pensoft Publishers, Sofia-Moscow, 327 pp.
- Rucner Z (1994) Beitrag zur Entomofauna einiger waldassoziationen Kroatiens. Natura Croatica 3: 1–22. <https://www.hrcak.srce.hr/file/16607>
- Schneider-Jacoby M (1994) Sava and Drava ecological value and future of the two main rivers in Croatia. Periodicum Biologorum 96: 348–356.
- Sisojević P, Čepelak J, Slamečkova M (1989) Contribution to the fauna of Sarcophagidae (Diptera) of Yugoslavia. Biosistematiка 15: 71–79. [In Croatian with English abstract]
- Strukan D (1964) Contribution to the knowledge of Sarcophaginae species and types in Yugoslavia. PhD dissertation, University of Zagreb, Zagreb, Croatia [In Croatian with English abstract]

- Strukan D (1967) Remarques sur deux Sarcophaginae peu connus de Yougoslavie (Diptera Sarcophagidae). Cahiers des Naturalistes, Bulletin NP, n.s. 23: 45–47.
- Strukan D (1970) Les Parasarcophaginae de Yougoslavie (Sarcophagidae – Diptera). Matica Srpska Zbornik za Prirodne Nauke 38: 90–114. [In Serbian with French abstract]
- Sukontason KI, Sanit S, Klong-Klaew T, Tomberlin JK, Sukontason K (2014) *Sarcophaga (Liosarcophaga) dux* (Diptera: Sarcophagidae): A flesh fly species of medical importance. Biological Research 47(1): 14. <https://doi.org/10.1186/0717-6287-47-14>
- Szpila K, Mądra A, Jarmusz M, Matuszewski S (2015) Flesh flies (Diptera: Sarcophagidae) colonising large carcasses in Central Europe. Parasitology Research 114(6): 2341–2348. <https://doi.org/10.1007/s00436-015-4431-1>
- Verves Yu G, Barták M (2021) New faunistic data on Sarcophagidae (Diptera) from Croatia. The Kharkov Entomological Society Gazette 29(1): 71–76. <https://doi.org/10.36016/KhESG-2021-29-1-6>
- Whitmore D (2009) A review of the *Sarcophaga (Heteronychia)* (Diptera: Sarcophagidae) of Sardinia. In: Cerretti P, Mason F, Minelli A, Nardi G, Whitmore D (Eds) Research on the Terrestrial Arthropods of Sardinia. Zootaxa 2318: 566–588. <https://doi.org/10.11646/zootaxa.2318.1.24>
- Whitmore D (2010) Systematics and phylogeny of *Sarcophaga (Heteronychia)* (Diptera, Sarcophagidae). PhD dissertation. Università degli Studi di Roma “La Sapienza”, Dipartimento di Biologia Animale e dell’Uomo, Rome, Italy, [v +] 257 pp.
- Whitmore D (2011) New taxonomic and nomenclatural data on *Sarcophaga (Heteronychia)* (Diptera: Sarcophagidae), with description of six new species. Zootaxa 2778(1): 1–57. <https://doi.org/10.11646/zootaxa.2778.1.1>
- Whitmore D, Pape T, Cerretti P (2013) Phylogeny of *Heteronychia*: the largest lineage of *Sarcophaga* (Diptera: Sarcophagidae). Zoological Journal of the Linnean Society 169(3): 604–639. <https://doi.org/10.1111/zoj.12070>
- Whitmore D, Plummer S, Clemons L (2020) First records of *Sarcophaga (Heteronychia) bulgarica* (Enderlein) (Diptera, Sarcophagidae) from Great Britain. Dipterists Digest 27: 223–229.