

The systematic position of *Cryptopotamon anacoluthon* (Kemp, 1918), with the description of a new species of *Sinolapotamon* Tai & Sung, 1975 (Crustacea, Decapoda, Brachyura, Potamidae) from southern China

Yuan-Biao Lu¹, Yi-Xuan Zhang¹, Jie-Xin Zou^{1,2} 

¹ Research Laboratory of Freshwater Crustacean Decapoda & Paragonimus, School of Basic Medical Sciences, Nanchang University, Nanchang City, Jiangxi Province, China

² Jiangxi Provincial Key Laboratory of Experimental Animals, Nanchang City, Jiangxi Province, China

Corresponding author: Jie-Xin Zou (jxzou@ncu.edu.cn)



Academic editor: Sameer Pati

Received: 10 February 2023

Accepted: 17 May 2023

Published: 8 June 2023

ZooBank: <https://zoobank.org/7BCF174D-E335-49B2-9799-4F7E99D01FB7>

Citation: Lu Y-B, Zhang Y-X, Zou J-X (2023) The systematic position of *Cryptopotamon anacoluthon* (Kemp, 1918), with the description of a new species of *Sinolapotamon* Tai & Sung, 1975 (Crustacea, Decapoda, Brachyura, Potamidae) from southern China. ZooKeys 1166: 155–173. <https://doi.org/10.3897/zookeys.1166.101737>

Copyright: © Yuan-Biao Lu et al.

This is an open access article distributed under terms of the Creative Commons Attribution License ([Attribution 4.0 International – CC BY 4.0](https://creativecommons.org/licenses/by/4.0/)).

Abstract

The systematics of the potamid freshwater crab *Cryptopotamon anacoluthon* (Kemp, 1918) is clarified, and its generic position in *Sinolapotamon* Tai & Sung, 1975, is confirmed based on morphological comparisons, geographical information and phylogenetic analyses. A new species of *Sinolapotamon*, *Sinolapotamon cirratum* **sp. nov.** is described from the Guangxi Zhuang Autonomous Region of China. *Sinolapotamon cirratum* **sp. nov.** is distinguished from its congeners by the combination of characters of its carapace, third maxilliped, anterolateral margin, and unique male first gonopod. Phylogenetic analyses based on partial COX1, 16S rRNA and 28S rRNA genes also support the species as new.

Key words: Freshwater crabs, molecular phylogeny, systematics, taxonomy

Introduction

Located in the southwest border region of China, with a warm climate, abundant precipitation, and a high percentage of forest coverage and karst landforms, Guangxi (Fig. 1) provides a suitable living environment for freshwater crabs. In China, which has the highest species richness of freshwater crabs globally (Cumberlidge et al. 2011), the species richness in Guangxi (43 species, including *S. cirratum* sp. nov.) is surpassing that of Taiwan (41 species) and only lower than that of Yunnan (74 species) (Chu et al. 2018; Wang et al. 2019; Cai et al. 2021). Rong County, situated in southeastern Guangxi and adjoining Guangdong Province, is the type locality of *Sinolapotamon cirratum* sp. nov. (Fig. 1). Hong Kong (Fig. 1), located in the south of China, consists of Hong Kong Island, Kowloon, the New Territories and 262 surrounding islands. The New Territories and Kowloon are connected to mainland China. It is worth noting that the New Territories is connected to Shenzhen, Guangdong Province.

The genus *Sinolapotamon* Tai & Sung, 1975, was described, with the type species as *Potamon (Geothelphusa) patellifer* Wu, 1934 (type locality, Luocheng).




Figure 1. Map showing collection sites of *Sinolapotamon* used in this study and the known collection sites of *Cryptopotamon anacoluthon*. Part of the studied region (Shenzhen and Hong Kong) is enlarged.

Zhu et al. (2010) described two species of *Sinolapotamon* from Guangxi: *S. auriculatum* Zhu, Naruse & Zhou, 2010, from Shanglin, and *S. palmatum* Zhu, Naruse & Zhou, 2010, from Liuzhou (Fig. 1). There has been disagreement regarding the generic position of *Cryptopotamon anacoluthon* (Kemp, 1918). Ng and Dudgeon (1992), while describing *Cryptopotamon* Ng & Dudgeon, 1992, included only *Potamon (Potamon) anacoluthon* Kemp, 1918, from Hong Kong (Fig. 1). Dai (1999) pointed out that *Cryptopotamon* is a synonym of *Sinolapotamon*, considering that the differences between these two genera could only be regarded as interspecific differences. Ng et al. (2008), however, continued to place *Potamon (Potamon) anacoluthon* in *Cryptopotamon*. Based on phylogenetic methods, we reconstructed the phylogenetic relationships of *Sinolapotamon*, which confirm the generic position of *Potamon (Potamon) anacoluthon* in *Sinolapotamon* and recognize a new species. *Sinolapotamon cirratum* sp. nov. is described here based on morphology and genetic data.

Materials and methods

Specimens were collected from the Duqiaoshan Forest Park and Silaochong, both in Rong County, Yulin City, Guangxi Zhuang Autonomous Region, China. The two sites are so close that they appear as one dot in Fig. 1 (about 5 km). In addition, the sites of the specimens of *C. anacoluthon* referred to in Ng and Dudgeon (1992) were added to the map (Fig. 1). The two sites are: Tai Po Kau Forest Reserve stream, New Territories, Hong Kong; and the stream at Wu Kwai

Sha, New Territories, Hong Kong. The linear distances from the ‘Shenzhen’ site to the two ‘Hong Kong’ sites are between 30–40 km (Fig. 1). Ethanol (95%) was used to preserve the collected specimens, which were deposited in the Department of Parasitology of the Medical College of the Nanchang University, Jiangxi, China (NCU MCP). Materials used herein, except for the new species and *S. anacoluthon*, are as follows: *S. patellifer*, 1 ♂, Yangshuo County, Guangxi Province, collection date not clear, NCU MCP 407301; *S. auriculatum*, 2 ♂♂, Shanglin County, Guangxi Province, July 2006, NCU MCP 72301, 72302; *S. palmatum*, 2 ♂♂, Liuzhou City, Guangxi Province, May 2018, NCU MCP 415301, 415302. Carapace width and length were measured in millimeters. The terminology used herein primarily follows that of Dai (1999) and Davie et al. (2015). The abbreviations used for the male first gonopod and male second gonopod are G1 and G2, respectively.

Approximately 50 mg of muscle tissue was excised from ambulatory legs. Total genomic DNA was extracted using the D3373-01 Mollusc DNA Kit (Omega Biotek, Inc., Norcross, USA). In our study, three fragments of target genes were amplified, including the mitochondrial COX1 and 16S rRNA genes and nuclear 28S rRNA gene. The primers and annealing temperatures used are presented in Table 1. Notably, the COX1 primers used were slightly modified based on the primers LCO1490 and HCO2198. A base T in the primer HCO2198 was replaced with a degenerate base Y (Folmer et al. 1994; Yang 2011). We performed phylogenetic analyses with the single-gene dataset (COX1) and 3-gene combined dataset (COX1, 16S rRNA and 28S rRNA). All molecular data are presented in Table 2. Sequences were aligned using ClustalW (Thompson et al. 2003), and the conserved regions were selected with Gblocks 0.91b (Castresana 2000) using the default settings. The optimal model for Bayesian inference (BI) analysis was determined using MrModeltest v. 2.3 (Nylander 2004) on the basis of the Akaike information criterion (AIC). The best-fitting model was GTR+G+I for both datasets. MrBayes v. 3.2.6 (Ronquist et al. 2012) was employed to perform BI analysis, and four Monte Carlo Markov chains of 2 000 000 generations were run with sampling every 1000 generations. The first 25% of generations were discarded as burn-in. Tracer v. 1.6 (Rambaut et al. 2013) was used to examine the sampling parameter. The optimal model, identified with MEGA X, for maximum likelihood (ML) analysis was also GTR+G+I for both datasets (Kumar et al. 2018). MEGA X was also employed to construct the ML tree based on 1000 bootstrap replicates and to calculate the pairwise distance based on the Kimura 2-parameter (K2P) model (Kumar et al. 2018). The map of the study area was prepared using ArcMap v. 10.2.

Table 1. Information on the primers used and annealing temperatures.

Gene	Primer name	Sequence 5'-3'	Product Length (bp)	Annealing Temperature (°C)	Reference
COI	LC01490	GGTCAACAAATCATAAAGATATTGG	~700	52	Folmer et al. (1994) and Yang (2011)
	HCO2198'	TAAACTTCAGGGTGACCAAAAAAYCA			
16S rRNA	1471	CCTGTTTANCAAAAACAT	~600	50	Crandall and Fitzpatrick (1996)
	1472	AGATAGAAACCAACCTGG			
28S rRNA	F28S	CAGCCCTAAGCAGGTGGTAAACT	~1000	53	Ji et al. (2016)
	R28S	CCATCTTTCGGGTCCCAACAT			

Table 2. Collection information and GenBank accession number of the species used for phylogenetic analyses.

Species	GenBank number	Voucher number	Locality	Reference
<i>Candidiopotamon okinawense</i> Minei, 1973	COI, MN737145	–	Okinawa, Japan	Zhang et al. 2020
<i>Candidiopotamon rathbunae</i> (De Man, 1914)	COI, AB433579	NCHU:Z00L:13146	Taiwan	Shih et al. 2011a
<i>Chinapotamon depressum</i> (Dai, Song, Li & Liang, 1980)	COI, MZ350918	–	Guangdong, China	Pan et al. 2021
<i>Geothelphusa dehaani</i> (White, 1847)	COI, AB187570	–	Tokyo, Japan	Segawa and Aotsuka 2005
<i>Geothelphusa minei</i> Shy & Ng, 1998	COI, AB625725	Gmn6	Ryukyu	Shih et al. 2011b
<i>Huananpotamon angulatum</i> (Dai, Chen, Song, Fan, Lin & Zeng, 1979)	COI, AB433576	NCHU:Z00L:13139	Fujian, China	Shih et al. 2011b
<i>Nanhaipotamon guangdongense</i> Dai, 1997	COI, MK226144	Ns7	Guangdong, China	Huang et al. 2018
<i>Nanhaipotamon hongkongense</i> (Shen, 1940)	COI, AB470509	Nh3	Hong Kong	Shih et al. 2011b
<i>Nanhaipotamon pingyuanense</i> Dai, 1997	COI, AB470513	Npy3	Guangdong, China	Shih et al. 2011b
<i>Neotiwariopotamon jianfengense</i> Dai & Naiyanetr, 1994	COI, MZ350933	–	Hainan, China	Pan et al. 2021
<i>Sinopotamon yaanense</i> (Chung & Ts'ao, 1962)	COI, LC155173	SC8	Sichuan, China	Shih et al. 2016
<i>Longpotamon nanlingense</i> (Dai & Jiang, 1991)	COI, LC155196	SPx173	Hunan, China	Shih et al. 2016
<i>Cantopotamon zhuhaiense</i> Huang, Ahyong & Shih, 2017	COI, LC342051	SYSBM:1439	Guangdong, China	Huang et al. 2017
<i>Parapotamon spinescens</i> (Calman, 1905)	COI, LC155209	PP4	Yunnan, China	Shih et al. 2016
<i>Tenuilapotamon latilum</i> (Chen, 1980)	COI, LC155206	TNL1	Hubei, China	Shih et al. 2016
<i>Tiwariopotamon edostilus</i> Ng & Yeo, 2001	COI, AB896762	TWs6	Haiphong, Vietnam	Shih and Do 2014
<i>Tiwariopotamon pingguoense</i> Dai & Naiyanetr, 1994	COI, LC145315	TWs13	Guangxi, China	Van et al. 2016
<i>Yarepotamon gracillipa</i> (Dai, Song, Li & Liang, 1980)	COI, AB433577	–	Guangxi, China	Direct Submission
<i>Apotamonautes hainanensis</i> (Parisi, 1916)	COI, MN737137	–	Hainan, China	Zhang et al. 2020
<i>Chinapotamon maolanense</i> Zou, Bai & Zhou, 2018	COI, MT134100	–	Guizhou, China	Cui et al. 2020
<i>Indochinamon chinghungense</i> (Dai, Song, He, Cao, Xu & Zhong, 1975)	COI, MZ350925	–	Yunnan, China	Pan et al. 2021
<i>Indochinamon daweshanense</i> (Dai, 1995)	COI, MZ350926	–	Yunnan, China	Pan et al. 2021
<i>Potamiscus yiwuensis</i> Dai & Cai, 1998	COI, MN737136	–	Yunnan, China	Zhang et al. 2020
<i>Qianguimon elongatum</i> Huang, 2018	COI, MZ350943	–	Guizhou, China	Pan et al. 2021
<i>Vadosapotamon sheni</i> (Dai & Chen in Dai, Chen, Liu, Luo, Yi, Liu, Gu & Liu, 1990)	COI, MZ350958	–	Sichuan, China	Pan et al. 2021
<i>Longpotamon yangtsekiense</i> (Bott, 1967)	COI, EU676302	TB5	China	Direct Submission
<i>Tenuilapotamon joshuiense</i> (Dai, Song, He, Cao, Xu & Zhong, 1975)	COI, MZ350951	–	Hunan, China	Pan et al. 2021
<i>Neotiwariopotamon whiteheadi</i> (Parisi, 1916)	COI, MZ350934	–	Hainan, China	Pan et al. 2021
<i>Huananpotamon lichuanense</i> Dai, Zhou & Peng, 1995	COI, MN737141	–	Jiangxi, China	Zhang et al. 2020
<i>Johora singaporensis</i> Ng, 1986	COI, MG010237	JSIN_BT01	Singapore	Tay et al. 2018
<i>S. cirratum</i> sp. nov.	COI, OP425670	Slsp01	Guangxi, China	This study
	COI, OP425672	Slsp02	Guangxi, China	This study
	COI, OP425671	Slsp03	Guangxi, China	This study
<i>S. auriculatum</i> Zhu, Naruse & Zhou, 2010	COI, OP425667	Slac01	Guangxi, China	This study
	COI, OP376822	Slpm02	Guangxi, China	This study
<i>S. anacoluthon</i> (Kemp, 1918)	COI, OP425668	Slal01	Guangdong China	This study
	COI, OP425669	Slal02	Guangdong China	This study
<i>S. patellifer</i> (Wu, 1934)	COI, MK883709	–	Guangxi, China	Ji et al. 2019
<i>Candidiopotamon okinawense</i> Minei, 1973	16S, AB208627	Co	Okinawa, Japan	Shih et al. 2006
<i>Candidiopotamon rathbunae</i> (De Man, 1914)	16S, AB208589	TPWL1	Taiwan	Shih et al. 2006
<i>Chinapotamon depressum</i> (Dai, Song, Li & Liang, 1980)	16S, KT586287	–	Guangxi, China	Ji et al. 2016
<i>Geothelphusa dehaani</i> (White, 1847)	16S, AB535460	Gd21	Kagoshima, Japan	Ng et al. 2010
<i>Geothelphusa minei</i> Shy & Ng, 1998	16S, AB625677	Gmn8	Ryukyu	Shih et al. 2011b
<i>Huananpotamon angulatum</i> (Dai, Chen, Song, Fan, Lin & Zeng, 1979)	16S, AB433555	NCHU:Z00L:13139	Fujian, China	Shih et al. 2011b
<i>Nanhaipotamon hongkongense</i> (Shen, 1940)	16S, AB212869	NHHK	Hong Kong	Shih et al. 2005
<i>Nanhaipotamon pingyuanense</i> Dai, 1997	16S, AB265237	NPy	Guangdong, China	Shih et al. 2007
<i>Neotiwariopotamon jianfengense</i> Dai & Naiyanetr, 1994	16S, KT586289	–	Hainan, China	Ji et al. 2016
<i>Sinopotamon yaanense</i> (Chung & Ts'ao, 1962)	16S, KT586263	02	Sichuan, China	Ji et al. 2016
<i>Longpotamon nanlingense</i> (Dai & Jiang, 1991)	16S, KT586180	01	Hunan, China	Ji et al. 2016
<i>Tenuilapotamon latilum</i> (Chen, 1980)	16S, AB428468	–	Hubei, China	Shih et al. 2009
<i>Longpotamon yangtsekiense</i> Bott, 1967	16S, KT586268	02	Jiangsu, China	Ji et al. 2016
<i>Tenuilapotamon joshuiense</i> (Dai, Song, He, Cao, Xu & Zhong, 1975)	16S, ON024657	NCU MCP 430301	Hunan, China	Direct Submission

Species	GenBank number	Voucher number	Locality	Reference
<i>S. cirratum</i> sp. nov.	16S, OP467587	Slsp01	Guangxi, China	This study
	16S, OP467588	Slsp02	Guangxi, China	This study
	16S, OP467584	Slsp03	Guangxi, China	This study
<i>S. auriculatum</i> Zhu, Naruse & Zhou, 2010	16S, OP467583	Slac01	Guangxi, China	This study
	16S, OP467590	Slpm02	Guangxi, China	This study
<i>S. anacoluthon</i> (Kemp, 1918)	16S, OP467585	Slal01	Guangdong, China	This study
	16S, OP467586	Slal02	Guangdong, China	This study
<i>S. patellifer</i> (Wu, 1935)	16S, MK883709	–	Guangxi, China	Ji et al. 2019
<i>Candidiopotamon okinawense</i> Minei, 1973	28S, AB503625	Co	Okinawa, Japan	Direct Submission
<i>Candidiopotamon rathbunae</i> (De Man, 1914)	28S, AB503628	Cr	Taiwan	Direct Submission
<i>Chinapotamon depressum</i> (Dai, Song, Li & Liang, 1980)	28S, KT586427	–	Guangxi, China	Ji et al. 2016
<i>Geothelphusa dehaani</i> (White, 1847)	28S, AB503607	Gdmms	Kagoshima, Japan	Direct Submission
<i>Geothelphusa minei</i> Shy & Ng, 1998	28S, AB503619	GmnlG	Okinawa, Japan	Direct Submission
<i>Huananpotamon angulatum</i> (Dai, Chen, Song, Fan, Lin & Zeng, 1979)	28S, AB576807	Hua2	Fujian, China	Shih et al. 2011b
<i>Nanhaipotamon hongkongense</i> (Shen, 1940)	28S, AB551401	Nh3	Hong Kong	Shih et al. 2011b
<i>Nanhaipotamon pingyuanense</i> Dai, 1997	28S, AB551405	Npy2	Guangdong, China	Shih et al. 2011b
<i>Neotiwariopotamon jianfengense</i> Dai & Naiyanetr, 1994	28S, KT586429	–	Hainan, China	Ji et al. 2016
<i>Sinopotamon yaanense</i> (Chung & Ts'ao, 1962)	28S, KT586416	04	Sichuan, China	Ji et al. 2016
<i>Longpotamon nanlingense</i> (Dai & Jiang, 1991)	28S, KT586368	01	Hunan, China	Ji et al. 2016
<i>Tenuilapotamon latilum</i> (Chen, 1980)	28S, MW540828	NCU MCP 66301	Hubei, China	Direct Submission
<i>Longpotamon yangtsekiense</i> (Bott, 1967)	28S, KT586417	02	Jiangsu, China	Ji et al. 2016
<i>Tenuilapotamon joshuiense</i> (Dai, Song, He, Cao, Xu & Zhong, 1975)	28S, ON033004	NCU MCP 430301	Hunan, China	Direct Submission
<i>S. cirratum</i> sp. nov.	28S, OP578215	Slsp01	Guangxi, China	This study
	28S, OP578219	Slsp02	Guangxi, China	This study
	28S, OP578212	Slsp03	Guangxi, China	This study
<i>S. auriculatum</i> Zhu, Naruse & Zhou, 2010	28S, OP578218	Slac01	Guangxi, China	This study
	28S, OP578217	Slpm02	Guangxi, China	This study
<i>S. anacoluthon</i> (Kemp, 1918)	28S, OP578213	Slal01	Guangdong, China	This study
	28S, OP578214	Slal02	Guangdong, China	This study
<i>S. patellifer</i> (Wu, 1936)	28S, OP578216	Slpl01	Guangxi, China	This study

Results

Systematics

Family Potamidae Ortmann, 1896

Sinolapotamon Tai & Sung, 1975

Cryptopotamon Ng & Dudgeon, 1992: 741, figs 3B, 4, 5.

Type species. *Potamon* (*Geothelphusa*) *patellifer* Wu, 1934, by original designation.

Sinolapotamon anacoluthon (Kemp, 1918)

Figs 2, 7C

Potamon (*Potamon*) *anacoluthon* Kemp, 1918: 243, fig. 5.

Cryptopotamon anacoluthon Ng & Dudgeon, 1992: 741, figs 3B, 4, 5. — Ng et al. 2008: 161 (list).

Sinolapotamon anacoluthon Dai, 1999: 150, fig. 79.

Material examined. CHINA • 4 ♂♂ (18.40 × 16.34 mm, 20.26 × 18.40 mm, 21.64 × 18.60 mm, 19.26 × 17.04 mm); Yangtaishan Forest Park, Shenzhen,




Figure 2. *Sinolapotamon anacoluthon*, male (25.84 × 22.76 mm) (NCU MCP 434101) **A** overall dorsal habitus **B** overall ventral habitus **C** frontal view of cephalothorax **D** outer view of chelipeds **E** sternopleonal cavity with G1. Scale bars: 1 cm.

Guangdong Province; 22.6587°N, 113.9837°E; July 2022; Sheng Yu leg.; NCU MCP 434001–434004 • 1 ♂ (25.84 × 22.76 mm); same collection data as above; NCU MCP 434101 • 3 ♀♀ (26.34 × 23.58 mm, 28.84 × 24.38 mm, 24.31 × 20.95 mm); same collection data as above; NCU MCP 434102–434104.

Diagnosis. Carapace gently convex, regions indistinct. Cervical groove shallow, indistinct; H-shaped groove depressed and distinct (Fig. 2A). Epigastric cristae weak, postorbital cristae flat, indistinct. External orbital angle triangular, with about 5 small granules. Epibranchial tooth sharp, distinctly separated with external orbital angle by V-shaped gap. Anterolateral margin of carapace cristate, with about 12 granules (Fig. 2A). Maxilliped 3 exopod reaching nearly 1/3 of merus length, with long flagellum (Fig. 2C). Chelipeds (pereiopod 1) strongly unequal (Fig. 2A, B, D). G1 slender, subterminal segment about 1.1 times as long as terminal segment; 2 lobes of terminal segment strongly unequal, dorsal lobe longitudinally extended, oval shaped, ventral lobe sharp and short, reaching 3/7 of terminal segment (Figs 2E, 7C).

Ecology. The species is usually inhabiting the clear hill streams at an altitude below 50 m. Stones could serve as shelter and leaf mould could serve as food (Dai, 1999).

Distribution. China: Shenzhen of Guangdong Province (present record) and Hong Kong.

Remarks. The specimens from Shenzhen, with gently convex dorsal surface of carapace, indistinct postorbital cristae, sharp epibranchial tooth, unequal lobes of the terminal segment of the G1 (Fig. 2), and other characteristics, agree well with the descriptions and illustrations in Ng and Dudgeon (1992) and Dai (1999). The ratio of the subterminal segment to the terminal segment of G1 calculated in this study is 1.1 (Fig. 7C), which is equal to that in Dai (1999) and slightly smaller than that in Ng and Dudgeon (1992) (1.17). Although the specimens are not from Hong Kong, they could still be determined as *S. anacoluthon* based on morphological examination and the proximity of their collection site to Hong Kong (Fig. 1).

Ng and Dudgeon (1992) listed the differences between *Cryptopotamon* and *Sinolapotamon*, including carapace, epigastric cristae, postorbital cristae, epibranchial tooth, and the ratio of the subterminal segment to the terminal segment of the G1. We, however, noticed that those differences are interspecific, while two or more species sharing the same character state with the remaining species is not. For instance, *S. anacoluthon* has a gently convex carapace similar to that of *S. cirratum* sp. nov. but different from the remaining congeners (Figs 2A, 3A). The weak epigastric cristae of *S. anacoluthon* are consistent with those of *S. auriculatum* and *S. cirratum* sp. nov. (Fig. 2A; see Zhu et al. 2010: figs 1a, 6a). The indistinct postorbital cristae of *S. anacoluthon* are comparable with those of *S. auriculatum* and *S. palmatum* (Fig. 2A; see Zhu et al. 2010: figs 1a, 6a). The sharp epibranchial tooth is consistent with that of *S. auriculatum* (Fig. 2A; see Zhu et al. 2010: fig. 1a). The different ratios of the subterminal segment to the terminal segment of the G1 could only be regarded as interspecific differences. Most importantly, all five species have accordant fundamental types of G1 (Fig. 7).

***Sinolapotamon cirratum* sp. nov.**

<https://zoobank.org/25AEC22F-BAAA-4DF5-8580-8334F8DBB9EC>

Figs 3–6, 7A, 8

Type material. Holotype: CHINA • ♂ (17.90 × 15.50 mm); Guangxi Zhuang Autonomous Region, Yulin City, Rong County, Duqiaoshan Forest Park; 22.8019°N, 110.6098°E; October 2022; Yi-Xuan Zhang leg.; NCU MCP 434201.

Paratypes: CHINA • 1 ♀ (17.42 × 15.45 mm); same collection data as for holotype; NCU MCP 433901 • 1 ♀ (23.74 × 20.30 mm); same collection data as for holotype; NCU MCP 433902 • 4 ♂♂ (22.54 × 19.55 mm, 22.92 × 19.80 mm, 23.05 × 20.24 mm, 18.00 × 16.02 mm); same collection data as for holotype; NCU MCP 433903–433906.

Other material. CHINA • 7 ♂♂ (21.96 × 19.12 mm, 13.36 × 11.93 mm, 15.76 × 13.18 mm, 17.34 × 14.99 mm, 17.26 × 14.94 mm, 17.93 × 14.91 mm, 20.73 × 17.62 mm); same collection data as for holotype; NCU MCP 434202–434208 • 8 ♂♂ (14.99 × 12.90 mm, 9.48 × 8.65 mm, 9.61 × 8.54 mm, 9.81 × 8.91 mm, 8.71 × 7.53 mm, 8.82 × 7.77 mm, 9.74 × 7.38 mm, 10.42 × 8.79 mm); Guangxi Zhuang Autonomous Region, Yulin City, Rong County, Silaochong, small stream; 22.8263°N, 110.6065°E; November 2018; Jie-Xin Zou et al. leg.; NCU MCP 416001–416008.

Diagnosis. Carapace subquadrate, regions indistinct; dorsal surface gently convex, anterolateral region weakly rugose. Cervical groove shallow and wide;




Figure 3. *Sinolapotamon cirratum* sp. nov., holotype male (17.90 × 15.50 mm) (NCU MCP 434201) **A** overall dorsal habitus **B** frontal view of cephalothorax. Scale bars: 1 cm.

H-shaped groove shallow (Figs 3A, 5A). Epigastric cristae distinct, separated from postorbital cristae by narrow gap; epibranchial region slightly depressed; mesogastric region gently convex. External orbital angle triangular, distinctly separated from anterolateral margin by wide notch. Anterolateral margin of carapace distinctly cristate, lined with approximately 20 granules (Figs 3A, 5A). Maxilliped 3 exopod reaching nearly 1/2 of merus length, with long flagellum, slightly longer than width of merus (Fig. 4B). Chelipeds (pereiopod 1) strongly




Figure 4. *Sinolapotamon cirratum* sp. nov., holotype male (17.90 × 15.50 mm) (NCU MCP 434201) **A** outer view of chelipeds **B** left third maxilliped **C** anterior thoracic sternum, pleonal somites 4–6 and telson **D** sternopleonal cavity with G1. Scale bars: 5 mm.

unequal in males, subequal in females (Figs 3A, 4A, 5A). G1 slender, subterminal segment about 1.7 times as long as terminal segment; 2 lobes of terminal segment strongly unequal, dorsal lobe longitudinally extended, oval shaped, ventral lobe blunt, reaching 1/2 of terminal segment (Figs 6A–D, 7A). Female vulvae ovate, medium-sized, occupying anterior 2/3 length of sternite 6 (Fig. 5B).

Description. Carapace subquadrate, nearly 1.2 times as wide as long; surface generally smooth, regions indistinct; dorsal surface slightly convex, with tiny pits, anterolateral region weakly rugose. Cervical groove shallow and wide; H-shaped groove shallow. Front gently deflexed; frontal margin slightly rimmed, weakly bilobed in dorsal view (Figs 3A, 5A). Epigastric cristae low and weak, separated from postorbital cristae by narrow gap; postorbital cristae laterally expanded, not fused with epibranchial tooth. Epibranchial region gently depressed; mesogastric region gently convex. External orbital angle bluntly triangular, distinctly separated from anterolateral margin by V-shaped wide gap (Figs 3A, 5A). Epibranchial tooth distinct, bluntly angular. Anterolateral margin of carapace distinctly cristate, lined with approximately 20 granules; bent inward posteriorly. Posterolateral surface smooth, with oblique striae, converging towards posterior carapace margin (Figs 3A, 5A). Orbits large; supraorbital and infraorbital margins cristate. Sub-orbital, pterygostomial, subhepatic regions covered with striae. Epistome posterior margin narrow longitudinally; median lobe triangular, lateral margins sinuous (Fig. 3B).




Figure 5. *Sinolapotamon cirratum* sp. nov., paratype female (17.42 × 15.45 mm) (NCU MCP 433901) **A** overall dorsal habitus **B** sternopleonal cavity with vulvae. Scale bars: 1 cm.

Maxilliped 3 exopod reaching nearly 1/2 of merus length, with long flagellum, slightly longer than width of merus. Merus subrectangular, 2 times as wide as long. Ischium subtrapezoidal, about 1.4 times as long as wide, with distinct sulcus (Fig. 4B).

Chelipeds (pereiopod 1) strongly unequal in males, subequal in females. Merus trigonal in cross section. Carpus surface gently depressed, with spine at inner distal angle and spinule at base in both males and females. Palm of larger chela about 1.3–1.5 times as long as high in males, 1.3–1.6 times in females. Dactylus of larger chela 0.6–1.0 times as long as palm in males, practically same proportion in females. Inner margin of fingers lined with granular teeth; fingers of larger chela leaving small gap while smaller one without gap when closed in both males and females (Figs 3A, 4A, 5A).

Ambulatory legs (pereiopod 2–5) slender, with setae; pereiopod 3 longest, merus 0.5–0.6 times as long as carapace length. Pereiopod 5 propodus about 2.0 times as long as broad in both males and females, 0.8–0.9 times as long as dactylus; dactylus gently curved (Figs 3A, 5A).




Figure 6. Gonopods of *Sinolapotamon cirratum* sp. nov., holotype male (17.90 × 15.50 mm) (NCU MCP 434201) **A** ventral view of left G1 **B** ventral view of terminal segment of left G1 **C** dorsal view of left G1 **D** dorsal view of terminal segment of left G1 **E** ventral view of left G2. Scale bars: 1 mm.



Figure 7. Left G1s (ventral view) of known species of *Sinolapotamon* **A** *Sinolapotamon cirratum* sp. nov., holotype (NCU MCP 434201) **B** *S. patellifer* (Yangshuo, NCU MCP 407301) **C** *S. anacoluthon* (Shenzhen, NCU MCP 434001) **D** *S. auriculatum* (Shanglin, NCU MCP 72301) **E** *S. palmatum* (Liuzhou, NCU MCP 415301).

Male thoracic sternum generally smooth, pitted; sternite 1 triangular; sternite 2–3 fused without visible sutures. Male sternopleonal cavity relatively deep, exceeding imaginary line connecting posterior edges of cheliped coxae base. Median longitudinal suture of sternites 7, 8 deep. Tubercle of abdominal lock positioned at approaching mid-length of sternite 5. Sutures between sternites in female indistinct (Fig. 4C, D). Female vulvae ovate, medium-sized, occupying anterior 2/3 length of sternite 6 (Fig. 5B).

Male pleon and telson triangular; pleonal somites 3–6 gradually narrowed longitudinally, lateral margins forming gently concave line with thoracic sternum; pleonal somite 6 about 2.2 times as wide as long; telson about 1.3 times as wide as long (Fig. 4C). Female pleon and telson broadly ovate (Fig. 5B).

G1 slender, tip of terminal segment exceeding beyond pleonal locking tubercle and suture between thoracic sternites 4 and 5 (Fig. 4D). Subterminal segment about 1.7 times as long as terminal segment; edges of dorsal lobe curled; 2 lobes of terminal segment strongly unequal, dorsal lobe longitudinal-




Figure 8. **A** *Sinolapotamon cirratum* sp. nov. in the wild **B** general habitat of *Sinolapotamon cirratum* sp. nov.

ly extended, oval, ventral lobe blunt, reaching 1/2 length of terminal segment (Fig. 6A–D). G2 slender, longer than G1 (Fig. 6A, E).

Remarks. Consistent with the diagnostic characters of *Sinolapotamon*, *Sinolapotamon cirratum* sp. nov. has a gently convex dorsal surface, long flagellum of the third maxilliped exopod and unequal lobes of the G1 terminal segment (Figs 3A, 4B, 7A). The dorsal lobe of the G1 terminal segment in *S. cirratum* sp. nov. is long and oval, which is similar to that of *S. anacoluthon*. The two species can nevertheless be distinguished by the ratio of the subterminal segment to the terminal segment of G1, which is 1.7 in *S. cirratum* sp. nov. and 1.1 in *S. anacoluthon* (Fig. 7A, C). When compared with *S. patellifer*, *S. auriculatum* and *S. palmatum*, the new species could be easily distinguished by the shape of the dorsal lobes and ventral lobes. The ventral lobe of *S. cirratum* sp. nov. is bluntly angular, while those of the other species in *Sinolapotamon* are pointed or shortly pointed (Fig. 7A, B, D, E). They also differ in comparative length of the ventral lobe relative to the terminal segment of the G1 (see Table 3). Additional differences among the known species of *Sinolapotamon* are provided in Table 3.

Etymology. The new species is named *Sinolapotamon cirratum* sp. nov. because of the curled edges of the dorsal lobe of the G1. In the Latin, ‘cirratus’ means ‘curled’.

Ecology. The specimens were collected from puddles in the Duqiaoshan Forest Park. These crabs live in the shallow water or under the wet stones (Fig. 8A, B).

Table 3. Morphological differences between five species of *Sinolapotamon*.

Species	<i>Sinolapotamon cirratum</i> sp. nov.	<i>S. patellifer</i> (cf. Dai 1999: fig. 78)	<i>S. anacoluthon</i> (cf. Ng and Dudgeon 1992: figs 4, 5)	<i>S. palmatum</i> (cf. Zhu et al. 2010: figs 6–9)	<i>S. auriculatum</i> (cf. Zhu et al. 2010: figs 1–4)
Flagellum of exopod of third maxilliped	slightly longer than width of merus (Fig. 4B)	slightly shorter than width of merus	exceeding width of merus	slightly longer than width of merus	shorter than width of merus
Anterolateral margin of carapace	distinctly cristate, lined with approximately 20 granules (Fig. 3A)	ridged, without conspicuous granules	cristate, lined with numerous small rounded granules	convex laterally, cristate, lined with fine granules	weakly convex laterally, cristate, lined with fine granules
Ratio of subterminal segment to terminal segment of G1	1.7 (Fig. 7A)	1.7 (Fig. 7B)	1.1 (Fig. 7C)	1.5 (Fig. 7E)	1.5 (Fig. 7D)
Terminal segment of G1	longitudinally extended oval (Fig. 7A)	oval (Fig. 7B)	longitudinally extended oval (Fig. 7C)	subcircular (Fig. 7E)	oval (Fig. 7D)
ventral lobe of G1	blunt, reaching 1/2 length of terminal segment (Fig. 7A)	sharp, reaching beyond proximal 2/3 length of terminal segment (Fig. 7B)	sharp and short, reaching 3/7 length of terminal segment (Fig. 7C)	sharp, reaching 5/6 length of terminal segment (Fig. 7E)	sharp, reaching proximal 1/2 length of terminal segment (Fig. 7D)

Distribution. China: Guangxi Zhuang Autonomous Region: Rong County, Yulin City.

Phylogenetic relationships

A single-gene dataset (COX1) and a 3-gene combined dataset (COX1, 16S rRNA, and 28S rRNA) were used to reconstruct the ML tree and BI tree, respectively. The topologies of the ML tree and BI tree based on the single-gene dataset and the 3-gene combined dataset were analogous. Both evolutionary trees based on the single-gene and 3-gene datasets offer strong evidence for the recognition of the new species as *Sinolapotamon cirratum* sp. nov., since it is clustered with the species of *Sinolapotamon* as a monophyletic clade. *Sinolapotamon patellifer* and *S. auriculatum* form a sister group. Notably, *S. anacoluthon* (previously *C. anacoluthon*) is in 'Clade *Sinolapotamon*', which provides supporting evidence for recognizing the species in *Sinolapotamon* (Figs 9, 10). The results show that pairwise genetic distances range from 0.0600–0.1106 within the genus *Sinolapotamon*, and the genetic distances between *Sinolapotamon cirratum* sp. nov. and its congeners range from 0.0728–0.0947 (Table 4). Phylogenetic analyses, therefore, provided evidence for the identification of *Sinolapotamon cirratum* sp. nov. as a new species.

Discussion

Previous studies on *Sinolapotamon* focused on morphological descriptions and lacked molecular evidence (Tai and Sung 1975; Ng and Dudgeon 1992; Zhu et al. 2010). For our study, we obtained sequences of the partial COX1, 16S rRNA and 28S rRNA genes of all the members of *Sinolapotamon*, thus compensating for this gap. Moreover, the taxonomic statuses of the new species and *S. anacoluthon* are demonstrated based on morphology, molecular phylogeny and geographical distribution.

Ng and Dudgeon (1992) listed the morphological differences between *Cryptopotamon* and *Sinolapotamon*, including a gently convex carapace against a strongly inflated carapace, and the extent of prominence of the epigastric and postorbital cristae. Dai (1999) stated that these differences could only be regarded as interspecific and that the fundamental types of G1 are accordant, thus considering *Cryptopotamon* as a synonym of *Sinolapotamon*. Ng et al. (2008), however, listed *S. anacoluthon* as belong to *Cryptopotamon*. We assessed the morphological differences among the five known species of *Sinolapotamon* (see Remarks above) (Table 3) and reconstructed the phylogenetic relationships in *Sinolapotamon*, in turn providing molecular evidence for transferring *C. anacoluthon* to *Sinolapotamon*. *Sinolapotamon anacoluthon* was previously recorded only from Hong Kong (Ng and Dudgeon 1992; Stanton et al. 2017), but we also collected this species in Shenzhen of the Guangdong Province. There is some geographical distance between *S. anacoluthon* (from Guangdong) and its congeners (from Guangxi). We, however, noticed that all the species of *Sinolapotamon* are distributed near the Pearl River Basin. We speculate that the Pearl River contributed to the spread of *Sinolapotamon*, but further surveys will be needed to validate this hypothesis.

Table 4. Pairwise genetic distances of known species of *Sinlapotamon*.

Species	1	2	3	4
<i>Sinlapotamon auriculatum</i>	–	–	–	–
<i>Sinlapotamon anacoluthon</i>	0.0802	–	–	–
<i>Sinlapotamon cirratum</i> sp. nov.	0.0890	0.0728	–	–
<i>Sinlapotamon palmatum</i>	0.1106	0.0692	0.0947	–
<i>Sinlapotamon patellifer</i>	0.0600	0.0863	0.0842	0.1067




Figure 9. Phylogenetic tree based on the COX1 gene. Topologies and branch lengths were obtained from BI analysis. Only values >50% are displayed.




Figure 10. Phylogenetic tree based on three genes (COX1, 16S rRNA and 28S rRNA). Topologies and branch lengths were obtained from BI analysis. Only values >50% are displayed.

Conclusion

In this study, a new species of *Sinolapotamon* is described from the Guangxi Zhuang Autonomous Region of China, based on its morphological characteristics, especially its unique G1 among congeners, and the results of phylogenetic analyses (phylogenetic tree based on COX1 and 3-gene combined datasets). In addition, the generic position of *Cryptopotamon anacoluthon* in *Sinolapotamon* is confirmed largely on the basis of its morphology, with further evidence from the genetic data. *Sinolapotamon* is now known by five species. Based on the geographical distributions of *Sinolapotamon*, there is still possibility to discover new species in Guangxi or Guangdong.

Acknowledgements

We thank Ruoying Cheng (Nanchang University) for giving advice in taking the photos. We thank Qihong Tan (Nanchang University) for helping with the morphological description of the new species. We thank Chao Huang (Australian Museum) for the suggestions for the revision of the manuscript. Finally, we thank the subject editor and reviewers for greatly improving our manuscript.

Additional information

Conflict of interest

No conflict of interest was declared.

Ethical statement

No ethical statement was reported.

Funding

This work was supported by the National Natural Science Foundation of China (no. 32060306 and 21866020) and the National Parasitic Resources Center (NPRC-2019-194-30).

Author contributions

Jiexin Zou: review and editing; Yuanbiao Lu: original draft; Yixuan Zhang: investigation.

Author ORCIDs

Jie-Xin Zou  <https://orcid.org/0000-0002-5549-2167>

Data availability

All of the data that support the findings of this study are available in the main text or Supplementary Information.

References

- Cai MR, Tan QH, Zou JX (2021) A new species of freshwater crab of the genus *Nanhaiapotamon* Bott, 1968 (Crustacea, Decapoda, Brachyura, Potamidae) from Longhai, Fujian Province, China. ZooKeys 1062: 11–30. <https://doi.org/10.3897/zookeys.1062.71171>

- Castresana J (2000) Selection of Conserved Blocks from Multiple Alignments for Their Use in Phylogenetic Analysis. *Molecular Biology and Evolution* 17(4): 540–552. <https://doi.org/10.1093/oxfordjournals.molbev.a026334>
- Chu KL, Ma XP, Zhang ZW, Wang PF, Lü LN, Zhao Q, Sun HY (2018) A checklist for the classification and distribution of China's freshwater crabs. *Shengwu Duoyangxing* 26(3): 274–282. <https://doi.org/10.17520/biods.2018062>
- Crandall KA, Fitzpatrick Jr JF (1996) Crayfish Molecular Systematics: Using a Combination of Procedures to Estimate Phylogeny. *Systematic Biology* 45(1): 1–26. <https://doi.org/10.1093/sysbio/45.1.1>
- Cui YY, Shi LB, Ji WB, Xu SX, Zhu CC, Zhou XM, Zou JX (2020) The complete mitochondrial genome of the freshwater crab *Chinapotamon maolanense* (Decapoda: Brachyura: Potamoidea). *Mitochondrial DNA, Part B, Resources* 5(3): 2357–2359. <https://doi.org/10.1080/23802359.2020.1773330>
- Cumberlidge N, Ng PKL, Yeo DCJ, Naruse T, Meyer KS, Esser LJ (2011) Diversity, endemism and conservation of the freshwater crabs of China (Brachyura: Potamidae and Gecarcinidae). *Integrative Zoology* 6(1): 45–55. <https://doi.org/10.1111/j.1749-4877.2010.00228.x>
- Dai AY (1999) *Fauna Sinica (Arthropoda. Crustacea. Malacostraca. Decapoda. Parathelphusidae. Potamidae)*. Science Press, Beijing, 501 pp. [In Chinese with English summary]
- Davie PJF, Guinot D, Ng PKL (2015) Anatomy and functional morphology of Brachyura. In: Castro P, Davie PJF, Guinot D, Schram FR, von Vaupel Klein JC (Eds) *Treatise on Zoology – Anatomy, Taxonomy, Biology. The Crustacea (Vol. 9) (Part C-I). Decapoda: Brachyura (Part 1)*. Brill, Leiden, The Netherlands, 11–163. https://doi.org/10.1163/9789004190832_004
- Folmer O, Black M, Hoeh W, Lutz R, Vrijenhoek R (1994) DNA primers for amplification of mitochondrial cytochrome c oxidase subunit I from diverse metazoan invertebrates. *Molecular Marine Biology and Biotechnology* 3: 294–299.
- Huang C, Ah Yong ST, Shih HT (2017) *Cantopotamon*, a new genus of freshwater crabs from Guangdong, China, with descriptions of four new species (Crustacea: Decapoda: Brachyura: Potamidae). *Zoological Studies* 56: e41. <https://doi.org/10.6620/zs.2017.56-41>
- Huang C, Wong KC, Ah Yong ST (2018) The freshwater crabs of Macau, with the description of a new species of *Nanhaipotamon* Bott, 1968 and the redescription of *Nanhaipotamon wupingense* Cheng, Yang, Zhong & Li, 2003 (Crustacea, Decapoda, Potamidae). *ZooKeys* 810: 91–111. <https://doi.org/10.3897/zookeys.810.30726>
- Ji YK, Sun YF, Gao W, Chu KL, Wang RC, Zhao Q, Sun HY (2016) Out of the Sichuan Basin: Rapid species diversification of the freshwater crabs in *Sinopotamon* (Decapoda: Brachyura: Potamidae) endemic to China. *Molecular Phylogenetics and Evolution* 100: 80–94. <https://doi.org/10.1016/j.ympev.2016.04.003>
- Ji WB, Cui YY, Xu SX, Jia XN, Zhu CC, Zhou XM, Zou JX (2019) The complete mitochondrial genome of the freshwater crab *Sinopotamon patellifer* (Decapoda: Brachyura: Potamoidea). *Mitochondrial DNA, Part B, Resources* 4(2): 2451–2453. <https://doi.org/10.1080/23802359.2019.1637791>
- Kumar S, Stecher G, Li M, Knyaz C, Tamura K (2018) MEGA X: Molecular Evolutionary Genetics Analysis across Computing Platforms. *Molecular Biology and Evolution* 35(6): 1547–1549. <https://doi.org/10.1093/molbev/msy096>
- Ng PKL, Dudgeon D (1992) The Potamidae and Parathelphusidae (Crustacea: Decapoda: Brachyura) of Hong Kong. *Invertebrate Systematics* 6(3): 741–768. <https://doi.org/10.1071/IT9920741>

- Ng PKL, Guinot D, Davie PJF (2008) Systema Brachyurorum: Part I. An annotated checklist of extant brachyuran crabs of the world. The Raffles Bulletin of Zoology 17 (Supplement): 1–286.
- Ng PKL, Shih HT, Naruse T, Shy JY (2010) Using molecular tools to establish the type locality and distribution of the endemic Taiwanese freshwater crab *Geothelphusa chiui* Minei, 1974 (Crustacea: Brachyura: Potamidae), with notes on the genetic diversity of *Geothelphusa* from Eastern Taiwan. Zoological Studies 49: 544–555.
- Nylander JAA (2004) MrModeltest v 2.3. Evolutionary Biology Centre, Uppsala University.
- Pan D, Shi BY, Du SY, Gu TY, Wang RX, Xing YH, Zhang Z, Chen JJ, Cumberlidge N, Sun HY (2021) Mitogenome phylogeny reveals Indochina Peninsula origin and spatiotemporal diversification of freshwater crabs (Potamidae: Potamiscinae) in China. Cladistics 38(1): 1–12. <https://doi.org/10.1111/cla.12475>
- Rambaut A, Drummond AJ, Suchard M (2013) Tracer v1. 6—MCMC trace analysis package. Institute of Evolutionary Biology, University of Edinburgh, UK.
- Ronquist F, Teslenko M, van der Mark P, Ayres DL, Darling A, Höhna S, Larget B, Liu L, Suchard MA, Huelsenbeck JP (2012) MrBayes 3.2: Efficient Bayesian Phylogenetic Inference and Model Choice Across a Large Model Space. Systematic Biology 61(3): 539–542. <https://doi.org/10.1093/sysbio/sys029>
- Segawa RD, Aotsuka T (2005) The mitochondrial genome of the Japanese freshwater crab, *Geothelphusa dehaani* (Crustacea: Brachyura): Evidence for its evolution via gene duplication. Gene 355: 28–39. <https://doi.org/10.1016/j.gene.2005.05.020>
- Shih HT, Do VT (2014) A new species of *Tiwaripotamon* Bott, 1970, from northern Vietnam, with notes on *T. vietnamicum* (Dang & Ho, 2002) and *T. edostilus* Ng & Yeo, 2001 (Crustacea, Brachyura, Potamidae). Zootaxa 3764(1): 26–38. <https://doi.org/10.11646/zootaxa.3764.1.2>
- Shih HT, Chen GX, Wang LM (2005) A new species of freshwater crab (Decapoda: Brachyura: Potamidae) from Dongyin Island, Matsu, Taiwan, defined by morphological and molecular characters, with notes on its biogeography. Journal of Natural History 39(31): 2901–2911. <https://doi.org/10.1080/00222930500214010>
- Shih HT, Hung HC, Schubart CD, Chen CA, Chang HW (2006) Intraspecific genetic diversity of the endemic freshwater crab *Candidiopotamon rathbunae* (Decapoda, Brachyura, Potamidae) reflects five million years of the geological history of Taiwan. Journal of Biogeography 33(6): 980–989. <https://doi.org/10.1111/j.1365-2699.2006.01472.x>
- Shih HT, Fang SH, Ng PKL (2007) Phylogeny of the freshwater crab genus *Somannia-thelphusa* Bott (Decapoda: Parathelphusidae) from Taiwan and the coastal regions of China, with notes on their biogeography. Invertebrate Systematics 21(1): 29–37. <https://doi.org/10.1071/IS06007>
- Shih HT, Yeo DCJ, Ng PKL (2009) The collision of the Indian plate with Asia: molecular evidence for its impact on the phylogeny of freshwater crabs (Brachyura: Potamidae). Journal of Biogeography 36(4): 703–719. <https://doi.org/10.1111/j.1365-2699.2008.02024.x>
- Shih HT, Ng PKL, Naruse T, Shokita S, Liu MY (2011a) Pleistocene speciation of freshwater crabs (Crustacea: Potamidae: Geothelphusa) from northern Taiwan and southern Ryukyus, as revealed by phylogenetic relationships. Zoologischer Anzeiger 250(4): 457–471. <https://doi.org/10.1016/j.jcz.2011.07.004>
- Shih HT, Zhou XM, Chen GX, Chien IC, Ng PKL (2011b) Recent vicariant and dispersal events affecting the phylogeny and biogeography of East Asian freshwater crab genus *Nanhaipotamon* (Decapoda: Potamidae). Molecular Phylogenetics and Evolution 58(3): 427–438. <https://doi.org/10.1016/j.ympev.2010.11.013>

- Shih HT, Huang C, Ng PKL (2016) A re-appraisal of the widely-distributed freshwater crab genus *Sinopotamon* Bott, 1967, from China, with establishment of a new genus (Crustacea: Decapoda: Potamidae). *Zootaxa* 4138(2): 309–331. <https://doi.org/10.11646/zootaxa.4138.2.5>
- Stanton DJ, Leven MR, Hui TCH (2017) Distribution of *Cryptopotamon anacoluthon* (Kemp, 1918) (Crustacea: Brachyura: Potamidae), a freshwater crab endemic to Hong Kong. *Journal of Threatened Taxa* 9(2): 9786–9794. <https://doi.org/10.11609/jott.3007.9.2.9786-9794>
- Tai AY, Sung YZ (1975) A preliminary study of the freshwater crabs as intermediate hosts of lung flukes from China. *Acta zoologica Sinica* 21: 169–178.
- Tay YC, Ng DJJ, Loo JB, Huang D, Cai Y, Yeo DCJ, Meier R (2018) Roads to isolation: Similar genomic history patterns in two species of freshwater crabs with contrasting environmental tolerances and range sizes. *Ecology and Evolution* 8(9): 4657–4668. <https://doi.org/10.1002/ece3.4017>
- Thompson JD, Gibson TJ, Higgins DG (2003) Multiple Sequence Alignment Using ClustalW and ClustalX. *Current Protocols in Bioinformatics* 00: 2.3.1–2.3.22. <https://doi.org/10.1002/0471250953.bi0203s00>
- Van Tu Do HTS, Huang C (2016) A new species of freshwater crab of the genus *Tiwaripotamon* Bott, 1970 (Crustacea, Brachyura, Potamidae) from northern Vietnam and southern China. *The Raffles Bulletin of Zoology* 64: 213–219.
- Wang SB, Huang C, Zou JX (2019) Description of a new species of freshwater crab of the genus *Qianguiamon* Huang, 2018 (Crustacea: Decapoda: Brachyura: Potamidae) from Yulin, Guangxi, Southern China. *Zoological Studies* 58: e31. <https://doi.org/10.6620/zs.2019.58-31>
- Yang FF (2011) Investigating the phylogenic relationship of ribbed-shelled and smooth-shelled snails of *Oncomelania hupensis* collected from Guangxi with these from neighborhood provinces, China. Master Thesis, Guangxi Medical University, Nanning, China.
- Zhang Z, Xing YH, Cheng JJ, Pan D, Lü LN, Cumberlidge N, Sun HY (2020) Phylogenetic implications of mitogenome rearrangements in East Asian potamiscine freshwater crabs (Brachyura: Potamidae). *Molecular Phylogenetics and Evolution* 143: e106669. <https://doi.org/10.1016/j.ympev.2019.106669>
- Zhu C, Naruse T, Zhou XM (2010) Two new species of freshwater crabs of the genus *Sinopotamon* Tai & Sung, 1975 (Decapoda, Brachyura, Potamidae) from Guangxi Zhuang Autonomous Region, China. *Crustaceana* 83(2): 245–256. <https://doi.org/10.1163/001121609X12603430877199>

Supplementary material 1

Phylogenetic tree (ML) based on the COX1 gene

Authors: Yuan-Biao Lu, Yi-Xuan Zhang, Jie-Xin Zou

Data type: phylogenetic (.mtsx file)

Copyright notice: This dataset is made available under the Open Database License (<http://opendatacommons.org/licenses/odbl/1.0/>). The Open Database License (ODbL) is a license agreement intended to allow users to freely share, modify, and use this Dataset while maintaining this same freedom for others, provided that the original source and author(s) are credited.

Link: <https://doi.org/10.3897/zookeys.1166.101737.suppl1>

Supplementary material 2

Phylogenetic tree (BI) based on the COX1 gene

Authors: Yuan-Biao Lu, Yi-Xuan Zhang, Jie-Xin Zou

Data type: phylogenetic (.tre file)

Copyright notice: This dataset is made available under the Open Database License (<http://opendatacommons.org/licenses/odbl/1.0/>). The Open Database License (ODbL) is a license agreement intended to allow users to freely share, modify, and use this Dataset while maintaining this same freedom for others, provided that the original source and author(s) are credited.

Link: <https://doi.org/10.3897/zookeys.1166.101737.suppl2>

Supplementary material 3

Phylogenetic tree (ML) based on 3 genes

Authors: Yuan-Biao Lu, Yi-Xuan Zhang, Jie-Xin Zou

Data type: phylogenetic (.mtsx file)

Copyright notice: This dataset is made available under the Open Database License (<http://opendatacommons.org/licenses/odbl/1.0/>). The Open Database License (ODbL) is a license agreement intended to allow users to freely share, modify, and use this Dataset while maintaining this same freedom for others, provided that the original source and author(s) are credited.

Link: <https://doi.org/10.3897/zookeys.1166.101737.suppl3>

Supplementary material 4

Phylogenetic tree (BI) based on 3 genes

Authors: Yuan-Biao Lu, Yi-Xuan Zhang, Jie-Xin Zou

Data type: phylogenetic (.tre file)

Copyright notice: This dataset is made available under the Open Database License (<http://opendatacommons.org/licenses/odbl/1.0/>). The Open Database License (ODbL) is a license agreement intended to allow users to freely share, modify, and use this Dataset while maintaining this same freedom for others, provided that the original source and author(s) are credited.

Link: <https://doi.org/10.3897/zookeys.1166.101737.suppl4>