

Revision of the genus *Heterosmylus* Krüger, 1913 from China (Neuroptera, Osmylidae)

Min Dong¹, Han Xu¹, Yongjie Wang², Chunfeng Jia¹, Zhiqi Liu¹

1 Department of Entomology, China Agricultural University, Beijing 100094, China **2** College of Life Sciences, Capital Normal University, Beijing 100048, China

Corresponding authors: Yongjie Wang (wangyjosmy@foxmail.com); Zhiqi Liu (liuzhiqi@cau.edu.cn)

Academic editor: S. Winterton | Received 27 August 2016 | Accepted 17 November 2016 | Published 2 December 2016

<http://zoobank.org/CB5E8347-7767-49FB-96F4-98E014846775>

Citation: Dong M, Xu H, Wang Y, Jia C, Liu Z (2016) Revision of the genus *Heterosmylus* Krüger, 1913 from China (Neuroptera, Osmylidae). ZooKeys 637: 107–128. <https://doi.org/10.3897/zookeys.637.10310>

Abstract

A new species of osmylid (*Heterosmylus processus* **sp. n.**) is described and the other species in the genus from mainland China are redescribed. *Heterosmylus zhamanus* Yang, 1987, **syn. n.** is identified as a junior synonym of *Heterosmylus yunnanus* Yang, 1986. A key is provided to differentiate Palaearctic and Oriental species of *Heterosmylus*.

Keywords

Heterosmylus, Oriental region, Osmylidae, Palaearctic

Introduction

Heterosmylus Krüger is a relatively small genus of lance lacewing (Osmylidae: Protosmylinae) described by Krüger (1913a) from the Oriental Region. There are presently nine species in the genus, including seven species in mainland China, one species in Taiwan (*Heterosmylus primus* Nakahara) and another species (*Heterosmylus aspersus* Krüger) in northern India. Krüger (1913a) established the genus just based on the comparison with other genera, but the detailed description of the type species *H. aspersus* was given in a subsequent publication (Krüger 1913b). As for the Chinese species, Nakahara (1955) described *Heterosmylus primus* from Taiwan. Subsequently, in a series of

publications, Yang (1986, 1987, 1992, 1997, 1999) described an additional six species from mainland China. However, all the early taxonomists ignored or poorly described the characters of genitalia in osmylids, which result in a vague definition of the genus.

Herein the genus *Heterosmylus* is revised with a focus on the species from mainland China, and detailed descriptions of genitalic structures are provided for the first time. A revised diagnosis of the genus is proposed based on both external morphology and genitalic characters. The new species, *Heterosmylus processus* sp. n., is described and *Heterosmylus zhamanus* Yang, 1987, syn. n. is identified as a junior synonym of *Heterosmylus yunnanus* Yang, 1986. The distribution of the genus in China is also discussed.

Material and methods

All the described specimens are deposited in the Entomological Museum of China Agricultural University (CAU), Beijing. Terminalia preparations were made by macerating the apex of the abdomen in hot 10% KOH for 3–5 min, neutralized with 10% acetic acid. The apex of the abdomen was then transferred to glycerol for further dissection and examination. After examination it was moved to fresh glycerol and stored in a microvial pinned below the specimen. Images of wings were taken with a Nikon D7000 digital camera. Drawings were made under a light microscope. The terminology for wing venation and genitalia follows Winterton and Wang (2016).

Taxonomy

Heterosmylus Krüger, 1913a

Type species. *Heterosmylus aspersus* Krüger 1913a: 37, original designation. Deposited in Stettiner Museum (National Museum Szczecin, Poland).

Type locality. India: Sikkim.

Diagnosis. Head brown or dark brown; compound eyes black; antennae shorter than half length of forewing; prothorax black and length longer than width, with yellow setae; meso- and metathorax dark brown with long setae; legs yellow with short brown setae; forewings oblong and subacute at apex, with few brown spots; nygmata clear surrounded by light brown spot; veins thickened; costal cross-veins simple and occasionally bifurcate; forewing Rs with 8–15 branches, distal to the base of wing; cross-veins among Rs branches forming two or three series of gradates; forewing M branching more basally than the divergence of basal branch of Rs; no more than four cross-veins present between the two branches of M; forewing Cu bifurcating near the base of wing; CuA and CuP with numerous pectinate branches; CuP longer than half length of CuA; hind wings similar to forewings in size and shape without any spot apart from pterostigma; hindwing M branching near the base of wing; hindwing CuA with numerous pectinate branches; CuP simple and shorter than half length of CuA; male

genitalia with 9th tergite narrow, and sternite approximately quadrate; ectoproct relatively large, callus cerci rounded and located at the middle or underside of ectoproct; male genitalia composed of gonarcus, entoprocesses and mediuncus, arched gonarcus similar to other genera of Protosmylinae; mediuncus attached with a membrane below gonarcus, mediuncus bent into C-shape laterally; female genitalia with 9th tergite narrow; spermathecae bent into n-shape with base expanded and apex columniform.

Included species. *Heterosmylus aspersus* Krüger, *H. curvagradata* Yang, *H. flavidus* Yang, *H. limulus* Yang, *H. primus* Nakahara, *H. processus* sp. n., *H. shennonganus* Yang, *H. wolonganus* Yang, *H. yunnanus* Yang.

Comments. Although the type species of *Heterosmylus* was not described in detail when the genus was established, Krüger provided a detailed description of the type species in a following paper later that year (Krüger 1913a, b). This subsequent work was overlooked by Nakahara (1955), and later Ghosh (2000) presented a brief description for the species without the genitalic characters. Consequently, the systematic status of *Heterosmylus* was not well defined, although it is clear that the monophyly of *Heterosmylus* is well supported in the recent phylogenetic work on Osmylidae by Winterton et al. (in press). *Heterosmylus* also can be distinguished from other genera (*Gryposmylus* Krüger, *Lysmus* Navás, *Paryphosmylus* Krüger) in Protosmylinae based mainly on wing venation. In *Heterosmylus*, the veins are thickened and branches of forewing M have no more than four *ma-mp* cross-veins. In both *Gryposmylus* and *Lysmus* these are slender and the M vein generally five *ma-mp* cross-veins. Moreover, the base of costal-field of forewing of *Heterosmylus* species is narrower compared with that of *Gryposmylus*. *Heterosmylus* differs from *Paryphosmylus* in that the wings are mostly hyaline and single cross-vein presents before the separation of basal branch of Rs (Martins et al. 2016).

Heterosmylus is mainly recorded in the Oriental Region and especially in China (Fig. 1), typically in warmer and humid environments. According to the distribution of the individual species, we find *H. wolonganus*, *H. shennonganus* and *H. yunnanus* with relatively wide distributions. It seems that *H. wolonganus* has a broad geographical distribution from central to western China, occurring in four geographically continuous provinces, Sichuan, Gansu, Shaanxi and Henan. Interestingly, most localities of *H. wolonganus* are along the boundary of Oriental and Palearctic regions in China. *Heterosmylus shennonganus* is principally distributed in central China, representing a typical Oriental species. *Heterosmylus yunnanus* is another widespread species, distributed in Tibet, Yunnan and Sichuan. Considering the similar environment in these localities, it is estimated that these species might be present in the whole southwest of China. The other five species, *H. curvagradata*, *H. limulus*, *H. flavidus*, *H. primus* and *H. processus* sp. n., are only recorded in a single region. *Heterosmylus curvagradata* is restricted to Fujian, while *H. flavidus* is a distinctive species restricted in the west of Yunnan. *Heterosmylus limulus* is limited to Yadong (Tibet) and it could be found at the same altitude as *H. yunnanus*. *Heterosmylus primus* is only recorded in Taiwan. We did not examine this species so we could not compare it with the other species, but *H. curvagradata* shows the similar appearance with this species, suggesting their potential close relationship. The new species *H. processus* sp. n. is highly distinctive (weakly de-



Figure 1. Distribution of *Heterosmylus* in China. ☆ = *H. curvagradata* ▲ = *H. flavidus* ■ = *H. limulus* ★ = *H. primus* ● = *H. processus* sp. n. △ = *H. shennonganus* ◆ = *H. wolonganus* ◇ = *H. yunnanus*.

fined perostigma and hyaline and colourless membrane) is found in Shaanxi near the Qinling Mountains.

Key to *Heterosmylus* species (males) in the Palearctic and Oriental regions

- 1 Head and thorax with spots or stripes 2
- Head and thorax without any spot and stripe; pronotum yellowish brown without any stripe, meso- and metanotum dark brown; cross-veins *rs-ma* with lance-shaped brown marks *H. aspersus*
- 2 Mediuncus with a process at base (Fig. 4); membrane of wings hyaline and veins not edged with spots (Fig. 2); pronotum with four brown spots along anterior margin, two round brown spots in middle; mesonotum with two brown spots in middle on both sides *H. processus* sp. n.
- Mediuncus without any process at base, membrane of wings only with a few spots 3
- 3 Forewings with many spots including along the gradate cross-veins 4
- Forewings with few spots in the membrane (Fig. 21); pronotum with two yellow narrow stripes in middle; metanotum with two brown spots on anterior margin *H. wolonganus*
- 4 More than 11 branches of Rs 5

- 11 or fewer branches of Rs.....7
- 5 An oblique brown stripe present from pterostigma to the outer margin, Cu with four brown spots; pronotum with two pale yellow longitudinal stripes, mesonotum brown anteriorly.....*H. limulus*
- No oblique stripes present from pterostigma to the outer margin6
- 6 Pronotum black with a black longitudinal stripe medially; meso- and metanotum black; Rs with 14–17 branches; Cu edged with 3–4 yellow spots; apex of gonarcus bent upwards and relative long in lateral view*H. yunnanus*
- Pronotum brown with a yellowish longitudinal stripe medially, meso- and metanotum brown; vertex with a Y-shaped mark..... *H. primus*
- 7 Membrane of wings yellow and gradate cross-veins not edged with spots (Fig. 29); pronotum with a dark brown longitudinal stripe medially...*H. flavidus*
- Membrane of wings hyaline and gradate cross-veins edged with brown8
- 8 Apex of mediuncus short, broad and flat in dorsal view (Figs 46–47); ectoproct without any process in lateral view; two yellow longitudinal stripes present from pronotum to mesonotum; mesoscutellum bright yellow, metanotum with a central yellow stripe*H. curvogradatus*
- Apex of mediuncus long, acute and protuberant in dorsal view (Figs 38–39); ectoproct with a dorsal coniform process in lateral view; pronotum with two narrow longitudinal dark brown marks medially..... *H. shennonganus*

***Heterosmylus processus* sp. n.**

<http://zoobank.org/18F14A57-CA9C-4329-BA1F-F885461CCCB6>

Figs 2–6

Material examined. Holotype Male. CHINA: Shaanxi (Province): Taibai, [33°55'N, 107°43'E] 09.v.1982, leg. Guojun Qi. [Verbatim label data translated from Chinese]: CHINA: Shaanxi, Taibai/ 09.v.1982/ Guojun Qi/ CAU. Terminalia cleared in KOH, and stored in a micro-vial pinned below the specimen.

Diagnosis. Pronotum with four brown spots at anterior margin, two round brown spots in middle and two brown spots at posterior margin; membrane hyaline, pterostigma yellow, without dark spots besides the pterostigma; veins light yellow at base but brown from middle to the end; mediuncus C-shaped in lateral view with a basal process and boat-shaped in dorsal view.

Description. Body length 8.6 mm. *Head.* Vertex yellow with brown setae; ocelli distinctively brown; compound eyes black; antennae yellow with a brown stripe at base; frons with two brown stripes, genae with two round brown spots; maxillary palpi yellow and thick, labium short and brown. *Thorax.* Yellow dorsally and dark brown ventrally; pronotum with four brown spots at anterior margin, two round brown spots in middle and two brown spots at posterior margin linking to the mesonotum; mesonotum with two brown spots in middle on both sides; metanotum similar to the mesonotum, with two spots in middle. *Legs.* Yellow with brown setae; claws brown





Figure 2. Wings of *Heterosmylus processus* sp. n. Abbreviations: ng, nygmata; pt, pterostigma.



Figures 3–6. *Heterosmylus processus* sp. n. Male: **3** terminalia, lateral view **4** mediuncus, lateral view **5** mediuncus, dorsal view **6** genitalia, lateral view.

with a small tooth. *Wings* (Fig. 2): Forewing length 15.4 mm, width 6.2 mm; membrane hyaline without any spot besides the yellow pterostigma; veins yellow; costal field broad, cross-veins simple without forks. Cross-veins in radial sector few besides the gradate cross-veins. Rs with 8 branches. The basal cross-vein between R_1 and M edged with a brown mark. Hind wing length 14.4 mm, width 5.3 mm; membrane hyaline, veins light yellow at base but brown from middle to the end. Costal field narrow, pterostigma yellow, and Rs with 7 branches. *Abdomen*. Yellow dorsally, dark brown ventrally, covered with yellow setae.

Male Terminalia (Figs 3–6). Ectoproct quadrate in lateral view; callus cerci rounded and small. Distal part of gonarcus bent upwards. Entoprocesses curved in middle and dilated apically. Mediuncus (Figs 4–5) C-shaped in lateral view with a basal process and boat-shaped in dorsal view; two mediuncus lobes fused at base and each one raised on both sides, the inner side larger than the outer one.

Distribution. China (Shaanxi).

Etymology. The specific name refers to the process at the base of the mediuncus.

Remarks. This species is known from its type locality, Shaanxi province. It is easily distinguished from other species in the genus because of the body coloration, morphology and the genital characters. *Heterosmylus processus* sp. n. has a smaller body size and hyaline wings, while the others usually have a patterned membrane. Moreover, the mediuncus within the new species possesses a distinct process at the base whereas it is absent in the other species of *Heterosmylus*.

Heterosmylus yunnanus Yang, 1986

Figs 7–13

Heterosmylus zhamanus Yang, 1988: 195. **syn. n.**

Material examined. Holotype male. CHINA: Yunnan (Province): Lushuixian Gangfang, [25°57'N, 98°52'E], 29.ix.1980, leg. Dejing Zou. [Verbatim label data (translated from Chinese)]: CHINA: Yunnan, Lushuixian, Gangfang, 29.ix.1980/ Dejing Zou. Terminalia cleared in KOH, and stored in a micro-vial pinned below the specimen. 3 males, 1 female (type specimens of *H. zhamanus*), Tibet: Zhangmu, 06.vi.1981, leg. Shengchang Hu; 3 males, 5 females, Tibet: Hanmi, 24.viii.2005, leg. Dakang Zhou; 2 males, Sichuan: Luding, Hailuogou, 26.vi.2006, leg. Xiaoshuan Bai.

Diagnosis. Pronotum black with two yellowish longitudinal stripes at anterior margin; mesonotum with two yellow spots in middle; apex of gonarcus bent upwards and relative long in lateral view.


Redescription. Body length 9–11 mm. *Head*. Vertex dark brown, frons bright yellow; ocelli yellow; compound eyes dark grey; antennae entirely dark; clypeus brownish, maxillary and labial palpi dark brown. *Thorax*. Pronotum black with two yellowish longitudinal stripes at anterior margin; mesonotum with two yellow spots



Figure 7. Wings of *Heterosmylus yunnanus* Yang, 1986.

in middle. *Wings* (Fig. 7). Forewings length 18–21 mm, width 6–7 mm; membrane hyaline, and veins dark brown; pterostigma dark brown with yellowish center. R_s with 15–17 branches; Cu edged with three or four yellow spots. Hind wing length 15–18 mm, width 5–6 mm; membrane hyaline, cross-veins edged with brown marks between Sc and R_1 .

Male Terminalia (Figs 8–11). Ectoproct quadrate in lateral view. Gonarcus rod-like in lateral view, the apex bent upwards and relatively long; entoprocesses distally expanded, lobe-shaped; mediuncus fused at base and curved into C-shape in lateral view; each mediuncus lobe raised on both sides, boat-shaped in dorsal view.



Figures 8–13. *Heterosmylus yunnanus* Yang, 1986. Male: **8** terminalia, lateral view **9** mediuncus, lateral view **10** mediuncus, dorsal view **11** genitalia, lateral view Female: **12** terminalia, lateral view **13** spermatheca, lateral view.

Female Terminalia (Figs 12–13). Ectoproct trapeziform in lateral view; 9th gonocoxite finger-shaped in lateral view, 9th gonostylus brown and long. Spermathecae bent into C-shaped and basal part longer than distal.

Distribution. China (Tibet, Yunnan, Sichuan)

Remarks. Although the original identification of both *H. yunnanus* Yang and *H. zhamanus* Yang was based on the colour pattern of the head, there are no convincing differences to distinguish them after comparison of their genitalic features. So we believe that *H. zhamanus* should be a synonym of *H. yunnanus*.

5 mm



Figure 14. Wings of *Heterosmylus limulus* Yang, 1987.


***Heterosmylus limulus* Yang, 1987**

Figs 14–20

Material examined. Holotype Male, CHINA: Tibet: Yadong, [27°31'N, 88°55'E], 24.viii.1978, leg. Fasheng Li. [Verbatim label data (translated from Chinese)]: CHINA: Tibet, Yadong/ 24.viii.1978/ Fasheng Li/ CAU. 1 male, same data as Holotype. 1 female, China: Tibet: Yadong, 30.viii.1984, leg. Yongxiang Zhao.

Diagnosis. Pronotum with two yellow stripes in middle, an oblique brown mark presenting from the pterostigma to the posterior outer margin of forewing; base of mediuncus slender and apex dilated as lobe-shape in lateral view.

Redescription. Body length 8.0 mm. *Head.* Vertex dark brown, frons brown. Ocelli yellowish, eyes dark grey. Antennae brown, encircled by a yellowish stripe at



Figures 15–20. *Heterosmylus limulus* Yang, 1987. Male: **15** terminalia, lateral view **16** mediuncus, lateral view **17** mediuncus, dorsal view **18** genitalia, lateral view. Female: **19** terminalia, lateral view **20** spermatheca, lateral view.

base. Clypeus yellow, maxillary palpi and labial palpi dark brown. *Thorax*. Dark brown. Pronotum with two light yellow stripes medially; mesonotum brown at anterior part with black setae. *Wings* (Fig. 14). Forewing length 16–17 mm, width 5–6 mm; membrane hyaline, veins brown; pterostigma yellow, but brown on both sides; an oblique brown stripe presenting from the pterostigma to the posterior outer margin; Cu with four brown spots. Outer gradate series of cross-veins and wing margin bordered with fuscous spots. Rs with 13–15 branches. Hind wing length 13–14 mm, width 4–5 mm; membrane hyaline with some light brown spots between Sc and R₁.

Male Terminalia (Figs 15–18). Ectoproct quadrate in lateral view; gonarcus rod-like in lateral view, distal part upswept and relatively small; mediuncus fused at base and curved into C-shape in lateral view; the base slender and apex dilated as lobe-shape in lateral view.

Female Terminalia (Figs 19–20). Ectoproct approximately oblong in lateral view; 9th gonocoxite fusiform in lateral view; 9th gonostylus brown and small; spermathecae bent into C-shape.

Distribution. China (Tibet).

Heterosmylus wolonganus Yang, 1992

Figs 21–28

Material examined. Holotype Male. CHINA: Sichuan (Province): Wolong, [31°01'N, 103°10'E], 25.vii.1993, leg. Shuyong Wang. [Verbatim label data (translated from Chinese)]: CHINA: Sichuan Prov., Wonglong/ 25.vii.1993/ Shuyong Wang/ CAU. 1 male, CHINA: Shaanxi (Province): Ningshan, 18.vi.1981, leg. Chikun Yang. 1 female, CHINA: Shaanxi (Province): Huangniupu, 15.viii.1981, leg. Weidong Wang. 1 male, CHINA: Shaanxi (Province): Taibaishan, 09.iii.1982, leg. Chikun Yang. 1 male, CHINA: Shaanxi (Province): Taibaishan, 09.v.1982, leg. Guojun Qi. 1 female, CHINA: Shaanxi (Province): Taibaishan, 15.vii.1982, leg. Jingruo Zhou, Lan Liu. 1 male, 1 female, CHINA: Shaanxi (Province): Ningshan, 18.vii.1982, leg. Shenghui Lei. 1 female, CHINA: Shaanxi (Province): Ningshan, Xiangyang, Fengqi, 18.vii.1982, leg. Deqing Wang. 2 males, CHINA: Shaanxi (Province): Nanzheng, Yuanba, 27.v.1983, leg. Dahan He. 1 male, CHINA: Gansu (Province): Wenxian, Bikou, 25.vii.1998, leg. Jun Chen. 2 males, CHINA: Henan (Province): Songxian, Baiyunshan, 19.vii.1996, leg. Xiaocheng Shen. 10 males, 12 females, CHINA: Henan (Province): Songxian, Baiyunshan, 14–18.vii.2004, leg. Bingzhen Yan.

Diagnosis. Pronotum with two yellow narrow stripes in middle; metanotum with two brown spots on anterior margin; gonarcus sclerotized distally and bent upward and hook-shaped in lateral view; base of mediuncus approximately finger-shaped in lateral view.

Redescription. Body length 9–11 mm. *Head*. Vertex yellowish brown; frons yellow with one dark brown spot near antennae; ocelli brown, eyes blackish brown; antennae blackish brown; clypeus yellow, maxillary and labial palpi dark brown

5 mm




Figure 21. Wings of *Heterosmylus wolonganus* Yang, 1992.

Thorax. Pronotum with two yellow narrow stripes in middle with some dark setae on both sides; mesonotum with two yellow longitudinal spots; metanotum with two brown spots on anterior margin. *Wings* (Fig. 21). Forewing length 18–20 mm, width 6–7 mm; membrane hyaline, veins mostly brown; pterostigma light yellow but brown on both sides; Rs with 10 branches. Four cross-veins present between MA and MP. Hind wing length 16–17 mm, width 5–6 mm; Rs with 11–12 branches, without any distinct spot besides the pterostigma.

Male Terminalia (Figs 22–25). Ectoproct quadrate in lateral view; gonarcus sclerotized distally and bent upwards as hook in lateral view; gonocoxite bent and distally dilated as lobe-shaped in lateral view; mediuncus curved into C-shape in lateral view and spoon-shaped in dorsal view; base approximately finger-shaped in lateral view.

Female Terminalia (Figs 26–28). Anterior part of 8th sternite reduced and short finger-shaped, posterior part broad; ectoproct broad and trapezoid in lateral view; 9th



Figures 22–28. *Heterosmylus wolonganus* Yang, 1992. Male: **22** terminalia, lateral view **23** mediuncus, lateral view **24** mediuncus, dorsal view **25** genitalia, lateral view. Female: **26** terminalia, lateral view **27** spermatheca, lateral view **28** 8th sternite, ventral view.

gonocoxite fusiform in lateral view; 9th gonostylus long in lateral view; spermatheca bent into C-shape.

Distribution. China (Sichuan, Shaanxi, Gansu, Henan).

Heterosmylus flavidus Yang, 1992

Figs 29–35

Material examined. Holotype Female. CHINA: Yunnan (Province): Lushui, Yaojiapin, [25°58'N, 98°42'E], 04.vi.1981, leg. Shuyong Wang. [Verbatim label data (translated

5 mm




Figure 29. Wings of *Heterosmylus flavidus* Yang, 1992.

from Chinese]): CHINA: Yunnan Prov., Lushui, Yaojiaping/ 04.vi.1981/ Shuyong Wang/ CAU. Paratype Male, CHINA: Yunnan (Province): Lushui, Pianma, 29.v.1981. leg. Xuezhong Zhang. [Verbatim label data (translated from Chinese)]: CHINA: Yunnan Prov., Lushui, Pianma/ 29.v.1981/ Xuezhong Zhang/ CAU.

Diagnosis. Pronotum with a dark brown longitudinal stripe medially; metanotum with a brown spot in middle; forewing light yellow; mediuncus C-shaped in lateral view and scoop-shaped in dorsal view.

Redescription. Body length 7–9 mm. *Head.* Vertex dark brown, frons yellow; ocelli yellowish; Compound eyes brown with some small spots; antennae fuscous except for the yellow scape; clypeus yellow, maxillary and labial palpi black. *Thorax.* Mostly yellow. Pronotum with a brown longitudinal stripe, and with some long setae



Figures 30–35. *Heterosmylus flavidus* Yang, 1992. Male: **30** terminalia, lateral view **31** mediuncus, lateral view **32** mediuncus, dorsal view **33** genitalia, lateral view. Female: **34** terminalia, lateral view **35** spermatheca, lateral view.

on both sides; mesonotum brown on margin; metanotum dark brown with a brown spot medially. *Wings* (Fig. 29). Forewing length 16–17 mm, width 5–6 mm; membrane light yellow, veins brown and thickened; pterostigma light yellow but brown on

both sides; R_1 edged with some brown spots, Rs with 11 branches, cross-veins among branches of Rs forming three series of gradates. Hind wing length 12–14 mm, width 4–6 mm; membrane hyaline with small spots between Sc and R_1 .

Male Terminalia (Figs 30–33). Ectoproct quadrate in lateral view; gonarcus rod-like, but apex not bent upwards in lateral view; gonocoxite bent in the middle; medianus C-shaped in lateral view and scoop-shaped in dorsal view.

Female Terminalia (Figs 34–35). 8th sternite finger-shaped in lateral view; ectoproct broad and approximately elliptical in lateral view; 9th gonocoxite finger-shaped in lateral view; 9th gonostylus relatively long and finger-shaped in lateral view; spermathecae bent into C-shape, base thicker than apex.

Distribution. China (Yunnan).

Heterosmylus shennonganus Yang, 1997

Figs 36–43

Material examined. Holotype Male. CHINA: Hubei (Province): Shennongjia, [31°42'N, 110°38'E], 21.viii.1985, leg. Xiaoyuan Mao. [Verbatim label data (translated from Chinese)]: CHINA: Hubei Prov., Shennongjia, 21.viii.1985/ Xiaoyuan Mao/ CAU. Paratype 4 males, 3 females, same data as the holotype. 1 female, Henan: Luanchuan, Longyuwan, 1997.08.14, leg. RLS. 1 male, CHINA: Henan (Province): Luanchuan, Longyuwan, 18.viii.1997, leg. RLS. 1 male, CHINA: Henan (Province): Luanchuan, Longyuwan, 19.vii.2004, leg. Bingzhen Yan. 1 female, CHINA: Henan (Province): Neixiang, Baotianman, 23.vii.2004, leg. Bingzhen Yan. 1 male, CHINA: Henan (Province): Neixiang, Baotianman, 24.vii.2004, leg. Bingzhen Yan. 1 male, CHINA: Chongqing: Jiangjin, Simianshan, 21.ix.2007, leg. Weiwei Zhang. 1 female, CHINA: Shaanxi (Province): Xiangyangba, 24.viii.1982, leg. Shenghui Lei.

Diagnosis. Pronotum with two narrow longitudinal dark brown marks in middle; ectoproct with a dorsal coniform process in lateral view; gonarcus with a short finger-like process distally in lateral view.

Redescription. Body length 8–10 mm. *Head.* Vertex with a brown cross-stripe; frons yellow but brown on both sides; ocelli large and prominent, compound eyes shiny black; antennae dark brown; clypeus fulvous, maxillary and labial palpi brown.

Thorax. Pronotum with two narrow longitudinal dark brown marks in the middle; mesonotum with dark setae; metanotum without spots. *Wings* (Fig. 36). Forewing length 16–17 mm, width 6.4 mm; membrane hyaline, veins mainly brown; pterostigma yellowish but brown on both sides; costal field with three or four brown spots; $r1-rs$ edged with brown marks; Rs with 8–9 branches, cross-veins among branches of Rs forming two series of gradates; the outer gradate cross-veins edged with brown spots. Hind wing length 15.5 mm, width 5.4 mm. Membrane hyaline with few spots.

Male Terminalia (Figs 37–40). Ectoproct approximately quadrate with a dorsal coniform process in lateral view; gonarcus approximately rod-like and distally sclerotized with a short finger-like process in lateral view; gonocoxite curved as ancon and distally




Figure 36. Wings of *Heterosmylus shennonganus* Yang, 1997.

dilated as lobe; mediuncus curved and thickened in lateral view and boat-shaped with a cone-shaped apex in dorsal view.

Female Terminalia (Figs 41–43). 8th sternite approximately finger-shaped in lateral view. Ectoproct trapeziform in lateral view; 9th gonocoxite approximately finger-shaped in lateral view; 9th gonostylus finger-shaped and brown; spermathecae bend into C-shape.

Distribution. China (Henan, Shaanxi, Hubei, Chongqing).



Figures 37–43. *Heterosmylus shennonganus* Yang, 1997. Male **37** terminalia, lateral view **38** mediuncus, lateral view **39** mediuncus, dorsal view **40** genitalia, lateral view. Female: **41** terminalia, lateral view **42** spermatheca, lateral view **43** 8th sternite, ventral view.

***Heterosmylus curvagratus* Yang, 1999**

Figs 44–50

Material examined. Holotype Male. CHINA: Fujian (Province): Wuyishan, Huanghuacong, [27°48'N, 117°42'E], 13.x.1980, leg. Fan Jiang. [Verbatim label data (translated from Chinese)]: CHINA: Fujian Prov., Wuyishan, Huanghuacong/ 13.x.1980/ Fan Jiang/ CAU. Paratype Female. CHINA: Fujian (Province): Wuyishan, Xianfening, 19.ix.1987. leg. Jiashe Wang.

Diagnosis. Two yellow longitudinal stripes present from pronotum to mesonotum. mesoscutellum bright yellow, metanotum with a central yellow stripe; apex of gonarcus slightly dilated, short and curved dorsally.




Figure 44. Wings of *Heterosmylus curvagratus* Yang, 1999.

Redescription. Body length 7–10 mm. *Head.* Vertex shiny yellow with a brown round spot in middle and a greyish yellow transverse band near antennae; frons bright yellow with a black brown stripe; ocelli grey but black at base; compound eyes grey and glossy; antennae black; clypeus yellow, maxillary and labial palpi dark brown. *Thorax.* Dark brown with two yellow longitudinal stripes from pronotum to mesonotum. mesoscutellum bright yellow, metanotum with a central yellow stripe. *Wings* (Fig. 44). Forewing length 15–17 mm, width 5–6 mm; membrane hyaline, veins fuscous with numerous long setae; pterostigma yellow but brown on both sides. Crossveins *r1-rs* edged with dark brown spots. *Rs* with 8–9 branches, gradates cross-veins with brown marks. Hind wing length 13–14 mm, width 4–6 mm; membrane hyaline without spots besides the pterostigma.

Male Terminalia (Figs 45–48). Ectoproct approximately pentagonal in lateral view; gonarcus rod-like in lateral view, apex slightly inflated, short and curved dorsad; mediuncus fused at base and scoop-shaped in dorsal view and apex flat-bottomed in dorsal view.

Female Terminalia (Figs 49–50). 8th sternite reduced into finger-shape in lateral view; ectoproct approximately trapezoid in lateral view; 9th gonocoxite approximately finger-shaped in lateral view; 9th gonostylus brown and short; spermathecae bend into C-shape.

Distribution. China (Fujian).



Figures 45–50. *Heterosmylus curvagradataus* Yang, 1999. Male: **45** terminalia, lateral view **46** mediuncus, lateral view **47** mediuncus, dorsal view **48** genitalia, lateral view. Female: **49** terminalia, lateral view **50** spermatheca, lateral view.

Acknowledgements

We wish to express our gratitude to Timothy R. New and Valdimir Makarkin for providing literature necessary to this project. This research was supported by the National Natural Science Foundation of China (Gants 31272352, 31301905 and 41372013), the Beijing Natural Science Foundation (Grant 5132008), and Ph.D. Programs Foundation of Ministry of Education of China (Grant 20131108120005).

References

- Adams PA (1969) A new genus and species of Osmylidae (Neuroptera) from Chile and Argentina, with a discussion of Planipennian genitalic homologies. *Postilla* 141: 1–11. doi: 10.5962/bhl.part.24591
- Krüger L (1913a) Osmylidae. Beiträge zu einer Monographie der Neuropteren-Familie der Osmylidaen. II. Charakteristik der Familie, Unterfamilien und Gattungen auf Grund des Geäders. *Stettiner Entomologische Zeitung* 74: 3–123.
- Krüger L (1913b) Osmylidae. Beiträge zu einer Monographie der Neuropteren-Familie der Osmyliden. IV. Beschreibung der Arten. *Stettiner Entomologische Zeitung* 74: 225–294.
- Kuwayama S (1962) A revisional synopsis of the Neuroptera in Japan. *Pacific Insects* 4: 325–412.
- Martins CC, Ardila-Camacho A, Aspöck U (2016) Neotropical osmylids (Neuroptera, Osmylidae): Three new species of *Isostenosmylus* Krüger, 1913, new distributional records, redescriptions, checklist and key for the Neotropical species. *Zootaxa* 4149(1): 1–66. doi: 10.11646/zootaxa.4149.1.1
- Nakahara W (1955) Formosan Neuroptera collected by the late Dr. T. Kano. *Kontyû* 23: 6.
- New TR (1983) A revision of the Australian Osmylidae: Kempyninae (Insecta: Neuroptera). *Australian Journal of Zoology* 31: 393–420. doi: 10.1071/ZO9830393
- Winterton SL, Wang YJ (2016) Revision of the genus *Gryposmylus* Krüger, 1913 (Neuroptera, Osmylidae) with a remarkable example of convergence in wing disruptive patterning. *ZooKeys* 617: 31–45. doi: 10.3897/zookeys.617.10165
- Winterton SL, Garzón-Orduña IJ, Zhao J, Wang YJ, Liu ZQ (2016) Phylogeny of the lacewing family Osmylidae (Insecta: Neuroptera). *Systematic Entomology*, (submitted).
- Yang CK (1986) Thirty new species and four new genera of Neuroptera from Yunnan, and the family Nemopteridae new to China. *Pei ching nung yeh ta hsueh hsueh pao= Acta Agriculturae Universitatis Pekinensis* 12: 153.
- Yang CK (1987) Neuroptera. In: Zhang S (Ed.) *Agricultural insects, spiders, plant diseases and weeds of Xizang*. Vol. 1. Xizang Renmin Press House, Tibet, 191–194.
- Yang CK (1992) Neuroptera. *Insects of the Hengduan Mountains Region* Science press, Beijing, 438–454.
- Yang CK (1997) Neuroptera: Osmylidae. In: Yang XK (Ed.) *Insects of the Three Gorge Reservoir Area of Yangtze River*. Chongqing Press, Chongqing, 580–581.
- Yang CK (1999) Osmylidae. In: Huang BK (Ed.) *Fauna of Insects Fujian Province of China* Vol. 3. Fujian Science and Technology Press, Fuzhou, 96–98.