

Exploring the relationship between bats (Mammalia, Chiroptera) and ectoparasitic flies (Diptera, Hippoboscoidea) of the Orinoquia Region in South America

Erika M. Ospina-Pérez^{1,2}, Fredy A. Rivera-Páez², Héctor E. Ramírez-Chaves^{2,3}

1 Doctorado en Ciencias – Biología, Facultad de Ciencias Exactas y Naturales, Universidad de Caldas, Calle 65 No. 26-10, 170004, Manizales, Caldas, Colombia

2 Grupo de Investigación en Genética, Biodiversidad y Manejo de Ecosistemas (GEBIOME), Departamento de Ciencias Biológicas, Facultad de Ciencias Exactas y Naturales, Universidad de Caldas, Calle 65 No. 26-10, 170004, Manizales, Caldas, Colombia

3 Centro de Museos, Museo de Historia Natural, Universidad de Caldas, Calle 58 No. 21-50, 170004, Manizales, Caldas, Colombia

Corresponding author: Héctor E. Ramírez-Chaves (hector.ramirez@ucaldas.edu.co)



This article is part of:
Biology of Pangolins and Bats

Academic editor: Liping Yan
Received: 14 March 2023
Accepted: 18 August 2023
Published: 8 September 2023

ZooBank: <https://zoobank.org/53C5FB34-A6B1-48BF-8B16-6C86E5F7865E>

Citation: Ospina-Pérez EM, Rivera-Páez FA, Ramírez-Chaves HE (2023) Exploring the relationship between bats (Mammalia, Chiroptera) and ectoparasitic flies (Diptera, Hippoboscoidea) of the Orinoquia Region in South America. ZooKeys 1179: 1–34. <https://doi.org/10.3897/zookeys.1179.103479>

Copyright: © Erika M. Ospina-Pérez et al.
This is an open access article distributed under terms of the Creative Commons Attribution License ([Attribution 4.0 International – CC BY 4.0](https://creativecommons.org/licenses/by/4.0/)).

Abstract

Bat flies (Nycteribiidae and Streblidae) have been used to study co-evolutionary patterns between ectoparasites and bats. In the world, Nycteribiidae and Streblidae are represented by approximately 276 and 237 species, respectively. In regions such as the Orinoquia located in the north of South America (Colombia and Venezuela), the richness of bats is high (more than 100 documented species), but studies on Nycteribiidae and Streblidae are scarce and discontinuous. To contribute to the knowledge of ectoparasitic flies in the Orinoquia, records of flies and their interactions with bats were reviewed, including new records and associations using interaction networks. We documented 124 species of Streblidae and only 12 of Nycteribiidae for the Orinoquia in approximately 102 bat species reported in Colombia and Venezuela. New records for six species of bat flies in Colombia were found (*Mastoptera guimaraesi*, *Noctiliostrebla maai*, *Paradyschiria parvuloides*, *Trichobius jubatus*, *Trichobius parasiticus*, and *Basilia ferrisi*) associated with six species of bats (*Cynomops planirostris*, *Desmodus rotundus*, *Myotis handleyi*, *Molossus rufus*, *Noctilio albiventris*, and *Phyllostomus hastatus*). The bat-ectoparasite interaction networks in the Orinoquia revealed a pattern of antagonistic relationships, with high specialization, modularity, and low connectivity and nesting. The identified networks are between bat fly species belonging to different ecomorphological groups with unique host species. This supports the idea of ecological niche partitioning among ectoparasitic bat flies and hosts. Our study expanded the knowledge of the distribution of some fly species and the associations with bat hosts in Colombia, by presenting morphological descriptions and new observations, which are key to understanding the ecology, diversity, and distribution of these species.

Key words: Bat flies, ecological networks, ecomorphology, Neotropics, Nycteribiidae, parasitism, Streblidae

Introduction

Bat flies (Diptera: Hippoboscoidea) are obligate blood-feeding ectoparasites of bats (Chiroptera) and an interesting evolutionary system for studying co-evolutionary patterns between hosts and ectoparasites (Porter et al. 2021).

Taxonomically bat flies are divided into two cosmopolitan families, Nycteribiidae and Streblidae (Wenzel et al. 1966; Dick and Patterson 2006; Patterson et al. 2007), but the latter is considered paraphyletic with New World and Old-World lineages (Dittmar et al. 2006; Petersen et al. 2007). Both families of bat flies show greater diversity in tropical latitudes and are less diverse in subtropical and temperate regions (Dick and Dittmar 2014). However, Nycteribiids (three subfamilies, 11 genera and ~ 276 species for 2018) are more numerous in the Eastern Hemisphere, while, for the Streblids (five subfamilies, 33 genera, and ~ 237 species for 2018) the richness is greater in the Western Hemisphere (Dick and Patterson 2006; Dick and Miller 2010; Soares et al. 2013; Graciolli and Dick 2018). Based on dispersion-vicariance analysis, Dittmar et al. (2006) suggested that the Neotropical region is the ancestral area for all New World Streblidae, while the Oriental region is considered the ancestral area of Nycteribiidae and Old-World Streblidae.

Both Nycteribiidae and Streblidae present a wide variety of morphological and physiological adaptations for their ectoparasitic lifestyle, among which adenotrophic viviparity stands out (Hagan 1951; Lehane 2005; Dick and Miller 2010; Dick and Dittmar 2014). In this process, the larvae develop individually in the female oviduct until the third instar (called prepupa) (Hagan 1951). Later, when the development of the prepupa is complete, female bat flies of both families deposit a single prepupa on substrates in the roost (Petersen et al. 2007; Dick and Dittmar 2014). Once the prepupa is deposited, it immediately forms a puparium, which, after a pupal stage, emerges as an unfed adult fly (teneral) and must locate and colonize a host (Ching and Marshall 1968).

In general, the life strategy of bat flies reflects their association with bats, and it has been found that some Streblidae flies can form specific parasite assemblages for each bat species due to their high host specificity (Wenzel 1976; Hiller et al. 2018). Each community can consist of two to five fly species, and each of them shows a preference for a specific part of the bat's body (Wenzel 1976; Ter Hofstede et al. 2004; Dick and Gettinger 2005; Tello et al. 2008; Patterson et al. 2009). Dick (2005) defined three ecomorphological groups of bat flies based on behavioral observations and morphological traits: 1) "wing crawler" primarily live on wing membranes, with non-flattened bodies and legs of the same size, 2) "fur runners" that live on the hairy membranes of the body and move by running on the fur, with well-developed wings and long legs (especially the hind legs), and 3) "fur swimmers" that inhabit areas of long fur such as the neck and have compressed bodies and heads, and possessing Ctenidia. Hiller et al. (2018) supported this classification, focusing on differences in the morphology and size of the hind legs, and found evidence of density-dependent competition among species within the same ecomorphological group.

In Neotropical countries with a high number of bat species such as Colombia (~ 217 species) and Venezuela (172 species) (Delgado-Jaramillo et al. 2016; Ramírez-Chaves et al. 2021), studies on Streblidae and Nycteribiidae are few and discontinuous (e.g., Tamsitt and Fox 1970; Marinkelle and Grose 1981; Guerrero 1994). In recent years, new research has expanded the knowledge about these ectoparasites in both countries (Herrera-Sepúlveda 2013; Tarquino-Carbonell 2014; Tarquino-Carbonell et al. 2015; Dick et al. 2016; Durán et al. 2018; Calonge-Camargo and Pérez-Torres 2018; Guerrero 2019; Liévano-Romero et al. 2019; Raigosa Álvarez et al. 2020; Cañizales and Guerrero 2022). Nonetheless, there are very few studies in the Orinoquia Region, a hydrographic basin that converges into the

Orinoco River, covering an area of almost 989,000 km² (Domínguez 1998), and shared by both countries (Guimarães 1972; Wenzel 1976; Dick et al. 2016; Liévanó-Romero et al. 2019; López Rivera et al. 2022). Approximately 65% of this area is located in Venezuela, while the remaining 35% is in Colombia (León 2005). The Venezuelan portion extends from the Andes and the Cordillera de la Costa to the north-western bank of the Orinoco River, forming most of the Venezuelan plains and the delta Orinoco (Domínguez 1998; León 2005). The Orinoquia is one of the most diverse areas in terms of mammal diversity with 318 species, including 150 species of Chiroptera (Ferrer Pérez et al. 2009; Pardo and Rangel-Ch. 2014).

Of the 125 Streblidae species reported in Colombia and Venezuela (81 and 107 respectively; Wenzel 1976; Tarquino-Carbonell et al. 2015; Dick et al. 2016; Duran et al. 2017; Calonge-Camargo and Pérez-Torres 2018; Guerrero 2019; Liévanó-Romero et al. 2019; Ascuntar-Osnas et al. 2020; Raigosa Álvarez et al. 2020; López Rivera et al. 2022), 124 have been documented for the Orinoquia in ~ 96 bat species of the families Emballonuridae, Molossidae, Mormoopidae, Natalidae, Noctilionidae, Phyllostomidae, and Vespertilionidae (Wenzel 1976; Dick et al. 2016; Guerrero 2019; Liévanó-Romero et al. 2019; López Rivera et al. 2022). Similarly, of the 16 Nycteribiidae reported in Colombia and Venezuela (11 and 13, respectively; Wenzel and Tipton 1966; Graciolli et al. 2016; Pastrana-Montiel et al. 2019; Raigosa Álvarez et al. 2020; López Rivera et al. 2022), 12 have been found in the Orinoquia (two in Colombia, 12 in Venezuela; Wenzel and Tipton 1966; Graciolli et al. 2016; López Rivera et al. 2022). Besides the high number of bats and bat flies documented in the Orinoquia, the interaction structure that might be evaluated using ecological or interaction networks (Dormann et al. 2009), has not been addressed. Interaction networks are formed by interactions (links) between species (nodes) that make up a community (network); they are considered as a synthesis tool in the study of ecological interactions that allow understanding the functioning of megadiverse systems (Blüthgen et al. 2006). The study of the host-parasite specificity interactions is essential to understand the mechanisms behind parasitism and its relationship with biodiversity functioning (Frainer et al. 2018) since parasites play an important role in the regulation of populations of host species (Poulin et al. 2006). For those reasons, and to contribute to the knowledge of ectoparasitic flies of the Orinoquia, we present novel records (including morphological description of ectoparasitic flies) and new associations with bats from this region in Colombia, and a review of records of ectoparasitic flies and the interactions with hosts bats in the whole Orinoquia (Colombia-Venezuela).

Materials and methods

Study Area

The Orinoquia Region shared by Colombia and Venezuela is a hydrographic basin, with waters that converge in the Orinoco River (Domínguez 1998). The Orinoco basin covers an area of almost 989,000 km² of which 643,480 km² (65%) are located in Venezuela and 35% in Colombia (Fig. 1). In Colombia, the Orinoquia extends from the eastern slopes of the Cordillera Oriental of Colombia. In Venezuela, the Orinoquia extends from the Venezuelan Andes and the Cordillera de la Costa to the north-western bank of the Orinoco River, forming most of the

Venezuelan plains and the delta Orinoco (León 2005). The Orinoco basin features isothermal climates with minimal temperature fluctuations throughout the year. This is typical of the inter-tropical zone and is characterized by a mere 3 °C difference between the average temperatures of the warmer and cooler months. In the lowlands, which reach up to 800 m above sea level, there are five primary climate types, including jungle, savanna, semi-desert, and desert proper (León 2005).




Figure 1. Locality records of Nycteriidae and Streblidae in the Orinoquia Region in South America **A** Orinoquia **B** South America highlighting Colombia (pink) and Venezuela (blue). The red line delimits the Orinoco basin according to Domínguez (1998); the yellow circles indicate 58 localities reported in the literature (Table 1) and the red star indicate the new records for the Department of Arauca, Colombia (Table 2). Detailed information is found in the Suppl. material 1: table S1.

Richness of bats and ectoparasite flies in the Orinoco Region and new records from Colombia

To gather the information on ectoparasitic flies associated with bats from Orinoquia Region (Fig. 1), we reviewed the information available in the literature retrieved from Science Direct, Web of Science, SciELO Scopus, and Google Scholar search engines using the keywords ((Fly*) OR (Flies*) AND ((Streblidae*) OR (Nycteribiidae*) AND (bat*)) AND (Colombia*) OR (Venezuela*)), without temporal restrictions. We also analyzed the sources cited and referenced in the publications to obtain more data and information for the compilation of the interaction networks. To update the bats taxonomy, we used recent check list and online resources (e.g., Ramírez-Chaves et al. 2021; Mammal Diversity Database 2022). To include an article in our study, we used the following criteria: bat species found in contact with ectoparasitic dipterans from a locality (including latitude and longitude) within the Orinoquia Region. It is relevant to highlight that when reviewing the associations, we consider previously reported bat fly species complexes (*sensu* Wenzel 1976), which are believed to have a generalist behavior. In publications lacking geographic coordinates we extrapolated this information using Google Earth (<https://www.google.com/intl/es/earth/>), when possible. To assess the quality of the published bat flies-host associations, we collected information on the abundances of flies found on the hosts, as well as the specific location and date of each report. To identify possible non-primary associations due to contamination transfers, two important criteria were applied (Dick 2005): 1) non-primary associations that represented $\leq 5\%$ of the total records, and 2) ectoparasitic flies had to be collected in the presence of the primary host at the same location and date (Table 1 and Suppl. material 1: table S2). Additionally, we reviewed the non-primary associations reported for the Department of Arauca in previous works (e.g., López Rivera et al. 2022), based on specimens deposited in the Collection of Ectoparasites (**Ec**) at the Museo de Historia Natural of the Universidad de Caldas (**MHN-UCa**).

We also included additional and noteworthy records of ectoparasitic bat flies collected from bats captured using four mist nets (12×2.5 m; with 36 mm mesh size) during November 2021 (8 days) in two localities of the Orinoquia Region. For this we performed field work in two localities of the municipality of Arauca in the Department of Arauca (Fig. 1), Colombia (1: Vereda El Socorro, Finca Los Trompillos, and 2: Vereda El Socorro, Finca Marsella; Table 2). The Department of Arauca is located in the Orinoquia Region bordering to the north and east with Venezuela, and occupies an area of $23,818 \text{ km}^2$ dominated by herbaceous plains and chaparral (Mosquera Guerra et al. 2019). The region shows a typical savanna climate with a well-defined wet season between June and July and a very dry season between December and April (López Rivera et al. 2022). The mist nets were placed randomly in both localities and operated between 18:00 and 22:30 hours. We examined each captured bat to search for ectoparasites by placing a clean white blanket on them and in some cases moistening the fur with 70% alcohol, to facilitate handling. We collected the flies and other ectoparasites using entomological tweezers and stored in Eppendorf tubes with 96% ethanol. The flies were collected and handled under the permission granted by the Autoridad Nacional de Licencias Ambientales (ANLA) to the Universidad de Caldas (Resolution 02497 of 31 December 2018) and by approval of the bioethics committee of the Facultad de Ciencias Exactas y Naturales of the

Table 1. Records of flies in bats in Orinoquia from 1911 to June 2023. Localities represents the number of localities in which the species has been documented. Letters in parentheses following the fly species indicate host-specificity (HS) behavior: M: monoxene (found in only one host species); O: oligoxene (found in different species of bat hosts, but of the same genus); Pl: pleioxene (found in different species and genera of bat hosts, but from the same family); Po: polyxene (found in different host bat species and genera and in different families). The hosts were arranged in descending order from primary hosts, transitional to incidental, the hosts marked with an asterisk are not primary ($\leq 5\%$ of the total records) and those followed by the letter (C) are possible associations resulting from contamination by manipulation when they are collected in the presence of the primary host in the same place and date (Dick 2005), however, as they are congeners of the primary host, this definition should be considered. The other associations are difficult to define if they are ecological or contamination due to the low number of records.

Bat-flies taxa	Bats	Country (Department/State)	Localities	References
Family Streblidae				
Subfamily Nycterophilinae				
<i>Nycterophilia coxata</i> (Pl)	<i>Pteronotus parnellii</i> (probably <i>P. fuscus</i>), * <i>Phyllostomus elongatus</i> , * <i>Artibeus planirostris</i> , * <i>Eumops glaucinus</i> , * <i>Pteronotus davyi</i> (C) and * <i>Pteronotus gymnonotus</i> (C)	Venezuela (Amazonas, Bolívar, Guárico, Monagas, Trujillo and Yaracuy)	8	Wenzel 1976
<i>Nycterophilia fairchildi</i> (O)	<i>Pteronotus davyi</i> , <i>Pteronotus gymnonotus</i> , * <i>Pteronotus parnellii</i> (probably <i>P. fuscus</i>) (C) and * <i>Platyrrhinus helleri</i> (C)	Venezuela (Yaracuy)	1	Wenzel 1976
<i>Nycterophilia mormoopsis</i>	<i>Mormoops megalophylla</i>	Venezuela (Yaracuy)	1	Wenzel 1976
<i>Nycterophilia parnelli</i> (M)	<i>Pteronotus parnellii</i> (probably <i>P. fuscus</i>), * <i>Lonchorhina orinocensis</i> (C), * <i>Lonchorhina aurita</i> , * <i>Pteronotus rubiginosus</i> and * <i>Sturnira tildae</i> (C)	Colombia (Vichada) and Venezuela (Amazonas, Apure, Bolívar Carabobo and Yaracuy)	14	Wenzel 1976; Dick et al. 2016
<i>Phalcophila puliciformes</i>	<i>Lonchophylla robusta</i> and <i>Artibeus planirostris</i>	Venezuela (Barinas)	1	Wenzel 1976
Subfamily Trichobiinae				
<i>Anatrichobius scorzai</i> (O)	<i>Myotis oxytus</i> , <i>Myotis keaysi</i> and * <i>Lonchophylla robusta</i>	Venezuela (Barinas, Bolívar and Carabobo)	4	Wenzel 1976
<i>Aspidoptera falcata</i> (O)	<i>Sturnira lilium</i> (probably <i>S. giannae</i>), <i>Sturnira ludovici</i> , <i>Sturnira tildae</i> , * <i>Dermanura cinerea</i> (C), * <i>Artibeus obscurus</i> (C), * <i>Artibeus planirostris</i> (C), * <i>Carollia perspicillata</i> (C), * <i>Phyllostomus hastatus</i> (C), and * <i>Uroderma bilobatum</i> (C)	Venezuela (Amazonas, Apure, Barinas, Carabobo, Guárico, Monagas, Trujillo and Yaracuy)	22	Wenzel 1976
<i>Aspidoptera phyllostomatis</i> (O)	, <i>Artibeus planirostris</i> , * <i>Artibeus lituratus</i> , * <i>Phyllostomus discolor</i> (C) and * <i>Sturnira giannae</i> (C)	Colombia (Arauca, Meta, Vichada and Casanare) and Venezuela (Amazonas, Apure, Barinas, Bolívar, Carabobo, Guárico, Monagas, Trujillo, Yaracuy)	24	Wenzel 1976; Dick et al. 2016; Liévano-Romero et al. 2019; López Rivera et al. 2022
<i>Aspidoptera delatorrei</i> (Pl)	<i>Sturnira giannae</i> and <i>Sturnira cf. parvidens</i> and * <i>Artibeus planirostris</i> ,	Colombia (Arauca and Meta)	2	Dick et al. 2016; López Rivera et al. 2022
<i>Exastinion clovisi</i> (O)	<i>Anoura caudifer</i> , <i>Anoura geoffroyi</i> , <i>Anoura latidens</i> , * <i>Artibeus planirostris</i> (C) and * <i>Pteropteryx macrotis</i> (C)	Venezuela (Amazonas, Barinas, Bolívar, Carabobo, Guarico, Monagas) and Colombia (Arauca)	14	Wenzel 1976; López Rivera et al. 2022
<i>Exastinion oculatum</i> (M)	<i>Anoura cultrata</i>	Venezuela (Aragua)	1	Wenzel 1976
<i>Exastinion deceptivum</i> (M)	<i>Anoura geoffroyi</i>	Venezuela (Merida and Monagas)	1	Wenzel 1976
<i>Mastoptera guimaraesi</i> (M)	<i>Phyllostomus hastatus</i>	Colombia (Arauca) and Venezuela (Apure, Barinas, Trujillo and Yaracuy)	6	This study; Wenzel 1976
<i>Mastoptera minuta</i> (O)	<i>Lophostoma silvicola</i> and <i>Tonatia saurophila</i>	Colombia (Meta, Casanare and Vichada) and Venezuela (Amazonas and Trujillo)	7	Wenzel 1976; Dick et al. 2016; Liévano-Romero et al. 2019
<i>Mastoptera minuta</i> s.l. (Pl)	<i>Lophostoma brasiliense</i> , <i>Lophostoma carrikeri</i> , <i>Lophostoma silvicola</i> , <i>Phyllostomus hastatus</i> , <i>Phyllostomus elongatus</i> , * <i>Anoura</i> sp. (C), * <i>Artibeus obscurus</i> (C), <i>Artibeus planirostris</i> (C) and * <i>Artibeus lituratus</i> (C)	Venezuela (Amazonas, Trujillo and Yaracuy)	10	Wenzel et al. 1976
<i>Megistopoda proxima</i> (Po)	<i>Sturnira giannae</i> , <i>Sturnira parvidens</i> , <i>Sturnira</i> sp., * <i>Carollia perspicillata</i> (C) and <i>Noctilio albiventris</i>	Colombia (Arauca)	2	López Rivera et al. 2022
<i>Megistopoda aranea</i> (O)	<i>Artibeus lituratus</i> and <i>Artibeus planirostris</i>	Colombia (Arauca and Casanare) and Venezuela (Amazonas, Apure, Barinas, Bolívar, Carabobo, Guarico, Monagas, Trujillo and Yaracuy).	28	Wenzel 1976; Liévano-Romero et al. 2019; López López-Rivera et al. 2022

Bat-flies taxa	Bats	Country (Department/State)	Localities	References
<i>Neotrichobius bisetosus</i> (O)	<i>Artibeus obscurus</i> and * <i>Artibeus planirostris</i> (C)	Colombia (Meta) and Venezuela (Amazonas and Bolívar)	9	Wenzel 1976; Dick et al. 2016
<i>Neotrichobius delicatus</i> (M)	<i>Vampyressa thyone</i>	Colombia (Meta) and Venezuela (Barinas, Carabobo and Yaracuy)	16	Wenzel 1976; Dick et al. 2016
<i>Neotrichobius delicatus</i> s.l. (Pl)	<i>Dermanura cinerea</i> , <i>Rhinophylla pumilio</i> , * <i>Artibeus</i> sp., * <i>Artibeus planirostris</i> , * <i>Platyrrhinus helleri</i> (C) and * <i>Uroderma magnirostrum</i>	Venezuela (Amazonas, Apure, Barinas, Bolívar, Carabobo and Guárico)	16	Wenzel 1976
<i>Neotrichobius ectophyllae</i> (M)	<i>Mesophylla macconnelli</i>	Venezuela (Amazonas)	1	Wenzel 1976
<i>Neotrichobius stenopterus</i> (Po)	<i>Dermanura cinerea</i>	Venezuela (Trujillo)	1	Wenzel 1976
<i>Noctiliostrebla aitkeni</i> (M)	<i>Noctilio leporinus</i> and * <i>Saccopteryx bilineata</i>	Colombia (Meta) and Venezuela (Amazonas, Bolívar, Monagas, Yaracuy)	6	Wenzel 1976; Dick et al. 2016
<i>Noctiliostrebla maai</i> (Po)	<i>Noctilio albiventris</i> , * <i>Molossus rufus</i> (C), and <i>Sturnira giannae</i>	Colombia (Arauca) and Venezuela (Amazonas, Apure, Bolívar, Monagas, Yaracuy)	9	This study; Wenzel 1976; López Rivera et al. 2022
<i>Noctiliostrebla dubia</i> (M)	Unidentified host	Colombia (Meta)	1	Dick et al. 2016
<i>Noctiliostrebla traubi</i> (M)	<i>Noctilio leporinus</i>	Venezuela (Guárico and Yaracuy)	2	Wenzel 1976
<i>Paradyschiria curvata</i> (M)	<i>Noctilio albiventris</i> , * <i>Desmodus rotundus</i> (C), * <i>Molossus rufus</i> (C) and * <i>Trachops cirrhosus</i> (C)	Venezuela (Apure)	2	Wenzel 1976
<i>Paradyschiria fusca</i> (M)	<i>Noctilio leporinus</i>	Colombia (Meta and Casanare) and Venezuela (Amazonas, Bolívar, Monagas)	7	Wenzel 1976; Dick et al. 2016
<i>Paradyschiria lineata</i> (M)	<i>Noctilio leporinus</i> , * <i>Noctilio albiventris</i> (C) and * <i>Pteronotus parnelli</i> (probably <i>P. fuscus</i>) (C)	Venezuela (Guarico and Yaracuy)	2	Wenzel 1976
<i>Paradyschiria parvula</i> (M)	<i>Noctilio albiventris</i> , * <i>Molossus aztecus</i> (C) and * <i>Molossus rufus</i> (C)	Colombia (Vichada) and Venezuela (Amazonas, Apure, Monagas and Yaracuy)	5	Wenzel 1976; Dick et al. 2016
<i>Paradyschiria parvuloides</i> (M)	<i>Noctilio albiventris</i> and * <i>Cynomops planirostris</i> (C)	Colombia (Arauca) and Venezuela (Apure and Trujillo)	3	This study; Wenzel 1976; López Rivera et al. 2022
<i>Parastrebla handleyi</i> (M)	<i>Trinycteris nicefori</i>	Venezuela (Bolívar)	1	Wenzel 1976
<i>Paratrichobius dunni</i> (O)	<i>Uroderma bilobatum</i> , <i>Uroderma magnirostrum</i> and * <i>Desmodus rotundus</i> (C),	Venezuela (Amazonas, Apure, Barinas, Bolívar, Trujillo and Yaracuy)	12	Wenzel 1976
<i>Paratrichobius longicrus</i> (M)	<i>Artibeus lituratus</i>	Colombia (Meta) and Venezuela (Amazonas, Apure, Aragua, Barinas, Bolívar, Carabobo, Trujillo and Yaracuy)	16	Wenzel 1976; Dick et al. 2016
<i>Paratrichobius longicrus</i> s.l.	<i>Platyrrhinus aurarius</i> , * <i>Platyrrhinus umbratus</i> , * <i>Platyrrhinus vittatus</i> , * <i>Desmodus rotundus</i> and * <i>Carollia perspicillata</i>	Venezuela (Amazonas, Apure, Barinas, Bolívar, Monagas)	5	Wenzel 1976
<i>Paratrichobius lowei</i> (O)	<i>Dermanura cinerea</i>	Venezuela (Bolívar)	1	Wenzel 1976
<i>Paratrichobius sanchezi</i> (M)	<i>Enchisthenes hartii</i>	Venezuela (Carabobo, Guárico and Monagas)	3	Wenzel 1976
<i>Paratrichobius salivini</i> (M)	<i>Chiroderma salvini</i>	Venezuela (Carabobo and Monagas)	1	Wenzel 1976
<i>Paratrichobius salivini</i> s.l. (Pl)	<i>Chiroderma trinitatum</i> , <i>Chiroderma villosum</i> , <i>Platyrrhinus helleri</i> , <i>Vampyriscus bidens</i> and * <i>Vampyrodes caraccioli</i>	Venezuela (Amazonas, Apure, Barinas, Carabobo and Yaracuy)	9	Wenzel 1976
<i>Pseudostrebla greenwelli</i>	<i>Lophostoma brasiliense</i> and <i>Tonatia maresi</i>	Colombia (Casanare) and Venezuela (Amazonas)	2	Wenzel 1976; Liévan Romero et al. 2019
<i>Pseudostrebla ribeiroi</i>	<i>Lophostoma silvicola</i>	Venezuela (Amazonas)	2	Wenzel 1976
<i>Pseudostrebla sparsisetis</i> (M)	<i>Lophostoma carrikeri</i>	Colombia (Meta) and Venezuela (Amazonas)	3	Wenzel 1976; Dick et al. 2016
<i>Speiseria ambigua</i> (O)	<i>Carollia perspicillata</i> and * <i>Carollia brevicauda</i> (C)	Colombia (Arauca, Meta and Casanare) and Venezuela (Amazonas Apure, Barinas Bolívar, Carabobo, Guárico, Monagas, Trujillo and Yaracuy)	33	Wenzel 1976; Liévan Romero et al. 2019; López Rivera et al. 2022
<i>Speiseria magnioculus</i> (M)	<i>Trachops cirrhosus</i>	Venezuela (Amazonas and Bolívar)	5	Wenzel 1976
<i>Speiseria peytoni</i> (M)	<i>Carollia brevicauda</i>	Colombia (Meta) and Venezuela (Apure, Barinas, Bolívar and Carabobo)	7	Wenzel 1976; Dick et al. 2016
<i>Stizostrebla longirostris</i>	<i>Lophostoma carrikeri</i>	Colombia (Meta) and Venezuela (Amazonas)	2	Wenzel 1976; Dick et al. 2016

Bat-flies taxa	Bats	Country (Department/State)	Localities	References
<i>Trichobius affinis</i> (M)	<i>Lophostoma brasiliense</i>	Venezuela (Amazonas and Apure)	2	Wenzel 1976
<i>Trichobius anducei</i> (M)	<i>Carollia perspicillata</i>	Colombia (Arauca)	1	López Rivera et al. 2022
<i>Trichobius angulatus</i> (M)	<i>Platyrrhinus auraritus</i>	Colombia (Meta) and Venezuela (Amazonas and Bolívar)	3	Wenzel 1976; Dick et al. 2016
<i>Trichobius assimilis</i> (M)	<i>Artibeus planirostris</i> and <i>*Platyrrhinus auraritus</i> (C)	Venezuela (Amazonas and Bolívar)	4	Wenzel 1976
<i>Trichobius bilobus</i> (M)	<i>Pteronotus gymnonotus</i>	Venezuela (Trujillo and Yaracuy)	2	Wenzel 1976
<i>Trichobius caecus</i> (M)	<i>Pteronotus parnellii</i> (probably <i>P. fuscus</i>), <i>Pteronotus rubiginosus</i> , <i>*Anoura geoffroyi</i> (C), <i>*Artibeus obscurus</i> (C), <i>*Artibeus lituratus</i> (C), <i>*Desmodus rotundus</i> (C), <i>*Macrophyllum macrophyllum</i> (C), <i>Myotis keaysi</i> , <i>*Pteronotus davyi</i> (C), <i>*Rhynchoycteris naso</i> (C), <i>*Trachops cirrhosus</i> (C) and <i>*Uroderma bilobatum</i> (C)	Colombia (Vichada) and Venezuela (Amazonas, Apure, Aragua, Bolívar, Carabobo, Guárico, Monagas and Yaracuy)	26	Wenzel 1976; Dick et al. 2016
<i>Trichobius costalimai</i> (M)	<i>Phyllostomus discolor</i> , <i>*Phyllostomus elongatus</i> and <i>*Eptesicus orinocensis</i> (C)	Colombia (Arauca and Meta) and Venezuela (Amazonas, Aragua, Barinas, Bolívar, Guárico, Carabobo, Monagas and Trujillo)	19	Wenzel 1976; Dick et al. 2019; López Rivera et al. 2022
<i>Trichobius diaemi</i> (M)	<i>Diaeus youngii</i>	Colombia (Guainía) and Venezuela (Amazonas)	4	Wenzel 1976; Dick et al. 2016
<i>Trichobius dybasi</i>	Unidentified host	Colombia (Meta)	1	Dick et al. 2016
<i>Trichobius diphyliae</i>	<i>Diphylla ecaudata</i>	Colombia (Vaupes) and Venezuela (Aragua)	1	Wenzel 1976; Dick et al. 2016
<i>Trichobius dugesii</i> (O)	<i>Glossophaga longirostris</i> , <i>Glossophaga soricina</i> , <i>*Carollia brevicauda</i> , <i>*Trinycteris nicefori</i> and <i>*Platyrrhinus helleri</i> (C)	Colombia (Meta and Vichada) and Venezuela (Amazonas, Apure, Barinas, Bolívar, Guárico, Monagas, Trujillo and Yaracuy)	25	Wenzel 1976; Dick et al. 2016
<i>Trichobius dugesioides dugesioides</i> (Pl)	<i>Trachops cirrhosus</i> , <i>*Chrotopterus auritus</i> (C), <i>*Desmodus rotundus</i> (C), <i>*Macrophyllum macrophyllum</i> (C), <i>Phyllostomus discolor</i> (C), <i>*Sphaeronycteris toxophyllum</i> (C) and <i>*Lophostoma silvicola</i> (C)	Colombia (Meta) and Venezuela (Amazonas, Apure, Barinas, Bolívar, Guárico, Trujillo and Yaracuy)	29	Wenzel 1976; Dick et al. 2016
<i>Trichobius dugesioides phyllostomus</i>	<i>Phyllostomus elongatus</i> and <i>*Phyllostomus hastatus</i>	Venezuela (Amazonas and Bolívar)	6	Wenzel 1976; Guerrero 1998
<i>T. dugesioides</i> (probably <i>Trichobius anducei</i>) (O)	<i>Carollia perspicillata</i> and <i>*Carollia brevicauda</i> (C)	Venezuela (Amazonas, Apure, Barinas, Bolívar and Yaracuy)	10	Wenzel 1976; Guerrero 1998
<i>Trichobius ethophallus</i> (M)	<i>Lonchorhina orinocensis</i>	Venezuela (Amazonas and Apure)	5	Wenzel 1976
<i>Trichobius flagellatus</i> (O)	<i>Lonchorhina aurita</i> and <i>*Lonchorhina orinocensis</i>	Venezuela (Amazonas, Barinas, Bolívar and Trujillo)	6	Wenzel 1976
<i>Trichobius furmani</i>	Unidentified host	Colombia (Meta)	1	Dick et al. 2016
<i>Trichobius galei</i> (M)	<i>Natalus tumidirostris</i>	Venezuela (Aragua and Bolívar)	2	Wenzel 1976
<i>Trichobius handleyi</i> (M)	<i>Micronycteris minuta</i> * and <i>*Phyllostomus elongatus</i> (C)	Venezuela (Amazonas, Apure, Bolívar, Guárico, Monagas and Trujillo)	7	Wenzel 1976
<i>Trichobius hispidus</i> (M)	<i>Sturnira bidens</i>	Venezuela (Mérida)	1	Wenzel 1976
<i>Trichobius imitator</i>	<i>Anoura</i> sp.	Venezuela (Bolívar)	1	Wenzel 1976
<i>Trichobius joblingi</i> (Pl)	<i>Carollia perspicillata</i> , <i>Phyllostomus elongatus</i> , <i>Carollia brevicauda</i> , <i>*Desmodus rotundus</i> (C), <i>*Phyllostomus hastatus</i> (C) and <i>*Platyrrhinus fusciventer</i>	Colombia (Arauca, Casanare, Meta and Vaupes) and Venezuela (Amazonas, Apure, Barinas, Bolívar, Guárico, Monagas, Trujillo and Yaracuy)	53	Wenzel 1976; Dick et al. 2016; Liévan-Romero et al. 2019
<i>Trichobius johnsonae</i> (O)	<i>Pteronotus gymnonotus</i> , <i>Pteronotus davyi</i> , <i>Pteronotus personatus</i> and <i>*Noctilio albiventris</i> (C)	Venezuela (Aragua, Bolívar and Yaracuy)	4	Wenzel 1976
<i>Trichobius jubatus</i> (M)	<i>Molossus rufus</i> , <i>Molossus pretiosus</i> and <i>*Molossus molossus</i>	Colombia (Arauca) and Venezuela (Amazonas, Apure and Monagas)	8	This study; Wenzel 1976; López Rivera et al. 2022
<i>Trichobius keemani</i>	<i>Micronycteris megalotis</i> , <i>Micronycteris microtis</i> and <i>Carollia perspicillata</i>	Venezuela (Amazonas, Apure and Barinas)	4	Wenzel 1976
<i>Trichobius leiomotus</i> (M)	<i>Mormoops megalophylla</i>	Venezuela (Bolívar and Yaracuy)	2	Wenzel 1976
<i>Trichobius lioni</i> (M)	<i>Lionycteris spelaea</i> , <i>*Carollia perspicillata</i> (C), <i>*Molossus aztecus</i> (C), <i>*Sturnira lilium</i> (probably <i>S. giannae</i>) (C) and <i>*Platyrrhinus helleri</i> (C)	Colombia (Meta) and Venezuela (Amazonas, Apure and Bolívar)	11	Wenzel 1976; Dick et al. 2019
<i>Trichobius lonchophyllae</i> (O)	<i>Lonchophylla robusta</i> , <i>Lonchophylla orienticollina</i> , <i>*Myotis handleyi</i> , <i>*Anoura latidens</i> (C) and <i>*Sturnira lilium</i> (probably <i>S. giannae</i>) (C)	Colombia (Arauca) and Venezuela (Barinas)	5	Wenzel 1976; López Rivera et al. 2022

Bat-flies taxa	Bats	Country (Department/State)	Localities	References
<i>Trichobius longipes</i> (O)	<i>Phyllostomus hastatus</i> , <i>Phyllostomus elongatus</i> , * <i>Artibeus planirostris</i> (C), * <i>Desmodus rotundus</i> (C), * <i>Molossus rufus</i> , * <i>Rhynchonycteris naso</i> (C) and * <i>Uroderma bilobatum</i> (C)	Colombia (Arauca) and Venezuela (Amazonas, Apure, Aragua, Barinas, Bolívar, Carabobo, Guárico, Monagas, Trujillo and Yaracuy)	23	Wenzel 1976; López Rivera et al. 2022
<i>Trichobius longipilis</i> (M)	<i>Pteropteryx macrotis</i>	Venezuela (Bolívar)	2	Wenzel 1976
<i>Trichobius macrophylli</i> (M)	<i>Macrophyllum macrophyllum</i>	Venezuela (Amazonas, Apure, Bolívar and Guárico)	6	Wenzel 1976
<i>Trichobius pallidus</i>	<i>Furipterus horrens</i>	Venezuela (Amazonas)	1	Wenzel 1976
<i>Trichobius parasparsus</i> (M)	<i>Pteronotus parnellii</i> (probably <i>P. fuscus</i>), * <i>Carollia perspicillata</i> (C), * <i>Sturnira lilium</i> (probably <i>S. giannae</i>) (C), * <i>Sturnira tildae</i> (C) and * <i>Uroderma convexum</i> (C)	Venezuela (Amazonas, Apure, Bolívar and Yaracuy) and Colombia (Vichada)	17	Wenzel 1976; Dick et al. 2016
<i>Trichobius parasiticus</i> (Pl)	<i>Desmodus rotundus</i> , <i>Carollia perspicillata</i> , <i>Chiroderma villosum</i> and <i>Platyrrhinus umbratus</i>	Colombia (Meta and Arauca) and Venezuela (Amazonas, Apure, Barinas, Bolívar, Carabobo, Guárico, Monagas, Trujillo and Yaracuy)	39	This study; Wenzel 1976; Dick et al. 2016
<i>Trichobius permilis</i> (O)	<i>Carollia brevicauda</i> , <i>Carollia perspicillata</i> and * <i>Phyllostomus elongatus</i>	Venezuela (Apure, Barinas, Bolívar, Carabobo and Monagas)	10	Wenzel 1976
<i>Trichobius petersoni</i> (O)	<i>Sturnira erythromos</i> and * <i>Sturnira bogotensis</i>	Venezuela (Barinas, Mérida and Monagas)	4	Wenzel 1976
<i>Trichobius propinquus</i> (O)	<i>Anoura geoffroyi</i>	Venezuela (Bolívar)	1	Wenzel 1976
<i>Trichobius silvicola</i> (M)	<i>Lophostoma silvicola</i> and * <i>Phyllostomus hastatus</i>	Venezuela (Amazonas and Bolívar)	2	Wenzel 1976
<i>Trichobius sparsus</i> (M)	<i>Pteronotus parnellii</i> and * <i>Natalus tumidirostris</i> (C)	Venezuela (Amazonas, Bolívar and Guárico)	10	Wenzel 1976
<i>Trichobius strictisternus</i>	<i>Lophostoma carrikeri</i> *	Venezuela (Amazonas)	1	Wenzel 1976
<i>Trichobius tiptoni</i> (M)	<i>Anoura caudifer</i> , * <i>Carollia perspicillata</i> (C), * <i>Desmodus rotundus</i> (C), * <i>Sturnira ludovici</i> (C) and * <i>Platyrrhinus helleri</i> (C)	Venezuela (Barinas, Bolívar, Carabobo and Yaracuy)	4	Wenzel 1976
<i>Trichobius tuttlei</i>	<i>Lampronycteris brachyotis</i>	Venezuela (Amazonas)	1	Wenzel 1976
<i>Trichobius uniformis</i> (O)	<i>Glossophaga longirostris</i> , <i>Glossophaga soricina</i> , * <i>Artibeus lituratus</i> (C), * <i>Carollia perspicillata</i> (C) and * <i>Platyrrhinus helleri</i> (C)	Venezuela (Amazonas, Apure, Barinas, Bolívar, Guárico, Monagas, Trujillo and Yaracuy)	21	Wenzel 1976
<i>Trichobius urodermae</i> (M)	<i>Uroderma bilobatum</i>	Venezuela (Amazonas and Trujillo)	5	Wenzel 1976
<i>Trichobius vampyrops</i> (O)	<i>Platyrrhinus vittatus</i>	Venezuela (Barinas)	1	Wenzel 1976
<i>Trichobioides perspicillatus</i> (M)	<i>Phyllostomus discolor</i>	Colombia (Arauca, Meta) and Venezuela (Amazonas, Aragua, Barinas, Bolívar, Carabobo, Guárico, Monagas and Trujillo)	13	Wenzel 1976; Dick et al. 2016; López Rivera et al. 2022
<i>Xenotrichobius noctilionis</i>	<i>Noctilio albiventris</i> and <i>Noctilio leporinus</i>	Venezuela (Amazonas and Apure)	2	Wenzel 1976
Subfamily Streblinae				
<i>Anastrebla caudiferae</i> (M)	<i>Anoura caudifer</i>	Venezuela (Amazonas, Barinas and Bolívar)	3	Wenzel 1976
<i>Anastrebla modestini</i> (M)	<i>Anoura geoffroyi</i> and <i>Anoura</i> sp.	Venezuela (Amazonas, Barinas, Bolívar, Carabobo, Guárico, Mérida and Monagas)	12	Wenzel 1976
<i>Anastrebla nycteridis</i> (O)	<i>Lonchophylla robusta</i> and <i>Lonchophylla orienticollina</i>	Colombia (Arauca) and Venezuela (Barinas)	2	Wenzel 1976; López Rivera et al. 2022
<i>Anastrebla spurrelli</i> (M)	<i>Lionycteris spurrelli</i> and * <i>Ametrida centurio</i> (C)	Venezuela (Amazonas and Bolívar)	7	Wenzel 1976
<i>Metalasmus pseudopterus</i> (Pl)	<i>Artibeus planirostris</i> , * <i>Artibeus lituratus</i> , * <i>Chiroderma villosum</i> , * <i>Myotis nigricans</i> , * <i>Pteropteryx macrotis</i> , * <i>Phyllostomus hastatus</i> and * <i>Uroderma magnirostrum</i>	Colombia (Meta) and Venezuela (Amazonas, Apure, Barinas, Bolívar, Guarico, Lara, Monagas, Trujillo and Yaracuy)	18	Wenzel 1976; Dick et al. 2016
<i>Metalasmus</i> sp.	<i>Sturnira ludovici</i>	Venezuela (Barinas)	1	Wenzel 1976
<i>Paraeuctenodes longipes</i> (M)	<i>Glossophaga soricina</i> , * <i>Artibeus lituratus</i> and * <i>Nyctinomops laticaudatus</i> (C)	Venezuela (Amazonas, Bolívar and Yaracuy)	7	Wenzel 1976
<i>Paraeuctenodes similis</i> (M)	<i>Carollia perspicillata</i>	Venezuela (Bolívar)	2	Wenzel 1976
<i>Strebla altmani</i> (O)	<i>Lonchorhina aurita</i> , <i>Lonchorhina orinocensis</i> and * <i>Macrophyllum macrophyllum</i>	Colombia (Meta) and Venezuela (Amazonas, Aragua, Apure, Barinas, Bolívar, Trujillo and Yaracuy)	15	Wenzel 1976; Dick et al. 2016
<i>Strebla alvarezi</i> (O)	<i>Micronycteris microtis</i> , <i>Micronycteris megalotis</i> , <i>Carollia brevicauda</i> , and <i>Lonchophylla thomasi</i>	Colombia (Meta) and Venezuela (Amazonas, Bolívar and Yaracuy)	5	Wenzel 1976; Dick et al. 2016

Bat-flies taxa	Bats	Country (Department/State)	Localities	References
<i>Strebla asternalis</i> (M)	<i>Saccopteryx bilineata</i> and <i>Saccopteryx</i> sp.	Venezuela (Amazonas)	2	Wenzel 1976
<i>Strebla christinae</i> (M)	<i>Phylloderma stenops</i> , * <i>Eumops glaucinus</i> (C) and * <i>Uroderma bilobatum</i> (C)	Venezuela (Amazonas, Apure and Bolívar)	8	Wenzel 1976
<i>Strebla choropteri</i> (M)	<i>Chrotopterus auritus</i>	Venezuela (Amazonas and Bolívar)	4	Wenzel 1976
<i>Strebla consocia</i> (O)	<i>Phyllostomus elongatus</i> , <i>Phyllostomus hastatus</i> , * <i>Desmodus rotundus</i> (C), * <i>Trachops cirrhosus</i> (C) and * <i>Platyrrhinus helleri</i> (C)	Colombia (Meta) and Venezuela (Amazonas, Apure, Barinas, Bolívar, Carabobo, Monagas, Trujillo, Yaracuy)	21	Wenzel 1976; Dick et al. 2016
<i>Strebla cormurae</i>	<i>Cormura brevirostris</i>	Venezuela (Amazonas)	1	Wenzel 1976
<i>Strebla curvata</i> (O)	<i>Glossophaga longirostris</i> , <i>Glossophaga soricina</i> , * <i>Carollia brevicauda</i> , * <i>Carollia perspicillata</i> and * <i>Noctilio albiventris</i> (C)	Venezuela (Amazonas, Apure, Barinas, Bolívar and Monagas)	12	Wenzel 1976
<i>Strebla diaemi</i> (M)	<i>Diaemus youngii</i>	Colombia (Guainía) and Venezuela (Amazonas)	4	Wenzel 1976; Dick et al. 2016
<i>Strebla diphyllae</i>	<i>Diphylla ecaudata</i>	Colombia (Vaupés) and Venezuela (Aragua)	1	Wenzel 1976; Dick et al. 2016
<i>Strebla galindoi</i> (M)	<i>Tonatia bidens</i>	Venezuela (Amazonas, Apure and Bolívar)	4	Wenzel 1976
<i>Strebla guajiro</i> (O)	<i>Carollia brevicauda</i> and <i>Carollia perspicillata</i>	Colombia (Casanare, Meta and Vichada) and Venezuela (Amazonas, Apure, Aragua, Barinas, Bolívar, Guárico, Monagas, Trujillo and Yaracuy)	50	Wenzel 1976; Dick et al. 2016; Liévano et al. 2019
<i>Strebla harderi</i> (M)	<i>Anoura geoffroyi</i> and <i>Anoura</i> sp.	Venezuela (Amazonas and Bolívar)	4	Wenzel 1976
<i>Strebla hertigi</i> (O)	<i>Phyllostomus discolor</i> and * <i>Phyllostomus elongatus</i> (C)	Colombia (Arauca) and Venezuela (Amazonas, Barinas, Bolívar, Guarico, Monagas and Trujillo)	14	Wenzel 1976; López Rivera et al. 2022
<i>Strebla kohlsi</i>	<i>Lophostoma silvicola</i>	Venezuela (Amazonas)	2	Wenzel 1976
<i>Strebla machadoi</i> (M)	<i>Micronycteris minuta</i>	Venezuela (Amazonas, Apure, Bolívar and Monagas)	4	Wenzel 1976
<i>Strebla matsoni</i> (M)	<i>Macrophyllum macrophyllum</i> and * <i>Rhynchonycteris naso</i> (C)	Venezuela (Amazonas, Apure and Bolívar)	5	Wenzel 1976
<i>Strebla mirabilis</i> (M)	<i>Trachops cirrhosus</i> , * <i>Phyllostomus hastatus</i> (C), * <i>Phyllostomus elongatus</i> (C), * <i>Artibeus planirostris</i> , * <i>Chrotopterus auritus</i> (C) and * <i>Diphylla ecaudata</i>	Colombia (Meta) and Venezuela (Amazonas, Apure, Aragua, Bolívar, Guarico and Yaracuy)	19	Wenzel 1976; Dick et al. 2016
<i>Strebla obtusa</i> (M)	<i>Trinycteris nicefori</i> and * <i>Phyllostomus elongatus</i> (C)	Venezuela (Amazonas and Bolívar)	5	Wenzel 1976
<i>Strebla paramirabilis</i> (Pl)	<i>Artibeus planirostris</i> , <i>Platyrrhinus aurarius</i> and * <i>Anoura geoffroyi</i> (C)	Colombia (Meta) and Venezuela (Amazonas and Bolívar)	5	Wenzel 1976; Dick et al. 2016
<i>Strebla proxima</i> (O)	<i>Peropteryx macrotis</i> and <i>Peropteryx trinitatis</i>	Venezuela (Amazonas and Yaracuy)	2	Wenzel 1976
<i>Strebla tonatiae</i> (O)	<i>Lophostoma brasiliense</i> , * <i>Tonatia saurophila</i> , * <i>Lophostoma carrikeri</i> (C) and * <i>Sturnira lilium</i> ("probably <i>S. giannae</i> ") (C)	Colombia (Casanare) and Venezuela (Amazonas, Apure, Bolívar, Monagas, Trujillo and Yaracuy)	9	Wenzel 1976; Liévano-Romero et al. 2019
<i>Strebla wiedemannii</i> (M)	<i>Desmodus rotundus</i>	Colombia (Meta) and Venezuela (Amazonas, Apure, Barinas, Bolívar, Guárico, Lara, Monagas, Trujillo and Yaracuy)	28	Wenzel 1976; Dick et al. 2016
Family Nycteribiidae				
<i>Basilia anceps</i>	Unidentified host	Venezuela (Bolívar and Amazonas)	2	Graciolli et al. 2007
<i>Basilia bequaerti</i>	Unidentified host	Venezuela (Amazonas)	1	Guimarães 1972
<i>Basilia constricta</i>	Unidentified host	Venezuela (Mérida)	1	Guimarães 1972
<i>Basilia dunnii</i>	Unidentified host	Venezuela (Amazonas)	1	Graciolli et al. 2007
<i>Basilia dubia</i> (M)	<i>Myotis albescens</i> and * <i>Saccopteryx bilineata</i> (C)	Venezuela (Amazonas and Apure)	3	Guimarães 1972
<i>Basilia ferrisi</i> (O)	<i>Myotis albescens</i> , <i>Myotis handleyi</i> , <i>Myotis nigricans</i> , <i>Myotis riparius</i> , * <i>Molossus pretiosus</i> (C), * <i>Desmodus rotundus</i> , * <i>Noctilio albiventris</i> (C), * <i>Phyllostomus elongatus</i> (C) and <i>Platyrrhinus helleri</i> (C)	Colombia (Arauca and Meta) and Venezuela (Amazonas, Apure, Aragua, Bolívar and Monagas)	8	Guimarães 1972; Graciolli et al. 2007; López Rivera et al. 2022
<i>Basilia juquiensis</i> (O)	<i>Myotis riparius</i>	Venezuela (Apure)	1	Guimarães 1972
<i>Basilia ortizi</i> (O)	<i>Eptesicus brasiliensis</i> , <i>Eptesicus furinalis</i> , <i>Eptesicus orinocensis</i> and * <i>Myotis riparius</i>	Colombia (Arauca), Venezuela (Amazonas, Bolívar and Monagas)	8	Guimarães 1972 López Rivera et al. 2022
<i>Basilia tiptoni</i> (M)	<i>Gardnerycteris crenulatum</i>	Venezuela (Apure and Trujillo)	2	Guimarães 1972
<i>Basilia typhlops</i>	<i>Myotis oxyotus</i>	Venezuela (Bolívar)	1	Guimarães 1972
<i>Basilia tuttlei</i>	<i>Myotis nigricans</i>	Venezuela (Amazonas)	1	Guimarães 1972
<i>Basilia wenzeli</i>	<i>Eptesicus fuscus</i> and <i>Lonchorhina aurita</i>	Venezuela (Aragua)	1	Guimarães 1972

Table 2. Ectoparasitic flies collected on bats during 2021 in the Department of Arauca, Colombia. * New records of bat fly species for Colombia. ** New association between bat flies and bats.

Taxon	No individuals	Host	Prevalence	Locality	Coordinates	Voucher
Streblidae						
<i>Mastoptera guimaraesi</i>	7♀ and 5♂	<i>Phyllostomus hastatus</i> (1♂)	1.00	1	06°46'47"N, 70°42'59.3"W	MHN-UCa-Ec 555
<i>Noctiliostrebla maai</i>	12♀ and 9♂	<i>Noctilio albiventris</i> (1♀ and 2♂)	0.26	1	06°46'46.4"N, 70°43'00"W; 06°46'46.7"N, 70°43'02.1"W	MHN-UCa-Ec 555, 561 and 562
	37♀ and 27♂	<i>Noctilio albiventris</i> (♀ and ♂)	0.26	2	06°46'43.2"N, 70°43'36.1"W	MHN-UCa-Ec 565, 568, 570–573, 580, 583, 586 and 588
<i>Paradyschiria parvuloides</i>	1♀	<i>Cynomops planirostris</i> (1♀)**	0.1	1	06°46'43.2"N, 70°43'36.1"W	MHN-UCa-Ec 509
	27♀ and 15♂	<i>Noctilio albiventris</i> (1♀ and 3♂)	0.32	1	06°46'46.4"N, 70°43'00"W; 06°46'46.7"N, 70°43'02.1"W	MHN-UCa-Ec 557, 559, 560 and 563
	39♀ and 32♂	<i>Noctilio albiventris</i> (♀ and ♂)	0.32	2	06°46'43.2"N, 70°43'36.1"W	MHN-UCa-Ec 566, 567, 569, 572, 574–579, 581, 582, 585 and 589
<i>Trichobius jubatus*</i>	1♂	<i>Molossus rufus</i> (♂)	0.05	2	06°46'43.2"N, 70°43'36.1"W	MHN-UCa-Ec 564
<i>Trichobius parasiticus</i>	1♂	<i>Desmodus rotundus</i> (1♂)	1.00	1	06°46'46.3"N, 70°42'59.2"W	
Nycteribiidae						
<i>Basilia ferrisi</i>	6♀ and 5♂	<i>Myotis handleyi</i> (1♀ and 1♂)	0.25	1	06°46'47"N, 70°42'59.3"W	MHN-UCa-Ec 552

Universidad de Caldas (2 June 2017). Bats and ectoparasites were deposited at the Mammals (M) and Ectoparasites (Ec) collections of the MHN-UCa. Of these records, we calculated the general prevalence for host species defined as the number of individuals of a host species infected with a particular parasite species divided by the number of hosts examined (Margolis et al. 1982).

For the taxonomic identification of the captured bats, we took morphometric measurements including the total length, tail length, forearm length, ear length and foot length (Nagorsen and Peterson 1980) and used taxonomic keys (e.g., Gardner 2008). For the identification of the specimens of Streblidae and Nycteribiidae, we used the dichotomous keys of Wenzel et al. (1966), Wenzel (1976), Guerrero (1994, 1995, 1998), Autino et al. (1999), and Alcantara et al. (2019) using morphological features to distinguish between different species (Wenzel 1976; Autino et al. 1999). These traits include the shape and size of the body, wing, mesonotum, head, eyes and female and male reproductive structures. Identifications were done with the help of a stereomicroscope and compared with additional specimens deposited in the MHN-UCa-Ec collection.

Bat-ectoparasite network structure and complex network metrics

We used the new records reported in the present study and the records collected in the literature to build bipartite interaction networks for the Streblidae and Nycteribiidae of the Orinoquia. In the networks, bat and ectoparasite species are represented by nodes and interacting species are linked by lines, with the width of the line proportional to the frequency of each interaction. We created a net that includes all raw records, as well as other nets that exclude records that could be the product of incidental transfer or contamination. In addition, we carried out an analysis of the Streblidae species present in the interaction networks, classifying them into the three ecomorphological groups proposed by Dick (2005): wing crawler, fur runners, and fur swimmers. This allowed us to infer possible niche partitioning between host species across the networks.

To evaluate the properties of the network we used the index of specialization by communities (H_2), the quantitative modularity QuanBiMo (Q), the connectance (C), and the nestedness (wNODF) (Dormann et al. 2009; Fortuna et al. 2010; Mello et al. 2016). The standardized two-dimensional entropy index (H_2) measures both the degree of niche complementarity among species and the specialization at species level (Blüthgen et al. 2006). This index varies from 0 (unspecialized network) to 1 (perfectly specialized network). We calculated Quantitative Modularity (QuanBiMo) that allows determining the existence of sets or groupings within the complete network, that is, when there are species that interact more closely, forming modules (Fortuna et al. 2010). The modularity ranges from 0 (non-modular) to 100 (fully modular) and were estimated using the algorithm QuanBiMo (Dormann and Strauss 2014). The C index represents the number of interactions or links observed in the network, between bats and their ectoparasitic flies considering the total number of potential interactions. It takes values from 0 to 1 where 0 indicates that there are no connections and 1 which denotes that most of the nodes in the network interact with each other (Blüthgen et al. 2006). Then we calculated the Weighted Nestedness (wNODF) ranging from 0 (non-nested) to 100 (fully nested), to measure how strongly species interactions of seldom connected species were nested within those of highly connected species (Almeida-Neto and Ulrich 2011).

Also, we assessed the role of bat and fly species using one centrality metric. The degree centrality (DC) measured the number of interactions of a given species, reflecting its degree of generalization versus specialization and the (González et al. 2010). All calculations were carried out using the R software (R Core Team 2022) and the vegan (diversity analysis), iNEXT (interaction accumulation curve), network and bipartite (interaction networks and metrics) packages (Butts 2008; Dormann et al. 2008; Hsieh et al. 2016; Oksanen et al. 2020).

Results

Richness of bats and ectoparasite flies in the Orinoco Region

Our review showed 1046 records of 129 species of ectoparasitic flies associated with 102 species of bats (Table 1) in the Orinoquia Region (Suppl. material 1: table S1), and seven species (*Basilia anceps*, *Basilia bequaerti*, *Basilia constricta*, *Basilia dunnii*, *Noctiliostrebla dubia*, *Trichobius dybasi* and *Trichobius furmani*) in which the associated host were not recorded (Table 1, Suppl. material 1: table S2). The flies were mainly associated with bats of the families Phyllostomidae (69 species, ~ 79% of the records), followed by Vespertilionidae (10 species), Molossidae (seven species), Emballonuridae (six species), Mormoopidae (six species), Noctilionidae (two species), Furipteridae (one species), and Natalidae (one species) (Table 1). Species of Nycteribiidae were found associated mainly with species of Vespertilionidae (27 records of seven species of flies associated with nine species of bats), followed by Phyllostomidae (six records of three species of flies associated with five species of bats) (Table 1). 119 species of Streblidae were reported associated mainly with species of Phyllostomidae (821 records of 97 species of bat flies associated with 68 species of bats) followed by Mormoopidae (89 records of 11 species of bat flies associated with six species of bats) (Table 1).

The evaluation of the quality of the associations between flies and hosts showed that 9.8% of them could be the result of incidental transfers or contamination (Table 1). For example, reported associations of *Trichobius caecus* were found with 11 bat species, of which nine are considered non-primary associations and might represent incidental transfers or contamination. Similarly, reported associations of *Aspidoptera falcata* with nine bat species were recorded, of which six were identified as possible contaminations. We also detected unresolved non-primary associations where a significant number of individuals (> 5% of records) were found in the absence of the primary hosts. For example: *Mastoptera minuta* is primarily associated with *Lophostoma silvicola* but Liévano-Romero et al. (2019) reported 16 individuals of this species parasitizing *Tonatia maresi* in Casanare, Colombia. Furthermore, *Megistopoda proxima* is primarily associated with *Sturnira* species, but López Rivera et al. (2022) found three individuals parasitizing *Noctilio albiventris* and seven individuals of *Noctiliostrebla maai* on a *Sturnira gigantea* in the absence of the primary host (*N. albiventris*) in Arauca, Colombia (Table 1). Distinguishing non-primary associations due to contamination among ecological associations were also challenging for species complexes, given their taxonomic complexity. For instance, *Mastoptera minuta* s.l. were recorded associated with eight bat species belonging to four genera (*Anoura*, *Artibeus*, *Lophostoma*, *Phyllostomus*). *Paratrichobius salvini* s.l. were reported associated with five bat species from three genera (*Chiroderma*, *Platyrrhinus*, and *Vampyriscus*). *Neotrichobius delicatus* s.l. were found associated with two bat species from different genera (*Dermanura cinerea* and *Rhynophylla pumilio*). Furthermore, the specificity of other ectoparasitic fly species could not be determined due to limited or unique records, such as *Neotrichobius ectophyllae*, *Neotrichobius stenopterus*, *Strebla cormurae*, *Trichobius pallidus*, and *Trichobius tuttlei* (Table 1). Regarding the morphological confirmation of some non-primary associations previously reported for the Department of Arauca, Colombia, we reidentified samples of *Trichobius mendezi* found in association with *Molossus molossus* and *Molossus pretiosus* as *Trichobius jubatus*. Similarly, *T. mendezi* found in non-primary association with *Phyllostomus hastatus* were morphologically reidentified as *Trichobius longipes*.

New records for Colombia

We captured 141 bats of 16 species (*Carollia brevicauda*, *Cynomops milleri*, *Cynomops planirostris*, *Desmodus rotundus*, *Eptesicus orinocensis*, *Eumops glaucinus*, *Eumops nanus*, *Molossops temminckii*, *Molossus coibensis*, *M. molossus*, *M. pretiosus*, *Molossus rufus*, *Myotis handleyi*, *N. albiventris*, *P. hastatus* and *Uroderma magnirostrum*) of which 24 individuals of six bat species had ectoparasitic flies (*C. planirostris*, *D. rotundus*, *M. handleyi*, *M. rufus*, *N. albiventris*, and *P. hastatus*). In total, we obtained 223 flies belonging to five species of Streblidae: *Mastoptera guimaraesi* (Fig. 2), *N. maai* (Fig. 3), *Paraduschiria parvuloides* (Fig. 4), *Trichobius jubatus* (Fig. 5A–C), and *T. parasiticus* (Fig. 5D, F), and one species of Nycteribiidae: *Basilia ferrisi* (Fig. 6). Our study reports for the first time the presence *T. jubatus* in Colombia previously reported in Venezuela (Table 2). The association between *Paraduschiria parvuloides*

and *C. planirostris*, although novel, could be related to possible contamination, since only one individual was collected in the presence of the primary host *N. albiventris* (see Table 2).

Bat-ectoparasite network structure and complex network metrics

The Streblidae-bat interaction network for the Orinoquia was made up of 121 species of ectoparasitic flies and 91 species of bats (Suppl. material 2). The quantitative modularity QuanBiMo and specialization calculated for the interaction network was high ($H_2' = 0.94$ and $Q = 0.85$, respectively), which indicates a high niche (bats) differentiation in the network. We observed low connectance ($C = 0.02$) and nestedness ($wNODF = 2.39$). The degree of centrality (DC) shows us 66 highly specialized bat flies interacting with a single bat species (Suppl. material 1: table S3), and ten relatively generalist species interacting with more than four species of bats: *Aspidoptera delatorrei*, *Exastinion clovisi*, *M. proxima*,




Figure 2. Micrographs of *Mastoptera guimaraesi*: female **A** dorsal and **B** ventral views (minute species, body 0.73–1.4 mm long, with short legs) **C** lateral lobes of tergum 1+2, these longer and heavier, dorsal view; male **D** dorsal and **E** ventral views **F** male genital apparatus, lateral view. Abbreviations: ae: aedeagus; gon: gonopods; mga: male genital apparatus; ms: mesonoto; St2: Sternum 2; t1+2: tergum 1 + 2, w: wings.




Figure 3. Micrographs of *Noctiliostrebla maai*: female **A** dorsal and **B** ventral views (body 3.0–3.2 mm long) **C** median and transverse mesonotal sutures united to form an inverted Y and median wing vein usually with 1 or 2 setae (arrowed), dorsal view **D** terminal cone, female abdomen, ventral view **E** male, ventral views **F** male genital apparatus, lateral view and **G** male and female copulating. Abbreviations: ae: aedeagus; gon: gonopods; mga: male genital apparatus; ms: mesonoto; St2: Sternum 2; sst: surstylus; t1+2: tergum I + 2; tc: termina cone.

Nycterophilus coxata, *Nycterophilus parnelli*, *Strebla alvarezi*, *Strebla curvata*, *Trichobius dugesii*, *Trichobius johnsonae* and *Trichobius parasiticus* (DC = 4) and one highly generalist species parasitizing eight bat species: *Metalasmus pseudopterus* (DC = 8). In addition, as expected, the species complexes also proved to be the most general, with associations encompassing up to five host species: *M. minuta* s.l., *N. delicatus* s.l., *P. longicrus* s.l., and *Paratrichobius salvini* s.l. (Suppl. material 1: table S3). We also observed that 39 of the 91 bat species were parasitized by a single fly species (Suppl. material 1: table S4) and 11 bat species were parasitized by more than five bat fly species: *Anoura geoffroyi*, *Artibeus lituratus*, *Lophostoma silvicola*, *Noctilio leporinus* and *Pteronotus parnellii*



Figure 4. Micrographs of *Paradyschiria parvuloides*: male **A** dorsal view (body 2.5–3.0 mm long) **B** mesonotum usually with a short seta on each side anterior and lateral to the long posterior macroseta (arrow), dorsal view **C** male genital apparatus, lateral view; female **D** ventral view **E** seventh sternites each with 3 or -4, rarely 2, spine-like setae on distal margin (arrows), ventral view **F** supra-anal plate as wide as long, the basal (anterior) margin roundly angulate, dorsal view. Abbreviations: ae: ae-deagus; gon: gonopods; ms: mesonoto; sap: supra-anal plate; St2: Sternum 2; St7: Sternum 7; sst: surstylus; t1+2: tergum I + 2.

(probably *P. fuscus*) ($DC = 5$), *C. brevicauda*, *N. albiventris*, and *P. hastatus* ($DC = 7$), *Phyllostomus elongatus* ($DC = 9$), *Artibeus planirostris* ($DC = 10$), and *Carollia perspicillata* ($DC = 11$) (Suppl. material 1: table S4). These bat species are key hosts, as they act as connectors between the different bat fly species that make up the web. This means that these species play a fundamental role in the interaction between different species of ectoparasites, therefore, their presence and characteristics can be determinant for the survival and dispersal of ectoparasites in the network.

The analysis of the interaction network between Streblidae and bats in the Orinoquia revealed a high modularity and specialization and three different modules. The first module or group was composed of nine species of flies belonging to four genera (*Anastrebla*, *Exastinion*, *Trichobius*, and *Strebla*) exclusive to bats of the genus *Anoura* (Fig. 7). In this group, *E. clovisi* were observed to be the most generalist species, associated with three *Anoura* species, but showing a stronger association with *Anoura geoffroyi*. *Anoura geoffroyi* was the bat species with the




Figure 5. A–C Micrographs of *Trichobius parasiticus* male: A dorsal view (body 5.4 mm long) B head (latero vertices and occipital lobes well defined) and mesonotum (prescutum with very short discal setae immediately in front of the transverse suture (black arrow) and scutum posteriorly with an irregular W-shaped row of short setae (white arrow)), dorsal view C gonopods, lateral view D–F micrographs of *Trichobius jubatus* male: D dorsal views (body 3.8 mm long) E head (occipital lobes of head densely setose, eyes multifaceted) and mesonotum (mesonotum essentially setose throughout, and median and transverse sutures not united), dorsal view F metasternal lobe united with metepimeron (arrows), ventral view. Abbreviations: cx.2: mesocoxa; cx.3: metacoxa; e: eyes; epm: metepimeron; gon: gonopods; mga: male genital apparatus; ms: mesonoto; mts: metasternum; mts.l.: metasternal lobe; St2: Sternum 2; t1+2: tergum I + 2; w: wings.

highest number of associated fly species ($DC = 5$). In module 1, two ecomorphological groups were mainly found: the “wing crawler” (represented by *Exastinion* and *Trichobius*) and the “fur swimmer” (represented by *Anastrebla* and *Strebla*). In general, the species of both groups were associated with the same hosts, indicating the possibility of a niche partition within the host bat species (Fig. 7).

The second module (the largest group) was composed of 91 species of flies of the three subfamilies (Nycterophiliinae, Streblinae and Trichobiinae) associated with 64 species of bats of the families Emballonuridae, Mormoopidae, Molossidae, Noctilionidae, Phyllostomidae, and Vespertilionidae, (Fig. 8). In this group, highly interconnected hosts are highlighted, hosting several species of ectoparasitic flies. Among them there are two species of Noctilionidae (*N. leporinus* and *N. albiventris*), one Mormoopidae (*Pteronotus parnellii*; probably *P. fuscus*), and seven of Phyllostomidae (*A. planirostris*, *Trachops cirrhosus*, *Phyllostomus elongatus*, *P. hastatus*, *Phyllostomus discolor*, *C. brevicauda*, and *C. perspicillata*).



Figure 6. Micrographs of *Basilia ferrisi*: female **A** ventral and **B** dorsal views: body 1.8–2.0 mm long, tibiae with three rows of setae (circle) **C** female with: tergal plate II transformed into two elongate lobes with short and long setae or with posterior margin arcuate, dorsal view; male **D** ventral and **E** dorsal views **F** male genital apparatus, lateral view. Abbreviations: ae: aedeagus; gon: gonopods; mga: male genital apparatus; tpl: tergal plate I and tpll: tergal plate II. Circles **A**, **D** indicate tibiae with three rows of setae.

Additionally, we confirmed primary associations supported by their abundances, such as *Paradyschiria parvula* with *N. albiventris*, *T. caecus* with *P. parnellii* (probably *P. fuscus*), *Trichobius joblingi* with *C. perspicillata*, *Trichobius costalimai* with *Phyllostomus discolor*, and *T. parasiticus* and *Strebla wiedemanni* with *D. rotundus*. For the second module the three ecomorphological groups were identified. Wing crawlers were the most representative, comprising ~ 57% of the fly species, followed by fur swimmers, representing ~ 27.5% of the fly species, and finally, in smaller proportion, fur runners (Fig. 8). In general, in this module, species from different ecomorphological groups were associated with the same host species. In some cases, only two ecomorphological groups were present, while for more generalist host species such as *A. planirostris* and *C. perspicillata*, all three ecomorphological groups were found together. Although the majority of the studies analyzed in this work do not specify which fly specimens were collected on the same hosts, these associations may indicate niche partitioning within the same host bat species (Fig. 8).

The third module of Strebidae interaction network is represented by the most specific species or unique associations. It consists of 25 species of bat flies and 21 species of bats of the families Emballonuridae, Furipteridae, Mormoopidae, Natalidae, and Phyllostomidae (Fig. 9). We confirmed primary associations based on the abundance of records between *Neotrichobius bisetosus* and *Artibeus obscurus*; *Strebla diaemi* and *Trichobius diaemi* with *Diaemus youngii*; *Anastrebla spurrelli* and *Trichobius lonycteridis* with *Lionycteris spurrelli*; *Strebla christinae* with *Phylloderma stenops*, and finally, *A. falcata* with three species of *Sturnira*




Figure 7. Bipartite bat-fly quantitative network (Module 1 - Streblidae). The size of the bar on the left (orange) represents the abundance (number of individuals) of bat flies per species observed and the size of the bar on the right (green) represents the abundance of bats for which the bat fly sample was obtained. The width of the black lines/bars indicates the frequency of interactions.

(*S. cf. lilum*, *S. ludovici*, and *S. tildae*). Like modules 1 and 2 of this network, the incidence of different ecomorphological types within the same host bat species suggests niche partitioning pattern across the entire network (Fig. 9).

For Nycteribiidae, the interaction network was composed of eight species of flies of the genus *Basilia* and 17 species of bats mainly of the families Vesperilionidae (nine species) and Phyllostomidae (five species), with additional re-




Figure 8. Quantitative bipartite bat-fly network (Module 2 - Streblidae). The size of the left bar (orange) represents the abundance (number of individuals) of bat flies per observed species and the size of the right bar (green) represents the abundance of bats for which the sample was obtained. Bat fly. The width of the black lines/bars indicates the frequency of interactions.




Figure 9. Quantitative bipartite bat-fly network (Module 3 - Streblidae). The size of the left bar (orange) represents the abundance (number of individuals) of bat flies per observed species and the size of the right bar (green) represents the abundance of bats for which the sample was obtained. The width of the black lines/bars indicates the frequency of interactions.




Figure 10. Quantitative bipartite bat-fly network (Nycteribiidae). The size of the left bar (red) represents the abundance (number of individuals) of bat flies per observed species and the size of the right bar (blue) represents the abundance of bats for which the sample was obtained. The width of the black lines/bars indicates the frequency of interactions.

cords of unique species of the families Emballonuridae (*Saccopteryx bilineata*), Molossidae (*M. pretiosus*), and Noctilionidae (*N. albiventris*) (Fig. 10). The quantitative modularity QuanBiMo for the interaction network was also high but lower compared to the findings for Streblidae ($Q = 0.67$). The specialization

(H_2' = 0.88) was higher, indicating a high niche differentiation among the bats in the network. Furthermore, we observed low connectivity (C = 0.15); however, nestedness ($wNODF$ = 16.31) was higher compared to Streblidae. The degree of centrality (DC) was notable for *B. ferrisi* (DC = 9) and *Basilia ortizi* (DC = 4), which were found parasitizing multiple species of host bats. Particularly, *Basilia ortizi* showed strong interaction with the genus *Eptesicus*, while *Basilia ferrisi* exhibited strong interaction with *Myotis nigricans*.

Discussion

Our study highlights the high diversity of bat flies that inhabit the Orinoco Region in northern South America. In other neotropical countries such as Brazil, Paraguay, and Panama, several studies have found a directly proportional relationship between the richness of Streblidae flies and bats (Dick and Gettinger 2005; Graciolli and Bianconi 2007). Considering that Colombia and Venezuela, are among the countries with the greatest diversity of bats in the Neotropics (with 217 and 172 species, respectively; Delgado-Jaramillo et al. 2016; Ramírez-Chaves et al. 2021), the finding of 136 species of bat flies in the Orinoquia Region was expected and will increase with additional studies in poorly sampled areas the Andean foothills of the Orinoco basin were endemic bat species such as *Vampyressa voragine* inhabit (Morales-Martínez et al. 2021). Besides the elevated number of the species documented in the Orinoquia, there are still gaps of information about these flies confirming the need for greater sampling efforts, particularly in some states/departments and biomes such as Portuguesa, Cojedes and Delta Amaruco in Venezuela and Guaviare in Colombia. The richness of bat flies in Colombia and Venezuela is also higher and surpasses the number of species documented in countries like Brazil, which despite having the largest area in South America (48%), has 181 species of bats (Garbino et al. 2022) and > 100 species of Streblidae and 26 of Nycteribiidae (da Silva et al. 2023).

Although the high specialization of some species of Streblidae has been controversial and some researchers have previously described them as mostly host-nonspecific (Dittmar et al. 2006), the evaluation of the quality of the non-primary associations reported in the literature for Orinoquia showed that several of them (~ 10% of the records) were probably due to accidental transfers or contamination of the samples during collection. These results support more recent studies suggesting that most species are host-specific (de Vasconcelos et al. 2016; Estrada-Villegas et al. 2018), a change attributable to methodological improvements in sample collection and taxonomic updates of flies and hosts. In addition, we have found controversial non-primary associations that could not be classified as possible cases of contamination. Some of these associations occurred in congeneric species, as is the case of the reports of *M. minuta* (primary host: *L. silvicola*) on *T. maresi* in Casanare, Colombia (Liévano-Romero et al. 2019). Considering that some unusual interaction occurred in non-congeneric species of the primary hosts, explaining the reasons behind is complicated. This is the case of *N. albiventris* parasitized by *M. proxima* (primary host: *Sturnira* species), and *S. giannae* parasitized by *N. maai* (primary host: *N. albiventris*) (López Rivera et al. 2022). Furthermore, the nature of the specificity in some non-primary associations for certain fly species complexes could not be determined (Table 1). In this sense, we suggested that the determination of the true specificity of ectoparasitic

flies will only be achieved when: 1) large samples of hosts and parasites are available, 2) sampling protocols strictly control cross-contamination between hosts, and 3) association cases are evaluated statistically (Dick 2005).

Although most studies on the interaction between bats and ectoparasites have been descriptive, these are crucial for understanding host-parasite dynamics in different environments (Patterson et al. 2007; Fagundes et al. 2017; Salinas-Ramos et al. 2018). Previous studies focused on interactions networks only covered few localities in the Colombian Orinoquia (López Rivera et al. 2022) or were limited to enumerating species and abundances in specific locations in Colombia (departments of Arauca, Casanare, and Meta) and Venezuela (states of Anzoátegui, Apure, Aragua, Barinas, Carabobo, Guárico, Lara, Mérida, Monagas, Trujillo and Yaracuy) (Wenzel and Tipton 1966; Wenzel 1976; Marinkelle and Grose 1981; Dick et al. 2016; Liévano-Romero et al. 2019; López Rivera et al. 2022). Considering that, our study represents the first attempt to define the ecological interactions between ectoparasitic flies and their host bats in the Orinoquia Region of Colombia and Venezuela. However, previously reported modularity and specialization values for the interaction networks in the Department of Arauca, in Colombia ($Q = 0.61$ and 0.69 ; $H_2' = 0.78$ and 0.91) (López Rivera et al. 2022), were similar to the values found for Nycteribiidae in the network of the entire Orinoquia ($Q = 0.67$ and $H_2' = 0.88$), but lower compared to the values found in the Streblidae network of the entire Orinoquia ($Q = 0.85$ and $H_2' = 0.94$). These findings match suggestions in the literature where the size of the network can influence the modularity values, being higher for larger networks (Júnior et al. 2020). This can be explained by the highly specialized nature of the interaction between bats and bat flies (Dick and Patterson 2007), where most fly species are associated with a single host (monoxenic), and even species that parasitize multiple bat species (oligoxenic and polyxenic) are restricted to phylogenetically close hosts, leading to the formation of groups with a similar composition of ectoparasitic fly species (Falcão 2015; López Rivera et al. 2022).

Ectoparasites typically display some host specificity, implying that some ectoparasitic fly species are adapted to parasitize only one or a few bat species (Wenzel 1976; Dick and Patterson 2007). The 11 bat species that were parasitized by more than five bat fly species might have characteristics or behaviors that make them prone to being parasitized by multiple bat fly species. Several studies have shown the common presence of various bat fly species on the same bat (Wenzel 1976; Dick 2005; Hiller et al. 2018). According to Wenzel (1976), 63% of the infested Venezuelan bat species harbored two to four species of bat flies. Several species of Streblidae coexist on the same hosts, most of these associations are negatively correlated in abundance, but competition is not strong enough to lead to local extinction (Dick 2005). In fact, some of the highly connected bat host species within the interaction network for the Orinoquia were reported to be involved in some of the previously documented cases of positive co-occurrence between bat flies (Dick 2005), such as: *E. clovisi* and *Trichobius propinquus* on *Anoura geoffroyi* (Fig. 7), *Trichobius joblingi* and *Strebla guajiro* on *C. perspicillata* (Fig. 8), and *Trichobius lonycteridis* with *A. spurrelli* on *L. spurrelli* (Fig. 9).

In a lesser extent, positive correlations have also been reported in abundance, indicating mutualistic relationships (Dick 2005; Hiller et al. 2018; Alcantara et al. 2022). It is possible that the presence and greater abundance of one species of parasite facilitates the presence and abundance of the other species, reciprocally eliminating the pressure of grooming (one of the main

causes of mortality of ectoparasites) in each population (Marshall 1981). Our interaction networks for the Orinoquia also revealed positive co-occurrences previously reported for ectoparasitic flies from Venezuela (Dick 2005). For example, a relatively equivalent association in the size of links and nodes is observed between *T. parasiticus* and *Strebla wiedemanni*, both associated with *D. rotundus* (Fig. 8). Similarly, an equivalent relationship between *S. diaemi* and *T. diaemi* is presented in its primary host *D. youngii* (Fig. 9). Cooccurrences of Neotropical Streblids have been reported mainly between species belonging to different genera (Dick 2005; Hiller et al. 2018) that differ in their general morphology facilitating the coexistence of species by utilizing different regions on the host's body surface (Wenzel 1976; ter Hofstede et al. 2004; Dick 2005; Hiller et al. 2018). Our interaction networks reveal the separation of the three ecomorphological groups (wing crawler – WC, fur runner – FR, and fur swimmer – FS), associated in a single bat species (Dick 2005; Hiller et al. 2018). Some notable examples of these associations include: i) *Metalasmus pseudopterus* (FS) with *A. phyllostomatis* (WC) or with *M. aranea* (FR), and *Aspidoptera phyllostomatis* (WC) with *Megistopoda aranea* (FR) in *Artibeus planirostris*; ii) the association of *Speiseria ambigua* (FR) with *S. guajiro* (FW) or with *Trichobius joblingi* (WC) on *Carollia perspicillata*, and the association of *Strebla guajiro* (FS) with *Trichobius joblingi* (WC) on the same host; iii) the presence of *T. parasiticus* (WC) and *Strebla wiedemanni* (FS) on *D. rotundus*, and iv) the association of *Strebla hertigi* (FS) with *Trichobius costalimai* (WC) or with *Trichobioides perspicillatus* (WC) on *Phyllostomus discolor* (Fig. 8). Therefore, our results suggest ecological niche partitioning of ectoparasitic flies on bat hosts in the Orinoco region. However, due to the lack of a detailed phylogeny of bat flies, it is unclear whether these morphological differences reflect the evolutionary history of bat flies or represent convergent adaptations to host habitat type (Hiller et al. 2018).

In this study, new records are presented that describe the co-occurrence of *N. maai* and *P. parvuloides* in *N. albiventris* in the Department of Arauca, Colombia, an association previously reported by López Rivera et al. (2022) (Table 2). The positive interactions between *Noctiliostrebla* and *Paradyschiria* species in *Noctilio* spp. are well known and documented (Wenzel 1976; Guerrero 1995; Guerrero 1998; Moura et al. 2003; Schad et al. 2012). For example, in the case of *N. albiventris*, *N. maai* coexists with *Paradyschiria curvata* or *Paradyschiria parvula*, and in the case of *Noctilio leporinus*, each individual host is infested with *Noctiliostrebla aitkeni* and *Paradyschiria fusca* or *Noctiliostrebla traubi* and *Paradyschiria lineata* (Wenzel 1976; Dick 2005). In this sense, *Paradyschiria* almost always presents the highest values of prevalence and abundance, while *Noctiliostrebla* occurs more commonly in the presence of *Paradyschiria* (Moura et al. 2003; Presley 2007; Schad et al. 2012; Guerrero 2019). Our results agree with this pattern since we observed a higher abundance of *P. parvuloides* compared to *N. maai* in *N. albiventris* (Fig. 8, Table 2).

Paradyschiria species are fur runners, while *Noctiliostrebla* spp. are wing-crawlers, again supporting the niche partitioning of these species on *Noctilio* spp. The ecomorphological classification for *Paradyschiria* has been controversial, as they had previously been classified as wing-crawlers (Dick 2005) for not having longer hind legs, a morphological feature associated with the fur runner microhabitat (Ter Hofstede et al. 2004). However, *Paradyschiria* spp. have an extremely strong forefemur compared to other streblid species, which allows

them greater agility to move in the fur and hold on the host's hair (Wenzel et al. 1966; Wenzel 1976; Alcantara et al. 2022). *Noctilio* bats are considered to be among the shortest-haired of Neotropical and Nearctic bats (Dick and Miller 2010), so *Paradyschiria* flies may not need longer hind legs, otherwise a strong forefemur can be used to hold on to the host's hair tightly if it cannot avoid grooming with rapid movements (Alcantara et al. 2022).

Finally, the analysis of ecological networks has been fundamental in the understanding of complex biological systems, providing information on how species are organized and connected in a community (Bascompte 2009; Butts 2009; Thébault and Fontaine 2010). In this sense, the study of host-parasite specificity interactions is essential to understand the mechanisms behind parasitism and its relationship with the functioning of biodiversity (Frainer et al. 2018), since parasites play an important role in the regulation of populations of host species (Poulin et al. 2006). Our results indicated that highly connected bat species in the Orinoquia interaction network act as hosts for a diversity of ectoparasitic fly species belonging to different ecomorphological groups. These groups differ in how they interact and are located on the bat's body. Therefore, the presence of these bats is not only important for ectoparasitic fly species in general, but also for maintaining the coexistence and interaction of different ecomorphological groups of ectoparasitic flies in the network.

Conclusions

The Orinoco Region, located in northern South America, harbors a remarkable diversity of bat flies. This diversity is largely attributed to the rich bat fauna found in Colombia and Venezuela, two countries known for their high bat diversity in the Neotropical region. In the Orinoco Region, associations have been identified between bat fly species belonging to different ecomorphological groups and unique host species. This supports the idea of a potential niche partitioning among ectoparasitic bat flies on their bat hosts. However, due to the lack of a detailed phylogeny of bat flies, it is unclear whether the observed morphological differences are the result of evolutionary history or convergent adaptations to different host habitat types. In this study, we present new records of associations between bat flies and their bat hosts, thereby expanding our knowledge of these interactions in the Orinoco Region. Overall, this study contributes to our understanding of the diversity, specificity, and ecological interactions between bat flies and their host bats in this region. These findings underscore the need for further research and sampling efforts to fill knowledge gaps in this field.

Acknowledgments

We thank the Museo de Historia Natural de la Universidad de Caldas, for allowing the work in the facilities of the MHN-UCa collections. This project was funded by the Vicerrectoría de Investigaciones, Universidad de Caldas (code 0318322 and 0277620). We thank the Convocatoria de subvención del Fondo de Ciencia, Tecnología e Innovación del Sistema General de Regalías para componer una lista de proyectos elegibles para su viabilidad, priorización y aprobación por parte de la OCAD como parte del Programa de Becas de Excelencia Doctoral cohorte 1 – 2019.. We thank the Ecology Laboratory of the Universi-

dad de Caldas for lending us their space and stereomicroscope with camera. We thank Liping Yan, Gustavo Graciolli, and two anonymous reviewers for comments on an early version of this manuscript and for the editorial support.

Additional information

Conflict of interest

The authors have declared that no competing interests exist.

Ethical statement

No ethical statement was reported.

Funding

Museo de Historia Natural de la Universidad de Caldas. Vicerrectoría de Investigaciones, Universidad de Caldas (code 0318322 and 0277620). Grant call of the Science, Technology and Innovation Fund of the General Royalties System to compost a list of eligible projects for viability, prioritization, and approval by the OCAD as part of the Scholarship Program of Doctoral Excellence cohort 1 – 2019.

Author contributions

Conceptualization: EMOP. Data curation: EMOP, HERC. Formal analysis: EMOP. Funding acquisition: FARP. Investigation: EMOP, HERC. Methodology: EMOP. Project administration: FARP. Supervision: FARP, HERC. Writing – original draft: EMOP. Writing – review and editing: FARP.

Author ORCIDs

Erika M. Ospina-Pérez  <https://orcid.org/0000-0001-5784-6216>

Fredy A. Rivera-Páez  <https://orcid.org/0000-0001-8048-5818>

Héctor E. Ramírez-Chaves  <https://orcid.org/0000-0002-2454-9482>

Data availability

All of the data that support the findings of this study are available in the main text or Supplementary Information.

References

- Alcantara DMC, Graciolli G, Nihei SS (2019) Revision of *Noctiliostrebla* (Diptera: Streblidae), parasites of bulldog bats (Chiroptera: Noctilionidae: Noctilio). Zootaxa 4560(3): 483–521. <https://doi.org/10.11646/zootaxa.4560.3.4>
- Alcantara DMC, Graciolli G, Toma R, Souza CS (2022) Sex-biased parasitism, host mass and mutualistic bat flies: An antagonistic individual-based network of bat-bat fly interactions. International Journal for Parasitology 52(4): 217–224. <https://doi.org/10.1016/j.ijpara.2021.10.010>
- Almeida-Neto M, Ulrich W (2011) A straightforward computational approach for measuring nestedness using quantitative matrices. Environmental Modelling & Software 26(2): 173–178. <https://doi.org/10.1016/j.envsoft.2010.08.003>
- Ascuntar-Osnas O, Montoya-Bustamante S, González-Chávez B (2020) Registros de Streblidae (Diptera: Hippoboscoidea) en un fragmento de bosque seco tropical en Colombia. Biota Colombiana 21(1): 16–27. <https://doi.org/10.21068/c2020.v21n01a02>

- Autino A, Claps GL, Barquez RM (1999) Insectos ectoparásitos de murciélagos de las yugas de la Argentina. *Acta Zoológica Mexicana* 78(78): 119–169. <https://doi.org/10.21829/azm.1999.78781921>
- Bascompte J (2009) Disentangling the web of life. *Science* 325(5939): 416–419. <https://doi.org/10.1126/science.1170749>
- Blüthgen N, Menzel F, Blüthgen N (2006) Measuring specialization in species interaction networks. *BMC Ecology* 6(1): 9. <https://doi.org/10.1186/1472-6785-6-9>
- Butts CT (2008) Network: A package for managing relational data in R. *Journal of Statistical Software* 24(2): 1–36. <https://doi.org/10.18637/jss.v024.i02>
- Butts CT (2009) Revisiting the foundations of network analysis. *Science* 325(5939): 414–416. <https://doi.org/10.1126/science.1171022>
- Calonge-Camargo B, Pérez-Torres J (2018) Ectoparasites (Polyctenidae, Streblidae, Nycteribiidae) of bats (Mammalia: Chiroptera) from the Caribbean Region of Colombia. *Therya* 9(2): 171–178. <https://doi.org/10.12933/therya-18-492>
- Cañizales I, Guerrero R (2022) Parasitofauna en Murciélagos de los Llanos Orientales de Venezuela. *Boletín del Centro de Investigaciones Biológicas* 56(2): 198–224. <https://doi.org/10.5281/zenodo.7469694>
- Ching LM, Marshall AG (1968) The breeding biology of the bat-fly *Eucampsipoda sundaticum* Theodor, 1955 (Diptera: Nycteribiidae). *Malayan Nature Journal* 21: 171–180.
- da Silva ADS, Barbier E, Júnior WRT, da Silva LAM, de Oliveira JB (2023) Parasitism on bats by bat flies in remnants of a humid forest enclave area in Northeastern Brazil. *Veterinary Parasitology. Regional Studies and Reports* 37: 100826. <https://doi.org/10.1016/j.vprsr.2022.100826>
- de Vasconcelos PF, Falcão LAD, Graciolli G, Borges MAZ (2016) Parasite-host interactions of bat flies (Diptera: Hippoboscoidea) in Brazilian tropical dry forests. *Parasitology Research* 115(1): 367–377. <https://doi.org/10.1007/s00436-015-4757-8>
- Delgado-Jaramillo M, García FJ, Machado M (2016) Diversidad de murciélagos (Mammalia: Chiroptera) en las áreas de protección estricta de Venezuela. *Ecotrópicos* 29(1–2): 28–42.
- Dick C (2005) Ecology and host specificity of Neotropical bat flies (Diptera: Streblidae) and their chiropteran hosts. PhD Dissertation. Texas Tech University, Lubbock, TX, USA.
- Dick CW, Dittmar K (2014) Parasitic bat flies (Diptera: Streblidae and Nycteribiidae): host specificity and potential as vectors. In: Klipple S, Mehlhorn H (Eds) *Bats (Chiroptera) as vectors of diseases and parasites*. Springer, Heidelberg, 131–155. https://doi.org/10.1007/978-3-642-39333-4_6
- Dick CW, Gettigner D (2005) A faunal survey of Streblid flies (Diptera: Streblidae) associated with bats in Paraguay. *The Journal of Parasitology* 91(5): 1015–1024. <https://doi.org/10.1645/GE-536R.1>
- Dick CW, Miller JA (2010) STREBLIDAE (bat flies). In: Brown BV, Borkent A, Cumming JM, Wood DM, Zumbado M (Eds) *Manual of Central American Diptera*, NRC Research Press, Ottawa 2: 1249–1260.
- Dick CW, Patterson BD (2006) Bat flies-obligate ectoparasites of bats. In: Morand S, Krasnov BR, Poulin R (Eds) *Micromammals and macro-parasites: from evolutionary ecology to management*. Springer-Verlag, Tokyo, 179–194. https://doi.org/10.1007/978-4-431-36025-4_11
- Dick CW, Patterson BD (2007) Against all odds: Explaining high host specificity in dispersal-prone parasites. *International Journal for Parasitology* 37(8–9): 871–876. <https://doi.org/10.1016/j.ijpara.2007.02.004>

- Dick CW, Graciolli G, Guerrero R (2016) Family Streblidae. *Zootaxa* 4122(1): 784–802. <https://doi.org/10.11646/zootaxa.4122.1.67>
- Dittmar K, Porter ML, Murray S, Whiting MF (2006) Molecular phylogenetic analysis of nycteriibiid and streblid bat flies (Diptera: Brachycera, Calyptratae): implications for host associations and phylogeographic origins. *Molecular Phylogenetics and Evolution* 38(1): 155–170. <https://doi.org/10.1016/j.ympev.2005.06.008>
- Domínguez C (1998) La gran cuenca del Orinoco. In: Domínguez C (Ed.) Colombia Orinoco. Instituto de Estudios Orinocenses, 39–67.
- Dormann CF, Strauss R (2014) A method for detecting modules in quantitative bipartite networks. *Methods in Ecology and Evolution* 5(1): 90–98. <https://doi.org/10.1111/2041-210X.12139>
- Dormann CF, Gruber B, Fründ J (2008) Introducing the bipartite Package: Analysing ecological networks. *R News* 8: 8–11.
- Dormann CF, Fründ J, Blüthgen N, Gruber B (2009) Indices, graphs and null models: Analyzing bipartite ecological networks. *The Open Ecology Journal* 2(1): 7–24. <https://doi.org/10.2174/1874213000902010007>
- Durán AA, Álvarez García DM, Graciolli G (2017) Ectoparasitic flies (Diptera, Streblidae) on bats (Mammalia, Chiroptera) in a dry tropical forest in the northern Colombia. *Papéis Avulsos De Zoología* 57(8): 105–111. <https://doi.org/10.11606/0031-1049.2017.57.08>
- Durán AA, Saldaña-Vázquez RA, Graciolli G, Peinado LC (2018) Specialization and modularity of a bat fly antagonistic ecological network in a dry tropical forest in northern Colombia. *Acta Chiropterologica* 20(2): 503–510. <https://doi.org/10.3161/15081109ACC2018.20.2.020>
- Estrada-Villegas S, Halczok TK, Tschapka M, Page RA, Brändel SD, Hiller T (2018) Bats and their bat flies: Community composition and host specificity on a Pacific island archipelago. *Acta Chiropterologica* 20(1): 161–176. <https://doi.org/10.3161/15081109ACC2018.20.1.012>
- Fagundes R, Antonini Y, Aguiar LM (2017) Overlap in cave usage and period of activity as factors structuring the interactions between bats and ectoparasites. *Zoological Studies* (Taipei, Taiwan) 56: 1–11. <https://doi.org/10.6620/ZS.2017.56-22>
- Falcão LAD (2015) Morcegos em florestas tropicais secas brasileiras. PhD Thesis, Management of the Federal University of Minas Gerais, Belo Horizonte.
- Ferrer Pérez A, Beltrán M, Díaz-Pulido AP, Trujillo F, Mantilla-Meluk H, Herrera O, Payán E (2009) Lista de los mamíferos de la cuenca del río Orinoco. *Biota Colombiana* 10(1–2): 179–207. <https://doi.org/10.15472/5v3n0h>
- Fortuna MA, Stouffer DB, Olesen JM, Jordano P, Mouillot D, Krasnov BR, Poulin R, Bascompte J (2010) Nestedness versus modularity in ecological networks: Two sides of the same coin? *Journal of Animal Ecology* 79: 811–817. <https://doi.org/10.1111/j.1365-2656.2010.01688.x>
- Frairer A, McKie BG, Amundsen PA, Knudsen R, Lafferty KD (2018) Parasitism and the biodiversity-functioning relationship. *Trends in Ecology & Evolution* 33(4): 260–268. <https://doi.org/10.1016/j.tree.2018.01.011>
- Garbino GST, Gregorin R, Lima IP, Loureiro L, Moras L, Moratelli R, Nogueira MR, Pavan AC, Tavares VC, Nascimento MC, Novaes RLM, Peracchi AL (2022) Updated checklist of Brazilian bats: versão 2020. Comitê da Lista de Morcegos do Brasil—CLMB. Sociedade Brasileira para o Estudo de Quirópteros. <https://www.sbeq.net/lista-de-especies>
- Gardner LA (2008) Mammals of South America. Marsupials, xenarthrans, shrews, and bats. University of Chicago Press, Chicago 2, 669 pp. <https://doi.org/10.7208/chicago/9780226282428.001.0001>

- González AM, Dalsgaard B, Olesen JM (2010) Centrality measures and the importance of generalist species in pollination networks. *Ecological Complexity* 7(1): 36–43. <https://doi.org/10.1016/j.ecocom.2009.03.008>
- Graciolli G, Bianconi GV (2007) Moscas ectoparasitas (Diptera, Streblidae e Nycteribiidae) em morcegos (Mammalia, Chiroptera) em área de floresta com Araucária no estado do Paraná, sul do Brasil. *Revista Brasileira de Zoologia* 24(1): 246–249. <https://doi.org/10.1590/S0101-81752007000100033>
- Graciolli G, Dick CW (2018) Checklist of the world Nycteribiidae (Diptera: Hippoboscoidea). Report for the Natural Science Foundation (DBI-0545051, DEB-1003459): 1–9.
- Graciolli G, Autino AG, Claps GL (2007) Catalogue of American Nycteribiidae (Diptera, Hippoboscoidea). *Revista Brasileira de Entomologia* 51(2): 142–159. <https://doi.org/10.1590/S0085-56262007000200004>
- Graciolli G, Dick CW, Guerrero R (2016) Familia Nycteribiidae. *Zootaxa* 4122(1): 780–783. <https://doi.org/10.11646/zootaxa.4122.1.66>
- Guerrero R (1994) Catálogo de los Streblidae (Diptera: Pupipara) parásitos de murciélagos (Mammalia: Chiroptera) del Nuevo Mundo. II. Los grupos: *pallidus*, *caucus*, *major*, *uniformis*, y *longipes* del género *Trichobius* Gervais, 1844. *Acta Biologica Venezolica* 15: 1–18.
- Guerrero R (1995) Catálogo de los Streblidae (Diptera: Pupipara) parásitos de murciélagos (Mammalia: Chiroptera) del Nuevo Mundo. V. Trichobiinae con alas reducidas o ausentes y miscelaneos. *Boletín de Entomología Venezolana* 10: 135–160.
- Guerrero R (1998) Notes on Neotropical batflies (Diptera, Streblidae). I. The genus *Trichobius*, with description of two new species and one new subspecies from Venezuela. *Acta Parasitologica* 43: 86–93.
- Guerrero R (2019) Streblidae (Diptera: Pupipara) de Venezuela: Sistemática, Ecología y Evolución. Editorial Académica Española, Muritius.
- Guimarães LR (1972) Venezuelan nycteribiid batflies (Diptera: Nycteribiidae). Brigham Young University Science Bulletin, Biological Series, 17(1): 1. <https://scholarsarchive.byu.edu/byuscib/vol17/iss1/>
- Hagan HR (1951) Embryology of the viviparous insects. Ronald, New York, 472 pp.
- Herrera-Sepúlveda MT (2013) Comparación de la carga de ectoparásitos entre harenas y grupos mixtos de la población de *Carollia perspicillata* en la cueva Macaregua (Santander, Colombia). B.Sc. Thesis, Facultad de Ciencias, Pontificia Universidad Javeriana, Bogotá, Colombia, 45 pp.
- Hiller T, Honner B, Page RA, Tschapka M (2018) Leg structure explains host site preference in bat flies (Diptera: Streblidae) parasitizing neotropical bats (Chiroptera: Phyllostomidae). *Parasitology* 145(11): 1475–1482. <https://doi.org/10.1017/S0031182018000318>
- Hsieh TC, Ma KH, Chao A (2016) iNEXT: An R package for rarefaction and extrapolation of species diversity (Hill numbers). *Methods in Ecology and Evolution* 7(12): 1451–1456. <https://doi.org/10.1111/2041-210X.12613>
- Júnior LD, De Araújo WS, Falcão LA (2020) Structure of the interaction networks between bats (Mammalia: Chiroptera) and ectoparasite flies (Diptera: Streblidae, Nycteribiidae) on a latitudinal gradient. *Acta Chiropterologica* 22(1): 187–196. <https://doi.org/10.3161/15081109ACC2020.22.1.018>
- Lehane MJ (2005) Biology of Blood-Sucking Insects. Cambridge University Press, 254–256. <https://doi.org/10.1017/CBO9780511610493>
- León GS (2005) La cuenca del río Orinoco: Visión hidrográfica y balance hídrico. *Revista Geográfica Venezolana* 46(1): 75–108.

- Liévano-Romero KS, Rodríguez-Posada ME, Cortés-Vecino JA (2019) Nuevos registros de ectoparásitos de murciélagos en sabanas inundables de la Orinoquía colombiana. Mastozoología Neotropical 26(2): 377–389. <https://doi.org/10.31687/saremMN.19.26.2.0.13>
- López Rivera C, Florez Padilla JM, Méndez Urbano F, Ospina-Pérez EM, Velásquez-Guarín D, Mejía-Fontecha IY, Ossa-López PA, Rivera-Páez FA, Ramírez-Chaves HE (2022) Interaction networks between bats (Mammalia: Chiroptera) and ectoparasitic flies (Diptera: Hippoboscoidea): a specificity relationship in the Colombian Orinoquia Region. Acta Chiropterologica 24(2): 379–394. <https://doi.org/10.3161/15081109ACC2022.24.2.008>
- Mammal Diversity Database (2022) Mammal Diversity Database (Version 1.10). Zenodo. <https://doi.org/10.5281/zenodo.7394529>
- Margolis L, Esch GW, Holmes JC, Kuris AM, Schad G (1982) The use of ecological terms in parasitology (report of an ad hoc committee of the American Society of Parasitologists). The Journal of Parasitology 68(1): 131–133. <https://doi.org/10.2307/3281335>
- Marinkelle CJ, Grose ES (1981) A list of ectoparasites of Colombian bats. Revista de Biología Tropical 29(1): 11–20.
- Marshall AG (1981) The ecology of ectoparasitic insects. Academic Press, London, 459 pp.
- Mello MAR, Muylaert R, Pereira R, Felix R (2016) Guia para análise de redes ecológicas. Belo Horizonte, 111 pp.
- Morales-Martínez DM, Rodríguez-Posada ME, Ramírez-Chaves HE (2021) A new cryptic species of the yellow-eared bat *Vampyressa melissa* species complex (Chiroptera: Phyllostomidae) from Colombia. Journal of Mammalogy 102(1): 90–100. <https://doi.org/10.1093/jmammal/gyaa137>
- Mosquera Guerra F, Trujillo F, Aya Cuero C, Bolívar L, Valencia K, Vargas Arboleda AF, Mantilla Meluk H (2019) Mamíferos. In: Trujillo F, Anzola F (Eds) Biodiversidad en el departamento de Arauca. Gobernación de Arauca, Fundación Omacha y Fundación Ecollano, Bogotá, 235–267.
- Moura MO, Bordignon MO, Graciolli G (2003) Host characteristics do not affect community structure of ectoparasites on the fishing bat *Noctilio leporinus* (L., 1758) (Mammalia: Chiroptera). Memorias do Instituto Oswaldo Cruz 98(6): 811–815. <https://doi.org/10.1590/S0074-02762003000600017>
- Nagorsen DW, Peterson RL (1980) Mammal collectors' manual: a guide for collecting, documenting, and preparing mammal specimens for scientific research. Life Sciences Miscellaneous Publications, Royal Ontario Museum, Toronto, Ontario. <https://doi.org/10.5962/bhl.title.60776>
- Oksanen J, Blanchet FG, Kindt R, Legendre P, Minchin PR, O'hara RB, Oksanen MJ (2020) Package 'vegan'. Community ecology package 2(9): 1–295. <https://cran.rsm.ac.jp/web/packages/vegan/vegan.pdf>
- Pardo A, Rangel-CH JO (2014) Mamíferos de la Orinoquia. In: Rangel-CH JO (Ed.) Colombia diversidad biótica XIV. Instituto de Ciencias Naturales, Facultad de Ciencias, Universidad Nacional de Colombia, Bogotá, 751–784.
- Pastrana-Montiel MR, Ballesteros-Corra J, Chacón-Pacheco J (2019) First record of the parasite bat fly *Basilia mimoni* Theodor & Peterson, 1964 (Diptera: Nycteribiidae) in Colombia. Oecologia Australis 23(3): 685–689. <https://doi.org/10.4257/oeco.2019.2303.27>
- Patterson DB, Dick WC, Dittmar K (2007) Roosting habits of bats affect their parasitism by bat flies (Diptera: Streblidae). Journal of Tropical Ecology 23(2): 177–189. <https://doi.org/10.1017/S0266467406003816>
- Patterson BD, Dick CW, Dittmar K (2009) Nested distribution of bat flies (Diptera: Streblidae) on Neotropical bats: artifact and specificity in host parasite studies. Ecography 32(3): 48–487. <https://doi.org/10.1111/j.1600-0587.2008.05727.x>

- Petersen FT, Meier R, Kutty SN, Wiegmann BM (2007) The phylogeny and evolution of host choice in the Hippoboscoidea (Diptera) as reconstructed using four molecular markers. *Molecular Phylogenetics and Evolution* 45(1): 111–122. <https://doi.org/10.1016/j.ympev.2007.04.023>
- Porter ML, Cronin TW, Dick CW, Simon N, Dittmar K (2021) Visual system characterization of the obligate bat ectoparasite *Trichobius frequens* (Diptera: Streblidae). *Arthropod Structure & Development* 60: 101007. <https://doi.org/10.1016/j.asd.2020.101007>
- Poulin R, Krasnov BR, Morand S (2006) Patterns of host specificity in parasites exploiting small mammals. In: Morand S, Krasnov BR, Poulin R (Eds) *Micromammals and macroparasites*. Springer, Tokyo, 233–256. https://doi.org/10.1007/978-4-431-36025-4_13
- Presley SJ (2007) Streblid bat fly assemblage structure on Paraguayan *Noctilio leporinus* (Chiroptera: Noctilionidae): nestedness and species co-occurrence. *Journal of Tropical Ecology* 23(4): 409–417. <https://doi.org/10.1017/S0266467407004245>
- Raigosa Álvarez JR, Osorio CG, Autino AG, Dias LG (2020) First records of ectoparasitic insects (Diptera: Hippoboscoidea) of bats in the Department of Caldas, Colombia. *Papéis Avulsos de Zoologia* 60: e20206018. <https://doi.org/10.11606/1807-0205/2020.60.18>
- R Core Team (2022) R: A Language and Environment for Statistical Computing. R Foundation for Statistical Computing, Vienna. <https://www.R-project.org/>
- Ramírez-Chaves HE, Morales-Martínez DM, Rodríguez-Posada ME, Suárez-Castro AF (2021) Checklist of the mammals (Mammalia) of Colombia: Taxonomic changes in a highly diverse country. *Mammalogy Notes* 7(2): 253. <https://doi.org/10.47603/mano.v7n2.253>
- Salinas-Ramos VB, Zaldívar-Riverón A, Rebollo-Hernández A, Herrera-M LG (2018) Seasonal variation of bat-flies (Diptera: Streblidae) in four bat species from a tropical dry forest. *Mammalia* 82(2): 133–143. <https://doi.org/10.1515/mammalia-2016-0176>
- Schad J, Dechmann DK, Voigt CC, Sommer S (2012) Evidence for the ‘good genes’ model: Association of MHC class II DRB alleles with ectoparasitism and reproductive state in the neotropical lesser bulldog bat, *Noctilio albiventris*. *PLoS ONE* 7(5): e37101. <https://doi.org/10.1371/journal.pone.0037101>
- Soares FAM, Graciolli G, Alcântara DMC, Ribeiro CEB, Valença GC, Ferrari SF (2013) Bat flies (Diptera: Streblidae) ectoparasites of bats at an Atlantic Rainforest site in northeastern Brazil. *Biota Neotropica* 13(2): 242–246. <https://doi.org/10.1590/S1676-06032013000200024>
- Tamsitt JR, Fox I (1970) Records of bat ectoparasites from the Caribbean Region (Siphonaptera, Acarina, Diptera). *Canadian Journal of Zoology* 48(5): 1093–1097. <https://doi.org/10.1139/z70-193>
- Tarquino-Carbonell A (2014) Ectoparásitos asociados a la quirópterofauna en la vereda chorrillo municipio de Ambalema, Tolima, Colombia. Thesis. Programa de Biología, Universidad del Tolima, Tolima, Colombia, 123 pp.
- Tarquino-Carbonell A, Gutiérrez-Díaz KA, Galindo-Espinosa EY, Reinoso-Flórez G, Solari S, Guerrero R (2015) Ectoparasites associated with bats in northeastern Tolima, Colombia. *Mastozoología Neotropical* 22: 349–358.
- Tello SJ, Stevens RD, Dick CW (2008) Patterns of species co-occurrence and density compensation: A test for interspecific competition in bat ectoparasite infracommunities. *Oikos* 117(5): 693–702. <https://doi.org/10.1111/j.0030-1299.2008.16212.x>

- Ter Hofstede HM, Fenton MB, Whitaker Jr JO (2004) Host and host-site specificity of bat flies (Diptera: Streblidae and Nycteribiidae) on Neotropical bats (Chiroptera). Canadian Journal of Zoology 82(4): 616–626. <https://doi.org/10.1139/z04-030>
- Thébaud E, Fontaine C (2010) Stability of ecological communities and the architecture of mutualistic and trophic networks. Science 329(5993): 853–856. <https://doi.org/10.1126/science.1188321>
- Wenzel RL (1976) The streblid batflies of Venezuela (Diptera: Streblidae). Brigham Young University Science Bulletin. Biological Series 20 (4): 1–175. <https://doi.org/10.5962/bhl.part.5666>
- Wenzel RL, Tipton VJ (1966) Ectoparasites of Panama. Chicago Field Museum of Natural History, 861 pp. <https://doi.org/10.5962/bhl.title.2633>
- Wenzel RL, Tipton VJ, Kiewlicz A (1966). The Streblid bat flies of Panama (Diptera: Calypterae: Streblidae). In: Wenzel RL, Tipton VJ (Eds) Ectoparasites of Panama. Chicago Field Museum of Natural History, 405–675. <https://doi.org/10.5962/bhl.title.2633>

Supplementary material 1

Supplementary tables

Authors: Erika M. Ospina-Pérez, Fredy A. Rivera-Páez, Héctor E. Ramírez-Chaves

Data type: docx

Explanation note: **table S1:** Information and coordinates of the 63 sampling locations in the Orinoquia Region in South America shown on the map in Fig. 1; **table S2:** Localities of the records of flies in bats in Orinoquia from 1911 to 2023; **table S3:** Centrality metric of the species of ectoparasitic flies; **table S4:** Centrality metrics of the species of bats; **table S5:** Metrics of the interaction networks for the study areas, the first landscape: Herbaceous plains and chaparral – HPC and Orinoco-Amazonian Forests – OAF in the Colombian and Venezuelan Orinoquia Region; **table S6:** Centrality metric of the species of ectoparasitic flies, the first landscape: Herbaceous plains and chaparral – HPC and Orinoco-Amazonian Forests – OAF in the Colombian and Venezuelan Orinoquia Region; **table S7:** Centrality metric of the species of bats, the first landscape: Herbaceous plains and chaparral – HPC and Orinoco-Amazonian Forests – OAF in the Colombian and Venezuelan Orinoquia Region.

Copyright notice: This dataset is made available under the Open Database License (<http://opendatacommons.org/licenses/odbl/1.0/>). The Open Database License (ODbL) is a license agreement intended to allow users to freely share, modify, and use this Dataset while maintaining this same freedom for others, provided that the original source and author(s) are credited.

Link: <https://doi.org/10.3897/zookeys.1179.103479.suppl1>

Supplementary material 2

Bipartite bat-fly quantitative network of the Herbaceous plains and chaparral – HPC

Authors: Erika M. Ospina-Pérez, Fredy A. Rivera-Páez, Héctor E. Ramírez-Chaves

Data type: pdf

Explanation note: **fig. S1:** Bipartite bat-fly quantitative network of the Herbaceous plains and chaparral – HPC. The size of the bar on the left (red) represents the abundance (number of individuals) of bat flies per species observed and the size of the bar on the right (blue) represents the abundance of bats for which the bat fly sample was obtained. The width of the black lines/bars indicates the frequency of interactions.

Copyright notice: This dataset is made available under the Open Database License (<http://opendatacommons.org/licenses/odbl/1.0/>). The Open Database License (ODbL) is a license agreement intended to allow users to freely share, modify, and use this Dataset while maintaining this same freedom for others, provided that the original source and author(s) are credited.

Link: <https://doi.org/10.3897/zookeys.1179.103479.suppl2>

Supplementary material 3

Bipartite bat-fly quantitative network of the Orinoco-Amazonian Forests – OAF

Authors: Erika M. Ospina-Pérez, Fredy A. Rivera-Páez, Héctor E. Ramírez-Chaves

Data type: pdf

Explanation note: **fig. S2:** Bipartite bat-fly quantitative network of the Orinoco-Amazonian Forests – OAF. The size of the bar on the left (red) represents the abundance (number of individuals) of bat flies per species observed and the size of the bar on the right (blue) represents the abundance of bats for which the bat fly sample was obtained. The width of the black lines/bars indicates the frequency of interactions.

Copyright notice: This dataset is made available under the Open Database License (<http://opendatacommons.org/licenses/odbl/1.0/>). The Open Database License (ODbL) is a license agreement intended to allow users to freely share, modify, and use this Dataset while maintaining this same freedom for others, provided that the original source and author(s) are credited.

Link: <https://doi.org/10.3897/zookeys.1179.103479.suppl3>