

Two new Oriental species of *Eumorphus* Weber (Coleoptera, Endomychidae)

Ling-Xiao Chang¹, Guo-Dong Ren¹

¹ College of Life Sciences, Hebei University, Baoding 071002, China

Corresponding author: Guo-Dong Ren (gdren@hbu.edu.cn)

Academic editor: M. Thomas | Received 4 September 2016 | Accepted 9 May 2017 | Published 26 May 2017

<http://zoobank.org/A1AB0C19-A751-4372-90B0-E51E79C2C913>

Citation: Chang L-X, Ren G-D (2017) Two new Oriental species of *Eumorphus* Weber (Coleoptera, Endomychidae). ZooKeys 677: 1–9. <https://doi.org/10.3897/zookeys.677.10399>

Abstract

Two new species of *Eumorphus* from Asia, *E. falcifasciatus* **sp. n.** and *E. qiujianyuei* **sp. n.** are described and illustrated.

Keywords

Coccinelloidea, Coleoptera, Lycoperdininae, new species, Oriental Region, taxonomy

Introduction

The genus *Eumorphus* was established by Weber (1801) with *Eumorphus sumatrae* Weber, 1801 (= *Erotylus quadriguttatus* Illiger, 1800) as the type species. This genus is classified in the largest subfamily of Endomychidae, Lycoperdininae, the monophyly of which was tested and confirmed by the phylogenetic studies of Tomaszewska (2000, 2005). Robertson et al. (2015) presented a large-scale phylogenetic study for the Cucujoidea, using molecular evidence to rebuild the relationship tree of this superfamily and established one new superfamily, Coccinelloidea, with Endomychidae placed within it. This study further confirmed the monophyly of the subfamily Lycoperdininae and established its sister relationship with the subfamily Epipocinae (Robertson et al. 2015).

Tomaszewska (2005) recognized five generic groups among 38 genera of Lycoperdininae known at that time. The 23 genera known then from the Oriental Region have been classified in four of five generic groups (*Lycoperdina*-, *Amphix*-, *Amphisternus*- and

Eumorphus-groups). Since then two new genera of Lycoperdininae have been described from the Oriental Region: *Stroheckeria* Tomaszewska, 2006 and *Humerus* Chang & Ren, 2013. Both, indicated by the authors as belonging to *Amphisternus*-group (Tomaszewska 2006, Chang and Ren 2013), sister group of *Eumorphus*-group which includes the genus *Eumorphus* Tomaszewska (2005).

The *Eumorphus*-group includes 14 genera, five of which are distributed in the Oriental Region: *Avencymon* Strohecker, 1971, *Encymon* Gerstaecker, 1857, *Eumorphus* Weber, 1801, *Platindalmus* Strohecker, 1979 and *Parindalmus* Achard, 1922.

Strohecker (1968) listed 73 species (including subspecies) in his synopsis of the genus *Eumorphus*, of which *E. convexus*, *E. cryptus*, *E. elegans*, *E. eurynotus*, *E. leptocerus*, *E. micans*, and *E. parvus* were described as new species, and *E. austerus indianus*, *E. bipunctatus crucifer*, *E. bipunctatus mirus*, and *E. murrayi carinensis* were introduced as new subspecies. In addition, eleven nominal species were reduced to subspecies: *E. assamensis subsinuatus*, *E. bulbosus arrowi*, *E. coloratus vitalisi*, *E. cyanescens thomsoni*, *E. dilatatus turritus*, *E. eburatus guerini*, *E. fryanus festivus*, *E. fryanus quadripustulatus*, *E. quadriguttatus andamanensis*, *E. quadriguttatus convexcicollis*, and *E. sybarita consobrinus*. Subsequently two species were removed from *Eumorphus* and transferred to other genera, *E. calcaratus* Arrow, 1920 to *Platindalmus* (Strohecker 1979) and *E. nanus* Arrow, 1920 to *Indalmus* (Strohecker 1971); and one species was been transferred into *Eumorphus*: *Engonius bicoloripedoides* (Mader, 1955) by Strohecker (1968).

In 2007, Ren and Wang described two new species of *Eumorphus*, *E. dentatus* and *E. letilimarginatus* from China. *Eumorphus* is the largest genus of the subfamily Lycoperdininae and prior to the present study, this genus included 76 species (including subspecies) (Shockley et al. 2009).

During the examination of Endomychidae collected in China and Borneo, two new species were recognized and are described here.

Materials and methods

Type specimens of the new species described here are deposited in the following institutions or private collections:

MHBU Museum of Heibei University, Baoding, China

CCLX Collection of Lingxiao Chang, Beijing, China

The specimens were examined and described using a Nikon[®] SMZ800 dissecting microscope. The following measurements were made using a Leica[®] M205 A dissecting microscope: body length from apical margin of clypeus to apex of elytra; width across both elytra (at widest part); elytral length along suture, including scutellum. The abdomen was boiled in 10% NaOH solution, cleaned, and finally aedeagus was dissected in distilled water. Habitus photos were taken using a Canon[®] Eos 5D III SLR camera and Canon[®] MP-E 65mm macro lens. All photographs were modified in Adobe Photoshop[®] CC 2015.

Taxonomy

Eumorphus Weber, 1801

Eumorphus Weber, 1801: 31.

Type species. *Erotylus quadriguttatus* Illiger, 1800.

Eumorphoides Guérin-Méneville, 1858: 12.

Type species. *Eumorphus tetraspilotus* Hope, 1832.

Enaisimus Guérin-Méneville, 1858: 16.

Type species. *Eumorphus quadrinotatus* Gerstaecker, 1857.

Haplomorphus Guérin-Méneville, 1858: 18.

Type species. *Eumorphus bipunctatus* Perty, 1831.

Heterandrus Guérin-Méneville, 1858: 26.

Type species. *Eumorphus confusus* Guérin-Méneville, 1857.

Diagnosis. The species of *Eumorphus* are most similar to those of *Platindalmus* and *Gerstaeckerus*. However, *Eumorphus* can be distinguished from these other genera by the following combination of characters: 1) lateral margin of pronotum with a tendency to form irregularly broken lines, inconstant and often asymmetrical; 2) mandible is narrowly chisel-shaped at its apex; 3) elytra with basal margin simple; 4) intercoxal process of mesoventrite with lateral margins subparallel; 5) male femora lacking fringes of long hairs on inner edges (after Tomaszewska 2005).

Eumorphus falcifasciatus sp. n.

<http://zoobank.org/1F8715EC-E053-4819-829C-54B9A59E769C>

Figs 1, 3

Type material. Holotype, male, Borneo, Sabah, Keningau district, Jungle Girl Camp, 1215 m, 2016-IV-26, Chang L.X. leg (CCLX).

Diagnosis. *Eumorphus falcifasciatus* is a very unique species by its colouration, differing from all others in having the anterior elytral maculae falciform and posterior maculae dentate.

Description. Length 16.1 mm. Body pyriform, approximately 1.9 times as long as wide; moderately convex; subopaque. Colour black brown with two yellow maculae on elytra.

Head. Antenna composed of 11 antennomeres, long, rather stout, nearly 1/2 body length, with antennomeres 1 and 3–8 distinctly longer than wide; scape approximately 5.5 times as long as pedicel; pedicel very short, nearly as long as wide; antennomere 3 longer than 4–5 combined; antennomere 4 as long as 5; antennomeres 5 slightly longer than 6; antennomeres 6–8 subequal in length; club composed of three antennomeres, very broad, approximately 4.0 times as wide as antennomere 8, moderately flat and compact.




Figure 1. Dorsal and ventral habitus of *E. falcifasciatus* sp. n. male. **a** dorsal view **b** ventral view. Scale bar 1 mm.

Thorax. Pronotum 3.5 mm long, 5.7 mm wide; widest at base; finely, rather densely punctate; lateral and anterior margins narrowly bordered; anterior edge with small stridulatory membrane; sides undulate, abruptly widened basally from $1/4$ length; anterior angles produced, rather acute; posterior angles strongly, acutely produced backwards, distinctly curved basally and overlapping most of humeri; disc weakly convex, surface uneven with one large transverse oval raised area posteromedially and two small round raised areas anterolaterally; median furrow absent; lateral sulci shallow, linear, extending to $1/2$ pronotal length; basal sulcus weakly sinuate, moderately deep. Prosternal process moderately widely separating the procoxae; gradually widening to apical $1/4$, thence abruptly converging towards apex. Mesoventral process nearly quadrate, disc weakly convex, sides subparallel. Elytra 11.8 mm long, 8.3 mm wide; 1.4 times as long as pronotum; 1.5 times as wide as pronotum, sides curved, widest near behind $1/2$ length of elytron; lateral flattened margins abruptly widening from basal $1/6$ to apex, nearly $1/5$ of elytral width; sides distinctly converging from apical $1/3$ towards apex; finely, densely punctate; humeri weakly prominent. Each elytron with two large irregular maculae. Anterior elytral macula falciform, occupies about $4/5$ of elytral width and $2/3$ of elytral length, outer sides touching elytral lateral margin, inner margin of macula placed closely to elytral suture. Posterior macula crown-shaped, located at apical $1/3$, its anterior margin

tridentate, posterior margin widely emarginate medially. Protibiae slender basally, abruptly widening from basal 1/4 to apex; outer edge strongly sinuate; dorsal edge with S-shaped longitudinal ridge; inner edge with large, sharp tooth near 1/2 length; mesotibiae weakly curved from about 1/3 length to apex; metatibiae simple throughout its length, acutely produced apically.

Abdomen with five ventrites. Ventrite 1 almost as long as three following ventrites combined; ventrites 2-4 subequal in length. Ventrite 5 with lateral margins strongly converging posteriorly, posterior margin deeply, narrowly emarginate medially. Aedeagus (Fig. 3) long, heavily sclerotized, straight. Median lobe hook-shaped at apex, and branched latero-apically; branch rather long and strongly reflexed upwardly. Tegmen basal, comparatively large, ring-shaped.

Etymology. The name refers to the anterior elytral macula falciform.

***Eumorphus qiujiangyuei* sp. n.**

<http://zoobank.org/DDECE70A-4C04-4C30-BC67-5758A2CF8899>

Figs 2, 4

Type material. Holotype, male, Hainan, Wuzhishan, 21.V.2014, Jian-Yue Qiu leg. (MHBU).

Diagnosis. *Eumorphus qiujiangyuei* is similar to *Eumorphus austerus austerus* in appearance, but can be differentiated based on the following combination of characters: posterior angles of pronotum strongly and acutely produced, with tips curved inwardly (in *E. austerus austerus* posterior angles of pronotum weakly produced); sides of pronotum undulate (in *E. austerus austerus* rather smooth); and mesotibiae gently curved distally from near 1/2 length (in *E. austerus austerus* abruptly and strongly curved distally from near 1/2 length).

Description. Length 12.3 mm. Body broadly oval, approximately 1.8 times as long as wide; moderately convex; shiny. Colour brown with four yellow maculae on elytra. Antenna with scape red. Femora at apical 1/2 or 1/3 red.

Head. Antenna composed of 11 antennomeres, long, rather slender, nearly 1/2 body length, with antennomeres 3-8 distinctly longer than wide; scape approximately 4.5 times as long as pedicel; pedicel short, subquadrate; antennomere 3 distinctly longer than 4-5 combined; antennomere 4 slightly longer than 5, antennomeres 5-8 subequal in length; club composed of three antennomeres, moderately broad, flat. Maxilla with terminal palpomere prolonged, nearly 2.0 times as long as palpomere 3, cylindrical, weakly curved distally.

Thorax. Pronotum 2.4 mm long, 4.9 mm wide; widest at base; finely, rather densely punctate; lateral and anterior margins narrowly bordered; anterior edge with small stridulatory membrane; sides undulate, distinctly converging from apical 1/3 to apex, abruptly widened basally from 1/5 length; anterior angles distinctly produced, rather acute; posterior angles strongly, acutely produced, with tips curved inwardly; disc weakly convex; median furrow absent; lateral sulci linear, deep, ex-




Figure 2. Dorsal and ventral habitus of *E. qiujianyuei* sp. n. male. **a** dorsal view **b** ventral view. Scale bar 1 mm.

tending to basal $\frac{1}{4}$ length; basal sulcus nearly straight, deep. Prosternal process moderately widely separating procoxae; subparallel, weakly widening before apex then abruptly converging apically. Mesoventral process transverse rectangle, parallel sided, flat; posterior margin nearly straight. Elytra 8.9 mm long, 6.7 mm wide; 1.3 times as long as pronotum; 1.4 times as wide as pronotum, sides curved, widest near $\frac{1}{2}$ length of elytron; lateral margins moderately widely flattened, nearly $\frac{1}{5}$ of elytral width; distinctly converging from apical $\frac{1}{3}$ to apex; finely, densely punctate; humeri weakly prominent. Each elytron with two small round spots. Anterior elytral spot occupies about $\frac{1}{4}$ of elytral width, located posterior to humerus. Posterior spot of the same size as anterior one, located at apical $\frac{1}{4}$. Protibiae in male with one large, sharp tooth near $\frac{1}{2}$ length at inner edge, strongly expanded basally; mesotibiae distinctly curved distally from near $\frac{1}{2}$ length; metatibiae simple, acutely produced apically.

Abdomen with five ventrites. Ventrite 5 with lateral margins strongly converging posteriorly, posterior margin deeply, narrowly emarginate medially. Aedeagus (Fig. 4) rather long, heavily sclerotized, weakly curved basally, abruptly widened from basal $\frac{1}{3}$ to apex. Median lobe branched apically; the long branch abruptly raised at basal $\frac{1}{3}$, strongly reflexed apically. Tegmen basal, comparatively large, ring-shaped.



Figures 3–4. Aedeagi. **3** *E. falcifasciatus* sp. n. **4** *E. qiujianyuei* sp. n. Abbreviations: **a** lateral view **b** apical view. Scale bars 1 mm.

Etymology. This new species is dedicated to Ms. Jian-Yue Qiu, an insect researcher from Chongqing, who has been working on classification of insects for many years, collecting and providing many specimens of Endomychidae used in our studies.

Acknowledgements

We are indebted to Dr. W. Tomaszewska (Museum and Institute of Zoology, Warsaw, Poland) and Dr. F.W. Shockley (Smithsonian Institution, Washington, DC, USA) for providing useful comments which improved this manuscript. We thank Mr. Wen-Xuan Bi (Shanghai, China) very much for he discussed the genus *Eumorphus* with us, and make his viewpoint about the some suspected new species if belong to this genus.

References

- Achard J (1922) Description de nouveaux endomychides (Col. Endomychidae). Fragments Entomologiques (Prague: Smichov), 1922: 28–30.

- Arrow GJ (1920a) A contribution to the classification of the Coleopterous family Endomychidae. Transactions of the Entomological Society of London 1920: 1–83.
- Chang L-X, Ren G-D (2013) *Humerus medogicus*, new genus and new species from Xizang, China (Coleoptera: Endomychidae). Annales Zoologici 63: 95–100. <https://doi.org/10.3161/00-0345413X666138>
- Gerstaecker A (1857) Versuch einer systematischen Auseinandersetzung der Gattungen *Eumorphus* Web. und *Endomychus*. Payk. Archiv für Naturgeschichte 23: 211–243.
- Guérin-Ménéville FE (1857c) Matériaux pour une Monographie des Coléoptères du groupe des Eumorphides, et plus. spécialement du genre *Eumorphus*. Archives Entomologiques, 1: 237–280 + pl. 13.
- Guérin-Ménéville FE (1858) Matériaux pour une Monographie des Coléoptères du groupe des Eumorphides, et plus. spécialement du genre *Eumorphus*. Revue et Magasin de Zoologie Pure et Appliquée (2)10: 10–29.
- Hope FW (1832) [Untitled. *Eumorphus tetraspilotus*]. In: Griffith E (Ed) The animal kingdom arranged in conformity with its organization, by the Baron Cuvier, with supplementary additions to each order by Edward Griffith FLS, AS, and others. Volume the Fifteenth [= Class Insecta, Volume 2]. Whittaker, Treacher and Co., London, 796 pp.
- Illiger JKW (1800) Vierzig neue Insekten aus der Helwigischen Sammlung in Braunschweig. Beschriben von Karl Illiger im Anfänge des Februars 1800. Mit einigen Abbildungen. In: Wiedemann CPW (Ed.) Archiv für Zoologie und Zootomie. 1(2). Reichard, Braunschweig, 103–150.
- Mader L (1955) Neue Coleopteren aus Fukien (China). Helotidae, Languriidae, Erotylidae, Endomychidae, Coccinellidae. Koleopterologische Rundschau 33: 62–78.
- Perty JAM (1831) Observationes nomullae in Coleoptera Indiae orientalis. Dissertatio philosophico-entomologica, quam unacum praemissis thesibus auctoritate et consensu illustris philosophorum ordinis in Academia Ludovico-Maximilianeae facultatem legendi rite adepturus. M. Lindauer, Monachii, 44 pp.
- Ren G-D, Wang J-L (2007) Two new species of the genus *Eumorphus* Weber from China (Coleoptera, Endomychidae). Acta Zootaxonomica Sinica 32(3): 700–704.
- Robertson JA, Slipinski A, Moulton M, Shockley FW, Giorgi A, Lord NP, McKenna DD, Tomaszewska W, Forrester J, Miller KB, Whiting MF, McHugh JV (2015) Phylogeny and classification of Cucujoidea and the recognition of a new superfamily Coccinelloidea (Coleoptera: Cucujiformia). Systematic Entomology 40: 745–778. <https://doi.org/10.1111/syen.12138>
- Shockley FW, Tomaszewska KW, McHugh JV (2009) An annotated checklist of the handsome fungus beetles of the world (Coleoptera: Cucujoidea: Endomychidae). Zootaxa 1999: 1–113.
- Strohecker HF (1968) A synopsis of the genus *Eumorphus* (Coleoptera: Endomychidae). Pacific Insect 10(1): 79–112.
- Strohecker HF (1971a) The genera *Engonius* and *Parindalmus* (Coleoptera: Endomychidae). Pacific Insects, 13: 13–25.
- Strohecker HF (1971b) Synopsis of genus *Encymon* and *Avencymon* new genus (Coleoptera: Endomychidae). Pacific Insects 13: 27–39.

- Strohecker HF (1979) The genus *Indalmus* in Asia, New Guinea and Australia, with description of a new genus, *Platindalmus* (Coleoptera: Endomychidae). Pacific Insects 20: 279–292.
- Tomaszewska KW (2000) Morphology, phylogeny and classification of adult Endomychidae (Coleoptera: Cucujoidea). Annales Zoologici 50(4): 449–558.
- Tomaszewska KW (2005) Phylogeny and generic classification of the subfamily Lycoperdinae with a reanalysis of the family Endomychidae (Coleoptera: Cucujoidea). Annales Zoologici 55 (suppl. 1): 1–172.
- Tomaszewska KW (2006) *Stroheckeria quadrimaculata*, new genus and new species of Lycoperdinae from Vietnam (Coleoptera: Endomychidae). Annales Zoologici 56: 465–470.
- Weber F (1801) Observationes Entomologicae, continens novorum quae condidit generum characters, et nuper detectarum specierum descriptions. Bibliopolii Academici Novi, Kiliae, 116 pp.