

A survey of pholcid spiders (Araneae, Pholcidae) from Guiyang, Guizhou Province, China

Lan Yang¹, Fangyu Zhao¹, Qiaoqiao He^{1,2,3}, Zhiyuan Yao^{1,2,3}

¹ College of Life Science, Shenyang Normal University, Shenyang 110034, Liaoning, China

² Liaoning Key Laboratory of Evolution and Biodiversity, Shenyang 110034, Liaoning, China

³ Liaoning Key Laboratory for Biological Evolution and Agricultural Ecology, Shenyang 110034, Liaoning, China

Corresponding authors: Qiaoqiao He (heqq@synu.edu.cn); Zhiyuan Yao (yaozy@synu.edu.cn)

Abstract

The family Pholcidae C.L. Koch, 1850 is highly diverse in Guizhou Province, southwestern China, and currently contains four genera and 22 species. Nevertheless, the distribution of pholcid spiders is conspicuously patchy in Guizhou. Species from Guiyang are poorly studied, and only *Pholcus spilis* Zhu & Gong, 1991 has been recorded. A survey was undertaken for the first time to study the pholcids in Guiyang. A total of four species are reported, comprising *Belisana yuhaoi* Yang & Yao, **sp. nov.** and three other species: *Leptopholcus tanikawai* Irie, 1999 (new record for Guiyang), *Pholcus spilis* Zhu & Gong, 1991 and *Spermophora senoculata* (Dugès, 1836) (new record for Guizhou).

Key words: Biodiversity, fauna, new record, new species, taxonomy



Academic editor: Fedor Konstantinov

Received: 29 April 2023

Accepted: 27 November 2023

Published: 13 December 2023

ZooBank: <https://zoobank.org/AB71200E-4EB6-40F9-B981-C78C0B3BD662>

Citation: Yang L, Zhao F, He Q, Yao Z (2023) A survey of pholcid spiders (Araneae, Pholcidae) from Guiyang, Guizhou Province, China. ZooKeys 1186: 175–184. <https://doi.org/10.3897/zookeys.1186.105736>

Copyright: © Lan Yang et al.

This is an open access article distributed under terms of the Creative Commons Attribution License ([Attribution 4.0 International – CC BY 4.0](https://creativecommons.org/licenses/by/4.0/)).

Introduction

The species-rich spider family Pholcidae currently contains 97 genera and 1937 species (World Spider Catalog 2023). It comprises five subfamilies: Arteminae Simon, 1893; Modisiminae Simon, 1893; Ninetinae Simon, 1890; Smeringopinae Simon, 1893; and Pholcinae C.L. Koch, 1850 (Huber 2011a; Dimitrov et al. 2013; Eberle et al. 2018), and has a worldwide distribution. Recently, a series of surveys of pholcid spiders have been undertaken in northern China. For instance, the expedition to Changbai Mountains revealed 26 species recorded from Liaoning Province (Lu et al. 2021; Yao et al. 2021; World Spider Catalog 2023; Zhao et al. 2023a). The expeditions to Yanshan-Taihang Mountains and Lüliang Mountains brought the fauna of pholcids from Hebei Province, Beijing and Shanxi Province to 31 species, six species, and 21 species, respectively (Lu et al. 2022a, b; World Spider Catalog 2023; Zhao et al. 2023b). To date, 17 genera and 271 species of pholcids have been recorded from China (World Spider Catalog 2023), of which nine genera and 99 species were collected in caves or at cave entrance ecotones from karst regions.

Guizhou Province, in the southwest of China, is one of the most spectacular examples of humid subtropical karst landscapes, and also exhibits high diversity of pholcids. Currently, four genera (*Belisana* Thorell, 1898, *Khorata* Huber, 2005, *Leptopholcus* Simon, 1893, *Pholcus* Walckenaer, 1805) and 22 species have been record-

ed (World Spider Catalog 2023). Nevertheless, only one species, *Pholcus spilis* Zhu & Gong, 1991, has been recorded from Guiyang, the provincial capital of Guizhou. In this paper, we undertook a survey in Guiyang for the first time and report four species, comprising a new species from a cave and three known species (Figs 1, 2).

Material and methods

Specimens were examined and measured with a Leica M205 C stereomicroscope. The left male palp was photographed. The epigyne was photographed before dissection. The vulva was treated in a 10% warm solution of potassium hydroxide (KOH) to dissolve soft tissues before illustration. Images were captured with a Canon EOS 750D wide zoom digital camera (24.2 megapixels) mounted on the stereomicroscope mentioned above and assembled using Helicon Focus v.3.10.3 image stacking software (Khmelik et al. 2005). All measurements are given in millimeters (mm). Leg measurements are shown as: total length (femur, patella, tibia, metatarsus and tarsus). Leg segments were measured on their dorsal sides. The distribution map was generated with ArcGIS v. 10.2 (ESRI Inc. 2002). The specimens studied are preserved in 75% ethanol and deposited in the College of Life Science, Shenyang Normal University (SYNU) in Liaoning, China and Guizhou Normal University (GZNU) in Guizhou, China.

Terminology and taxonomic descriptions follow Huber (2005) and Yao et al. (2015). The following abbreviations are used in the descriptions: **ALE** = anterior lateral eye, **AME** = anterior median eye, **PME** = posterior median eye, **L/d** = length/diameter; used in the illustrations: **b** = bulb, **ba** = bulbal apophysis, **da** = distal apophysis, **e** = embolus, **ep** = epigynal pocket, **f** = flap, **pa** = proximo-lateral apophysis, **pp** = pore plate, **pr** = procursus.




Figure 1. Distribution of Pholcidae treated in this paper 1 *Belisana yuhai* sp. nov. 2 *Leptopholcus tanikawai* 3 *Pholcus spilis* 4 *Spermophora senoculata*.




Figure 2. Living specimens of Pholcidae treated in this paper **A, B** *Belisana yuhaoi* sp. nov. (♂♀) **C, D** *Sperophora senoculata* (♂) **E** *Pholcus spilis* (♂) **F** *Leptopholcus tanikawai* (juvenile). Photographs by Q Lu (Shenzhen).

Taxonomic accounts

Family Pholcidae C.L. Koch, 1850

Subfamily Pholcinae C.L. Koch, 1850

Genus *Belisana* Thorell, 1898

Type species. *Belisana tauricornis* Thorell, 1898.

***Belisana yuhaoi* Yang & Yao, sp. nov.**

<https://zoobank.org/35D4BC52-53E7-47D4-9E54-2FC5291E33DC>

Figs 3, 4

Type material. Holotype: ♂ (SYNU-Ar00301), cave without a name (27°5.40'N, 106°30.00'E, 1109 m), Liutong Town, Xiuwen County, Guiyang, **Guizhou, China**, 5 June 2022, H Yu & Q Lu leg. **Paratypes:** 2♂ (SYNU-Ar00302, Ar00303) and 3♀ (SYNU-Ar00304–00306), same data as for the holotype. 1♀ (SYNU-Ar00307), Duobing Cave (27°6.00'N, 106°30.00'E, 1026 m), other data as for the holotype.

Etymology. The specific name is a patronym in honor of the collector Hao Yu; noun (name) in genitive case.

Diagnosis. The new species resembles *B. galeiformis* Zhang & Peng, 2011 (Zhang and Peng 2011: 52, fig. 1A–F) with similar male chelicerae and bulbal apophysis (Fig. 4C, D), but it can be distinguished by prolatero-ventral lamella of procursus nearly round (arrow 2 in Fig. 3C; nearly angular in *B. galeiformis*), by distal membranous lamella of procursus laterally weak sclerotized (sclerotized part nearly half-round, arrow 3 in Fig. 3C; with triangular sclerite in *B. galeiformis*), by procursus with triangular retrolateral membranous flap (f in Fig. 3D; flap large and half-round in *B. galeiformis*), by epigynal plate nearly round, posteriorly strongly curved (Fig. 4A; hat-shaped, posteriorly straight in *B. galeiformis*), and by vulval pore plates nearly triangular (Fig. 4B; long elliptic in *B. galeiformis*).

Description. Male (holotype, SYNU-Ar00301): total length 1.93 (2.03 with clypeus), prosoma 0.74 long, 0.78 wide, opisthosoma 1.19 long, 0.96 wide. Legs I and IV missing, femur II: 3.92 (other segments missing), leg III: 10.18 (2.97, 0.33, 2.48, 3.56, 0.84). Eye interdistances and diameters: PME–PME 0.13, PME 0.10, PME–ALE 0.04, AME absent. Sternum width/length: 0.65/0.62. Habitus as in Fig. 4E, F. Dorsal shield of prosoma yellowish, with large, brown radiating marks; ocular area and clypeus yellowish, with brown marks; sternum yellowish, with triangular posterior brown marks. Legs whitish, without darker rings. Opisthosoma yellowish, with dorsal and lateral brown spots. Ocular area not elevated. Thoracic furrow absent. Clypeus unmodified. Chelicerae as in Fig. 4D, with a pair of proximo-lateral apophyses and a pair of curved distal apophyses (distance between tips of distal apophyses: 0.21). Palp as in Fig. 3A, B; trochanter with short retrolatero-ventral apophysis; femur with small retrolatero-proximal protrusion; procursus simple proximally but complex distally, with prolatero-subdistal sclerite (arrow 1 in Fig. 3C), sclerotized prolatero-ventral lamella (arrow 2 in Fig. 3C), distal membranous lamella (arrow 3 in Fig. 3C), curved distal spine (arrow 4 in Fig. 3C), sclerotized dorsal apophysis (arrow 5 in




Figure 3. *Belisana yuhaoi* sp. nov., holotype male **A, B** palp: **A** prolateral view **B** retrolateral view **C, D** distal part of procurus: **C** prolateral view, arrow 1 indicates prolatero-subdistal sclerite, arrow 2 indicates sclerotized prolatero-ventral lamella, arrow 3 indicates distal membranous lamella, arrow 4 indicates curved distal spine, arrow 5 indicates sclerotized dorsal apophysis **D** retrolateral view. Abbreviations: b = bulb, ba = bulbal apophysis, e = embolus, f = flap, pr = procurus. Scale bars: 0.10 mm (**A, B**); 0.02 mm (**C, D**).




Figure 4. *Belisana yuhaoi* sp. nov., holotype male (C–F) and paratype female (A, B, G, H) **A** epigyne, ventral view **B** vulva, dorsal view **C** bulbal apophyses, prolateral view **D** chelicerae, frontal view **E–H** habitus: **E**, **G** dorsal view **F** lateral view **H** ventral view. Abbreviations: b = bulb, ba = bulbal apophysis, da = distal apophysis, e = embolus, ep = epigynal pocket, pa = proximo-lateral apophysis, pp = pore plate. Scale bars: 0.05 mm (A–D); 0.20 mm (E–H).

Fig. 3C), and angular retrolateral membranous flap (f in Fig. 3D); bulb (Fig. 4C) with hooked apophysis and simple embolus.

Female (paratype, SYNU-Ar00304): similar to male, habitus as in Fig. 4G, H. Total length 2.03 (2.13 with clypeus), prosoma 0.83 long, 0.91 wide, opisthosoma 1.20 long, 1.04 wide; tibia I: 4.02; tibia I L/d: 50. Eye interdistances and diameters: PME–PME 0.12, PME 0.08, PME–ALE 0.03, AME absent. Sternum width/length: 0.58/0.51. Epigyne (Fig. 4A) simple and flat, with brown marks and a pair of posterior pockets, 0.17 apart (ep in Fig. 4A, B). Vulva (Fig. 4B) with ridge-shaped anterior arch and a pair of nearly triangular pore plates.

Variation. In one male paratype (SYNU-Ar00302), leg I: 23.28 (6.02, 0.39, 5.71, 9.36, 1.80); tibia I L/d: 60. Retrolateral trichobothrium of tibia I at 6% proximally; legs with short vertical setae on metatarsi, without spines and curved setae; tarsus I with 22 distinct pseudosegments. Tibia I in another male paratype (SYNU-Ar00303): 5.38. Tibia I in the other two female paratypes (SYNU-Ar00305, Ar00306): 3.28, 3.75 (leg I missing in SYNU-Ar00307).

Habitat. The species was found inside cave.

Distribution. China (Xiuwen County in Guizhou; type locality, Fig. 1).

Genus *Leptopholcus* Simon, 1893

Type species. *Leptopholcus signifer* Simon, 1893.

Leptopholcus tanikawai Irie, 1999

Leptopholcus tanikawai Irie, 1999: 37, figs 1–5 (♂♀).

Leptopholcus tanikawai Irie, 2009: 108, figs 9–11 (♂♀). Huber 2011b: 97, figs 228–231, 273, 274, 395–401, 426, 427 (♂♀). Fu and Chen 2017: 18, figs 2A–E, 3A–E (♂♀).

New material examined. 1♂ (GZNU), Taoyuanhe (26°31.80'N, 106°28.20'E, 1237 m), Xiuwen County, Guiyang, **Guizhou, China**, 4 June 2022, H Yu leg. 3♀ (GZNU), a forest park (26°33.60'N, 106°45.60'E, 1165 m), Nanming District, Guiyang, **Guizhou, China**, 10 August 2021, H Yu, H Zhang, D Wang, L Li & J Xin leg. 1 juvenile (GZNU), Guizhou Botanical Garden (26°37.80'N, 106°43.80'E, 1249 m), Yunyan District, Guiyang, **Guizhou, China**, 4 June 2022, H Yu & Q Lu leg.

Distribution. China (Xiuwen County, Nanming District and Yunyan District in Guizhou; Fig. 1).

Genus *Pholcus* Walckenaer, 1805

Type species. *Aranea phalangoides* Fuesslin, 1775.

Pholcus spilis Zhu & Gong, 1991

Pholcus spilis Zhu & Gong, 1991: 22, fig. 4A–G (♂♀).

Pholcus spilis Song, Zhu and Chen 1999: 59, fig. 24E–H (♂♀). Zhang and Zhu 2009: 83, fig. 47A–G (♂♀). Huber 2011b: 359, figs 1654, 1727, 1728 (♂). Yao and Li 2012: 33, figs 161A–D, 162A–C (♂♀). Yin et al. 2012: 171, fig. 35a–g (♂♀). Zhang 2018: 4, figs 1–3A–F, pl. 1 (♂♀).

New material examined. 1♂ (GZNU), Sanchahe (26°16.20'N, 106°48.00'E, 1162 m), Gaopo Town, Huaxi District, Guiyang, **Guizhou, China**, 20 May 2022, H Yu & Q Lu leg.

Distribution. China (Huaxi District in Guizhou; Fig. 1).

Genus *Spermophora* Hentz, 1841

Type species. *Spermophora meridionalis* Hentz, 1841.

Spermophora senoculata (Dugès, 1836)

See World Spider Catalog 2023

New material examined. 2♂ (GZNU) and 2♀ (GZNU), Sanchahe (26°16.20'N, 106°48.00'E, 1162 m), Gaopo Town, Huaxi District, Guiyang, **Guizhou, China**, 20 May 2022, H Yu & Q Lu leg. 1♂ (GZNU), Laobanghe (26°20.40'N, 106°43.80'E, 1022 m), Qiantao Town, Huaxi District, Guiyang, **Guizhou, China**, 18 May 2022, H Yu & Q Lu leg.

Distribution. China (Huaxi District in Guizhou; Fig. 1).

Acknowledgements

The manuscript benefited greatly from comments by Gergin Blagoev, Fedor Konstantinov and Kadir B Kunt. Joseph KH Koh kindly checked the English.

Additional information

Conflict of interest

The authors have declared that no competing interests exist.

Ethical statement

No ethical statement was reported.

Funding

This study was supported by the National Natural Science Foundation of China (NSFC-32170461, 31872193) and the Research Fund for the Doctoral Program of Shenyang Normal University (BS201841). Field work and part of the laboratory work was supported by the Project of Biodiversity Survey and Assessment in Guiyang (GZZC-2021-018).

Author contributions

ZY and QH designed and funded the study. LY and ZY performed morphological species identification. LY and FZ finished the species descriptions and took the photos. LY and ZY drafted the manuscript. QH and ZY revised the manuscript. All authors read and approved the final version of the manuscript.

Author ORCIDs

Lan Yang  <https://orcid.org/0000-0002-7754-9275>

Fangyu Zhao  <https://orcid.org/0000-0002-1005-8471>

Qiaoqiao He  <https://orcid.org/0000-0002-9381-7444>

Zhiyuan Yao  <https://orcid.org/0000-0002-1631-0949>

Data availability

All of the data that support the findings of this study are available in the main text.

References

- Dimitrov D, Astrin JJ, Huber BA (2013) Pholcid spider molecular systematic revisited, with new insights into the biogeography and the evolution of the group. *Cladistics* 29(2): 132–146. <https://doi.org/10.1111/j.1096-0031.2012.00419.x>
- Eberle J, Dimitrov D, Valdez-Mondragón A, Huber BA (2018) Microhabitat change drives diversification in pholcid spiders. *BMC Evolutionary Biology* 18(1): 1–141. <https://doi.org/10.1186/s12862-018-1244-8>
- ESRI Inc (2002) ArcView GIS 10.2. Environmental Systems Research Institute, Redlands, California. <http://www.esri.com/software/arcgis/arcview/> [Accessed 18 April 2023]
- Fu Y, Chen J (2017) A newly recorded species of the genus *Leptopholcus* from China (Araneae: Pholcidae). *Acta Arachnologica Sinica* 26(1): 18–21. <https://doi.org/10.3969/j.issn.1005-9628.2017.01.004>
- Huber BA (2005) High species diversity, male-female coevolution, and metaphyly in Southeast Asian pholcid spiders: The case of *Belisana* Thorell 1898 (Araneae, Pholcidae). *Zoologica* 155: 1–126.
- Huber BA (2011a) Phylogeny and classification of Pholcidae (Araneae): An update. *The Journal of Arachnology* 39(2): 211–222. <https://doi.org/10.1636/CA10-57.1>
- Huber BA (2011b) Revision and cladistic analysis of *Pholcus* and closely related taxa (Araneae, Pholcidae). *Bonner Zoologische Monographien* 58: 1–509.
- Irie T (1999) A new species of the genus *Leptopholcus* (Araneae: Pholcidae) from the Yaeyama Islands, Japan. *Acta Arachnologica* 48(1): 37–39. <https://doi.org/10.2476/asjaa.48.37>
- Irie T (2009) Pholcidae. In: Ono H (Ed.) *The Spiders of Japan with Keys to the Families and Genera and Illustrations of the Species*. Tokai University Press, Kanagawa, 106–111.
- Khmelik VV, Kozub D, Glazunov A (2005) *Helicon Focus* 3.10.3. <https://www.heliconsoft.com/heliconsoft-products/helicon-focus/> [Accessed 10 April 2023]
- Lu Y, Yang F, He Q (2021) *Pholcus maxian* sp. nov., the fifth endemic spider species of *Pholcus phungiformes* species-group (Araneae: Pholcidae) at the border between Jilin, China and North Korea. *Biodiversity Data Journal* 9(e72464): 1–7. <https://doi.org/10.3897/BDJ.9.e72464>
- Lu Y, Chu C, Zhang X, Li S, Yao Z (2022a) Europe vs. China: *Pholcus* (Araneae, Pholcidae) from Yanshan-Taihang Mountains confirms uneven distribution of spiders in Eurasia. *Zoological Research* 43(4): 532–534. [Supplement 1–78] <https://doi.org/10.24272/j.issn.2095-8137.2022.103>
- Lu Y, Yao Z, He Q (2022b) A new species of *Pholcus yichengicus* species-group (Araneae, Pholcidae) from Hebei Province, China. *Biodiversity Data Journal* 10(e81800): 1–7. <https://doi.org/10.3897/BDJ.10.e81800>
- Song D, Zhu M, Chen J (1999) *The Spiders of China*. Hebei Science and Technology Publishing House, Shijiazhuang, 640 pp.

- World Spider Catalog (2023) World Spider Catalog, Version 24.5. Natural History Museum Bern. <http://wsc.nmbe.ch> [Accessed 22 November 2023]
- Yao Z, Li S (2012) New species of the spider genus *Pholcus* (Araneae: Pholcidae) from China. *Zootaxa* 3289(1): 1–271. <https://doi.org/10.11646/zootaxa.3289.1.1>
- Yao Z, Pham DS, Li S (2015) Pholcid spiders (Araneae: Pholcidae) from northern Vietnam, with descriptions of nineteen new species. *Zootaxa* 3909(1): 1–82. <https://doi.org/10.11646/zootaxa.3909.1.1>
- Yao Z, Wang X, Li S (2021) Tip of the iceberg: species diversity of *Pholcus* spiders (Araneae, Pholcidae) in the Changbai Mountains, Northeast China. *Zoological Research* 42(3): 267–271. [Supplement 1–60] <https://doi.org/10.24272/j.issn.2095-8137.2021.037>
- Yin C, Peng X, Yan H, Bao Y, Xu X, Tang G, Zhou Q, Liu P (2012) Fauna Hunan: Araneae in Hunan, China. Hunan Science and Technology Press, Changsha, 1590 pp.
- Zhang F (2018) Pholcidae, Miturgidae, Clubionidae, Eutichuridae. In: Zhang F, Xue X (Eds) Fauna of Tianmu Mountain (Vol. II). Arachnida. Araneae Eriophyoidea. Zhejiang University Press, Hangzhou, 1–5. [92–101, 105–107, pl. 1, 19–22.]
- Zhang F, Peng Y (2011) Eleven new species of the genus *Belisana* Thorell (Araneae: Pholcidae) from south China. *Zootaxa* 2989(1): 51–68. <https://doi.org/10.11646/zootaxa.2989.1.2>
- Zhang F, Zhu M (2009) A review of the genus *Pholcus* (Araneae: Pholcidae) from China. *Zootaxa* 2037(1): 1–114. <https://doi.org/10.11646/zootaxa.2037.1.1>
- Zhao F, Jiang T, Yang L, He Q, Zheng G, Yao Z (2023a) Pholcid spiders of the *Pholcus phungiformes* species group (Araneae, Pholcidae) from Liaoning Province, China: An overview, with description of a new species. *ZooKeys* 1156: 1–14. <https://doi.org/10.3897/zookeys.1156.98331>
- Zhao F, Yang L, Zou Q, Ali A, Li S, Yao Z (2023b) Diversity of *Pholcus* spiders (Araneae: Pholcidae) in China's Lüliang Mountains: an integrated morphological and molecular approach. *Insects* 14: e364. [pp. 1–34] <https://doi.org/10.3390/insects14040364>
- Zhu M, Gong L (1991) Four new species of the genus *Pholcus* from China (Araneae: Pholcidae). *Acta Zootaxonomica Sinica* 16(1): 18–26.