

# A taxonomic review of the Hydraenidae in South Korea (Coleoptera)

Dae-Hyun Lee<sup>1</sup>, Kee-Jeong Ahn<sup>2</sup>

**1** Institute of Ecosystem Restoration Planning, 97 Deokmyeong-ro, Yuseong-gu, Daejeon 34155, Republic of Korea **2** Department of Biology, Chungnam National University, Daejeon 34134, Republic of Korea

Corresponding author: Kee-Jeong Ahn (kjahn@cnu.ac.kr)

---

Academic editor: M. Michat | Received 26 September 2016 | Accepted 3 November 2016 | Published 21 November 2016

<http://zoobank.org/A32A88C2-C671-4301-A4D6-E02A40A1F41A>

---

**Citation:** Lee D-H, Ahn K-J (2016) A taxonomic review of the Hydraenidae in South Korea (Coleoptera). ZooKeys 634: 57–78. doi: 10.3897/zookeys.634.10640

---

## Abstract

A taxonomic study of the South Korean Hydraenidae is presented. Eight species in two genera are recognized, one of which is reported for the first time in the Korean peninsula, *Ochthebius marinus* (Paykull). It was also found that *Hydraena riparia* Kugelann and *O. inermis* Sharp previously recorded in South Korea were incorrect identifications of *H. puetzi* Jäch and *O. lobatus* Pu, respectively. Habitus and SEM photographs, line drawings of aedeagus, distribution maps, keys, and redescriptions of the species are provided.

## Keywords

Coleoptera, Hydraenidae, South Korea, taxonomy

## Introduction

The Hydraenidae, relatively small water beetles with unique habitus, are usually found at margins of running water, slowly flowing water, and stagnant water in the vicinity of lotic biotopes (Jäch et al. 2016). Approximately 1600 species in 42 genera have been recorded in the world (Ślipiński et al. 2011); 941 species in nine genera in the Palaearctic (Jäch and Skale 2015); 100 species in seven genera in China, 35 species in three genera in Japan; and 11 species in three genera in the Far East of Russia (Jäch and Skale 2015).

Members of the Hydraenidae are characterized by combination of the following features: separation of the gula and submentum by confluent genae; presence of an anterior plate-like premento-hypopharyngeal extension; large labrum with a deep medial incision in some species; and palpigers connected by a transverse sclerotized bar (Beutel et al. 2003).

Kwon and Suh (1986) first recorded *Hydraena riparia* Kugelann in South Korea. Later, Lee (1995) added two species (*Ochthebius inermis* Sharp and *O. satoi* Nakane) with descriptions and habitus illustrations. Recently, Jäch and Delgado (2014) found that a marine littoral species of *Neochthebius granulosus* (Satô) previously recorded by Park and Ahn (2008) was an incorrect identification of *O. ahni* Jäch and Delgado and *O. parki* Jäch and Delgado. Jäch and Skale (2015) added four species [*H. miyatakei* Satô, *H. puetzi* Jäch, *O. hasegawai* Nakane and Matsui, and *O. lobatus* Pu]. Accordingly, nine species in two genera have been recorded in South Korea.

In this paper, one species is reported for the first time in the Korean peninsula, *O. marinus* (Paykull). It was also found that *H. riparia* and *O. inermis* previously recorded in South Korea were incorrect identifications of *H. puetzi* and *O. lobatus*, respectively. Habitus and scanning electron microscopy photographs, keys, redescriptions, and diagnostic characters with illustrations of the species are provided.

## Material and methods

To identify South Korean hydraenid species more reliably, they were compared with voucher specimens in the Natural History Museum (NHM, London, United Kingdom), Naturhistorisches Museum (NMW, Wien, Austria) and Ehime University Museum (EUMJ, Matsuyama, Japan). The specimens used in this study are deposited in Chungnam National University Insect Collection (CNUIC, Daejeon, Korea). Habitus and scanning electron microscope (SEM) photographs were prepared based on a former study (Lee and Ahn 2015). See Jäch and Delgado (2014) for detailed descriptions of *O. ahni* and *O. parki*. The terms of taxonomic characters and measurements of specimens mainly followed Perkins (2001) and Jäch et al. (2016). The geographical subdivision of China and Russia was based on the standards of Löbl and Löbl (2015).

## Results

### Hydraenidae Mulsant, 1844

#### Key to the genera of South Korean Hydraenidae

- |   |  |                   |
|---|--|-------------------|
| 1 | Anterior margin of mentum (Fig. 8) dentate; maxillary palpi longer than antenna.....       | <i>Hydraena</i>   |
| - | Anterior margin of mentum (Fig. 18) not dentate; maxillary palpi shorter than antenna..... | <i>Ochthebius</i> |

## Genus *Hydraena* Kugelann, 1794

*Hydraena* Kugelann, 1794: 578. Type species *Hydraena riparia* Kugelann, 1794.

**Diagnosis.** Labral-mandibular interlocking device present. Mentum (Fig. 8) with acute median projection on anterior part. Sensilla various and complex. Features of the secretion delivery system specialized (Jäch et al. 2000).

### Key to the subgenera of South Korean *Hydraena*

- |   |   |                     |
|---|---|---------------------|
| 1 | Body brown to dark blue; longitudinal median genal suture and longitudinal inner genal carina complete .....          | <i>Hydraena</i>     |
| - | Body yellowish brown; longitudinal median genal suture and longitudinal inner genal carina incomplete or absent ..... | <i>Hydraenopsis</i> |

## Subgenus *Hydraena* Kugelann, 1794

*Hydraena* Kugelann, 1794: 578. Type species: *Hydraena riparia* Kugelann, 1794. See Jäch and Skale (2015) for more detailed synonymy and references.


**Diagnosis.** Lateral margin of labrum (Fig. 6) abruptly constricted posteriorly. Setae of labial palpomere 2 closely set. Anterior apex of modified longitudinal hypomeral carina with antennal pocket setae. Elytra (Fig. 11) with 15 or more striae. Mesoventral process (Fig. 14) obtuse, between 100° and 130° angle. Sternite III (Fig. 17) without pubescence behind coxal pits. Sternite VII (Fig. 17) pubescent with semicircular glabrous posterior region. Gonocoxite with a pair of subapical tufts (Jäch et al. 2000).

## *Hydraena* (*Hydraena*) *puetzi* Jäch, 1994

Figs 1, 6–17, 50, 55


*Hydraena* (*s. str.*) *puetzi* Jäch, 1994: 43; Jäch and Skale 2015: 139.

**Specimens examined. SOUTH KOREA:** Gangwon Prov.: 1♀, Goseong-gun, Gangeong-eub, Jinburyeong, N38°15'59" E128°23'04", 640m, 21.VII.2004, KM Yang, JS Park, leaf litter; 1♂ 2♀, Hongcheon-gun, Nae-myeon, Changchon-ri, Unduryeong-hill, 10.VIII.2012, SW Jeong, mountain stream; 1♂, Hongcheon-gun, Seo-myeon, Bangok-ri, Bangokgyo, 17.IX.2009, DH Lee, mountain stream (1♂, on slide); 1♂ 3♀, Pyeongchang-gun, Daegwanryeong-eub, Yucheon-ri, Haewon-temple, 15.IX.2009, YJ Park, under stone on stream margin (1♂, on slide); 1♀, Pyeongchang-gun, Jinbu-myeon, Dongsan-ri, Mt. Odaesan, Sangwon-temple, 16.VIII–15.IX. 2001, SJ Park, CW Shin, FIT; 1♂, same data as former except for, 8.V.2004, DH



**Figures 1–5.** Habitus. **1** *Hydraena puetzi*, 2.2 mm **2** *Ochthebius hasegawai*, 1.8 mm **3** *O. lobatus*, 2.2 mm **4** *O. marinus*, 2.0 mm **5** *O. satoi*, 1.8 mm.

Lee, stream; 1♂, same data as former except for, 8.V.2004, DH Lee, stream; 9♂♂ 11♀♀, same data as former except for, Mt. Odaesan, 22.V.2012, DH Lee, springfed pool; 1♂, Samcheok-si, Gagok-stream, 25.VI.1985. SH Lee; 3♀♀, Yeongwol-gun, Suju-myeon, Mt. Baehyang-san, 17.VII.2010, SW Jeong, mountain stream; Gyeongbuk Prov.: 1♀, Uljin-gun, Buk-myeon, Mt. Eungbongsan, 6.VI.1995, SH Lee; 1♂, Uljin-gun, Seo-myeon, Wangpi-ri, Golan-bridge, N $36^{\circ}54'27.1''$  E $129^{\circ}14'34.2''$ , 385m, 26.IV.2012, DH Lee, mountain stream; Gyeonggi Prov.: 1♀, Gapyeong-gun, Buk-myeon, Jeongmok-ri, Garim-bridge, 16.IX.2010, SW Jeong, stream; 1♂, Paju-si, Jangdan-myeon, Nosang-ri, 21.IX.2012, SW Jeong, stream.



**Figures 6–17.** *Hydraena puetzi*. **6** head (dorsal aspect) **7** head (ventral aspect) **8** mentum (ventral aspect) **9** maxillary palpus (ventral aspect) **10** pronotum (dorsal aspect) **11** anterior part of elytron (dorsal aspect) **12** posterior part of elytron (dorsal aspect) **13** prosternum (ventral aspect) **14** mesoventrite (ventral aspect) **15** mesotibia in male (dorsal aspect) **16** metaventrite (ventral aspect) **17** abdominal sternites (ventral aspect).

**Published South Korean records.** *Hydraena puetzi*: Jäch and Skale (2015: 103). *Hydraena riparia*: Kwon and Suh (1986: 99); Kim et al. (1994: 134); Lee (1995: 13); Han et al. (2007: 271); Han et al. (2008: 263); Cho and Park (2010: 96) [misidentification].

**Redescription.** Length 2.0–2.3 mm. Head black; pronotum and elytra brown to dark blue; ventral surface dark brown. Head (Fig. 6) trapezoidal, widest cross eyes, ventral side (Fig. 7) with sparse setae. Anterior margin of labrum (Fig. 6) nearly straight except antero-medial part deeply excised. Clypeus (Fig. 6) with relatively small punctures; antero-medial margin broadly rounded; antero-lateral part

protruded. Frontoclypeal suture (Fig. 6) bisinuate. Mentum (Fig. 8) subquadrate, widest at posterior corner, with large punctures on anterior and lateral parts; anterior margin broadly rounded, with a row of thick setae; posterior part protruded laterally. Submentum (Fig. 7) semicircular; antero-medial margin nearly straight, dentate; antero-lateral margin protruded and acute. Antenna with nine antennomeres; 1 longest, approx. 2.5 times as long as 2; 2 approx. 6.0 times as long as 3, bulbous at basal part; 3 bulbous at apical part; 4 dish-shaped; 5–9 clubbed and with pubescence. Maxillary palpomere (Fig. 9) 1 smallest; 2 longest, approx. 3.0 times as long as 3, few setae present on ventral part, with transverse imbricate reticulation except apical part; 3 bulbous at apical part, few setae present on dorsal and ventral parts, with imbricate reticulation except anterior third; 4 approx. 2.0 times as long as 3, many setae present on dorsal and ventral parts, apical part acute, with imbricate reticulation except anterior to middle. Pronotum (Fig. 10) hexagonal, widest at middle and narrowest posterior part, medial part protruded and flattened; anterior margin sinuate; antero-median margin straight; anterior and posterior corners rectangular; lateral margin rounded and serrated. Elytra (Figs 11, 12) widest at middle; antero-lateral and postero-lateral parts serrated. Prosternum (Fig. 13) transverse, with longitudinal carina on medial part; anterior and posterior corners acute. Mesoventrite (Fig. 14) reverse pentagonal, with two vertical carina on each side of midline. Mesotibia (Fig. 15) serrate on apico-lateral part, with thick spines. Metaventrite (Fig. 16) flattened, with two glabrous parts medially. Sternite VII (Fig. 17) with many long setae on medial part in female. Male terminal sternite semicircular; postero-medial part bifid; posterior part rounded and with many setae. Median lobe of aedeagus (Fig. 50) longer than paramere, strongly curved at middle; a seta present on anterior third; apical part acute and protruded; flagellum bisinuate. Left paramere (Fig. 50) as long as right; apical part long-oval; many long setae present on apical margin. Apical part of right paramere (Fig. 50) semicircular; many long setae present on apical margin.

**Distribution.** Korea, China (Liaoning, Shandong), Russia (Far East).

**Biological note.** They are found under leaves or stones in margins of mountain streams. Some specimens were collected from submerged branch and leaves in seepage on small cliff.

**Remarks.** Kwon and Suh (1986) first recorded this species as *H. riparia* in Korea. After that, many entomologists [Kim et al. (1994); Lee (1995); Han et al. (2007); Han et al. (2008); Cho and Park (2010)] reported the species, only in the local fauna without any taxonomic comments. After studying the specimen (1♂, Samcheok-si, Gagok-stream, 25.VI.1985. SH Lee) previously examined by Kwon and Suh (1986), we found that it had been incorrectly identified and actually is *H. puetzi*. This species can be distinguished from *H. riparia* by left paramere (Fig. 7A) parallel at middle, and apical part of right paramere (Fig. 7A) semicircular, with compact long setae on apex.

## Subgenus *Hydraenopsis* Janssens, 1972

*Hydraenopsis* Janssens, 1972: 254. Type species: *Hydraenopsis vietnamensis* Janssens, 1972.

**Diagnosis.** Transverse anterior inner gular carina absent; mesosternal intercoxal process protruding from mesosternal disc at an angle of 150°–180°; at least one of the two parameres shifted (Jäch et al. 2000).

### *Hydraena (Hydraenopsis) miyatakei* Satô, 1959

Fig. 55

*Hydraena miyatakei* Satô, 1959: 62; *Hydraena (Hydraenopsis) miyatakei*: Jäch and Skale 2015: 143.

**Specimens examined.** SOUTH KOREA: Gangwon Prov.: 1♀, Pyeongchang-gun, Baesujang, 29 VII 1994, SH Lee.

**Published South Korean record.** *Hydraena (Hydraenopsis) miyatakei*: Jäch and Skale (2015: 143).

**Distribution.** Korea, China (Jilin, Liaoning, Shandong), Japan, Russia (Far East).

**Remarks.** Only one specimen of this species was collected, not suitable for description or illustration.

## Genus *Ochthebius* Leach, 1815

Figs 18, 19, 22, 23, 29, 30, 31, 37, 38, 45, 46

*Ochthebius* Leach, 1815: 95. Type species: *Helophorus marinus* Paykull, 1798.

**Diagnosis.** Body (Fig. 26) with distinct pronoto-elytral angle. Head (Fig. 27) with a transverse groove separating clypeus and frons, the latter on each side with an interocular pit-like depression and a short longitudinal depression or groove on postero-medial part. Anterior margin of labrum truncate, sometimes with a small medial emargination. Pronotum (Figs 21, 30, 37, 45) with a narrow hyaline membranous cuticula, middle pronotal portion raised, and with longitudinal or transversal groove. Elytra (Figs 22, 31, 38, 46) with rows of punctures (Chiesa 1959; Hansen 1987; Jäch 1992).

### Key to the species of South Korean *Ochthebius*

- |   |  |                          |
|---|--|--------------------------|
| 1 | Pronotum widest at anterior third to three seventh part, protruded laterally.... | <b>2</b>                 |
| - | Pronotum widest at middle part, rounded laterally .....                          | <b>5</b>                 |
| 2 | Elytra (Fig. 31) with long setae .....   | <b><i>O. lobatus</i></b> |
| - | Elytra (Figs 22, 38, 46) with short setae.....                                   | <b>3</b>                 |

- 3 Maxillary palpomere 3(Fig. 20) approx. 4.0 times as long as 4... *O. hasegawai*  
 – Maxillary palpomere 3 (Figs 36, 44) approx. 2.0 times as long as 4..... 4  
 4 Medial part of metaventrite (Fig. 41) with pubescence; apical part of distal lobe of aedeagus acute..... *O. marinus*  
 – Medial part of metaventrite (Fig. 48) without pubescence; apical part of distal lobe of aedeagus round..... *O. satoi*  
 5 Pronotal longitudinal groove distinct (Jäch and Delgado 2014: Fig. 6); distal lobe of aedeagus curved in lateral view (Jäch and Delgado 2014: Figs 8, 15h)..... *O. abni*  
 – Pronotal longitudinal groove indistinct (Jäch and Delgado 2014: Fig. 7); distal lobe of aedeagus bisinuate in lateral view (Jäch and Delgado 2014: Figs 13, 15a) ..... *O. parki*

### Subgenus *Ochthebius* Leach, 1815

Figs 22, 27, 29, 31, 37

*Ochthebius* Leach, 1815: 95. Type species: *Helophorus marinus* Paykull, 1798.  
 See Jäch and Skale (2015) for more detailed synonymy and references.

**Diagnosis.** Lateral margin of pronotum (Figs 21, 30, 37, 45) with hyaline membranous cuticula. Marginal elytral ridge (Fig. 39) disappearing posteriorly. Pedicel (Fig. 19, 35) not enlarged distally (Hansen 1987; Jäch 1992).

### *Ochthebius (O.) ahni* Jäch & Delgado, 2014

*Ochthebius (s. str.) ahni* Jäch & Delgado, 2014: 85; Jäch and Skale 2015: 153.

**Specimens examined.** Holotype: 1♂ (CNUIC), with labels as follows: "KOREA: Gyeongbuk prov.: Kugae, 6 VII 1991, K. J. Ahn, ex., rock crevice., Holotype *Ochthebius ahni* sp. n. Jäch and Delgado 2014". Paratypes: 1♂1♀, same data as holotype. 3♂♂ 2♀♀, same data as holotype.

**Published South Korean records.** *Ochthebius ahni*: Jäch and Delgado (2014: 85); Jäch and Skale (2015: 153). *Neochthebius granulosus*: Park and Ahn (2008: 2506) [misidentification].

**Diagnosis.** See Jäch and Delgado (2014).

**Distribution.** Korea.

***Ochthebius (O.) hasegawai* Nakane & Matsui, 1986**

Figs 2, 18–25, 51, 56

*Ochthebius (s. str.) hasegawai* Nakane & Matsui, 1986: 81; Jäch 1998: 186; Jäch and Skale 2015: 156.

*Ochthebius mamagri* Shatrovsky, 1989: 263.


**Specimens examined. SOUTH KOREA:** Chungnam Prov.: 1♀, Nonsan-si, Beolgok-myeon, Sajeong-ri, N36°11'41.77" E127°16'18.77" 118 m, 24 V 2014, IS Yoo, under stone in stream; Gyeongbuk Prov.: 6♂♂ 8♀♀, Uljin-gun, Giseong-myeon, Dacheon-ri, Giseong-stream 17 VII 1995, SH Lee, mountain stream (3♂ 2♀, on slides); Gyeonggi Prov.: 1♀, Gapyeong-gun, Buk-myeon, Dodae-ri, Seungcheon-temple, 5 VII 2013, SW Jeong, HJ Park, stream.

**Published South Korean record.** *Ochthebius hasegawai*: Jäch and Skale (2015: 156).

**Redescription.** Length 1.6–2.0 mm. Body dark blue to black. Mentum (Fig. 18) 1.2 times as wide as long and with sparse setae; protruded and widest at anterior third; with a row of long setae on anterior margin. Anterior gena without long setae on posterior part in ventral view. Antenna (Fig. 19) with nine antennomeres; 1 longest, approx. 6.0 times as long as 2, two long setae present on lateral part; 2 widest at base; 3 bulbous at apical part; 4 semicircular; 5–9 clubbed and with pubescence. Maxillary palpomere (Fig. 20) 1 small, approx. 2.0 times as long as wide; 2 bulbous at apical part, 2.0 times as long as 1; 3 largest, bulbous at apical part, slightly longer than 2; 4 slender and paralleled, 0.25 times as long as 3. Pronotum (Fig. 21) reverse trapezoidal, widest at anterior fourth, with indistinct longitudinal groove on medial part, two small oval grooves on anterior part, relatively large oval groove on posterior part; anterior margin bisinuate; antero-medial margin protruded; anterior corner rectangular; lateral margin protruded at anterior third; posterior corner obtuse; postero-medial part rounded. Elytra (Fig. 22) widest at middle, with setae. Mesoventrite (Fig. 23) pentagonal, with T-shaped carina on anterior part; anterior margin transverse. Metaventrite (Fig. 24) with pubescence on medial part. Sternite VIII (Fig. 25) with more or less long setae on posterior part. Male terminal sternite (Fig. 25) semicircular and with a row of compact setae on posterior part. Median lobe of aedeagus (Fig. 51) long, slender, very weakly curved at middle; apical part acute; distal lobe slender, slightly acute apically. Paramere (Fig. 51) very short, with long setae on apical part.

**Distribution.** Korea, Japan, Russia (Far East).

**Biological note.** The specimens were collected under boulders in a fast-flowing stream.



**Figures 18–25.** *Ochthebius hasegawai*. **18** mentum (ventral aspect) **19** antennae (lateral aspect) **20** maxillary palpus (ventral aspect) **21** pronotum (dorsal aspect) **22** anterior part of elytron (dorsal aspect) **23** mesovenitrite (ventral aspect) **24** metaventrite (ventral aspect) **25** male terminal sternite (ventral aspect).

***Ochthebius (O.) lobatus* Pu, 1958**

Figs 3, 26–33, 52, 56

*Ochthebius (s. str.) lobatus* Pu, 1958: 256; Jäch, 1995:178; 2003: 351; Jäch and Skale, 2015: 157.


**Specimens examined.** SOUTH KOREA: Gangwon Prov.: 4♂♂ 2♀♀, Samcheok-si, Wondeok-eub, Wolcheon-ri, Gagok-steam, 1 V 1994, SH Lee, beside stream (4♂ 1♀, on slides); 5♂♂ 7♀♀, same data as former except for, 1 VI 1995; 1♀, same data as former except for, 25 VI 1995; Gyeongbuk Prov.: 2♂♂ 3♀♀, Uljin-gun, Namdae-stream, 17 VII 1995, SH Lee.

**Published South Korean records.** *Ochthebius lobatus*: Jäch and Skale (2015: 157). *Ochthebius inermis*: Lee (1995: 13) [misidentification].

**Redescription.** Length 2.2–2.6 mm. Head dark brown; pronotum and elytra brown to dark brown; ventral surface mostly brown. Dorsum (Figs 27, 30, 31) with whitish long setae. Mentum (Fig. 28) 1.2 times as wide as long and with sparse setae, widest anterior third; antero-lateral part protruded. Anterior gena without long setae on posterior part in ventral view. Antenna with nine antennomeres; 1 longest, approx. 6.0 times as long as 2, two long setae present on lateral part; 2 bulbous at medial part; 3 bulbous at apical part; 4 semicircular; 5–9 clubbed and with pubescence. Maxillary palpomere (Fig. 29) 1 small, approx. 2.0 times as long as wide; 2 bulbous at apical part, 3.5 times as long as 1; 3 bulbous at apical part, as long as 2; 4 slender and parallel-sided, 0.5 times as long as 3. Pronotum (Fig. 30) reverse trapezoidal, widest at anterior three seventh, with distinct longitudinal groove on medial part, two rounded grooves on anterior part, two oval grooves on posterior part; anterior margin bisinuate; antero-median margin straight; anterior corner rectangular; posterior corner obtuse; postero-medial part rounded. Elytra (Fig. 31) widest at middle, with long setae. Mesoventrite (Fig. 32) pentagonal with T-shaped carina on anterior part; anterior margin transverse. Metaventrite (Fig. 32) with pubescence on medial part. Sternite VIII (Fig. 33) with more or less long setae on posterior part. Male terminal sternite semicircular with few spines on posterior part. Median lobe of aedeagus (Fig. 52) weakly slender, curved; subapical part vented, inflated at anterior third; apical part acute with few setae; distal lobe cylindrical, approx. 0.25 times as long as median lobe. Paramere (Fig. 52) slender, shorter than median lobe, with few setae on apical part.

**Distribution.** Korea, China (Chongqing, Jilin, Liaoning, Sichuan, Yunnan).

**Remarks.** Lee (1995) first reported *O. inermis* Sharp in South Korea but we found that this was a misidentification of *O. lobatus*, based on our examination of his voucher specimens (4♂♂ 2♀♀, Samcheok-si, Wondeok-eub, Wolcheon-ri, Gagok-steam, 1 V 1994, SH Lee, ex. beside stream). This species can be distinguished from *O. inermis* by elytra brown and distal lobe of aedeagus robust.



**Figures 26–33.** *Ochthebius lobatus*. **26** body (dorsal aspect) **27** head (dorsal aspect) **28** mentum (ventral aspect) **29** maxillary palpus (dorsal aspect) **30** pronotum (dorsal aspect) **31** anterior part of elytron (dorsal aspect) **32** meso- and metaventrite (ventral aspect) **33** sternites VIII–IX (ventral aspect).

***Ochthebius (O.) marinus* (Paykull, 1798)**

Figs 4, 34–41, 53, 56

*Helophorus marinus* Paykull, 1798: 245.*Ochthebius pallidus* Mulsant, 1844: 61.*Ochthebius holmbergi* Mannerheim, 1853: 166.*Ochthebius subabruptus* Rey, 1885: 23.*Ochthebius (s. str.) marinus* Jäch, 1992: 112; 2003: 330; Jäch and Skale 2015: 157.


**Specimens examined.** SOUTH KOREA: Gyeonggi Prov.: 2♂♂ 3♀♀, Ansan-si, Danwon-gu, Daebudo-dong, Dongju-salt farm, N37°14'05.60" E126°36'19.55" 4 m, 7 XI 2013, DH Lee, IS Yoo, SG Lee, saline pond near salt farm (1♂, on slide).

**Redescription.** Length 1.7–2.0 mm. Head metallic dark green; pronotum mostly pale yellowish brown and medial part brown; elytra pale yellow and rows of serial punctures dark brown; ventral surface mostly dark brown. Mentum (Fig. 34) as long as wide, widest anterior third and with sparse setae; antero-medial part broadly excised; antero-lateral part protruded. Anterior gena (Fig. 35) with long setae on posterior part in ventral view. Gula (Fig. 35) with spiny microreticulation. Antenna (Fig. 35) with nine antennomeres; 1 longest, approx. 6.0 times as long as 2, with two long setae on lateral part; 2 widest at base; 3 bulbous at apical part; 4 dish-shaped; 5–9 clubbed with pubescence. Maxillary palpomere (Fig. 36) 1 small, approx. 2.0 times as long as wide; 2 bulbous at apical part, 4.0 times as long as 1; 3 largest, bulbous at apical part, slightly longer than 2; 4 slender and paralleled, 0.5 times as long as 3. Pronotum (Fig. 37) reverse trapezoidal, widest at anterior third, with indistinct longitudinal groove on median part; anterior margin nearly straight; anterior corner rectangular; lateral margin protruded at anterior third; posterior corner obtuse; postero-medial part rounded. Elytra (Figs 38, 39) widest at middle, with setae. Mesoventrite (Fig. 40) pentagonal, with transverse thick carina on anterior part; antero-lateral part with two grooves. Metaventrite (Fig. 41) with pubescence on medial part. Sternite VIII with more or less long setae on posterior part. Male terminal sternite semicircular and with long setae on posterior margin. Median lobe of aedeagus (Fig. 53) slightly curved; apical part acute; distal lobe approx. 0.1 times as long as median lobe. Paramere (Fig. 53) shorter than median lobe; apical part oval with setae.

**Distribution.** Asia: Korea, China (Beijing, Heilongjiang), Russia (East Siberia); Europe: Denmark, Estonia, Finland, France, Great Britain, Germany, Ireland, Latvia, The Netherlands, Norway, Poland, Russia, Spain, Sweden; North America.

**Biological note.** The specimens were collected in saline pond (salinity 32.84‰) with algae and plentiful vegetation near salt farm. They were found with *Hydroglyphus coreanus* Lee & Ahn (Dytiscidae), *Berosus lewisius* Sharp, *B. spinosus* (Steven), *Enochrus simulans* (Sharp), and *Paracymus aeneus* (Germar) (Hydrophilidae).

**Remarks.** This species is recorded for first time in Korea.



**Figures 34–41.** *Ochthebius marinus*. **34** mentum (ventral aspect) **35** head (ventral aspect) **36** maxillary palpus (ventral aspect) **37** pronotum (dorsal aspect) **38** anterior part of elytron (dorsal aspect) **39** posterior part of elytron (dorsal aspect) **40** mesoventrite (ventral aspect) **41** metaventrite (ventral aspect).

### *Ochthebius (O.) parki* Jäch & Delgado, 2014

*Ochthebius (s. str.) parki* Jäch & Delgado, 2014: 88; Jäch and Skale 2015: 158.

**Specimens examined.** Holotype: 1♂ (CNUIC), with labels as follows: "Geoje City, Gabae-ri, 1 VII 2000, K.-J. Ahn, H.-J. Kim, M.-J. Jeon, on barnacles, Holotype *Ochthebius parki* sp. n. Jäch and Delgado 2014". Paratypes: 2♂♂, same data as holotype. **SOUTH KOREA:** Chungnam Prov.: 1♂, Boryeong-si, Ungcheon-eup, Doksan-ri, Holmoi beach, 6 IX 2003, K.J. Ahn, J.S. Park, under seaweeds; Gangwon Prov.: 1♀, Sokcho-si, Dongmyeong-dong, N38°12'48.0" E128°36'05.9" -1 m, 28 V 2012, J.H. Song, under stone in beach; Gyeongbuk Prov.: 3♂♂ 4♀♀, Yongyeon [=Pohang-si, Buk-gu, Heunghae-eub, Yonggok-ri], 20 VII 1991, K.J. Ahn, , in rock crevice; Gyeonggi Prov.: 1♂ 1♀, Incheon-City, Is. Baekryeongdo, 7 VIII 2000, C.-W. Shin, on rock (near sea); Gyeongnam Prov.: 5♂♂ 2♀♀, Geoje City, Gabae-ri, 1 VII 2000, K.-J. Ahn, H.-J. Kim, M.-J. Jeon, ex barnacles; 2♂♂ 3♀♀, Is. Geojedo, Dongbu-myeon, Gabae-ri, 30 VI 2001, S.-J. Park, inside barnacles; 1♂ 1♀, Koje, Kabae, Korean Marine Biological Laboratory, 3 VII 1998, J.-Y. Lyu, H.-J. Kim, empty barnacle; 1♂ 1♀, Kōje-City, Gabae-ri, 30 VII 1999, K.-J. Ahn, on rock; Jeju Prov.: 1♂, Gwagji beach [=Jeju-si, Aewol-eub, Gwakji-ri, Gwakji beach], 3 VII 1991, K.J. Ahn, , on rock with clam; 2♂♂ 1♀, Namjeju-gun, Seongsan-eub, Ilchulbong, 11 VI 2005, S.I. Lee, S.J. Park, K.J. Ahn, M.J. Jeon, D.H. Lee, ex stones; Jeonnam Prov.: 2♂♂ 5♀♀, Jindo, Imhoe-myeon, Geumgab beach, 21 VIII 2001, S.-J. Park, on barnacles; 3♂♂, Wando, Sinji-myeon, Myeongsasimri beach, 23 VIII 2001, K.-J. Ahn, J.-H. Ahn, on barnacles.

**Published South Korean records.** *Ochthebius parki*: Jäch and Delgado (2014: 88); Jäch and Skale (2015:158). *Neochthebius granulosus*: Park and Ahn (2008: 2506) [misidentification].

**Diagnosis.** See Jäch and Delgado (2014).


**Distribution.** Korea.

### *Ochthebius (O.) satoi* Nakane, 1965


Figs 5, 42–49, 54, 56

*Ochthebius (s. str.) satoi* Nakane, 1965: 51; Jäch 1991: 77; 1998: 177; 2003: 328; Jäch and Skale 2015: 159.


**Specimens examined.** **SOUTH KOREA:** Chungbuk Prov.: 2♂♂ 4♀♀, Okcheon-gun, Okcheon-eub, Gyodong-ri, 4 V 1990, SH Lee, beside stream; Gangwon Prov.: 3♂♂ 5♀♀, Samcheok-si, Wondeok-eub, Wolcheon-ri, Gagok-steam, 1 VI 1995, SH Lee, beside stream (3♂♂ 1♀, on slides); Gyeongbuk Prov.: 10♂♂ 7♀♀, Uljin-gun, Giseong-myeon, Dacheon-ri, Giseong-stream 17 VII 1995, SH Lee, ex. mountain stream (2♂♂ 2♀♀, on slides); 2♂♂ 1♀, Uljin-gun, Wonnam-myeon, Gilgok-ri, Maehwa-stream, beside stream (2♂♂, on slides); Gyeonggi Prov.: 1♀, Yeoncheon-



**Figures 42–49.** *Ochthebius satoi*. **42** mentum (ventral aspect) **43** gena and gula (ventral aspect) **44** maxillary palpus (dorsal aspect) **45** pronotum (dorsal aspect) **46** anterior part of elytron (dorsal aspect) **47** mesoventrite (ventral aspect) **48** metaventrite (ventral aspect) **49** male terminal sternite (ventral aspect).



**Figures 50–54.** Aedeagus (lateral aspect). **50** *Hydraena puetzi* **51** *Ochthebius basegawai* **52** *O. lobatus* **53** *O. marinus* **54** *O. satoi*. Scale bars 0.1 mm.



**Figures 55–56.** Distribution maps. **55** *Hydraena puetzi* (circle), *H. miyatakei* (triangle) **56** *Ochthebius ahni* (trapezoid), *O. lobatus* (circle), *O. marinus* (diamond), *O. parki* (star), *O. satoi* (triangle), *O. hasegawai* (square).

gun, Jangnam-myeon, Wondal-ri, Samicheon-bridge, 5 VII 2014, SW Jeong, stream; Jeonbuk Prov.: 1♀, Buan-gun, Byeonsan-myeon, Junggye-ri, Jikso-fall, 11 VII 2014, SW Jeong, mountain stream.

**Published South Korean records.** *Ochthebius satoi*: Lee (1995: 14); Jäch and Skale (2015: 159).

**Redescription.** Length 1.6–1.8 mm. Head dark brown; pronotum and elytra yellowish brown; ventral surface mostly brown. Mentum (Fig. 42) slightly longer than wide and with sparse setae; antero-lateral part protruded. Anterior gena (Fig. 43) without long setae on posterior part in ventral view. Antenna with nine antennomeres; 1 longest, approx. 6.0 times as long as 2, two long setae present on lateral part; 2 bulbous at middle; 3 oval, widest at middle; 4 dish-shaped; 5–9 clubbed and with pubescence. Maxillary palpomere (Fig. 44) 1 small, approx. 2.0 times as long as wide; 2 bulbous at apical part, 3.0 times as long as 1; 3 largest, bulbous at apical part, slightly longer than 2; 4 slender and paralleled, 0.5 times as long as 3. Pronotum (Fig. 45) reverse trapezoidal, widest at anterior third, with indistinct longitudinal groove on medial part, two small oval grooves on anterior part, relatively large oval groove on posterior part; anterior margin bisinuate; antero-medial margin straight; anterior corner rectangular; lateral side protruded at anterior third; posterior corner obtuse; postero-medial part rounded. Elytra (Fig. 46) widest at middle, with setae. Mesoventrite (Fig. 47) pentagonal, with

T-shaped carina on anterior part; anterior margin transverse. Metaventrite (Fig. 48) without pubescence on medial part. Metatrochanter with a row of setae. Sternite VIII (Fig. 49) with more or less long setae on posterior part. Male terminal sternite (Fig. 49) semicircular, with sparse setae, relatively long setae on posterior part. Median lobe of aedeagus (Fig. 54) slender, curved; apical part strongly vented; distal lobe rounded. Paramere (Fig. 54) nearly parallel-sided, apical margin rounded, with few setae.

**Distribution.** Korea, Japan, China (Henan, Jilin, Liaoning, Nei Mongol, Shaanxi, Shandong, Taiwan), Russia (Far East).

## Discussion

This study revises the number and taxonomic status of hydraenid species known to occur in South Korea. In total, eight species are recognized. Two species (*O. ahni* and *O. parki*) are apparently endemic to the South Korean fauna (Jäch and Delgado 2014). Of the remaining species, five are widespread in the East Palaearctic region (*H. puetzi*, *H. miyatakei*, *O. lobatus*, *O. hasegawai*, *O. satoi*) and one occurs in the Holarctic region (*O. marinus*).

Diversity of South Korean Hydraenidae is very low compared to other adjacent countries, such as China (82 endemic species out of 100 species), Japan (29 endemic species out of 35 species), and the Far East Russia (no endemic species out of 11 species) (Jäch 2003; Jäch and Díaz 2003; 2004; 2005; 2006; 2012; Jäch and Delgado 2014; Jäch and Skale 2015). They show high diversity and endemism because of their small size and limited dispersal abilities (Jäch and Balke 2008). Further collecting efforts in the Korean Peninsula will probably add more hydraenid species to the Korean fauna such as the Palaearctic species, *H. riparia* Kugelann and *O. angusi* Jäch, and more species.

## Acknowledgements

We thank Dr. M. A. Jäch (NMW, Wien), Dr. S. W. Jung (National Science Museum, Daejeon), Dr. H. Yoshitomi (EUMJ, Matsuyama), Mr. S. H. Lee (Acheon Elementary School, Gimcheon) and Ms. C. Taylor (NHM, London) for providing many specimens including voucher specimens and the literature. Comments by H. Frank (Univ. of Florida, Gainesville) and M. A. Jäch significantly improved the manuscript. This work was supported by the research fund of 2016 Chungnam National University.

## References

- Beutel RG, Anton E, Jäch MA (2003) On the evolution of adult head structures and the phylogeny of Hydraenidae (Coleoptera, Staphyliniformia). Journal of Zoological Systematics and Evolutionary Research 41: 256–275. doi: 10.1046/j.1439-0469.2003.00224.x

- Chiesa A (1959) Hydrophilidae Europae (Coleoptera, Palpicornia). A. Forni, Bologna, 199 pp.
- Cho YB, Park SW (2010) Coleoptera. In: Paek MK (Ed.) Checklist of Korean Insects. Nature and Ecology, Seoul, 83–157.
- Han MS, Na YE, Bang HS, Kim MH, Kim KM, Roh KA, Lee JT (2007) The Fauna of Aquatic Invertebrates in Paddy Field. Korean Journal of Environmental Agriculture 26(3): 267–273. doi: 10.5338/KJEA.2007.26.3.267
- Han MS, Na YE, Bang HS, Kim MH, Kang KK, Hong HK, Lee JT, Ko BG (2008) An Illustrated book of Aquatic Invertebrates in Paddy Field (a Revised Edition). National Institute of Agricultural Science and Technology, RDA, Suwon, 416 pp. [In Korean]
- Hansen M (1987) The Hydrophiloidea (Coleoptera) of Fennoscandia and Denmark. Fauna Entomologica Scandinavica 18: 1–254.
- Jäch MA (1991) Revision of the Palearctic species of the genus *Ochthebius* VII. The *foveolatus* group (Coleoptera: Hydraenidae). Koleopterologische Rundschau 61: 61–94.
- Jäch MA (1992) Revision of the Palearctic species of the genus *Ochthebius* Leach VI. The *marinus* group (Coleoptera: Hydraenidae). Entomologica Basiliensia 14: 101–145.
- Jäch MA (1994) New and little known Palearctic species of the genus *Hydraena* (s. l.) Kugelann II (Coleoptera: Hydraenidae). Entomological Problems 25: 37–46.
- Jäch MA (1995) Hydraenidae (Coleoptera). In: Jäch MA, Ji L (Eds) Water Beetles of China, Volume I. Zoologisch-Botanische Gesellschaft in Österreich and Wiener Coleopterologenverein, Wien, 173–180.
- Jäch MA (1998) Hydraenidae: II. The Taiwanese and Japanese species of *Ochthebius* Leach (Coleoptera). In: Jäch MA, Ji L (Eds) Water Beetles of China, Volume II. Zoologisch-Botanische Gesellschaft in Österreich and Wiener Coleopterologenverein, Wien, 173–193.
- Jäch MA (2003) Hydraenidae: II. Synopsis of *Ochthebius* Leach from Mainland China, with descriptions of 23 new species In: Jäch MA, Ji L (Eds) Water Beetles of China, Volume III. Zoologisch-Botanische Gesellschaft in Österreich and Wiener Coleopterologenverein, Wien, 313–369.
- Jäch MA, Balke M (2008) Global diversity of water beetles (Coleoptera) in freshwater. Hydrobiologia 595: 419–442. doi: 10.1007/s10750-007-9117-y
- Jäch MA, Beutel RG, Díaz JA, Kodada J (2000) Subgeneric classification, description of head structures, and world check list of *Hydraena* Kugelann (Insecta: Coleoptera: Hydraenidae). Annalen des Naturhistorischen Museums in Wien 102B: 177–258.
- Jäch MA, Beutel RG, Delgado JA, Díaz JA (2016) Hydraenidae Mulsant, 1844. In: Beutel RG, Leschen RAB (Eds) Handbook of Zoology (Volume 4, Part 38) – Coleoptera, Beetles (Volume 1): Morphology and Systematics (Archostemata, Adephaga, Myxophaga, Polyphaga partim) (2<sup>nd</sup> Edn). Walter de Gruyter, Berlin & New York, 316–345.
- Jäch MA, Delgado JA (2014) Revision of the Palearctic species of the genus *Ochthebius* Leach XXIX. The Asian species of the *O. vandykei* group (Coleoptera: Hydraenidae). Koleopterologische Rundschau 84: 81–100.
- Jäch MA, Díaz JA (2003) Hydraenidae: I. *Edaphobates puetzi* gen. et sp. n. from Sichuan (Coleoptera: Hydraenidae). In: Jäch MA, Ji L (Eds) Water Beetles of China (Volume 3). Zoologisch-Botanische Gesellschaft in Österreich and Wiener Coleopterologenverein, Wien, 301–311.

- Jäch MA, Díaz JA (2005) Revision of the Chinese species of *Hydraena* Kugelann I. Descriptions of 15 new species of *Hydraena* s. str. from China (Coleoptera: Hydraenidae). *Koleopterologische Rundschau* 75: 53–104.
- Jäch MA, Díaz JA (2006) Revision of the Chinese species of *Hydraena* Kugelann II. *Hydraena* s. str. from Gansu and Sichuan (Coleoptera: Hydraenidae). *Koleopterologische Rundschau* 76: 79–95.
- Jäch MA, Díaz JA (2012) Descriptions of six new species of *Hydraena* s. str. Kugelann from Japan (Coleoptera: Hydraenidae). *Koleopterologische Rundschau* 82: 115–136.
- Jäch MA, Skale A (2015) Family Hydraenidae. In: Löbl I, Löbl D (Eds) Catalogue of Palaearctic Coleoptera (Volume 1, Revised and updated edition) – Hydrophiloidea–Staphylinoidea. Brill, Leiden, 130–162.
- Kim JI, Kwon YJ, Paek JC, Lee SM, An SR, Park HC, Chu HY (1994) Coleoptera. In: The Entomological Society of Korea and Korean Society of Applied Entomology (Eds) Check List of Insects from Korea. Kon-Kuk University Press, Seoul, 117–214.
- Kugelann JG (1794) Verzeichniss der in einigen Gegenden Preussens bis jetzt entdeckten Käfer-Arten, nebst kurzen Nachrichten von denselben. Neustes Magazin für die Liebhaber der Entomologie, herausgegeben von DH Schneider 1: 513–582.
- Kwon YJ, Suh SJ (1986) Check list of Water Beetles from Korea. *Insecta Koreana* 6: 91–113.
- Leach WE (1815) Entomology. In: Brewster D (Ed.) The Edinburgh Encyclopaedia (Volume 9, part 1). Balfour, Edinburgh, 57–172.
- Lee DH, Ahn KJ (2015) A taxonomic review of the Gyrinidae (Coleoptera) in Korea. *ZooKeys* 509: 87–107. doi: 10.3897/zookeys.509.9442
- Lee SH (1995) Water beetles of Bulyeong valley. *Corentomon* 6(1): 11–16.
- Löbl I, Löbl D (2015) Catalogue of Palaearctic Coleoptera (Volume 1, Revised and updated edition) – Hydrophiloidea–Staphylinoidea. Brill, Leiden, 1702 pp.
- Mannerheim CG (1853) Dritter Nachtrag zur Kaefer-Fauna der Nord-Amerikanischen Laender des Russischen Reiches. *Bulletin de la Société Impériale des Naturalistes de Moscou* 26(3–4): 95–273.
- Matsui E, Delgado JA (1997) A new species of the genus *Ochthebius* from Japan (Coleoptera, Hydraenidae). *Esakia* 37: 71–76.
- Mulsant E (1844) Histoire naturelle des coléoptères de France – Palpicornes. L. Maison, Paris [et Ch. Savy Jeune, Lyon], 196 pp.
- Nakane T (1965) New or little-known Coleoptera from Japan and its adjacent regions. XXIII. *Fragmента Coleopterologica Japonica* 13: 51–54.
- Nakane T, Matsui E (1986) Notes on some new Hydrophiloidea from Japan (Coleoptera). Papers on entomology presented to Prof. Takehiko Nakane in commemoration of his retirement. Japanese Society of Coleoptera, Tokyo, 81–90.
- Park SJ, Ahn KJ (2008) Redescription of the littoral genus and species of *Neochthebius graulosus* (Satô), and the first larval description for the genus (Coleoptera: Hydraenidae: Ochthebiinae). *Journal of Natural History* 42: 2501–2514. doi: 10.1080/00222930802354803
- Paykull G (1798) Fauna Suecica – Insecta (Tomus I). J.F. Edman, Upsaliae, 358 pp.
- Perkins PD (1980) Aquatic beetles of the family Hydraenidae in the western hemisphere: Classification, biogeography and inferred phylogeny. *Quaestiones Entomologicae* 16: 3–554.

- Perkins PD (2001) Hydraenidae Mulsant, 1844. In: Arnett RH Jr, Thomas MC (Eds) American beetles (Volume 1). CRC Press, Boca Raton, London, New York, Washington D.C., 228–232.
- Pu CL (1958) The genus *Ochthebius* Leach of China (Coleoptera, Palpicornia). *Acta Entomologica Sinica* 8: 247–265.
- Rey C (1885) Descriptions de coléoptères nouveaux ou peu connus de la tribu des Palpicornes. *Annales de la Société Linnéenne de Lyon* 31[1884]: 13–32.
- Satô M (1959) Notes on Japanese *Hydraena* (Coleoptera: Limnebiidae). *Transactions of the Shikoku Entomological Society* 6: 62–64.
- Satô M (1963) A new inter-tidal species of the genus *Ochthebius* Leach from Japan (Coleoptera: Hydraenidae). *Transactions of the Shikoku Entomological Society* 7: 129–132.
- Sharp D (1884) The water beetles of Japan. *The Transactions of the Entomological Society of London* 1884: 439–464.
- Shatrovsky AG (1989) Hydraenidae, Hydrophilidae. In: Ler PA (Ed.) Opredelitel' Nasekomykh Dal'nego Vostoka SSSR v shestitomakh. Volume 3. Zhestkokrylye (Part 1). Nauka, Leningrad, 260–293.
- Slipinski, SA, Leschen RAB, Lawrence JF (2011) Order Coleoptera Linnaeus, 1758. In: Zhang ZQ (Ed.) Animal biodiversity: An outline of higher level classification and survey of taxonomic richness. *Zootaxa* 3148: 203–208.
- Stephens JF (1829) Illustrations of British entomology; or, a synopsis of indigenous insects: containing their generic and specific distinctions; with an account of their metamorphoses, times of appearance, localities, food, and economy, as far as practicable – Mandibulata (Vol. II). Baldwin and Cradock, London, 113–200. [pls 13–15]