

Two new species of *Ismarus* Haliday (Hymenoptera, Ismaridae) from Yunnan, China

Cheng-Jin Yan¹, Yan-Qiong Peng², Hua-Yan Chen³

1 Southern Zhejiang Key Laboratory of Crop Breeding, Wenzhou Vocational College of Science and Technology, Wenzhou, 325006, China

2 CAS Key Laboratory of Tropical Forest Ecology, Xishuangbanna Tropical Botanical Garden, Chinese Academy of Sciences, Mengla 666303, China

2 Key Laboratory of Plant Resources Conservation and Sustainable Utilization, South China Botanical Garden, Chinese Academy of Sciences, Guangzhou 510650, China

Corresponding author: Hua-Yan Chen (huayanc@scbg.ac.cn)

Abstract

The genus *Ismarus* Haliday are rarely collected parasitoids in the small family Ismaridae. In this study, two new species are described from China's Yunnan Province: *Ismarus robustus* Chen & Yan, sp. nov. and *Ismarus unisulcus* Chen & Yan, sp. nov. An updated key to the Chinese species of the genus is provided.

Key words: Diaprioidea, hyperparasitoid, key, new species, taxonomy, wasp

Introduction



Academic editor: Norman Johnson
Received: 15 May 2023
Accepted: 21 July 2023
Published: 10 August 2023

ZooBank: <https://zoobank.org/B8FB8234-B222-4DE5-B142-F976446DD112>

Citation: Yan C-J, Peng Y-Q, Chen H-Y (2023) Two new species of *Ismarus* Haliday (Hymenoptera, Ismaridae) from Yunnan, China. ZooKeys 1174: 207–217. <https://doi.org/10.3897/zookeys.1174.106404>

Ismarus Haliday belongs to the small parasitoid wasp family Ismaridae, with 59 described species worldwide (Masner 1976; Johnson 1992; Liu et al. 2011; Comério et al. 2016; Kolyada and Chemyreva 2016; Kim et al. 2018a, b; Zhang et al. 2021). Previous studies found that species of *Ismarus* are hyperparasitoids of Dryinidae, which are primary parasitoids of leafhoppers, planthoppers and treehoppers (Chambers 1955, 1981; Nixon 1957; Wall 1967; Kozlov 1971; Masner 1976; Jervis 1979; Tussac and Tussac 1991; Olmi 2000). Studies also suggested that species of *Ismarus* prefer wooded areas at higher elevations in warmer climatic zones and at low elevations in cooler climatic zones (Masner 1976; Kim et al. 2018a, b), although some species have been found to occur in warm subtropical regions (Zhang et al. 2021).

The Chinese fauna of *Ismarus* has been extensively studied recently and nine species (Table 1) have been recorded (Liu et al. 2011; Kim et al. 2018b; Zhang et al. 2021). In this study, we describe another two new species from the mountainous region of Yunnan Province of southwest China.

Copyright: © Cheng-jin Yan et al.
This is an open access article distributed under terms of the Creative Commons Attribution License ([Attribution 4.0 International – CC BY 4.0](https://creativecommons.org/licenses/by/4.0/)).

Table 1. An updated list of the Chinese species of *Ismarus* with their distribution in China.

Species	Distribution in China	
	Province	Realm
<i>Ismarus apicalis</i> Kolyada & Chemyreva	Jilin	Palearctic
<i>Ismarus areolatus</i> Chen	Guangdong	Indomalayan
<i>Ismarus excavatus</i> Kim & Lee	Jilin	Palearctic
<i>Ismarus halidayi</i> Foerster	Sichuan, Guizhou, Yunnan, Tibet	Indomalayan
	Ningxia	Palearctic
<i>Ismarus longus</i> Liu, Chen & Xu	Yunnan	Indomalayan
<i>Ismarus nigritrochanter</i> Liu, Chen & Xu	Yunnan	Indomalayan
<i>Ismarus paradorsiger</i> Chen	Guangdong	Indomalayan
<i>Ismarus parvicellus</i> Liu, Chen & Xu	Hainan	Indomalayan
<i>Ismarus spinalis</i> Kolyada & Chemyreva	Heilongjiang	Palearctic
<i>Ismarus robustus</i> Chen & Yan, sp. nov.	Yunnan	Indomalayan
<i>Ismarus unisulcus</i> Chen & Yan, sp. nov.	Yunnan	Indomalayan

Material and method

This work is based on the specimens collected by Malaise trap set in Gaoligongshan National Nature Reserve, Yunnan Province, China. All the studied specimens are deposited in the insect collection of the South China Botanical Garden, Chinese Academy of Sciences, Guangzhou, China (SCBG).

Abbreviations and morphological terms used in the text: **A1, A2, ... A12**: antennomere 1, 2, ... 12; **POL**: posterior ocellar line (shortest distance between posterior ocelli); **OOL**: ocular ocellar line (shortest distance between posterior ocellus and compound eye); **T1, T2, ... T8**: metasomal tergite 1, 2, ... 8. Morphological terminology otherwise follows Masner (1976) and Zhang et al. (2021). Species of *Ismarus* from China have been thoroughly reviewed and keyed (Zhang et al. 2021), which provides us a useful tool to identify the species in this study.

Specimens were examined using a Nikon SMZ800N microscope. Images and measurements were made using a Nikon SMZ25 microscope with a Nikon DS-Ri 2 digital camera system. Image plates were post-processed with Adobe Photoshop CS6 Extended.

Taxonomy

Ismarus Haliday, 1835

Ismarus Haliday, 1835: 467. Type species *Cinetus dorsiger* Haliday, 1831, by monotypy.

Entomia Herrich-Schäffer, 1840: 127. Type species *Entomia campanulata* Herrich-Schäffer, 1840, by monotypy.

Agonophorus Dahlbom, 1858: 289. Type species *Ismarus rugulosus* Förster, 1850, designated by Muesebeck (1972).

Diagnosis. Low insertion of antennae; antennal shelf not developed; antenna of female 15-merous and male 14-merous; notauli absent; mesoscutual

suprathumeral sulcus absent or present as a single pit or several pits; mesoscutum strongly arched in lateral view; base of second tergite with median furrow (Masner 1976; Zhang et al. 2021).

Key to species of *Ismarus* from China (females)

- 1 Body mostly pale to bright yellow (fig. 12A in Zhang et al. 2021); suture separating T2 and T3 incomplete (fig. 12F in Zhang et al. 2021)
..... *Ismarus paradorsiger* Chen
- Body mostly black or dark brown (Figs 1A, 3A); suture separating T2 and T3 complete (Figs 1F, 3F) 2
- 2 Posterior surface of mesoscutellum areolate (fig. 2C in Zhang et al. 2021); lateral pronotal area with an oblique submedian carina (fig. 2D in Zhang et al. 2021) *Ismarus areolatus* Chen
- Posterior surface of mesoscutellum smooth (Figs 1C, 3C); lateral pronotal area without carina (Figs 1D, 3D) 3
- 3 Radial cell of fore wing distinctly shorter than marginal vein (figs 8A, 10A, 16A in Zhang et al. 2021) 4
- Radial cell of fore wing slightly shorter or as long as marginal vein (Figs 2A, 4A) 6
- 4 Mesoscutal suprathumeral sulcus present as a single pit that is longer than length of the mesoscutellar disc (fig. 15C in Zhang et al. 2021); radial cell of fore wing 0.3× length of marginal vein (fig. 16A in Zhang et al. 2021)....
..... *Ismarus parvicellus* Liu, Chen & Xu
- Mesoscutal suprathumeral sulcus present as four or five foveae of varying size (figs 7C, 9C in Zhang et al. 2021); anterior mesoscutellar pit distinctly shorter than length of the mesoscutellar disc (figs 7C, 9C in Zhang et al. 2021); radial cell of fore wing 0.6× length of marginal vein (figs 8A, 10A in Zhang et al. 2021) 5
- 5 Second flagellomere 5.0× as long as wide (fig. 8B in Zhang et al. 2021); radial cell of fore wing 3.0× as long as high (fig. 8A in Zhang et al. 2021); antenna black with scape brown, pedicel and first flagellomere dark brown (fig. 8B in Zhang et al. 2021); all trochanters brown (fig. 7A in Zhang et al. 2021) *Ismarus longus* Liu, Chen & Xu
- Second flagellomere 3.5× as long as wide (fig. 10B in Zhang et al. 2021); radial cell of fore wing 2.0× as long as high (fig. 10A in Zhang et al. 2021); antenna uniformly black (fig. 10B in Zhang et al. 2021); all trochanters black (fig. 9A in Zhang et al. 2021)..... *Ismarus nigritrochanter* Liu, Chen & Xu
- 6 Antenna uniformly bright yellow or only A15 brown (figs 22, 23 in Kolyada and Chemyreva 2016) 7
- Antenna not bright yellow, variable (Figs 2B, 4B) 8
- 7 Antenna uniformly bright yellow, including A15 (fig. 22 in Kolyada and Chemyreva 2016); anterior mesoscutellar pit with median keel; radial cell of fore wing as long as marginal vein (fig. 17 in Kolyada and Chemyreva 2016) *Ismarus spinalis* Kolyada & Chemyreva
- Antenna bright yellow, except A15 brown (fig. 23 Kolyada and Chemyreva 2016); anterior mesoscutellar pit without median keel; radial cell of fore wing 0.8× length of marginal vein (fig. 12 Kolyada and Chemyreva 2016)
..... *Ismarus apicalis* Kolyada & Chemyreva

- 8 Mesoscutal suprathumeral sulcus present as six small pits (Fig. 1C); antenna entirely black (Fig. 2B) *Ismarus robustus* Chen & Yan, sp. nov.
- Mesoscutal suprathumeral sulcus present as a single anterior pit (Fig. 3C); antenna brown to dark brown (Fig. 4B) 9
- 9 Median furrow of T2 long, half the length of T2 (fig. 4E in Zhang et al. 2021) *Ismarus halidayi* Förster
- Median furrow of T2 very short, distinctly less than half the length of T2 (Fig. 3E) 10
- 10 Base of T2 with several short costae forming several furrows; A10–A15 dark brown (fig. 5A in Kim et al. 2018b) *Ismarus excavatus* Kim & Lee
- Base of T2 with two costae forming a single furrow (Fig. 3E); A10–A15 black (Fig. 4B) *Ismarus unisulcus* Chen & Yan, sp. nov.

***Ismarus robustus* Chen & Yan, sp. nov.**

<https://zoobank.org/CC12F866-7B9F-4DB3-8426-AA29B70309A9>

Figs 1, 2

Material examined. **Holotype.** CHINA•1♀; Yunnan, Gaoligongshan National Nature Reserve, Dulong River, GLG12; 27°53'51.96"N, 98°20'11.89"E, 1496 m; May–Jun. 2020; Lang Yi leg.; Malaise trap; SCBG 3044338.

Diagnosis. This species can be easily distinguished from other *Ismarus* species by the following characters: largely black; mesoscutal suprathumeral sulcus present as six small pits; mesoscutellum with posterior rim excavate and slightly prominent posterolateral corners; hind tibia abruptly incrassate.

Description. Female. Body length 3.33 mm.

Colour. Body black; antenna entirely black; coxae, trochanters and basal femora of fore and mid legs dark, remainder brown to yellow, with tibia and tarsi becoming paler distally, hind leg mostly black with hind tibia laterally yellow-brown and hind tarsi pale yellow; wings hyaline, veins brown to black-brown.

Head. Head 2.0× as wide as long in dorsal view; vertex abruptly sloping behind ocelli in lateral view; POL as long as OOL; most of frons with scattered setae, except densely setose ventro-laterally; transverse facial carina convex ventrally; A3 as long as A4; A4 1.3× length of A5; A6–A14 with each segment approximately 1.4× longer than wide; A15 approximately 2.5× longer than wide.

Mesosoma. Dorsal pronotal area punctate and setose; lateral pronotal area rugose-punctate ventrally, smooth dorsally; mesoscutum smooth, shiny and convex; mesoscutal suprathumeral sulcus present as six small pits; mesoscutal humeral sulcus deep and finely crenulate, 1.6× length of tegula; mesoscutellum convex with scattered punctae, posterior rim excavate with slightly prominent posterolateral corners; anterior mesoscutellar pit large and deep, as long as the length of the mesoscutellar disc, distinctly crenulate medially, rugose-punctate posteriorly, median keel strong; mesopleuron smooth, with area below tegula rugulose; metapleuron rugose-punctate and covered with dense, long whitish setae.

Wings. Radial cell completely closed, moderately large, 5.0× as long as wide and 0.7× as long as marginal vein.

Legs. Fore and mid legs slender; hind tibia abruptly incrassate, its maximum width slightly wider than hind femur.




Figure 1. *Ismarus robustus* Chen & Yan, sp. nov., holotype, female (SCBG 3044338) **A** lateral habitus **B** head, anterior view **C** head and mesosoma, dorsal view **D** head and mesosoma, lateral view **E** apex of propodeum and basal metasoma, dorsal view **F** posterior half of T2 and T3 to T7, dorsal view.

Metasoma. Petiole slightly shorter than wide (8:9), with irregular longitudinal costae dorsally; tergites smooth with scattered fine punctures; base of T2 with several short costae and short median furrow, extending 0.27× length of T2; sutures between tergites complete and deeply impressed.

Male. Unknown.

Etymology. Named after the comparatively robust body of this species.

Distribution. China (Yunnan).



Figure 2. *Ismarus robustus* Chen & Yan, sp. nov., holotype, female (SCBG 3044338) **A** fore wing **B** antenna **C** hind tibia.

***Ismarus unisulcus* Chen & Yan, sp. nov.**

<https://zoobank.org/25B2BB9E-00E3-44AC-B9BB-CB6B5A7F1D60>

Figs 3, 4

Material examined. **Holotype.** CHINA•1♀; Yunnan Province, Gaoligongshan National Nature Reserve, Dulong River, GLG13; 27°50'55.81"N, 98°28'3.15"E, 2824 m; 2–15 Jul. 2020; Lang Yi leg.; Malaise trap; SCBG 3044337. **Paratype.** CHINA•1♀; same locality as holotype, but 2–16 Jul. 2020; Lang Yi leg.; Malaise trap; SCBG 3049369.




Figure 3. *Ismarus unisulcus* Chen & Yan, sp. nov., holotype, female (SCBG 3044337) **A** lateral habitus **B** head, anterior view **C** head and mesosoma, dorsal view **D** head and mesosoma, lateral view **E** apex of propodeum and basal metasoma, dorsal view **F** apical metasoma, dorsal view.




Figure 4. *Ismarus unisulcus* Chen & Yan, sp. nov., holotype, female (SCBG 3044337) **A** fore wing **B** antenna **C** hind tibia.

Diagnosis. This species is most similar to *I. halidayi* Förster but can be distinguished by the following characters: A4 slightly longer than A3 (A4 shorter than A3 in *I. halidayi*); median furrow of T2 very short, distinctly less than half the length of T2 (median furrow of T2 long, reaching half the length of T2 in *I. halidayi*); radial cell of fore wing 0.75× as long as marginal vein (radial cell as long as marginal vein in *I. halidayi*).

Description. Female. Body length 2.50–2.64 mm.

Colour. Body black; A1–A9 brown to dark brown, remainder of antenna dark brown; fore and mid legs yellow-brown, with tarsi becoming darker distally, hind leg mostly dark brown with basal coxae somewhat dark and hind femur, trochanter and basal tibia yellow-brown; tegulae yellow-brown; wings hyaline, veins light brown to black-brown.

Head. Head 2.0× as wide as long in dorsal view; vertex abruptly sloping behind ocelli in lateral view; POL as long as OOL; most of frons with scattered setae, except densely setose ventro-laterally; transverse facial carina slightly convex ventrally; A4 slightly longer than A3; A4 1.5× length of A5; A6–A14 with each segment less than 1.5× as long as wide; A15 approximately 2.0× longer than wide.

Mesosoma. Dorsal pronotal area rugose-punctate and setose; lateral pronotal area rugose-punctate ventrally, smooth dorsally; mesoscutum smooth, shiny and convex, posterior margin with scattered long setae; mesoscutal suprakumeral sulcus present as a single anterior pit; mesoscutal humeral sulcus as long as tegula, deep and crenulate; mesoscutellum smooth and slightly convex, posterior rim rounded; anterior mesoscutellar pit large and deep, shorter than length of the mesoscutellar disc, sparsely punctate posteriorly, median keel weakly defined; mesopleuron smooth, with area below tegula rugulose; metapleuron rugose-punctate and covered with dense, long whitish setae.

Wings. Radial cell closed, moderately large, 5.6× as long as wide and 0.94× as long as marginal vein.

Legs. Fore and mid legs slender; hind tibia gradually swollen, its maximum width slightly wider than hind femur.

Metasoma. Petiole slightly shorter than wide (8:9), with strong costae dorsally; tergites smooth with scattered fine punctures; base of T2 with two short costae and a short median furrow, extending 0.37× length of T2; sutures between tergites complete and deeply impressed.

Male. Unknown.

Etymology. The name refers to the single furrow present on the base of T2.

Distribution. China (Yunnan).

Acknowledgments

We thank Zachary Lahey (USDA-ARS, U.S. Vegetable Laboratory) for improving the manuscript and Sufeng Pan (Wenzhou Vocational College of Science and Technology, Wenzhou) for handling the photographs.

Additional information

Conflict of interest

The authors have declared that no competing interests exist.

Ethical statement

No ethical statement was reported.

Funding

This material is based upon work supported in part by the Insect Network of Sino-BON and the National Natural Science Foundation of China (32100360).

Author contributions

Conceptualization and methodology, all authors; resources, Yan-Qiong Peng; writing—original draft, Chengjin Yan and Hua-Yan Chen; writing—review and editing, all authors; supervision, Chengjin Yan and Hua-Yan Chen; funding acquisition, Chengjin Yan and Yan-Qiong Peng. All authors have read and agreed to the published version of the manuscript.

Author ORCIDs

Cheng-Jin Yan  <https://orcid.org/0000-0001-5723-9635>

Yan-Qiong Peng  <https://orcid.org/0000-0002-7453-9119>

Hua-Yan Chen  <https://orcid.org/0000-0002-0382-1635>

Data availability

All of the data that support the findings of this study are available in the main text.

References

- Chambers VH (1955) Some hosts of *Anteon* spp. (Hym. Dryinidae) and a hyperparasite *Ismarus* (Hymenoptera. Belytidae). Entomologist's Monthly Magazine 91: 114–115.
- Chambers VH (1981) A host for *Ismarus halidayi* Foerst. (Hym., Diapriidae). Entomologist's Monthly Magazine 117: 1–29.
- Comério EF, Perioto NW, Lara RIR (2016) A new species of *Ismarus* Haliday (Hymenoptera: Ismaridae) from Brazil and a New Occurrence Record for *Ismarus gracilis* Masner. EntomoBrasilis 9(3): 197–201. <https://doi.org/10.12741/ebrasilis.v9i3.589>
- Dahlbom AG (1858) Svenska små-ichneumonernas familjer och slægten. Öfversigt af Kongliga Vetenskaps-Akademiens Förfärlingar 14: 289–298.
- Haliday AH (1835) Essay on parasitic Hymenoptera of the Ichneumones Adsciti. Entomological Magazine 2: 458–468.
- Herrich-Schäffer GAW (1840) Nomenclator entomologicus. Verzeichniss der europäischen Insecten; zur Erleichterung des Tauschverkehrs mit Preisen versehen. Zweites Heft. Coleoptera, Orthoptera, Deratoptera und Hymenoptera. Friedrich Pustet, Regensburg.
- Jervis MA (1979) Parasitism of *Aphelopus* species (Hymenoptera: Dryinidae) by *Ismarus dorsiger* (Curtis) (Hymenoptera: Diapriidae). Entomologist's Gazette 30: 127–129.
- Johnson NF (1992) Catalog of world Species of Proctotrupoidea, exclusive of Platygastidae (Hymenoptera). Memoirs of the American Entomological Institute 51: 1–825.
- Kim CJ, Copeland RS, Notton DG (2018a) The family Ismaridae Thomson (Hymenoptera, Diaprioidea): First record for the Afrotropical region with description of fourteen new species. African Invertebrates 59(2): 127–163. <https://doi.org/10.3897/afrinvertebr.59.24403>
- Kim CJ, Notton DG, Ødegaard F, Lee JW (2018b) Review of the Palaearctic species of Ismaridae Thomson, 1858 (Hymenoptera: Diaprioidea). European Journal of Taxonomy 417(417): 1–38. <https://doi.org/10.5852/ejt.2018.417>

- Kolyada VA, Chemyreva VG (2016) Revision of species of the genus *Ismarus* Haliday, 1835 (Hymenoptera: Diaprioidea: Ismaridae) of the Russian fauna. Far Eastern Entomologist = Dal'nevostochnyi Entomolog 318: 1–19.
- Kozlov MA (1971) Proctotrupoids (Hymenoptera, Proctotruipoidea) of the USSR. Trudy Vsesoyuznogo Entomologicheskogo Obshchestva 54: 3–67.
- Liu J, Chen H, Xu Z (2011) Notes on the genus *Ismarus* Haliday (Hymenoptera, Diapriidae) from China. ZooKeys 108: 49–60. <https://doi.org/10.3897/zookeys.108.768>
- Masner L (1976) A revision of the Ismarinae of the New World (Hymenoptera, Proctotruipoidea, Diapriidae). Canadian Entomologist 108(11): 1243–1266. <https://doi.org/10.4039/Ent1081243-11>
- Muesebeck CFW (1972) On the identity of *Agonophorus* Dahlbom (Hymenoptera: Diapriidae). Proceedings of Entomological Society of Washington 74: e131.
- Nixon GEL (1957) Hymenoptera, Proctotruipoidea, Diapriidae subfamily Belytinae. Handbooks for the Identification of British Insects 8(3dii): 1–107.
- Olmi M (2000) Bio-ecologia degli Imenotteri Driinidi e loro impiego in programmi di lotta biologica. In: Lucchi A (Ed.) La Metcalfa negli ecosistemi italiani. ARSIA, Firenze, 93–117.
- Tussac H, Tussac M (1991) Récapitulatif d'une collecte de Dryinidae et Diapriidae (Hym. Chrysidoidea et Proctotrypoidea). Entomologiste 47(4): 189–194.
- Wall I (1967) Die Ismarinae und Belytinae der Schweiz. Entomologische Abhandlungen 35: 123–265.
- Zhang X, Chen H, Liu J, Luo S (2021) The genus *Ismarus* Haliday (Hymenoptera, Ismaridae) from China. Journal of Hymenoptera Research 82: 139–160. <https://doi.org/10.3897/jhr.82.62148>