

Amplicon metagenomics of dung beetles (Coleoptera, Scarabaeidae, Scarabaeinae) as a proxy for lemur (Primates, Lemuroidea) studies in Madagascar

Andrey V. Frolov¹, Lilia A. Akhmetova¹, Maria S. Vishnevskaya^{1,2}, Bogdan A. Kiriukhin³, Olivier Montreuil⁴, Fernando Lopes⁵, Sergei I. Tarasov⁵

¹ Zoological Institute, Russian Academy of Sciences, Saint Petersburg, Russia

² Department of Entomology, Saint Petersburg State University, Saint Petersburg, Russia

³ AquaBioSafe Laboratory, University of Tyumen, Tyumen, Russia

⁴ National Museum of Natural History, Paris, France

⁵ Finnish Museum of Natural History, University of Helsinki, Helsinki, Finland

Corresponding author: Andrey V. Frolov (afrolov@zin.ru)



Academic editor: Christopher Majka

Received: 4 June 2023

Accepted: 7 September 2023

Published: 28 September 2023

ZooBank: <https://zoobank.org/980DD85A-4DC9-4667-8B18-CDD3A39120E2>

Citation: Frolov AV, Akhmetova LA, Vishnevskaya MS, Kiriukhin BA, Montreuil O, Lopes F, Tarasov SI (2023) Amplicon metagenomics of dung beetles (Coleoptera, Scarabaeidae, Scarabaeinae) as a proxy for lemur (Primates, Lemuroidea) studies in Madagascar. ZooKeys 1181: 29–39. <https://doi.org/10.3897/zookeys.1181.107496>

Copyright: © Andrey V. Frolov et al.

This is an open access article distributed under terms of the Creative Commons Attribution License ([Attribution 4.0 International – CC BY 4.0](https://creativecommons.org/licenses/by/4.0/)).

Abstract

Dung beetles (Scarabaeidae, Scarabaeinae) are among the most cost-effective and informative biodiversity indicator groups, conveying rich information about the status of habitats and faunas of an area. Yet their use for monitoring the mammal species, that are the main providers of the food for the dung beetles, has only recently been recognized. In the present work, we studied the diet of four endemic Madagascan dung beetles (*Helictopleurus fissicollis* (Fairmaire), *H. giganteus* (Harold), *Nanos agaboides* (Boucomont), and *Epilissus splendidus* Fairmaire) using high-throughput sequencing and amplicon metagenomics. For all beetle species, the $\frac{2}{3}$ – $\frac{3}{4}$ of reads belonged to humans, suggesting that human feces are the main source of food for the beetles in the examined areas. The second most abundant were the reads of the cattle (*Bos taurus* Linnaeus). We also found lower but significant number of reads of six lemur species belonging to three genera. Our sampling localities agree well with the known ranges of these lemur species. The amplicon metagenomics method proved a promising tool for the lemur inventories in Madagascar.

Key words: Beetles, coprophagy, gut content analysis, Madagascar, next-generation sequencing, scarabaeines

Introduction

Madagascar is known for its unique biota characterized by an exceptionally high level of endemism at all taxonomic levels (Vences et al. 2009). Species-level endemism reaches 100% in many taxa, and there are many families and tribes that are found only in Madagascar. For example, all native amphibians and land mammals in Madagascar are endemic (Goodman and Benstead 2003). Due to its exceptional endemism and generally high species diversity, Madagascar is considered one of the most important biodiversity hotspots in the world (Myers et al. 2000), yet its biodiversity is highly underestimated (Vieites et al. 2009).

Dung beetles are a marquee focal group for global efforts to assess the status of biodiversity. They are among the most cost-effective and informative biodiversity indicator groups (Spector 2006; Nichols and Gardner 2011). Dung beetles provide rich information about the status of habitats and faunas of an area (Davis et al. 2004; Nichols et al. 2008; Scholtz et al. 2009). In Madagascar, they are important elements in forest food chains and ecosystems where they originally evolved to decompose lemur excrements (Orsini et al. 2007; Wirta and Montreuil 2008; Wirta et al. 2008; Viljanen et al. 2010; Wirta et al. 2010; Montreuil and Viljanen 2022). Increasing anthropogenic pressure reduces forest habitats, where bulk of dung beetles occurs, and population of lemurs, the original producers of food for dung beetles. This forces dung beetles to switch to other food sources including human feces and cattle dung. This drastically affects distribution, population size, and survival of dung beetles, and leads to a global rewiring in tropical food chains (Hanski et al. 2008; Rahagalala et al. 2009; Wirta et al. 2014).

Recently, attempts have been made to use DNA metabarcoding with High-Throughput, or Next-Generation, sequencing (NGS) to identify taxa in material samples, such as plant species in herbivore dung, prey species in the gut contents of predators, or soil samples (Taberlet et al. 2012; Kocher et al. 2017). Animal manure includes not only the DNA of the food, but also the DNA of the animal producing the manure (Shehzad et al. 2012; De Barba et al. 2014; Lopes et al. 2023). Three studies have been published using NGS to identify mammalian food sources for coprophagous beetles. Gillett et al. (2016) tested the intestinal contents of 10 specimens of different species of coprophagous beetles collected using window traps in the Mbuluzi Game Reserve, eastern Swaziland. In these samples, they found DNA from seven species of mammals, including two species of cattle, domestic mice, and humans. Kerley et al. (2018) used the flightless beetle *Circellium bacchus* (Fabricius) to study changes in its diet, identify beetle food sources using DNA metabarcoding, and compare with published larval feeding data. Traditionally, *C. bacchus* was thought to specialize in the dung of large herbivores as a food source for both larvae and adult beetles. Kerley et al. (2018) extracted mammalian DNA from 151 adult *C. bacchus* fecal samples and sequenced them from 16 mammalian species (ranging from elephants to small rodents), many of which were not yet known as a food source for this species. This approach also confirmed the presence of unknown taxa of mammals in the study area. These authors suggested that the data obtained could be used to study and monitor the biodiversity of mammals. Mouse-like rodent feces were the most common food source (77.5%) for adult *C. bacchus*, which differed markedly from dung-bearing large herbivores used to raise larvae. These results support the hypothesis of changes in the use of food resources in dung beetles associated with the life cycle and reveal a previously unknown but ecologically important role of these beetles in the disposal of rodent feces. Drinkwater et al. (2021) studied gut content of large dung-beetle species and communities of smaller beetles in lowland tropical rainforest in Sabah, Malaysia. They successfully identified six mammalian species, including bearded pig (*Sus barbatus* Müller) and sambar deer (*Rusa unicolor* (Kerr)).

The main goal of the present study was to evaluate the amplicon metagenomics methods to find out what are the major food providers for a few model

Madagascan dung-beetle species. We did not aim at studying the complete diet of these species, nor did we study differences in the diet among individuals or groups. Instead, the study was designed to include specimens of different taxa collected in a few different areas where both native and introduced animals are available as potential food producers for dung beetles. We were specifically interested in finding sequences of lemurs (Lemuroidea) because of their importance for nature conservation in Madagascar. For preparation of amplicon libraries, we used the primers for the 16S rRNA gene fragment, successfully utilized by Ji et al. (2022) to study trophic associations of leeches. Another primer pair, used by Ji et al. to amplify 12S rRNA gene fragment, was tested but not adopted, because, being highly degenerate, it is less specific for mammalian taxa and is not species-specific for some lemur species.

Materials and methods

Materials and collection methods

Examined material is housed in the collection of Muséum national d'Histoire naturelle, Paris (MNHN), and Zoological Institute, Saint Petersburg, Russia (ZIN). Beetles were collected in four localities in northern (Montagne d'Ambre National Park and Montagne d'Ambre Nature Reserve) and central (Ankaratra Andraraty Forest Reserve and Andasibe-Mantadia National Park) Madagascar (Fig. 1):




Figure 1. Madagascan Scarabaeidae dung beetles, general view of beetles (left, not to scale) and sampling localities map (right) **1** *Helictopleurus fissicollis* (Fairmaire) **2** *Nanos agaboides* **3** *H. giganteus* (Harold) **4** *Epilissus splendidus* Fairmaire.

Helictopleurus fissicollis (Fairmaire). Madagascar • 12 females and 5 males (ZIN), Montagne d'Ambre National Park, 12°31'29"S, 49°10'19"E, forest, 5–10.02.2022, A.V. Frolov leg.

Helictopleurus giganteus (Harold). Madagascar • 5 females and 4 males (ZIN), Ankaratra Andraraty, 19°21'20"S, 47°18'18"E PFT, cow dung 23.02.2022, A.V.Frolov leg.

Nanos agabooides Madagascar • 12 females and 6 males (MNHN), Montagne d'Ambre Nature Reserve, 12°28'15"S, 49°13'4"E, forest, 7–10.02.2022, A.V. Frolov leg.

Epilissus splendidus Fairmaire. Madagascar • 2 females and 4 males (MNHN), Andasibe-Mantadia National Park, 18°49'31"S, 48°26'5"E, 18–20.II.2022, A.V. Frolov leg.

The beetles we collected by standard pitfall traps baited with human feces (all except *H. giganteus*) and from cow dung pads (*H. giganteus*). The trap captured *H. fissicollis* contained a preservation solution with SDS and EDTA (Pokluda et al. 2014) and was exposed for 5 days until picking up. In other pitfall traps, we used funnels over the collecting jars, so the beetles attracted to the traps fell into the jars and stayed alive until picking up; these traps were exposed overnight. After picking up, the beetles were placed in containers with 96% ethanol and transported to the laboratory in two or three weeks at room temperature; the alcohol was changed the next day after collecting and after a week. At the laboratory, the beetles were placed in a freezer until dissection.

Gut content extraction

For the analysis, the preserved beetles were dissected under a stereomicroscope. Abdominal tergites were cut with micro-scissors and, if the gut had visible content, it was dissected and placed in Eppendorf micro-tube with 96% ethanol for DNA extraction. Gut content was extracted from eight specimens.

DNA extraction and sequencing

DNA was extracted by phenol-chloroform extraction method (Sambrook and Russell 2006) [*H. giganteus*] or using a diaGene extraction kit (Dia-M, Moscow) according to the manufacturer's protocol. The extracted DNA was quantified using a Qubit fluorimeter 4.0 with high-sensitivity reagents (Lumiprobe QuDye dsDNA HS Assay Kit) and 1 µl of DNA. Four samples from different species with highest concentration of the extracted DNA were selected for high throughput sequencing. For amplicon metagenomics sequencing with the following primer pairs were used: 16Smam1 (5'-CGGTTGGGGTGACCTCGGA-3') and 16Smam2 (5'-GCTGTTATCCCTAGGGTAACT-3') (Taylor 1996). Libraries were prepared using the NEBNext Ultra II DNA Library Prep Kit, checked with Qubit (high-sensitive reagents) and real-time PCR for quantification, and Bioanalyzer for size distribution detection. The amplicon paired-end libraries (PE250) targeting an insert size of 350 bp were sequenced on Illumina NovaSeq 6000 platform aiming for 30K raw tags per sample. DNA extraction was performed at "Chromas" Core Facility, Saint Petersburg State University (Peterhoff, Russia), and library preparation, quality control, and sequencing were performed

at Novogene (Novogene, Cambridge, UK). The data presented in the study are deposited in the NCBI Sequence Read Archive (SRA) database, accession number PRJNA958125.

Bioinformatics methods

Raw reads were trimmed from adapter reads using *bbduk*. *CutAdapt* 4.3 was used to trim primer reads and to discard reads without primers (Martin 2011). Quality filtering was applied with the *Dada2* (Callahan et al. 2016) command «*filterAndTrim*» with total expected errors = 1 for forward reads and 2 for reverse reads. For reads visualization and quality checking *FastQC* tool was used on each of previous steps. Standard *Dada2* pipeline was used for further ASV producing. Taxonomy assignment was carried out using the *BLASTN* (McGinnis and Madden 2004) algorithm using the NCBI nucleotide database as a reference. ASVs with fewer than 10 copies were removed to minimize impact of likely spurious reads. The number of reads of each sample was rarefied by the value of the sample with the smallest number using the *phyloseq* (McMurdie and Holmes 2013) function «*rarefy_even_depth*».

To validate the identification of the hosts, we recovered a maximum-likelihood (ML) phylogeny with the most representative ASVs from each sample and reads obtained from GenBank for closely related species under the same genera. Reads were aligned, manually trimmed, and the ML phylogeny was recovered with *MEGAX* (Kumar et al. 2018).

Results

We have successfully amplified 16S rDNA marker from all samples with a total 363558 reads; 326522 reads passed through quality filtering and preprocessing procedures. Filtered reads were denoised and formed 150 ASVs. Taxonomy assignments produced by *BLASTN* were manually checked to address possible “overclassification” due to similarity of the marker (for example in Hominidae species). The results are summarized in Table 1.

The total number of mammalian taxa identified includes 15 species with most of them belonging to primates (7 species including human). Small number of reads was classified as belonging to domestic animals (dog), and may also be considered contaminations, although dogs may occur in the studied areas. The *Nanos agaboides* sample yielded reads of *Arvicolinae* rodents, not occurring in Madagascar. The *H. fissicollis* sample yielded reads of *Bovidae*, having high but not 100% similarity with a number of ruminant taxa. The *H. giganteus* sample yielded reads of Norway lemming (*Lemmus lemmus* (Linnaeus)), which does not occur in Madagascar and is apparently a contamination (this sample was processed separately in the DNA extraction laboratory). All but *H. giganteus* samples yielded reads of northern mole vole (*Ellobius talpinus* (Pallas)). We consider these reads as contaminations, because these animals do not occur in Madagascar but they were studied in the past years in the molecular laboratory where the DNA was extracted.

The mammal species that we think were true sources of the food for the examined beetle species (Fig. 2) are discussed below.

Table 1. Results of the amplicon metagenomic analysis of the gut content of four Madagascan Scarabaeidae dung beetle species.

Mammal species	<i>Nanos agaboides</i>	<i>Helictopleurus fissicollis</i>	<i>Epilissus splendidus</i>	<i>Helictopleurus giganteus</i>
<i>Homo sapiens</i>	55074	44855	80293	56315
<i>Eulemur coronatus</i>	0	329	0	0
<i>Eulemur fulvus</i>	0	0	28	63
<i>Eulemur rubriventer</i>	9	13	1161	0
<i>Eulemur sanfordi</i>	2853	14803	0	0
<i>Hapalemur griseus</i>	0	0	212	0
<i>Propithecus diadema</i>	3	6	425	0
<i>Bos taurus</i>	13224	3895	12481	28376
Bovidae	0	166	0	1
<i>Canis lupus</i>	0	93	0	2
<i>Sus scrofa</i>	256	700	323	427
Arvicolinae	912	0	0	4
<i>Ellobius talpinus</i>	427	223	1180	6
<i>Lemmus lemmus</i>	8	7	0	3907
<i>Rattus norvegicus</i>	412	189	1	2860




Figure 2. Results of the amplicon metagenomic analysis of the gut content of four Madagascan Scarabaeidae dung-beetle species (contamination reads are excluded). Left, number of effective tags per beetle sample. Right, proportion of effective tags per beetle sample, excluding human and cattle sequences.

Discussion

The major challenge in studying dung-beetle gut content by molecular methods is that the DNA of the putative food producers is highly degraded and is generally in low quantity. At the same time, the total DNA is extracted from the samples of the gross guts dissected from the beetles, and it is hardly possible with the beetles of this size to separate gut content from the gut tissue. In addition, the gut

content is normally full of symbiotic microorganisms. Therefore, using shotgun NGS sequencing for this DNA will be economically and computationally suboptimal since the great majority of the reads would belong to non-target organisms.

Our results demonstrated that instead of shotgun approach, the amplicon metagenomic method could be successfully applied to investigate the diet of Madagascan dung beetles and the composition of the local communities of animals that provide food resources to them.

It should be noted that in all samples 2/3–4/5 of the reads belong to humans. Part of the reads might result from the contamination by the bait, although the traps were designed in such a way as to minimize possible contact of the beetles with bait. Even if the beetles manage to consume bait, it may contribute to the total reads but not substitute the reads from the dung consumed before. In addition, since the beetles are fixed immediately or shortly after they arrived to traps, the foodstuff from the bait may not go through digestive system to get to the hindgut. The contamination is also possible during the laboratory work. However, we believe the human contaminations can be small or negligible percent of all reads we encountered.

The second most abundant reads belong to cattle (*Bos taurus* Linnaeus). Not surprisingly, they are the most abundant in the *H. giganteus* sample, where the beetle was collected from a cattle dung pad. But a reasonable amount of reads were also found in the *H. fissicollis* sample from Amber Mountain National Park, collected by pitfall traps a few kilometers inside the park.

A high number of reads of humans (more than 2/3 in our analyses) and cattle, which was similar for different samples, regardless of the taxonomy of the beetles and collecting localities, shows that humans and cattle can be considered as the main food producers at least for some native Madagascar dung beetle taxa.

In all our samples (except for *Epilissus splendidus*) we revealed the sequences of Norway rat (*Rattus norvegicus* (Berkenhout)). However, with the data available we cannot be sure that in all these cases rats were the real food producers for the beetles. It is possible that in case of *H. giganteus* at least a part of sequences might be a result of contamination in the laboratory, which previously processed rodent samples. *Helictopleurus giganteus* sample also yielded reasonable number of lemming reads, which are definitely a result of contamination.

However, the revealed sequences of our main focal taxon, lemurs, cannot be contaminations because no experiments were carried out with Madagascan primates in the involved laboratories. Also, the known and predicted distribution of the six lemur species (Mittermeier et al. 2008; Brown and Yoder 2015; Kamilar and Tecot 2016) agrees well with our sampling localities. Crowned lemur (*Eulemur coronatus* (Gray)) and Sanford's brown lemur (*E. sanfordi* Archbold) found in *H. fissicollis* and *Nanos agaboides* samples, occur in extreme north of Madagascar including the Amber Mountains. Brown lemur (*E. fulvus* (Geoffroy)), red-bellied lemur (*E. rubriventer* (Geoffroy)), eastern lesser bamboo lemur (*Hapalemur griseus* (Link)), and diademed sifaka (*Propithecus diadema* Bennett) have wider ranges throughout much of the eastern Madagascar lowland and mountain rain forests. The later species is classified as Critically Endangered by the International Union for the Conservation of Nature.

Our results demonstrated that Madagascan dung beetles can be considered a promising indirect tool for monitoring lemurs. They can easily be collected by standardized pitfall traps baited with ready available human feces. They can

also be used in other regions, but more research is needed to assess the species-specificity of trophic relationships between beetles and mammals, as well as to evaluate the primers that are most suitable for the target mammal taxa.

With the development of amplicon metagenomics methods, they can be used for more complex analyses in the future. For example, as opposed to lemurs and other mammals, dung beetles are represented in natural history collections in large numbers from numerous localities, and they have been accumulated during decades. Thus, the rich dung beetle collections putatively contain information about occurrence of dung producer species in a particular area, sometimes not existing as a natural habitat any longer.

Conclusions

1. Although the DNA in the gut of dung beetles is highly degraded, even minor mammal dung producers can be reliably identified by amplicon metagenomics methods.
2. Due to high sensitivity of amplicon methods in comparison to shotgun, the issue of contamination should be specifically considered, but in case of qualitative rather than quantitative studies, it does not pose a problem.
3. At present, cattle and especially humans can be the main food producers for some Madagascar dung beetle taxa.
4. Despite the fact that lemur reads may be a small percent of total mammalian reads, they allow to reliably identify lemur species that served as a food source for beetles. The 16S rDNA gene fragment used in the present study showed high specificity being rather small sized.
5. Dung beetles can be a promising indirect tool for monitoring lemurs in Madagascar.

Acknowledgements

We are thankful to the Malagasy Institute of Conservation of Tropical Ecosystems (MICET) for the support in the acquisition of permits and logistic support during the expedition to Madagascar. Michele Rossini and Thomas Merrien (Finnish Museum of Natural History, Helsinki, Finland) helped with fieldwork. Aleš Bezděk (Institute of Entomology ASCR, České Budějovice, Czech Republic) commented the draft manuscript. This research was funded by Russian Science Foundation, grant number 22-24-00715 to A.F. The scarab beetle collection of the Zoological Institute is maintained under the Russian State research program 122031100272-3.

Additional information

Conflict of interest

The authors have declared that no competing interests exist.

Ethical statement

No ethical statement was reported.

Funding

Russian Science Foundation, grant number 22-24-00715.

Author contributions

Conceptualization: A.F., S.T., O.M., L.A.; methodology: A.F., S.T., F.L.; bioinformatics analysis: B.K.; field work: A.F. and S.T.; specimen identification: O.M.; specimen curation: L.A.; sample preparation, M.V.; draft preparation: A.F.; draft review and editing: all authors; project administration: A.F.; funding acquisition: A.F.

Author ORCIDs

Andrey V. Frolov  <https://orcid.org/0000-0002-6724-6828>

Lilia A. Akhmetova  <https://orcid.org/0000-0002-2151-1278>

Maria S. Vishnevskaya  <https://orcid.org/0000-0002-2634-2610>

Bogdan A. Kiriukhin  <https://orcid.org/0000-0001-8840-2976>

Olivier Montreuil  <https://orcid.org/0000-0003-2682-7594>

Fernando Lopes  <https://orcid.org/0000-0002-1246-2777>

Sergei I. Tarasov  <https://orcid.org/0000-0001-5237-2330>

Data availability

The data presented in the study are deposited in the NCBI Sequence Read Archive (SRA) database, accession number PRJNA958125.

References

- Brown JL, Yoder AD (2015) Shifting ranges and conservation challenges for lemurs in the face of climate change. *Ecology and Evolution* 5(6): 1131–1142. <https://doi.org/10.1002/ece3.1418>
- Callahan BJ, McMurdie PJ, Rosen MJ, Han AW, Johnson AJA, Holmes SP (2016) DADA2: High-resolution sample inference from Illumina amplicon data. *Nature Methods* 13(7): 581–583. <https://doi.org/10.1038/nmeth.3869>
- Davis AL, Scholtz CH, Dooley PW, Bham N, Kryger U (2004) Scarabaeine dung beetles as indicators of biodiversity, habitat transformation and pest control chemicals in agro-ecosystems. *South African Journal of Science* 100: 415–424.
- De Barba M, Miquel C, Boyer F, Mercier C, Rioux D, Coissac E, Taberlet P (2014) DNA metabarcoding multiplexing and validation of data accuracy for diet assessment: Application to omnivorous diet. *Molecular Ecology Resources* 14(2): 306–323. <https://doi.org/10.1111/1755-0998.12188>
- Drinkwater R, Williamson J, Clare EL, Chung AY, Rossiter SJ, Slade E (2021) Dung beetles as samplers of mammals in Malaysian Borneo—A test of high throughput metabarcoding of iDNA. *PeerJ* 9: e11897. <https://doi.org/10.7717/peerj.11897>
- Gillett CPDT, Johnson AJ, Barr I, Hulcr J (2016) Metagenomic sequencing of dung beetle intestinal contents directly detects and identifies mammalian fauna. *bioRxiv*. <https://doi.org/10.1101/074849>
- Goodman SM, Benstead JP (2003) *Natural History of Madagascar*. University of Chicago Press, 1709 pp. <https://doi.org/10.7208/chicago/9780226337609.001.0001>
- Hanski I, Wirta H, Nyman T, Rahagalala P (2008) Resource shifts in Malagasy dung beetles: Contrasting processes revealed by dissimilar spatial genetic patterns. *Ecology Letters* 11(11): 1208–1215. <https://doi.org/10.1111/j.1461-0248.2008.01239.x>
- Ji Y, Baker CCM, Popescu VD, Wang J, Wu C, Wang Z, Li Y, Wang L, Hua C, Yang Z, Yang C, Xu CCY, Diana A, Wen Q, Pierce NE, Yu DW (2022) Measuring protected-area effectiveness using vertebrate distributions from leech iDNA. *Nature Communications* 13(1): e1555. <https://doi.org/10.1038/s41467-022-28778-8>

- Kamilar JM, Tecot SR (2016) Anthropogenic and climatic effects on the distribution of Eulemur species: An ecological niche modeling approach. *International Journal of Primatology* 37(1): 47–68. <https://doi.org/10.1007/s10764-015-9875-8>
- Kerley GIH, Landman M, Ficetola GF, Boyer F, Bonin A, Rioux D, Taberlet P, Coissac E (2018) Diet shifts by adult flightless dung beetles *Circellium bacchus*, revealed using DNA metabarcoding, reflect complex life histories. *Oecologia* 188(1): 107–115. <https://doi.org/10.1007/s00442-018-4203-6>
- Kocher A, de Thoisy B, Catzeflis F, Huguin M, Valière S, Zinger L, Bañuls AL, Murienne J (2017) Evaluation of short mitochondrial metabarcodes for the identification of Amazonian mammals. *Methods in Ecology and Evolution* 8(10): 1276–1283. <https://doi.org/10.1111/2041-210X.12729>
- Kumar S, Stecher G, Li M, Knyaz C, Tamura K (2018) MEGA X: Molecular Evolutionary Genetics Analysis across computing platforms. *Molecular Biology and Evolution* 35(6): 1547–1549. <https://doi.org/10.1093/molbev/msy096>
- Lopes F, Rossini M, Losacco F, Montanaro G, Gunter N, Tarasov S (2023) Metagenomics reveals that dung beetles (Coleoptera: Scarabaeinae) broadly feed on reptile dung. Did they also feed on that of dinosaurs? *Frontiers in Ecology and Evolution* 11: e1132729. <https://doi.org/10.3389/fevo.2023.1132729>
- Martin M (2011) Cutadapt removes adapter sequences from high-throughput sequencing reads. *EMBnet.Journal* 17(1): 10–12. <https://doi.org/10.14806/ej.17.1.200>
- McGinnis S, Madden TL (2004) BLAST: At the core of a powerful and diverse set of sequence analysis tools. *Nucleic Acids Research* 32(Web Server): W20–W25. <https://doi.org/10.1093/nar/gkh435>
- McMurdie PJ, Holmes S (2013) phyloseq: An R package for reproducible interactive analysis and graphics of microbiome census data. *PLoS ONE* 8(4): e61217. <https://doi.org/10.1371/journal.pone.0061217>
- Mittermeier RA, Ganzhorn JU, Konstant WR, Glander K, Tattersall I, Groves CP, Rylands AB, Hapke A, Ratsimbazafy J, Mayor MI, Louis Jr EE, Rumpel Y, Schwitzer C, Rasoloarison RM (2008) Lemur diversity in Madagascar. *International Journal of Primatology* 29(6): 1607–1656. <https://doi.org/10.1007/s10764-008-9317-y>
- Montreuil O, Viljanen H (2022) Révision du genre *Epilissus* Dejean, 1836 (Coleoptera: Scarabaeidae: Epilissini). *Annales de la Société Entomologique de France (NS)*: 1–38. <https://doi.org/10.1080/00379271.2022.2049457>
- Myers N, Mittermeier RA, Mittermeier CG, Da Fonseca GA, Kent J (2000) Biodiversity hotspots for conservation priorities. *Nature* 403(6772): 853–858. <https://doi.org/10.1038/35002501>
- Nichols ES, Gardner TA (2011) Dung beetles as a candidate study taxon in applied biodiversity conservation research. In: Simmons LW, Ridsdill-Smith TJ (Eds) *Ecology and Evolution of Dung Beetles*. Blackwell Publishing, Oxford, 267–291. <https://doi.org/10.1002/9781444342000.ch13>
- Nichols E, Spector S, Louzada J, Larsen T, Amezcuita S, Favila M, Network TSR (2008) Ecological functions and ecosystem services provided by Scarabaeinae dung beetles. *Biological Conservation* 141(6): 1461–1474. <https://doi.org/10.1016/j.biocon.2008.04.011>
- Orsini L, Koivulehto H, Hanski I (2007) Molecular evolution and radiation of dung beetles in Madagascar. *Cladistics* 23(2): 145–168. <https://doi.org/10.1111/j.1096-0031.2006.00139.x>
- Pokluda P, Čížek L, Stříbrná E, Drag L, Lukeš J, Novotný V (2014) A goodbye letter to alcohol: An alternative method for field preservation of arthropod specimens and

- DNA suitable for mass collecting methods. *European Journal of Entomology* 111(2): 175–179. <https://doi.org/10.14411/eje.2014.024>
- Rahagalala P, Viljanen H, Hottola J, Hanski I (2009) Assemblages of dung beetles using cattle dung in Madagascar. *African Entomology* 17(1): 71–89. <https://doi.org/10.4001/003.017.0109>
- Sambrook J, Russell DW (2006) Purification of nucleic acids by extraction with phenol: chloroform. *Cold Spring Harbor Protocols* 2006: pdb. prot4455. <https://doi.org/10.1101/pdb.prot4455>
- Scholtz CH, Davis ALV, Kryger U (2009) *Evolutionary Biology and Conservation of Dung Beetles*. Pensoft, Sofia–Moscow, 569 pp.
- Shehzad W, Riaz T, Nawaz MA, Miquel C, Poillot C, Shah SA, Pompanon F, Coissac E, Taberlet P (2012) Carnivore diet analysis based on next-generation sequencing: Application to the leopard cat (*Prionailurus bengalensis*) in Pakistan. *Molecular Ecology* 21(8): 1951–1965. <https://doi.org/10.1111/j.1365-294X.2011.05424.x>
- Spector S (2006) Scarabaeine dung beetles (Coleoptera: Scarabaeidae: Scarabaeinae): an invertebrate focal taxon for biodiversity research and conservation. *Coleopterists Bulletin* 60(sp5): 71–83. [https://doi.org/10.1649/0010-065X\(2006\)60\[71:SD-BCSS\]2.0.CO;2](https://doi.org/10.1649/0010-065X(2006)60[71:SD-BCSS]2.0.CO;2)
- Taberlet P, Coissac E, Pompanon F, Brochmann C, Willerslev E (2012) Towards next-generation biodiversity assessment using DNA metabarcoding. *Molecular Ecology* 21(8): 2045–2050. <https://doi.org/10.1111/j.1365-294X.2012.05470.x>
- Taylor PG (1996) Reproducibility of ancient DNA sequences from extinct Pleistocene fauna. *Molecular Biology and Evolution* 13(1): 283–285. <https://doi.org/10.1093/oxfordjournals.molbev.a025566>
- Vences M, Wollenberg KC, Vieites DR, Lees DC (2009) Madagascar as a model region of species diversification. *Trends in Ecology & Evolution* 24(8): 456–465. <https://doi.org/10.1016/j.tree.2009.03.011>
- Vieites DR, Wollenberg KC, Andreone F, Köhler J, Glaw F, Vences M (2009) Vast underestimation of Madagascar's biodiversity evidenced by an integrative amphibian inventory. *Proceedings of the National Academy of Sciences of the United States of America* 106(20): 8267–8272. <https://doi.org/10.1073/pnas.0810821106>
- Viljanen H, Wirta H, Montreuil O, Rahagalala P, Johnson S, Hanski I (2010) Structure of local communities of endemic dung beetles in Madagascar. *Journal of Tropical Ecology* 26(5): 481–496. <https://doi.org/10.1017/S0266467410000325>
- Wirta H, Montreuil O (2008) Evolution of the Canthonini Longitarsi (Scarabaeidae) in Madagascar. *Zoologica Scripta* 37(6): 651–663. <https://doi.org/10.1111/j.1463-6409.2008.00352.x>
- Wirta H, Orsini L, Hanski I (2008) An old adaptive radiation of forest dung beetles in Madagascar. *Molecular Phylogenetics and Evolution* 47(3): 1076–1089. <https://doi.org/10.1016/j.ympev.2008.03.010>
- Wirta H, Viljanen H, Orsini L, Montreuil O, Hanski I (2010) Three parallel radiations of Canthonini dung beetles in Madagascar. *Molecular Phylogenetics and Evolution* 57(2): 710–727. <https://doi.org/10.1016/j.ympev.2010.08.013>
- Wirta HK, Hebert PDN, Kaartinen R, Prosser SW, Várkonyi G, Roslin T (2014) Complementary molecular information changes our perception of food web structure. *Proceedings of the National Academy of Sciences of the United States of America* 111(5): 1885–1890. <https://doi.org/10.1073/pnas.1316990111>