

An update to the taxonomy of *Serica* MacLeay, 1819 (sensu lato) from China (Coleoptera, Scarabaeidae, Sericinae, Sericini)

Xie Su¹ , Wei Li¹ , Wan-Gang Liu² , Dirk Ahrens³ , Ying-Juan Zheng⁴, Yang Liu¹ 

1 Key Laboratory of Resource Biology and Biotechnology in Western China (Northwest University), Shaanxi Key Laboratory for Animal Conservation, College of Life Science, Northwest University, Taibai North Road 229, Xi'an 710069, China

2 SKLLQG, Institute of Earth and Environment, Chinese Academy of Sciences, Yanxiang Road 97, Yanta District, Xi'an 710061, China

3 Zoologisches Forschungsmuseum A. Koenig, Adenauerallee 127, 53113 Bonn, Germany

4 Chinese Research Academy of Environmental Science, Beijing 100012, China

Corresponding authors: Wan-Gang Liu (liuwangang@ieecas.cn); Dirk Ahrens (ahrens.dirk_col@gmx.de); Yang Liu (liuyangent@nwu.edu.cn)

Abstract

In this paper we update the knowledge on the species of *Serica* McLeay, 1819 (sensu lato) occurring in Yunnan, Sichuan, and Shaanxi provinces, China. Three new species are described: *Serica allonanhuensis* Liu, Ahrens, Li & Su, **sp. nov.**, *S. breviantennalis* Liu, Ahrens, Li & Su, **sp. nov.**, and *S. fengensis* Liu, Ahrens, Li & Su, **sp. nov.**. The key to the species groups and species is updated. The habitus and male genitalia of the new species are illustrated, and a map showing their distribution is provided. New distributional data are given for four species.

Key words: Beetles, chafers, key, new species, new records, Yunnan



Academic editor: Andrey Frolov

Received: 7 July 2023

Accepted: 4 November 2023

Published: 28 November 2023

ZooBank: <https://zoobank.org/601D4A0B-BF21-4FB5-984EC5887EC39227>

Citation: Su X, Li W, Liu W-G, Ahrens D, Zheng Y-J, Liu Y (2023) An update to the taxonomy of *Serica* MacLeay, 1819 (sensu lato) from China (Coleoptera, Scarabaeidae, Sericinae, Sericini). ZooKeys 1185: 181–198. <https://doi.org/10.3897/zookeys.1185.109154>

Copyright: © Xie Su et al.

This is an open access article distributed under terms of the Creative Commons Attribution License ([Attribution 4.0 International – CC BY 4.0](https://creativecommons.org/licenses/by/4.0/)).

Introduction

The tribe Sericini Kirby, 1837 includes more than 200 genera and 4000 species. While the phylogenetic position of Sericini appears now to be resolved, its internal relationships still await refinement with additional sampling of taxa and genetic markers needed (Dietz et al. 2023).

The genus *Serica* was established by Macleay (1819), and the type species, by monotypy, is *Serica brunnea* (Linnaeus, 1758) (Ahrens 1999, 2005a). It is one of the most diverse sericine genera in the Palaearctic Region, with hundreds of species in the temperate regions of Eurasia and higher mountains of Asia (Ahrens 2005a, 2006b, 2007b; Ahrens and Bezděk 2016). Under currently understood, *Serica* is polyphyletic, as shown by a series of molecular phylogenetic studies (Ahrens and Vogler 2008; Liu et al. 2015b; Eberle et al. 2017).

Ahrens (1999, 2005a) established the subgenus *Serica* (*sensu stricto*) for the species most closely related to the type species, *S. brunnea* (Linnaeus, 1758). Molecular evidence suggests that many species with an elongate body shape, trilamellate antennal club, and narrower, longer legs have been placed incorrectly in *Serica* by many authors due to convergence of these characters (Ahrens 1999, 2005a). As a consequence of the often occurring strong homoplasy, many diagnostic characteristics of Sericini have been proven to be misleading for higher classification, including the number of antennomeres which traditionally has been prominently used for the definition of genera of Sericini (Ahrens and Vogler 2008). This had led to many unresolved problems in generic classification, especially in the “*Maladera–Neoserica–Serica*” clade (Eberle et al. 2017). Therefore, many of our recent taxonomic treatments of *Serica*, *Maladera* Mulsant & Rey, 1871, and *Neoserica* Brenske, 1894 divided these genera into compact species groups or subgenera, which are likely to represent monophyletic groups, as a basis for future systematic work (e.g. Ahrens and Fabrizi 2016; Fabrizi et al. 2021). These are generally summarized under “*sensu lato*” in association with the respective genus name, or as subgenera. The subgenus *Serica* (*sensu stricto*) has had a vast radiation in the Asian highlands, mainly the Himalayas, eastern Tibet, and the mountains of the eastern Palaearctic (Ahrens 2007c).

This study is another contribution to a comprehensive taxonomic revision of the Sericini of China and adjacent regions. Previous works have comprised taxonomic revisions of all genera occurring in China to which also already several updates have been added (Ahrens 1996, 2000a, b, c, 2001, 2003a, b, c, d, 2005a, 2005b, 2006a, b, c, 2007a, b, c, 2009, 2011, 2021, 2023a, b; Ahrens and Pacholátko 2003, 2007; Ahrens and Fabrizi 2009a, 2009b, 2011; Liu et al. 2011, 2014a, b, c, d, e, 2015a, 2016, 2017a, 2017b, 2019; Ahrens et al. 2014a, b, c, 2021a, 2021b; Bohacz and Ahrens 2020; Fabrizi et al. 2021; Ahrens et al. 2022, 2023; Zhao and Ahrens 2023). Recently, the species of *Serica* MacLeay, 1819 (*sensu lato*) from China and adjacent areas were revised comprehensively (Ahrens et al. 2022). In this paper, we provide further updates on *Serica* (s.l.) species in China, which include new records and the description of three new species.

Materials and methods

The terminology and methods used for measurements, specimen dissection, and genital preparation follow Ahrens (2004). Data from specimens examined are cited in the text with original label contents given in quotation marks; multiple labels for a single specimen are separated by a “/”. Descriptions and illustrations of new taxa are based on the holotype specimen if not otherwise stated, while the variation of specimens is given separately under “variation”. Male genitalia were glued to a small, pointed card and photographed in both lateral and dorsal views using an Olympus SZ61-TR stereomicroscope with an Mshot MD 50 digital camera. All the parameter adjustment and images acquisition were done in Mshot Image Analysis System v. 1.6.1. To obtain an focused image, several partly focused images were also focus-stacked with this software. The resulting images were subsequently digitally edited with Adobe Photoshop v. 21.0.2. Distribution maps were generated using ArcGIS v. 10.3, based on the geographical coordinates from the labels.

Abbreviations used in the text for collection depositories are as follows:

SENU	Laboratory of Systematic Entomology, Northwest University, Xi'an, Shaanxi
IZAS	Institute of Zoology, Chinese Academy of Sciences, Beijing, China
ZFMK	Zoologisches Forschungsinstitut und Museum A. Koenig, Bonn, Germany

Results

Key to males of *Serica* (*sensu lato*) species of continental South-East Asia

- | | | |
|---|---|---|
| 1 | Dorsal surface unicoloured..... | 2 |
| - | Dorsal surface, particularly elytra, multicoloured..... | 28 |
| 2 | Dorsal surface brown and glabrous..... | 3 |
| - | Dorsal surface darker and generally with erect setae on dorsal surface | 12 |
| 3 | Body reddish to dark brown. Antennal club at least twice as long as remaining antennomeres combined | 4 |
| - | Body yellowish brown. Antennal club at maximum 1.5 times as long as remaining antennomeres combined | 8 |
| 4 | Metatibia long and narrow, ratio width:length 1:4.2. Antenna with 10 antennomeres..... | 5 |
| - | Metatibia long and little wider, ratio width:length 1:3.7 | 6 |
| 5 | Metatibia long and narrow, ratio width:length 1:4.2. Apex of the phallobase strongly bent in lateral view | <i>S. solivaga</i> (Brenske, 1898) |
| - | Metatibia moderately long and wide, ratio width:length 1:3.2. Apex of the phallobase more straight (lateral view) | <i>S. mengsonana</i> Ahrens, Fabrizi & Liu, 2022 |
| 6 | Antenna with 9 antennomeres. Antennal club 5 times as long as remaining antennomeres combined. Basal lobe of the right paramere wide and large..... | <i>S. fashengi</i> Liu & Ahrens, 2014 |
| - | Antenna with 10 antennomeres. Antennal club 4 times as long as remaining antennomeres combined. Basal lobe of the right paramere normal or narrow | 7 |
| 7 | Right paramere narrower, without blunt dorsomedian extension, sharply pointed towards apex in lateral view. Dorsal lobe of left paramere externally reflexed..... | <i>S. falcifera</i> Ahrens & Fabrizi, 2009 |
| - | Right paramere wider, with a blunt dorsomedian extension, moderately pointed towards apex in lateral view. Dorsal lobe of left paramere internally reflexed..... | <i>S. segregata</i> Arrow, 1946 |
| 8 | Dorsal surface dull. Phallobase at right side with an apical lateral apophyses laterally | 9 |
| - | Dorsal surface shiny. Phallobase at right side without apical lateral apophyses laterally | <i>S. longwan</i> Ahrens, Fabrizi & Liu, 2022 |
| 9 | Basal lobe of left paramere extended only basomedially..... | 10 |
| - | Basal lobe of left paramere extended basomedially and distally | <i>S. longula</i> Frey, 1972 |

- 10 Basal lobe of left paramere strongly curved and directed distally
..... *S. tiammushan Ahrens, Fabrizi & Liu, 2022* 11
- Basal lobe of left paramere medially directed 11
- 11 Antennal club distinctly longer than remaining antennomeres combined.
Right paramere moderately dorsally angled
..... *S. daqiaoana Ahrens, Fabrizi & Liu, 2022*
- Antennal club distinctly equal to length remaining antennomeres combined. Right paramere strongly dorsally angled
..... *S. jiankouensis Ahrens, Fabrizi & Liu, 2022*
- 12 Labroclypeus distinctly medially emarginate 13
- Labroclypeus weakly medially emarginate 15
- 13 Dorsal surface shiny. Aedeagus in apical half dorsoventrally flattened
..... *S. excisa* (Frey, 1972)
- Dorsal surface dull. Aedeagus in apical half almost spherical in cross section 14
- 14 Labroclypeus deeply medially emarginate. Lateral margin of pronotum slightly concave *S. shiduensis Ahrens, Fabrizi & Liu, 2022*
- Labroclypeus moderately medially emarginate. Lateral margin of pronotum straight *S. klapperichi* Frey, 1972
- 15 Body more oval, pronotum subequal, a third of elytral length 16
- Body elongate, pronotum subequal, a quarter of elytral length 17
- 16 Antennal club in male long, three times as long as remaining antennomeres combined. Parameres with distinct basal lobes
..... *S. huangbaiyuanensis Ahrens, Fabrizi & Liu, 2022*
- Antennal club in male short, as long as remaining antennomeres combined. Parameres without distinct basal lobes
..... *S. guangnanensis Ahrens, Fabrizi & Liu, 2022*
- 17 Parameres at maximum almost subequal to one-half length of phallobase 18
- Parameres distinctly shorter 19
- 18 Left paramere almost perpendicular to right one; club longer than twice of remaining antennomeres combined *S. lupina* Arrow, 1946
- Left paramere almost parallel to or forming an acute angle with right one; club shorter than twice the length of remaining antennomeres combined *S. breviantennalis* Liu, Ahrens, Li & Su, sp. nov.
- 19 Parameres at maximum subequal to one-third of phallobase length 20
- Parameres at maximum subequal to one-quarter or less of phallobase length 21
- 20 Right paramere curved ventrally in lateral view, without preapical lateral tooth 22
- Right paramere straight in lateral view, with sharp and distinct preapical lateral tooth *S. yini* Ahrens, Fabrizi & Liu, 2022
- 21 Dorsal surface of body densely setose. Left paramere nearly straight, distinctly shorter than right one
..... *S. baishuitaiensis Ahrens, Fabrizi & Liu, 2022*
- Dorsal surface of body sparsely setose. Left paramere strongly curved and bent ventrally below ventral portion of phallobase, almost as long as right paramere *S. paralupina* Ahrens, Fabrizi & Liu, 2022

22	Apex of right paramere sharply pointed	<i>S. gaoligong Ahrens, Fabrizi & Liu, 2022</i>
-	Apex of right paramere rounded or bluntly truncate	23
23	Dorsal pilosity dense.....	24
-	Dorsal pilosity sparse.....	27
24	Left paramere weakly curved	25
-	Left paramere strongly curved. Right paramere weakly curved in dorsal view	<i>S. longipes Ahrens, Fabrizi & Liu, 2022</i>
25	Parameres shorter, at maximum one-sixth as long as phallobase. Left paramere strong, obviously shorter than right one. Right paramere weakly curved in dorsal view.....	<i>S. nanhua Ahrens, Fabrizi & Liu, 2022</i>
-	Parameres longer, at least one-quarter as long as phallobase. Left paramere narrower and slightly shorter than right one. Right paramere strongly curved in dorsal view.....	26
26	Parameres without basal lobes.....	
- <i>S. allonanhuai Liu, Ahrens, Li & Su, sp. nov.</i>	
-	Parameres with long and distinct basal lobes	
- <i>S. fengensis Liu, Ahrens, Li & Su, sp. nov.</i>	
27	Right paramere strongly curved. Left paramere wide, flattened in cross section over almost two-thirds of its entire length	
- <i>S. allolongipes Ahrens, Fabrizi & Liu, 2022</i>	
-	Right paramere weakly curved. Left paramere narrow, sickle-shaped, and almost circular in cross section over almost its entire length	
- <i>S. paralongipes Ahrens, Fabrizi & Liu, 2022</i>	
28	Parameres inserted strongly asymmetric; left paramere much more basally than the right one.....	29
-	Phallobase rather symmetric; parameres inserted almost at the same level	37
29	Right paramere with a narrow basal lobe directed medially	31
-	Right paramere without a narrow basal lobe directed medially. Phallobase narrowed towards apex in dorsal view	30
30	Phallobase concavely dorsoapically excised	<i>S. dathei Ahrens, 2005</i>
-	Phallobase with a process-like median extension at mesoapical margin. Antennal club 1.2 times as long as remaining antennomeres combined ...	
- <i>S. dissimillima Ahrens, Fabrizi & Liu, 2022</i>	
31	Phallobase truncate apex (dorsal view), its dorsal margin straight. Antennal club 1.5 times as long as remaining antennomeres combined ...	
- <i>S. nanjiangana Ahrens, 2005</i>	32
-	Phallobase concavely dorsoapically excised	33
32	Left paramere more than one-half length of right paramere	
- <i>S. zhenba Ahrens, Fabrizi & Liu, 2022</i>	
-	Left paramere less than one-half length of right paramere	
- <i>S. nanjiangana Ahrens, 2005</i>	
33	Right paramere very long and narrow, sharply pointed. Left paramere without any lateral tooth	<i>S. trapezicollis Ahrens, 2005</i>
-	Right paramere moderately long and wider, rounded at apex	34
34	Left paramere about one-third length of right paramere	35
-	Left paramere more than one-half length of right paramere	36

- 35 Right paramere abruptly narrowed behind middle. Left paramere not dorsoventrally flattened..... *S. jani Ahrens, Fabrizi & Liu, 2022*
- Right paramere evenly narrowed towards apex. Left paramere dorsoventrally flattened..... *S. jaroslavi Ahrens, Fabrizi & Liu, 2022*
- 36 Left paramere in distal portion with a lateral tooth.....
..... *S. albisetis Ahrens, 2005*
- Left paramere without lateral apical tooth, but its left paramere having its dorsomedial tooth at basal third ..
..... *S. shengtangshan Ahrens, Fabrizi & Liu, 2022*
- 37 Aedeagus robust, with a long double dorsal phallobasal process which exceeds almost parameres *S. adspersa Frey, 1972*
- Aedeagus elongate, without dorsal phallobasal process, but simply concavely emarginate between parameres 38
- 38 Elytral surface with moderately dense to dense setae 39
- Elytral surface with sparse setae 40
- 39 Pilosity of dorsal surface double, with at least two types of setae. Elytra with a large dark spot before apex..... *S. nigroguttata group*
- Pilosity of dorsal surface simple, with one type of setae only. Elytra without dark spot before apex. Metasternum anteriorly abruptly and strongly elevated..... *S. velutina group*
- 40 Parameres both subdivided into a long dorsal and ventral lobe. Metafemur, submarginal serrate line beside anterior margin present or absent....
..... *Serica (subgenus Taiwanoserica Nomura, 1974)*
- Parameres not subdivided into a long dorsal and ventral lobe, or only one of both subdivided into a long dorsal and ventral lobe. Metafemur, submarginal serrate line beside anterior margin always absent..... 41
- 41 Pronotum almost glabrous, with a few adpressed setae on posterior disc; surface evenly plane, in anterior part shiny, posteriorly dull. Frons shiny in anterior half..... *S. ziqingi Ahrens, Fabrizi & Liu, 2022*
- Pronotum densely setose on anterior disc, surface with shallow transverse impressions, entirely dull or iridescent shiny. Frons entirely dull 42
- 42 Pronotum narrow. Body more elongate. Parameres shorter, left one half as long as right..... *S. jirii Ahrens, Fabrizi & Liu, 2022*
- Pronotum more transverse. Body oblong and less convex. Parameres long, equal in length *S. pangwa Ahrens, Fabrizi & Liu, 2022*

Species descriptions

Serica breviantennalis Liu, Ahrens, Li & Su, sp. nov.

<https://zoobank.org/09579AA9-C713-4E38-8CBD-921A41747FCF>

Figs 1A–H, 3

Type materials. **Holotype:** ♂ [China] “Black Tiger Mountain, Tianxin Township, Wuding County, Chuxiong, Yunnan Province, 20.V.2022, light trap, leg. Zhao Yuchen, Lu Jinbo” (SENU). **Paratypes:** 28 ♂♂ [China] “Black Tiger Mountain, Tianxin Township, Wuding County, Chuxiong, Yunnan Province, 20.V.2022, light trap, leg. Zhao Yuchen, Lu Jinbo” (SENU, IZAS, ZFMK), 2 ♂♂ [China] “Gaojian-chou village, Yao ‘an County, Chuxiong, Yunnan Province, 24.V.2022, light trap, leg. Xu Rentao” (SENU), 2 ♂♂ [China] “Xiaogupu Village, Shangdianliang Road,

Wuding County, Chuxiong Autonomous Prefecture, Yunnan Province, 21.V.2022, light trap, 2092.6m, 25.6724°N, 102.2782°E, leg. Lu Jinbo" (SENU), 1 ♂ [China] "Zhangjiacun Mixing Town, Yao 'an County, Chuxiong City, Yunnan Province, 23.V.2022, light trap, 25.4041°N, 101.1683°E" (SENU).

Description of the holotype. Length 7.9 mm, elytral length 5.6 mm, width 4.1 mm. Body oblong, reddish brown; frons dark reddish brown, dull; legs yellowish brown; antenna yellow, with moderately long setae on head dorsal surface.

Labroclypeus wide, trapezoidal, with lateral margins moderately convex and convergent, producing a blunt angle with ocular canthus; anterior angles strongly convex; anterior margin distinctly emarginate medially; margins moderately reflexed; surface moderately convex, finely and densely punctate, with numerous long and erect setae; frontoclypeal suture distinct, moderately curved; ocular canthus moderately narrow (length slightly >1/3 ocular width), smooth and glabrous, with four terminal setae. Frons with dull toment, with fine, moderately dense punctures, anteriorly and laterally with a few long setae. Antenna with nine antennomeres, club with three antennomeres, 1.5 times as long as remaining antennomeres combined, distinctly reflexed. Eyes small, ratio diameter to interocular width 0.47. Mentum convexly elevated and flattened anteriorly.

Pronotum moderately wide, widest at base, lateral margins moderately convex and convergent anteriorly; anterior angles acute, moderately produced, posterior angles blunt; anterior margin weakly convex, marginal line fine but complete; basal margin without marginal line; surface with coarse, dense punctures and minute setae, anteriorly and laterally bearing erect long setae. Hypomeron not carinate ventrally. Scutellum narrow, long, triangular, with fine but dense punctuation and minute setae in punctures, and basally with a triangular smooth area.

Elytra elongate, narrow, widest in posterior third; striae distinctly impressed, finely punctate; intervals weakly convex, finely and densely punctate, with punctures concentrated along striae; elytra glabrous, lateral intervals with a few moderately long, adpressed or erect setae; epipleural margin robust, ending at strongly rounded external apical angle of elytra; epipleura sparsely setose; apex of elytra with a very fine, membranous rim of short microtrichomes.

Ventral surface finely and densely punctate, glabrous. Metacoxae laterally with a few longer setae. Mesosternum between mesocoxae slightly wider than mesofemur. Ratio of length of metepisternum to metacoxa 1:1.52. Abdominal sternites finely and moderately densely punctate, each with a row of coarse punctures, each puncture bearing each a robust seta. Pygidium almost flat and finely, densely punctate, with moderately dense, long setae.

Legs long, narrow, shiny; femora finely, sparsely punctate, with two longitudinal rows of setae. Metafemur with anterior margin acute, without adjacent serrate line; posterior ventral margin almost straight, weakly widened in apical half, ventrally and dorsally not serrate, sparsely setose. Metatibia long, narrow, widest at apex, ratio of width to length 1:3.9, sharply carinate dorsally, with three groups of spines, basal one at one-third, middle one at three-fifths, apical one at four-fifths of metatibial length, basally with a number of robust single setae; lateral face longitudinally convex, finely and moderately densely punctate, glabrous; apex shallowly, concavely truncate interiorly near tarsal articulation. Ventral edge finely serrate, with four robust equidistant setae.

Tarsomeres finely punctate dorsally, with moderately dense, fine setae ventrally; metatarsomeres ventrally with a strongly serrate carina, subventrally with a second, smooth longitudinal carina; first metatarsomere a little shorter than following two tarsomeres combined, and little shorter than dorsal tibial spur. Protibia long, bidentate; anterior claws symmetrical, interior tarsal claws sharply pointed.

Aedeagus: Fig. 1A–C, E–G. Habitus: Fig. 1D, H. Female unknown.

Diagnosis. *Serica breviantennalis* sp. nov. differs from *S. lupina* Arrow, 1946 and all other species of this species group in the short antenna and the shape of aedeagus; both parameres are long (subequal to half of the length of phallobase), nearly equal in length, parallel, and basally expanded to basal short lobes.

Etymology. The name of the new species is derived from the combined Latin words *brevis* (short) and *antennalis* (antenna), with reference on the short antenna. (An adjective in nominative singular case.)

Variation. Length 7.5–8.2 mm, elytral length 5.6–6.1 mm, width 4.0–4.3 mm. There is some variation of the parameres and lateral intervals of the elytra. Parameres of some specimens are closed, some are slightly open (Fig. 1A–C), and in others turned or rotated outside (Fig. 1E–G). Furthermore, the left paramere is sometimes curved externally (Fig. 1A–C) or sometimes dorsally (Fig. 1E–G). Lateral intervals of the elytra have a few moderately or distinctly dense, long, adpressed or erect setae.

***Serica allonanhuai* Liu, Ahrens, Li & Su, sp. nov.**

<https://zoobank.org/976AA8D8-1FEE-4297-9F9F-194390DDCFCD>

Figs 1I–L, 3

Type materials examined. Holotype: ♂ [China] “Madeng Village, Jianchuan County, Dali, Yunnan Province, 31.V.2022, light trap, leg. Zhao Yuchen” (SENU).

Paratypes: 1 ♀ [China] “Black Tiger Mountain, Tianxin Township, Wuding County, Chuxiong, Yunnan Province, 20.V.2022, light trap, leg. Zhao Yuchen, Lu Jinbo” (SENU), 1 ♂ “LW-1065 / Yunnan, Lijiang, 2750 m Yulongxueshan, 2011-V-17. 27.013°N, 100.206°E” (ZFMK).

Description of the holotype. Length 7.8 mm, elytral length 6.0 mm, width 4.2 mm. Body oblong, light reddish brown; frons reddish brown; some lateral and basal parts of elytra intervals dark reddish brown dull; legs yellowish brown; antenna yellow, with moderately long setae on dorsal surface.

Labroclipeus wide, trapezoidal; lateral margins moderately convex and convergent, producing a blunt angle with ocular canthus; anterior angles strongly convex; anterior margin distinctly emarginate medially; margins moderately reflexed; surface moderately convex and finely, densely punctate, with moderately long, erect setae; frontoclypeal suture distinct, moderately curved; ocular canthus moderately narrow (almost 1/3 of ocular width), smooth and glabrous, with one terminal seta. Frons with fine, moderately dense punctures, anteriorly and laterally with a few long setae. Antenna with nine antennomeres; club with three antennomeres, 1.8 times as long as remaining antennomeres combined, slightly reflexed. Eyes moderately large, ratio of diameter to interocular width 0.56. Mentum weakly elevated and flattened anteriorly.




Figure 1. *Serica* species **A–D** *S. breviantennalis* Liu, Ahrens, Li & Su, sp. nov. (holotype) **E–H** *S. breviantennalis* (paratype) **I–L** *S. allonanhuensis* Liu, Ahrens, Li & Su, sp. nov. (paratype). **A, E, I** aedeagus, left side view **C, G, K** aedeagus, right side view **B, F, J** aedeagus, dorsal view **D, H, L** habitus (not to scale). Scale bars: 0.5 mm (**A, B, C, E, F, G, I, J, K**).

Pronotum moderately wide, widest at base; lateral margins moderately convex and convergent anteriorly; anterior angles acute, moderately produced, posterior angles blunt; anterior margin weakly convex, marginal line fine but complete; basal margin without marginal line; surface with coarse, dense punctures and minute setae, laterally bearing erect, long setae. Hypomeron not ventrally carinate. Scutellum triangular, with fine but very dense punctuation and minute setae, with a triangular smooth area at middle.

Elytra elongate, narrow, widest in posterior third; striae distinctly impressed, finely punctate; intervals weakly convex, finely and densely punctate, punctures concentrated along striae; glabrous, lateral intervals with a few moderately long, adpressed or erect setae; epipleural margin robust, ending at strongly rounded external apical angle of elytra; epipleura sparsely setose; apex of elytra with a very fine, membranous rim of short microtrichomes.

Ventral surface finely and densely punctate, glabrous. Metacoxa laterally with several longer setae. Mesosternum between mesocoxae almost as wide as mesofemur. Ratio of length of metepisternum to metacoxa 1:1.92. Abdominal sternites finely and moderately densely punctate, each with a row of coarse punctures; each puncture bearing a robust seta. Pygidium almost flat and finely, densely punctate, with moderately dense, long setae.

Legs long, narrow, shiny; femora finely and sparsely punctate, with two longitudinal rows of setae, but anterior row reduced to a few setae. Metafemur with anterior margin acute, without adjacent serrate line; posterior ventral margin almost straight, weakly widened in apical half, ventrally and dorsally not serrate, sparsely setose. Metatibia long, narrow, widest at apex, ratio of width to length 1:3.5, moderately carinate dorsally, with three groups of spines, basal one at one-third, middle one at three-fifths, apical one at five-sixths of metatibial length, basally with a number of robust single setae; lateral face longitudinally convex, with fine, moderately dense punctae, glabrous; apex shallowly concavely truncate interiorly near tarsal articulation. Ventral edge finely serrate, with three robust, equidistant setae. Tarsomeres missing.

Aedeagus: Fig. 1I–K. **Habitus:** Fig. 1L.

Diagnosis. *Serica allonanhu* sp. nov. differs from *S. nanhua* and all other species of this species group in the light body colour and in the shape of aedeagus: the phallobase is strongly bent, the parameres are moderately long instead of only short (right and left paramere approximately one-third and one-fourth, respectively, as long as phallobase). Compared to *S. nanhua*, the parameres are distinctly longer, and in contrast to *S. baishuitaiensis* the parameres are straight in lateral view and narrow; the left paramere is slightly shorter than right one.

Etymology. The name of the new species is derived from the combination of the Greek prefix *allo-* (different, other) and *nanhua*, in reference to the similarity of the new species to *Serica nanhua*. (Used as a noun in apposition.)

Variation. Female Length 7.6 mm, elytral length 5.8 mm, width 4.1 mm. Antennal club short, slightly shorter than remaining antennomeres combined; eyes smaller than in male, ratio of diameter to interocular width 1:0.40; pygidium weakly convex, ratio of metepisternum to metacoxa 1:1.57; ratio of width to length of metatarsomeres 1:4.1. Tarsomeres dorsally finely punctate, ventrally with moderately dense, fine setae; metatarsomeres ventrally with a distinct serrate carina, subventrally with a second smooth, longitudinal carina; first metatarsomere a little shorter than following two tarsomeres combined and almost

as long as dorsal tibial spur. Protibia long, bidentate; anterior claws symmetrical; interior tarsal claws sharply pointed.

NCBI accession numbers. A paratype of the species (voucher number LW-1065) was sequenced as *Serica* sp. by Liu et al. (2015b) for two mitochondrial (CO1: [KJ959139](#), 16s: [KJ959212](#)) and one ribosomal marker (28S: [KJ959057](#)).

***Serica fengensis* Liu, Ahrens, Li & Su, sp. nov.**

<https://zoobank.org/49FA8835-21E3-4320-BFB0-7DD9BD547B99>

Figs 2A–E, 3

Type materials examined. Holotype: ♂ [China] “Feng County, Shaanxi, end of June.1974/ LW-174/ Asia Sericini 1325 spec.” (SENU). **Paratypes:** 1 ♂ “LW-1034 China: Shaanxi Huangbaiyuan” (ZFMK), 1 ♂ “LW-1273 China: Shaanxi Liuba Huoshaodian, 2012-VI-19” (ZFMK), 1 ♂ “LW-1177 China: Shaanxi, Liuba, 2012-VI-22” (ZFMK), 1 ♂ “LW-1280 China: Shaanxi Liuba Huoshaodian, 2012. VI.23, 33.576°N, 106.961°E, 966m” (ZFMK), 1 ♂ “LW-1273bis China: Shaanxi Liuba Huoshaodian, 2012-VI-19” (ZFMK), 1 ♂ “X-DA3562 – China, Sichuan prov., W of Heishui, 2500m, 32.0465°N, 103.0168°E, 3.–10.VI.2012 leg. M. Murzin” (ZFMK).

Description of the holotype. Length 8.4 mm, elytral length 6.0 mm, width 3.9 mm. Body oblong, narrow, dark brown; ventral surface, legs, and labroclypeus reddish brown; antenna yellow, with moderately dense, long setae on dorsal surface.

Labroclypeus wide, trapezoidal, shiny; lateral margins moderately convex and convergent, producing a blunt angle with ocular canthus; anterior angles moderately convex; anterior margin distinctly emarginate medially; margins moderately reflexed; surface flat, finely and densely punctate, with numerous long and erect setae; frontoclypeal suture distinct, moderately curved; ocular canthus long and narrow (ca 1/3 of ocular width), smooth and glabrous, with two terminal setae. Frons dull, with fine, dense punctures and dense, long, erect setae. Antenna with nine antennomeres; club with three antennomeres, 2.5 times as long as remaining antennomeres combined, strongly reflexed. Eyes moderately large, ratio of diameter to interocular width 0.67. Mentum weakly elevated and anteriorly flattened.

Pronotum moderately wide, widest at middle; lateral margins moderately convex and convergent anteriorly, slightly narrowed towards base; anterior angles rectangular, moderately produced, posterior angles blunt; anterior margin weakly convex, marginal line fine and complete; basal margin without marginal line; surface with fine, dense punctures and minute setae in punctures, with sparse, erect, long setae. Hypomeron not ventrally carinate. Scutellum triangular, with fine, dense punctuation and minute setae; basal midline with a triangular smooth area.

Elytra elongate, narrow, widest in posterior third; striae distinctly impressed, finely punctate; intervals weakly convex and finely, densely punctate; glabrous, lateral intervals with a few moderately long, erect setae, otherwise with only minute setae in punctures; epipleural margin robust, ending at convex external apical angle of elytra; epipleura sparsely setose; apex of elytra with a fine, membranous rim of short microtrichomes.




Figure 2. **A–E** *S. fengensis* Liu, Ahrens, Li & Su, sp. nov. (holotype) **A** aedeagus, left side view **C** aedeagus, right side view **B** aedeagus, dorsal view **D** habitus, dorsal view **E** habitus, lateral view. Scale bar: 0.5 mm (**A–C**).




Figure 3. The distribution map of *Serica breviantennalis* Liu, Ahrens, Li & Su, sp. nov., *S. allonanhuia* Liu, Ahrens, Li & Su, sp. nov., *S. fengensis* Liu, Ahrens, Li & Su, sp. nov., including updated records of *S. nanhua* Ahrens, Fabrizi & Liu, 2022 and *S. baishuitaiensis* Ahrens, Fabrizi & Liu, 2022.

Ventral surface finely, densely punctate, and metasternum sparsely setose. Metacoxa glabrous, laterally with numerous, robust, fine, longer setae. Mesosternum between mesocoxae almost as wide as mesofemur. Ratio of length of metepisternum to metacoxa 1:1.52. Abdominal sternites finely, moderately, and densely punctate, each sternite with a row of coarse punctures, each puncture bearing a robust seta. Pygidium almost flat and finely, densely punctate, with moderately dense, short and long setae.

Legs long, narrow, shiny; femora finely, sparsely punctate, with two longitudinal rows of setae, but anterior row reduced to a few setae. Hind legs missing in holotype. Tarsomeres dorsally finely punctate, ventrally with moderately dense, fine setae. Protibia long, bidentate; anterior claws symmetrical; interior tarsal claws sharply pointed.

Aedeagus: Fig. 2A–C. **Habitus:** Fig. 2D. **Female unknown.**

Diagnosis. *Serica fengensis* sp. nov. differs from the quite similar *S. allonanhuia* sp. nov. in the long basal lobes of the parameres; these lobes are absent in the latter species.

Etymology. The name of the new species is derived from the location of type locality in Feng County. (Used as an adjective in nominative singular case.)

Variation. Length 6.8–8.4 mm, elytral length 5.3–6.5 mm, width 3.8–4.0 mm. Most paratypes with dense pilosity on pronotum and elytra, suggesting that pilosity of holotype has been widely abraded. Metafemur with anterior margin acute, without adjacent serrate line; posterior ventral margin almost straight, weakly widened in apical half, ventrally and dorsally not serrate, sparsely setose. Metatibia long and moderately narrow, widest at middle, ratio of width to length

1:3.3, dorsally sharply carinate, with two groups of spines, basal one shortly behind middle, apical one at four-fifths of metatibial length, basally with a number of robust, single setae in line; lateral face longitudinally convex, coarsely and densely punctate, with a numerous short setae in basal half; apex shallowly concavely truncate interiorly near tarsal articulation. Ventral margin finely serrate, with five or six robust equidistant setae. Metatarsomeres ventrally with a strongly serrate carina, subventrally with a second, smooth, longitudinal carina; first metatarsomere distinctly shorter than following two tarsomeres combined and little longer than dorsal tibial spur.

NCBI accession numbers. Paratypes of the species have been sequenced as *Serica* sp. by Liu et al. (2015b) for two mitochondrial and one ribosomal marker (voucher number LW-1034: 16s: [KJ959207](#), CO1: [KJ959135](#), 28s: [KJ959052](#); LW-1273: 16s: [KJ959208](#), CO1: [KJ959136](#), 28s: [KJ959053](#)). The locality data of the LW-1273 specimen were erroneously given in the supplementary table 1 of Liu et al. (2015b) as “China: Yunnan, Pianma” instead of “Shaanxi Liuba Huoshaodian, 2012-VI-19”.

New records

Serica nanhua Ahrens, Fabrizi & Liu, 2022

Additional materials examined. 1 ♂ [China] “Qianchang town, Yao ‘an, Chuxiong, Yunnan Province, 23.V.2022, light trap, 2095 m, 101.37449(E), 25.52360(N), leg. Zhao Yuchen” (SENU), 1 ♂ [China] “Dali Jianchuan, Yunnan Province, 1.VI.2022, light trap, 2200m, 26.321291°N, 99.541980°E, leg. Zhao Yuchen” (SENU).

Serica baishuitaiensis Ahrens, Fabrizi & Liu, 2022

Additional materials examined. 3 ♂♂ [China] “Black Tiger Mountain, Tianxin Township, Wuding County, Chuxiong, Yunnan Province, 20.V.2022, light trap, 1160 m, 25.49412°N, 102.113091°E, leg. Zhao Yuchen, Lu Jinbo” (SENU).

Acknowledgements

We gratefully acknowledge Hu Li, Yuchen Zhao, Jinbo Lu, and Rentao Xu (Shaanxi University of Technology) for providing the specimens.

Additional information

Conflict of interest

The authors have declared that no competing interests exist.

Ethical statement

No ethical statement was reported.

Funding

This research was supported by Key Research and Development Project in Shaanxi Province (no. S2023-YF-YBSF-1112).

Author contributions

Conceptualization: YL, WGL. Data curation: WGL, DA. Funding acquisition: WGL. Investigation: WL. Methodology: DA, WGL. Project administration: WGL. Resources: YL. Software: XS, YJZ, WL. Supervision: YL. Validation: WGL, DA. Visualization: YJZ, XS, WL. Writing - original draft: XS, WL. Writing - review and editing: DA, WGL.

Author ORCIDs

Xie Su  <https://orcid.org/0009-0004-1723-2542>

Wei Li  <https://orcid.org/0009-0001-0195-7889>

Wan-Gang Liu  <https://orcid.org/0000-0003-4788-7967>

Dirk Ahrens  <https://orcid.org/0000-0003-3524-7153>

Yang Liu  <https://orcid.org/0000-0003-0301-0186>

Data availability

All of the data that support the findings of this study are available in the main text.

References

- Ahrens D (1996) Revision der *Sericini* des Himalaya und angrenzender Gebiete. Die Gattungen *Lasioserica* Brenske, 1896 und *Gynaecoserica* Brenske, 1896. Coleoptera-Schwanfelder Coleopterologische Mitteilungen 16: 1–48.
- Ahrens D (1999) Revision der Gattung *Serica* (s. str.) MacLeay des Himalaya-Gebiets (Coleoptera, Melolonthidae). Fragmenta Entomologica 31: 205–332.
- Ahrens D (2000a) *Serica korgei* Petrovitz, 1967 aus der Türkei – ein neues Synonym von *Autoserica castanea* Arrow, 1913 (Coleoptera: Melolonthidae: Sericini). Reichenbachia 33: 395–298.
- Ahrens D (2000b) Synopsis der Gattung *Gastroserica* Brenske, 1897 des ostasiatischen Festlandes (Coleoptera: Melolonthidae: Sericini). Entomologische Abhandlungen des Staatlichen Museums für Tierkunde Dresden 59: 73–121.
- Ahrens D (2000c) Weiteres Material der Gattung *Lasioserica* Brenske, 1896, mit der Beschreibung acht neuer Arten (Coleoptera: Melolonthidae: Sericini). Acta Musei Moraviae. Scientiae Biologicae 85: 1–28.
- Ahrens D (2001) Zwei neue *Chrysoserica*-Arten aus dem Himalaya. (Coleoptera Melolonthidae). Bollettino della Società Entomologica Italiana 133: 133–147.
- Ahrens D (2003a) Neue Arten aus der Gruppe der *Amiserica insperata* (Brenske, 1898) (Coleoptera, Scarabaeoidea, Sericini). Spixiana 26: 65–81.
- Ahrens D (2003b) On the identity of *Maladera infuscata* (Moser, 1915), with description of a new species from Taiwan (Coleoptera, Scarabaeidae, Sericini). Zootaxa 392(1): 1–8. <https://doi.org/10.11646/zootaxa.392.1.1>
- Ahrens D (2003c) Zur Identität der aus Madagaskar beschriebenen *Autoserica stupida* Brenske, 1900 (Coleoptera: Scarabaeidae: Sericini). Beiträge zur Entomologie = Contributions to Entomology 53: 231–238. <https://doi.org/10.21248/contrib.entomol.53.1.231-238>
- Ahrens D (2003d) Zur Identität der Gattung *Neoserica* Brenske, 1894, nebst Revision und Beschreibung neuer Arten. Koleopterologische Rundschau 73: 169–226.
- Ahrens D (2004) Monographie der Sericini des Himalaya (Coleoptera: Scarabaeidae). Dissertation.de-Verlag im Internet GmbH, Berlin, 534 pp.
- Ahrens D (2005a) A taxonomic review on the *Serica* (s. str.) MacLeay, 1819 species of Asian mainland (Coleoptera, Scarabaeidae, Sericini). Nova Supplementa Entomologica 18: 1–163.

- Ahrens D (2005b) Taxonomic revision of the genus *Anomalophylla* Reitter, 1887 (Coleoptera: Scarabaeidae: Melolonthinae: Sericini). Zootaxa 1076(1): 1–62. <https://doi.org/10.11646/zootaxa.1076.1.1>
- Ahrens D (2006a) The phylogeny of the genus *Lasioserica* inferred from adult morphology—implications on the evolution of montane fauna of the South Asian orogenic belt (Coleoptera: Scarabaeidae: Sericini). Journal of Zoological Systematics and Evolutionary Research 44(1): 34–53. <https://doi.org/10.1111/j.1439-0469.2005.00340.x>
- Ahrens D (2006b) Revision und phylogenetische Analyse der Gattung *Pachyserica* Brenske, 1897 (Coleoptera, Melolonthidae, Sericini). Revue Suisse de Zoologie 113: 487–557. <https://doi.org/10.5962/bhl.part.80360>
- Ahrens D (2006c) Sericinae. In: Löbl I, Smetana A (Eds) Catalogue of Palaearctic Coleoptera. Volume 3. Scarabaeoidea – Scirtoidea – Dascilloidea – Buprestoidea – Byrrhoidea. Apollo Books, Stenstrup, 229–248.
- Ahrens D (2007a) Beetle evolution in the Asian highlands: Insight from a phylogeny of the scarabaeid subgenus *Serica* (Coleoptera, Scarabaeidae). Systematic Entomology 32(3): 450–476. <https://doi.org/10.1111/j.1365-3113.2006.00373.x>
- Ahrens D (2007b) Revision der *Serica nigroguttata* Brenske, 1897-Gruppe (Coleoptera, Scarabaeidae, Sericini). Bulletin de l’Institut Royal des Sciences Naturelles de Belgique 77: 5–37.
- Ahrens D (2007c) Taxonomic changes and an updated catalogue for the Palaearctic Sericini (Coleoptera: Scarabaeidae: Melolonthinae). Zootaxa 1504(1): 1–51. <https://doi.org/10.11646/zootaxa.1504.1.1>
- Ahrens D (2009) A cladistic analysis reveals an eastern Tibetan occurrence of *Taiwanoserica* (Coleoptera: Scarabaeidae). Annales de la Société Entomologique de France (NS) 45(3): 285–296. <https://doi.org/10.1080/00379271.2009.10697612>
- Ahrens D (2011) A revision of the genus *Archeohomaloplia* Nikolaev, 1982. Coleoptera: Scarabaeidae: Sericini). Bonn Zoological Bulletin 60: 117–138.
- Ahrens D (2021) Additions to the fauna of *Tetraserica* Ahrens, 2004 of China and India. Koleopterologische Rundschau 91: 131–135.
- Ahrens D (2023a) Updates on the *Neoserica vulpes* group (Coleoptera, Scarabaeidae, Melolonthinae, Sericini): new species and records. European Journal of Taxonomy 888: 97–110. <https://doi.org/10.5852/ejt.2023.888.2231>
- Ahrens D (2023b) Two new species of *Archeohomaloplia* Nikolajev, 1972 from China (Coleoptera, Scarabaeidae, Sericini). Bonn Zoological Bulletin 72(1): 55–61. <https://doi.org/10.20363/BZB-2023.72.1.055>
- Ahrens D, Bezděk A (2016) Sericini. In: Löbl I, Löbl D (Eds) Catalogue of Palaearctic Coleoptera. Apollo Books, Stenstrup, 281–317.
- Ahrens D, Fabrizi S (2009a) New species of Sericini from the Eastern Himalaya and Tibet (Coleoptera, Scarabaeidae). In: Hartmann M, Weipert J (Eds) Biodiversität und Naturausstattung im Himalaya III. Verein der Freunde und Förderer des Naturkundemuseums Erfurt e.V., Erfurt, 249–284.
- Ahrens D, Fabrizi S (2009b) A review of the genus *Gynaecoserica* Brenske, (Coleoptera, Scarabaeidae, Sericini). Journal of Natural History 43(25–26): 1505–1584. <https://doi.org/10.1080/00222930902968809>
- Ahrens D, Fabrizi S (2011) New species of Sericini from the Himalaya and adjacent mountains (Coleoptera: Scarabaeidae). Bonn Zoological Bulletin 60: 117–138.
- Ahrens D, Fabrizi S (2016) A monograph of the Sericini of India (Coleoptera: Scarabaeidae). Bonn Zoological Bulletin 65: 1–355.

- Ahrens D, Pacholátko P (2003) New data on the distribution of species of *Gastroserica* Brenske, 1897, with descriptions of five new taxa from China and Laos (Coleoptera: Scarabaeidae: Sericini). Zootaxa 342(1): 1–18. <https://doi.org/10.11646/zootaxa.342.1.1>
- Ahrens D, Pacholátko P (2007) Two new species of *Gastroserica* Brenske, 1897 from China and Myanmar (Coleoptera: Scarabaeidae, Sericini). Entomologische Zeitschrift 117: 137–139.
- Ahrens D, Vogler AP (2008) Towards the phylogeny of chafers (Sericini): Analysis of alignment-variable sequences and the evolution of segment numbers in the antennal club. Molecular Phylogenetics and Evolution 47(2): 783–798. <https://doi.org/10.1016/j.ympev.2008.02.010>
- Ahrens D, Liu W-G, Fabrizi S, Bai M, Yang X-K (2014a) A revision of the species of the *Neoserica* (sensu lato) *vulpes* group (Coleoptera: Scarabaeidae: Sericini). Journal of Natural History 49(17–18): 1073–1130. <https://doi.org/10.1080/00222933.2014.974707>
- Ahrens D, Liu W-G, Fabrizi S, Bai M, Yang X-K (2014b) A taxonomic review of the *Neoserica* (sensu lato) *abnormis* group (Coleoptera, Scarabaeidae, Sericini). ZooKeys 439: 28–82. <https://doi.org/10.3897/zookeys.439.8055>
- Ahrens D, Liu W-G, Fabrizi S, Bai M, Yang X-K (2014c) A taxonomic review of the *Neoserica* (sensu lato) *septemlamellata* group (Coleoptera, Scarabaeidae, Sericini). ZooKeys 402: 76–102. <https://doi.org/10.3897/zookeys.402.7360>
- Ahrens D, Liu W-G, Fabrizi S, Bai M (2021a) Taxonomic review on the *Trioserica* Moser, 1922 species of China (Coleoptera: Scarabaeidae: Melolonthinae: Sericini). Zootaxa 4999(4): 343–355. <https://doi.org/10.11646/zootaxa.4999.4.4>
- Ahrens D, Liu W-G, Pham P, Fabrizi S (2021b) An overview on the genus *Amiserica* Nomura, 1974 (Coleoptera: Scarabaeidae: Melolonthinae: Sericini). Zootaxa 5050(1): 1–63. <https://doi.org/10.11646/zootaxa.5050.1.1>
- Ahrens D, Fabrizi S, Bai M, Liu W-G (2022) Taxonomic revision of *Serica* MacLeay, 1819 (sensu lato) from China and adjacent areas (Coleoptera: Scarabaeidae: Melolonthinae: Sericini), with updates on *Nipponoserica* Nomura, 1972. Zootaxa 5186(1): 1–83. <https://doi.org/10.11646/zootaxa.5186.1.1>
- Ahrens D, Lukic D, Pham P, Li W, Liu W (2023) *Tetraserica* Ahrens, 2004 of continental Southeast Asia: new records, new species, and an updated key to species (Coleoptera, Scarabaeidae: Sericini). Zootaxa 5374(4): 451–486. <https://doi.org/10.11646/zootaxa.5374.4.1>
- Bohacz C, Ahrens D (2020) A new species of the *Neoserica* (sensu lato) *vulpes* group from China, with a corrigendum on *Neoserica ailaoshanica* Liu, Fabrizi, Bai, Yang Ahrens, 2014 (Coleoptera: Scarabaeidae: Sericini). Journal of Natural History 54: 2991–2997. <https://doi.org/10.1080/00222933.2020.1758819>
- Dietz L, Seidel M, Eberle J, Misof B, Pacheco TL, Podsiadlowski L, Ranasinghe S, Gunter NL, Niehuis O, Mayer C, Ahrens D (2023) A transcriptome-based phylogeny of Scarabaeoidea confirms the sister group relationship of dung beetles and phytophagous pleurostict scarabs (Coleoptera). Systematic Entomology 48(4): 1–15. <https://doi.org/10.1111/syen.12602>
- Eberle J, Fabrizi S, Lago P, Ahrens D (2017) A historical biogeography of megadiverse Sericini—Another story “out of Africa”? Cladistics 33(2): 183–197. <https://doi.org/10.1111/cla.12162>
- Fabrizi S, Liu W-G, Bai M, Yang X-K, Ahrens D (2021) A monograph of the genus *Maladera* Mulsant & Rey, 1871 of China (Coleoptera: Scarabaeidae: Melolonthinae: Sericini). Zootaxa 4922(1): 1–400. <https://doi.org/10.11646/zootaxa.4922.1.1>

- Liu W-G, Ahrens D, Bai M, Yang X-K (2011) A key to species of the genus *Gastroserica* Brenske of the China (Coleoptera, Scarabaeidae, Sericini), with the description of two new species and two new records for China. ZooKeys 139: 23–44. <https://doi.org/10.3897/zookeys.139.1702>
- Liu W-G, Bai M, Yang X-K, Ahrens D (2014a) An update to the taxonomy of the genus *Gastroserica* Brenske (Coleoptera, Scarabaeidae, Sericini). ZooKeys 426: 87–110. <https://doi.org/10.3897/zookeys.426.7578>
- Liu W-G, Fabrizi S, Bai M, Yang X-K, Ahrens D (2014b) A taxonomic revision of the *Neoserica* (sensu lato) *calva* group (Coleoptera, Scarabaeidae, Sericini). ZooKeys 448: 47–81. <https://doi.org/10.3897/zookeys.448.8368>
- Liu W-G, Bai M, Yang X, Ahrens D (2014c) Towards an improved knowledge of Sericini of the Tibetan highland: new species and records (Coleoptera: Scarabaeidae). Bonn Zoological Bulletin 63: 157–172.
- Liu W-G, Fabrizi S, Bai M, Yang X, Ahrens D (2014d) A review of the *Tetraserica* species of China (Coleoptera, Scarabaeidae, Sericini). ZooKeys 448: 83–121. <https://doi.org/10.3897/zookeys.448.8429>
- Liu W-G, Fabrizi S, Bai M, Yang X, Ahrens D (2014e) A taxonomic revision of the *Neoserica* (s.l.) *pilosula* group (Coleoptera, Scarabaeidae, Sericini). ZooKeys 440: 89–113. <https://doi.org/10.3897/zookeys.440.8126>
- Liu W-G, Bai M, Yang X-K, Ahrens D (2015a) New species and records of the *Neoserica* (sensu stricto) group (Coleoptera: Scarabaeidae: Sericini). Journal of Natural History 49(39–40): 2379–2395. <https://doi.org/10.1080/00222933.2015.1034208>
- Liu W-G, Eberle J, Bai M, Yang X-K, Ahrens D (2015b) A phylogeny of Sericini with particular reference to Chinese species using mitochondrial and ribosomal DNA (Coleoptera: Scarabaeidae). Organisms, Diversity & Evolution 15(2): 343–350. <https://doi.org/10.1007/s13127-015-0204-z>
- Liu W-G, Fabrizi S, Bai M, Ahrens D (2016) A taxonomic revision of *Neoserica* (sensu lato): The species groups *N. lubrica*, *N. obscura*, and *N. silvestris* (Coleoptera, Scarabaeidae, Sericini). ZooKeys 635: 123–160. <https://doi.org/10.3897/zookeys.635.9915>
- Liu W-G, Bai M, Yang X, Ahrens D (2017a) New species and records of *Lasioserica* and *Gynaecoserica* from China (Coleoptera, Scarabaeidae, Sericini). Bonn Zoological Bulletin 66: 29–36.
- Liu W-G, Fabrizi S, Yang X, Bai M, Ahrens D (2017b) New species of *Nipponoserica* and *Paraserica* from China (Coleoptera, Scarabaeidae, Sericini). ZooKeys 721: 65–91. <https://doi.org/10.3897/zookeys.721.13918>
- Liu W-G, Fabrizi S, Bai M, Yang X, Ahrens D (2019) A taxonomic revision of Chinese *Neoserica* (sensu lato): final part (Coleoptera: Scarabaeidae: Sericini). Bonn Zoological Bulletin (Supplement 64): 1–71.
- Macleay WS (1819) Horae entomologicae: or, Essays on the annulose animals. S. Bagster, London, 1–160. <https://doi.org/10.5962/bhl.title.48636>
- Zhao M-Z, Ahrens D (2023) New species of the tribe Sericini Kirby, 1837 from China, with further updates on their taxonomy and distribution (Coleoptera: Scarabaeidae: Melolonthinae). Zootaxa 5361(2): 181–220. <https://doi.org/10.11646/zootaxa.5361.2.3>