

Description of a new species of *Parens* Fibiger, 2011 (Lepidoptera, Erebidae, Hypenodinae) from Korea

Ji-Young Lee¹, Bong-Kyu Byun¹¹ Department of Biological Science and Biotechnology, Hannam University, Daejeon, 34054, Republic of KoreaCorresponding author: Bong-Kyu Byun (bkbyun@hnu.ac.kr)

Abstract

The genus *Parens* comprises small moths, with a wingspan of 9–13 mm, belonging to the family Erebidae. Until now, only four species have been described worldwide. In Korea, only one species, *P. occi* (Fibiger & Kononenko, 2008) has been known to date. In this study, a new species from Korea, *P. fibigerina* Lee & Byun, **sp. nov.**, is described. As a result, two *Parens* species are now known from Korea. Figures of adults, male and female genitalia, and a key to the species of *Parens* in Korea are provided.

Key words: Checklist, identification key, Micronoctuini, new species, taxonomy

Introduction

The genus *Parens* Fibiger, 2011 belongs to the tribe Micronoctuini, subfamily Hypenodinae of the family Erebidae. Early authors considered *Parens* as a member of the family Micronoctuidae based the simple male genitalia without a uncus and the two-veined hindwing (Kononenko and Han 2007; Fibiger and Kononenko 2008; Fibiger et al. 2011; Fibiger 2011). More recently, Zahiri et al. (2012) proposed a new taxonomy of the Erebidae based on molecular phylogenetics. The family Micronoctuidae was downgraded to a tribe within the subfamily Hypenodinae, with its subfamilies given the rank of subtribes. This genus was established by Fibiger (2011) with *Parens paraocci* Fibiger, 2011 as the type species. *Parens* is a relatively small genus with only four recognized species worldwide. In Korea, only one species, *P. occi* (Fibiger & Kononenko, 2008) is known, which was first reported by Kononenko and Han (2007) as *Micronoctua* sp. In the present study, a new species, *P. fibigerina* Lee & Byun, sp. nov., is described from Korea.

Materials and methods

Terminology

We follow the general terminology proposed by Fibiger (2011).



Academic editor: Reza Zahiri

Received: 13 October 2023

Accepted: 6 January 2024

Published: 28 February 2024

ZooBank: <https://zoobank.org/C889C428-DB22-4DC9-BF7A-B12D16D62CD1>

Citation: Lee J-Y, Byun B-K (2024) Description of a new species of *Parens* Fibiger, 2011 (Lepidoptera, Erebidae, Hypenodinae) from Korea. ZooKeys 1193: 19–23. <https://doi.org/10.3897/zookeys.1193.113303>

Copyright: © Ji-Young Lee & Bong-Kyu Byun. This is an open access article distributed under terms of the Creative Commons Attribution License ([Attribution 4.0 International – CC BY 4.0](https://creativecommons.org/licenses/by/4.0/)).

Collection and preparation of specimens

The specimens were mainly collected using a bucket light trap with a 20 W black-light lamp and a LepiLED, standard model (WIF, Dr Gunnar Brehm, Sonnenblumenweg, Germany) and a 200 V / 400 W mercury-vapor lamp (220 V / 400 W).

All specimens were photographed before the dissection of their genitalia. Images of the adults were taken using a Canon EOS 600D digital camera (Canon Inc., Ota, Tokyo, Japan). Male and female genitalia were dissected and mounted in Euparal solution, following the procedure described in Holloway et al. (1987). The genitalia slides were photographed using a digital camera attached to a Leica M205C microscope (Leica Microsystems, Wetzlar, Germany) and refined with Photoshop CS5 (Adobe Systems Inc., San Jose, CA, USA).

Most specimens examined in this study were deposited in the Systematic Entomology Laboratory, Hannam University, Daejeon, Korea (**HNSUEL**). Additional specimens examined are from the Korea National Insect Collection, Korea National Arboretum, Korea (**KNA**). Abbreviations for localities in Korea are as follows: **GG** (Gyeonggi-do), **GW** (Gangwon-do), **CB** (Chungcheongbuk-do), **DJ** (Daejeon), **GB** (Gyeongsangbuk-do), **GN** (Gyeongsangnam-do), **JN** (Jeollanam-do), and **JJ** (Jeju-do). Other abbreviations are **TL** (type locality) and **TD** (type depository).

Systematic accounts

Family Erebidae Leech, [1815]

Subfamily Hypenodinae Forbes, 1954

Genus *Parens* Fibiger, 2011

Type species. *Parens paraocci* Fibiger, 2011.

Key to *Parens* species in Korea

- 1 In female genitalia, ductus bursae coiled at terminal margin ***Parens occi***
- In female genitalia, ductus bursae not coiled at terminal margin ***Parens fibigerina***

***Parens fibigerina* sp. nov.**

<https://zoobank.org/60C683F2-89B4-49D0-BBDA-FA1E8C3C1F84>

Figs 1A, 2A, B

Type material. **Holotype** Female, Korea, Donghae-si, GW, 12.viii.2021 (BK Byun), gen. slide no. HNUSEL-6442-coll. HNUSEL.

Diagnosis. This species is distinguished from *P. occi* by the shape of the signum in the corpus bursae. In *P. occi*, the cross-shaped signum in the corpus bursae, while *P. fibigerina* has a triangular signum. Additionally, the *P. fibigerina* has the forewing with a more rounded apex, a whitish-beige ground color, and a terminal margin covered with more blackish scales. The hindwing is grayish with mixed black scales. Also, the new species is distinguished from *P. chekiangi* Fibiger, 1911 by characters of the 8th abdominal segment and




Figure 1. Adults of *Parens* **A** *P. fibigerina* sp. nov. (gen. slide no. HNUSEL_6442) **B** *P. occi* (gen. slide no. HNUSEL_5931) Scale bars: 0.5 mm.




Figure 2. Male and female genitalia of Korean *Parens* species **A** ♀, *P. fibigerina* sp. nov., holotype (gen. slide no. HNUSEL_6442) **B** ditto, signum **C** ♂, *P. occi* (gen. slide no. HNUSEL_5931, 5937) **D** ♀, *P. occi* (gen. slide no. HNUSEL_6958) **E** ditto, signum. Scale bars: 0.1 mm.

the shape of the signum. The new species has the 8th abdominal segment is 2/3 length of the posterior apophyses and is well sclerotized. The antrum is strongly sclerotized. The new species has a triangular signum in the corpus bursae, while in *P. chekiangi* has a cross-shaped signum.

Description. Adult (Fig. 1A). Wingspan 11 mm. Head bend down, black; antenna filiform, black; frons rounded; labial palps porrect. Thorax and abdomen with sparse scales, beige; A8–10 dark brown. Forewing ground color whitish

beige, with black line from base to costal margin; base with half-round, blackish patch; apex rounded and with four yellowish blotches; antemedial and post-medial lines parallel, wavy, dark brown; spot reniform with whitish inner area and light brown outline. Hindwing ground color grayish brown, mixed black, and outline slightly curved to inner side with many cilia.

Male genitalia. Unknown.

Female genitalia (Fig. 2A, B). Papillae anales normal shape, rounded apex, with many short hairs, and well sclerotized. Posterior apophyses length equal to papillae anales; anterior apophyses extremely short. Ostium bursae placed in the median of abdomen. Antrum short, strongly sclerotized. Ductus bursae long, almost straight, narrow, membranous, and dilated at junction to corpus bursae (ca twice as broad as main tube). Corpus bursae globular and membranous, with triangular signum positioned slightly to left. Signum strongly sclerotized at base; upper side rather weakly sclerotized.

Distribution. Korea (endemic).

Etymology. This new species is dedicated to the memory of Michael Fibiger, a Danish entomologist, who was a renowned researcher of the tribe Micronoctuini.

***Parens occi* (Fibiger & Kononenko, 2008)**

Figs 1B, 2C–E

Micronoctua occi Fibiger & Kononenko, 2008: 52. TL: Russia, Primorye territory, Gornotaezhnoe.

Micronoctua sp.: Kononenko and Han 2007: 29.

Parens occi: Fibiger 2011: 19.

Parens occi: Lee and Byun 2022: 612.

Description. Adult (Fig. 1B). Wingspan 10–12 mm. See Lee and Byun (2022).

Male genitalia (Fig. 2C). See Lee and Byun (2022).

Female genitalia (Fig. 2D, E). See Lee and Byun (2022).

Materials examined. [GG] 1♂, Mt. Bongmisan, 03.xi.2008 (BW Lee, SY Park, DH Kwon), genitalia slide no. HNUSEL-5936-coll. KNAE; [GW] 1♂, Girin-myeon, 27.vi.2013 (BK Byun), genitalia slide no. HNUSEL-5931-coll. HNUSEL; 1♂, Yeongwol-gun, 26.viii.2021 (BK Byun), genitalia slide no. HNUSEL-6438-coll. HNUSEL; [CB] 2♂, Boeun-gun, 17.ix.2021 (BK Byun), genitalia slide no. HNUSEL-6309-coll. HNUSEL; Chungju-si, 10.vii.2020 (BK Byun), genitalia slide no. HNUSEL-6437-coll. HNUSEL; [DJ] 1♀, Masan-dong, 9.viii.2022 (BK Byun), genitalia slide no. HNUSEL-6958-coll. HNUSEL; [JN] 3♂, Isl. Jindo, 4.vi.2022 (BK Byun), genitalia slide no. HNUSEL-6763, 6913, 6914-coll. HNUSEL; [JJ] 1♂, Sanghyo-dong, 01.viii.2018 (BK Byun), genitalia slide no. HNUSEL-5937-coll. HNUSEL.

Distribution. Korea, China, Japan, Russia (Russian Far East).

Remarks. This species was reported first time from Korea by Kononenko and Han (2007).

Acknowledgements

We thank to Dr Thomas Pape and Dr Ole Karsholt for opportunity of a visiting study on the Hyphenodinae and allied groups in Natural History Museum of Denmark.

Additional information

Conflict of interest

The authors have declared that no competing interests exist.

Ethical statement

No ethical statement was reported.

Funding

This work was supported by a grant from the National Institute of Biological Resources (NIBR), funded by the Ministry of Environment (MOE) of the Republic of Korea (NIBR202333201).

Author contributions

Conceptualization: BKB. Data curation: JYL. Investigation: JYL. Visualization: JYL. Writing - original draft: JYL. Writing - review and editing: BKB.

Author ORCIDs

Ji-Young Lee  <https://orcid.org/0000-0001-8215-7957>

Bong-Kyu Byun  <https://orcid.org/0000-0003-0393-6464>

Data availability

All of the data that support the findings of this study are available in the main text.

References

- Fibiger M (2011) Revision of the Micronoctuidae (Lepidoptera: Noctuoidea). Part 4, taxonomy of the subfamilies Tentaxinae and Micronoctuinae. *Zootaxa* 2842: 1–188. <https://doi.org/10.11646/zootaxa.2583.1.1>
- Fibiger M, Kononenko VS (2008) Revision of the Micronoctuidae species occurring in the Russian Far East and neighbouring countries with description of a new species (Lepidoptera, Noctuoidea). *Zootaxa* 1890(1): 50–58. <https://doi.org/10.11646/zootaxa.1890.1.2>
- Fibiger M, Han HL, Kononenko VS (2011) Five new species and one new subspecies of Micronoctuidae from China, with a checklist of Chinese species, including Taiwan (Lepidoptera: Noctuoidea, Micronoctuidae). *Zootaxa* 2777(1): 41–53. <https://doi.org/10.11646/zootaxa.2777.1.3>
- Holloway JD, Bradley JD, Carger DJ (1987) CIE Guides to Insects of Importance to Man 1. Lepidoptera. CAB International, London, 262 pp.
- Kononenko VS, Han HL (2007) Atlas genitalia of the Noctuidae in Korea (Lepidoptera). In: Park KT (Ed.) *Insects of Korea Series 11*. Korea National Arboretum & Center for Insect Systematics, Pocheon, 464 pp.
- Lee JY, Byun BK (2022) Taxonomic review of the subfamily Hypenodinae (Lepidoptera: Erebidae) from Korea. *Journal of Asia-Pacific Biodiversity* 15(4): 603–612. <https://doi.org/10.1016/j.japb.2022.09.006>
- Zahiri R, Holloway JD, Kitching IJ, Lafontaine JD, Mutanen M, Wahlberg N (2012) Molecular phylogenetics of Erebidae (Lepidoptera, Noctuoidea). *Systematic Entomology* 37(1): 102–124. <https://doi.org/10.1111/j.1365-3113.2011.00607.x>