

# A new species of the genus *Parachemmis* Chickering, 1937 from Colombia (Araneae, Corinnidae, Corinninae)

Leonel Martínez<sup>1,2</sup>, Eduardo Villarreal<sup>1,2</sup>, Neis Martínez<sup>1,2</sup>

**1** Grupo de Investigación ECOSIN, Semillero de Investigación Artrópodos e Insectos Neoptera del Caribe Colombiano, Departamento de Biología, Universidad del Atlántico, Barranquilla, Colombia **2** Grupo de Investigación Biodiversidad del Caribe colombiano. Programa de Biología, Universidad del Atlántico, Barranquilla

Corresponding author: Leonel Martínez ([leonelmarbio@gmail.com](mailto:leonelmarbio@gmail.com))

Academic editor: I. Agnarsson | Received 24 February 2017 | Accepted 26 April 2017 | Published 8 June 2017

<http://zoobank.org/DB0C3FE4-C912-457D-B551-8D4CFD013490>

**Citation:** Martínez L, Villarreal E, Martínez N (2017) A new species of the genus *Parachemmis* Chickering, 1937 from Colombia (Araneae, Corinnidae, Corinninae). ZooKeys 679: 29–35. <https://doi.org/10.3897/zookeys.679.12421>

## Abstract

The spider genus *Parachemmis* Chickering, 1937 (Araneae: Corinnidae: Corinninae) is reported from Colombia for the first time. *Parachemmis julioblancoi* **sp. n.** Martínez-G & Villarreal is described and illustrated from the Sierra Nevada de Santa Marta, Magdalena department. The exclusive morphology of the short and apically truncated retrolateral tibial apophysis and club-like regular laminar process of the male palp indicates that the specimens described herein belong to a new species of *Parachemmis*. A map of the distribution of species in the genus is included.

## Keywords

Dionycha, distribution, Neotropical Region, Sierra Nevada de Santa Marta, taxonomy

## Introduction

The family Corinnidae Karsch, 1880 currently includes 754 species in 67 genera (World Spider Catalog 2017) in two subfamilies (Ramírez 2014). The subfamily Corinninae includes 17 genera, characterized by two synapomorphies: the male palpal reservoir primarily coiled and with a sclerotized distal sector, this last is share too with some Castianeirinae (Platnick and Baptista 1995, Bonaldo 2000, Ramírez 2014). The genus *Para-*

*chemmis* was proposed by Chickering (1937) with *Parachemmis fuscus* Chickering, 1937 as type species, and placed in the subfamily Micariinae (Clubionidae). Posteriorly Reiskind (1969) considered *Parachemmis* as a Liocraninae by the presence of four to six pairs of long ventral spines on the tibiae of the first pair of legs and setae simply on abdomen, however, Bonaldo and Brescovit (1994) examined this characters and transferred *Parachemmis* to the subfamily Corinninae, because several pairs of ventral spines on anterior tibiae are common among Corinninae and the photograph of the *Parachemmis hassleri* (Gertsch 1942) reveal feathery setae on abdomen and legs (Bonaldo and Brescovit 1994). Currently this genus includes three valid species, distributed in Brazil, Guyana, and Panamá. Nevertheless, these numbers are highly conservative, because in countries considered megadiverse as Colombia, only has reported 12 genera and 25 species for the family Corinnidae (World Spider Catalog 2017; William Galvis pers. comm.). Therefore, it is expected that with greater effort and review of material deposited in Natural History Museums our understanding of this diversity will increase considerably.

The species of the genus *Parachemmis* can be recognized by having serrula in the lateral edge of the endites, subovate carapace, abdominal traqueal tubercule, male palp with retrolateral tibial apophysis (RTA) entire and by the presence of an articulated ventral tibial apophysis (VTA), as well as by the presence of a prolateral laminar process (PLP) arising from the tegulum. In females, the epigyne have two copulatory openings, the copulatory ducts are large and the secondary spermathecae are poorly-developed. On the other hand, specimens of the genus also have anterior eye row strongly procurved, posterior eye row lesser procurved, and anterior median eyes clearly larger than the others, in addition to having the sternum with two external anterolateral excavations; the latter is also characteristic of the genera *Stethorrhagus* Simon, 1896 and *Tupirinna* Bonaldo, 2000 (Bonaldo 2000).

In this paper, both sexes of a new species of *Parachemmis* Chickering, 1937 from Colombia is described and illustrated: *P. julioblancoi* sp. n. from the Sierra Nevada de Santa Marta (SNSM), Magdalena, Colombia. Finally, a map of the distribution of the genus is included.

## Material and methods

The specimens examined are deposited in the Arachnological Collection of the Instituto de Ciencias Naturales of the Universidad Nacional de Colombia (ICN-Ar, Eduardo Flórez), Bogotá. The multifocal photographs of the copulatory structures and the measurements of the specimens were taken with a Leica MC-120 HD digital camera attached to a Leica S8AP0A stereomicroscope, the photographs were united by the image stacking software Leica Application Suite version 4.1.0. The illustrations of the palp and epigyne were made with a light camera attached to a Leica M125 stereomicroscope and the software Inkscape version 0.91. Platnick and Shadab (1975) is used as model for describing leg spination (with minor changes). For visualization of

female genitalia, the epigynal plate was dissected and cleared in KOH solution 10% concentration, following the guideline proposed by Platnick et al. (1999).

Abbreviations used in the text and figures are:

<b>AER</b>	anterior eye row	<b>PLP</b>	prolateral laminar process
<b>ALE</b>	anterior lateral eye	<b>PME</b>	posterior median eye
<b>AME</b>	anterior median eye	<b>po</b>	epigynal pocket
<b>c</b>	conductor	<b>pr</b>	proximal
<b>cd</b>	copulatory duct	<b>r</b>	retrolateral
<b>co</b>	copulatory opening	<b>RTA</b>	retrolateral tibial apophysis
<b>d</b>	dorsal	<b>sp</b>	spermathecal
<b>e</b>	embolus	<b>Spe</b>	spermophore
<b>fd</b>	fertilization duct	<b>ST I</b>	primary spermatheca
<b>m</b>	meters above mean sea level	<b>ST II</b>	secondary spermatheca
<b>p</b>	prolateral	<b>v</b>	ventral
<b>PER</b>	posterior eye row	<b>VTA</b>	ventral tibial apophysis
<b>PLE</b>	posterior lateral eye		

The map was prepared in the Geographic Information System QGIS “Las Palmas” (version 2.18.0, <http://www.qgis.org/es/site/>). The measurements are given in millimeters.

## Taxonomy

### *Parachemmis* Chickering, 1937

*Parachemmis* Chickering, 1937:38 (type species: *Parachemmis fuscus* Chickering, 1937)

**Diagnosis and description.** See Bonaldo 2000: 126.


### *Parachemmis julioblancoi* Martínez-G & Villarreal, sp. n.

<http://zoobank.org/EE03D67B-E20A-4DCB-98F3-4911195437DE>

Figs 1, 2, 3

**Holotype.** Male in ICN-Ar, from Colombia, Magdalena, Sierra Nevada de Santa Marta, San Pedro de la Sierra, 2104 m, 10.895138°N, 73.999611°W, 28 Mar.2017, L. Martínez (ICN-Ar 8420). **Paratypes.** 1♀, same data (ICN-Ar 8421); 2♂ (ICN-Ar 8422- 8423) and 4♀ (ICN-Ar 8424) from same locality; 1 ♂ and 1♀ (ICN-Ar 8327-8328) same locality. 6 May 2016.


**Etymology.** The specific epithet is a patronym in honor of Dr. Julio Enrique Blanco (Founder of the Universidad del Atlántico), for his many contributions to art and education in Colombia.



**Figure 1.** *P. julioblancoi* sp. n., holotype (ICN–Ar 8420), **A** habitus **C** sternum, paratype female (ICN–Ar 8421) **B** habitus **D** sternum. Scale bars 2 mm (**A**), 5 mm (**B**), 0.3 mm (**C**), 0.5 mm (**D**).


**Diagnosis.** Males of *P. julioblancoi* sp. n. can be distinguished from all remaining species of the genus by the large, club-like PLP, which extends towards the middle and distal part of the tegulum, wide conductor that ends next to embolus, a short, apically truncated RTA and a wide VTA (Fig. 2A–C), Females of *P. julioblancoi* sp. n. resembles those of *P. fuscus* by having the copulatory openings placed medially and by the non-coiled copulatory ducts, but can be diagnosed by the short, wide and rounded copulatory ducts and very short ST II (Fig. 2D–E).

**Description. Male (holotype, ICN–Ar 8327).** Total length: 7.77. Carapace brown, 3.20 long, 2.65 wide, 1.15 high. Eyes AER 1.18 wide, PER 1.32 wide. AME 0.25, PME 0.17, ALE 0.20, PLE 0.21. (Fig. 1A). Sternum brown 1.69 long, 1.65 wide (Fig. 1C). Legs 4123, I-femora 3.10/ patella 0.94/ tibiae 3.40/ Metatarsus 3.01/ tarsus 1.65/total = 12.10; II 3.29/0.99/2.75/2.85/1.63/11.51; III 2.33/0.70/2.32/2.05/1.21/8.61; IV 3.2/1.25/2.67/3.65/1.47/12.24. Chelicerae dark brown, with three promarginal and four retromarginal teeth. Labium brown. Legs yellow (Fig. 1A). Leg macrosetae: femora, I d 1-1-0, p 1 di; II d=I, p 1-1-1; III d 1-1-1, p 1-0-1, r 1-0-1; IV d=III, p 1 di, r 1 di. Tibiae, I v 1-2-2; II v 0-2-2; III v 2-2-2, p 1-0-1, r 1-0-1; IV v=III, p 1-1-0, r1-1-0. Metatarsus, I v 2-0-2; II v 2-0-1; III v 2-2-2, p 1 pr, r 1-1-0; IV v=III, p 1-1-0, r=III. Abdomen Dark gray, with an anterior yellow patch (Fig. 1A); ventrally gray. Spinnerets gray.



**Figure 2.** *P. julioblancoi* sp. n., holotype male (ICN–Ar 8420) **A** left male palp, prolateral view **B** same, ventral view **C** same, retrolateral view; paratype female (ICN–Ar 8421) **D** epigyne, dorsal view **E** same, ventral view. Abbreviations: c = conductor; cd = copulatory duct; co = copulatory opening; e = embolus; fd = fertilization duct; PLP = prolateral laminar process; po = epigynal; RTA = retrolateral tibial apophysis; sp = spermathecae; Spe = spermophore; VTA = ventral tibial apophysis. Scale bars 1 mm (**A–C**), 0.5 mm (**D–E**).

**Female. (paratype, ICN–Ar 8328).** Total length: 11.00. Carapace brown with yellowish posterior borders, 4.72 long, 3.84 wide, 1.97 high. Eyes AER 1.35 wide, PER 1.56 wide. AME 0.29, PME 0.24, ALE 0.27, PLE 0.19 (Fig. 1B). Sternum brown 2.42 long, 2.04 wide (Fig. 1D). Legs 4123. I-femora 4.26/ patella 1.46/ tibiae 4.41/ Metatarsus 3.77/ tarsus 1.01/total= 14.91; II 4.32/1.15/4.16/3.41/0.97/14.01; III 4.02/1.16/3.52/3.65/1.36/13.71; IV 5.17/1.49/4.45/5.32/1.61/18.04. Chelicerae dark brown with two promarginal and four retromarginal teeth. Labium brown. Legs yellow (Fig. 1B). Leg macrosetae: femora, I d 1-1-0 , p 0-1-1; II d 1-1-1, p 0-2-1; III d 1 di, p 2-1-1, 1, r 0-1-1; IV d 1-1-1, p 1 di, r 1 di. Tibiae, I-II v 2–2–2; III v 2-2-2-2, p



**Figure 3.** Distribution of the species of the genus *Parachemmis* in South America.

0-1-1, 0-1-1; IV v 2-2-2, p 1 di, 1-0-1. Metatarsus, I v 2-0-2; II v 2-1-2; III v 2-1-2-2, p 1-1-1, r 1-1-1; IV v 1-1-2-2, p 1-1-0, r 1-0-1. Abdomen dorsal gray, with several anterior yellow patches. (Fig. 1B); ventrally gray. Spinnerets gray.

**Natural history.** The type material was collected manually, on leaf litter, in a conserved high mountain wet forest ecosystem.

**Distribution.** Only known from the type locality (Fig. 3).

**Authors' contributions.** LM and EV collected, and identified the material. LM was responsible for species description. LM, NM and EV reviewed the literature, drafted the manuscript, and contributed to the critical discussion. EV and LM prepared the images. All authors read and approved the final manuscript.

## Acknowledgments

Thanks to Eduardo Flórez-Daza (ICN-Ar) for allowing us to examine the material here described, to Universidad del Atlántico for financial support. To Willian Galvis

(LAM-UN, Universidad Nacional de Colombia) and to Dr Alexandre Bonaldo (Museu Paraense Emílio Goeldi) for their support and comments on the manuscript; to the Neoptera team (Universidad del Atlántico) for their help and support, to Conchita Pinzón for help with the illustrations, and to the Laboratorio de Ecología y Entomología (LEE), Departamento de Biología, Facultad de Ciencias, Universidad del Atlántico, for the assistance in photographing the material.

## References

- Bonaldo AB (2000) Taxonomy of the subfamily Corinninae (Araneae, Corinnidae) in neotropical and nearctic regions. *Iheringia, série Zoologia* 89: 3–148. <https://doi.org/10.1590/S0073-47212000000200001>
- Bonaldo AB, Brescovit AD (1994) Revision of the Neotropical spider genus *Stethorrhagus* (Araneae, Corinnidae). *Andrias* 13: 33–64.
- Chickering AM (1937) The Clubionidae of Barro Colorado Island, Panama. *Transactions of the American Microscopical Society, Lancaster* 56: 1–47. <https://doi.org/10.2307/3222720>
- Gertsch WJ (1942) New American spiders of the family Clubionidae III. *American Museum Novitates* 1195: 1–18.
- Platnick NI, Baptista RLC (1995) On the spider genus *Attacobius* (Araneae, Dionycha). *American Museum Novitates* 3120: 1–9.
- Platnick NI, Grismado CJ, Ramírez MJ (1999) On the genera of the spider subfamily Otiotopinae (Araneae, Palpimanidae). *American Museum Novitates* 3257: 1–25.
- Platnick N, Shadab MV (1975) A revision of the spider genus *Gnaphosa* (Araneae, Gnaphosidae) in America. *Bulletin of the American Museum of Natural History* 155(1): 1–66.
- Ramírez MJ (2014) The morphology and phylogeny of dionychan spiders (Araneae: Araneomorphae). *Bulletin of the American Museum of Natural History* 390: 1–374. <https://doi.org/10.1206/821.1>
- Reiskind J (1969) The spider subfamily Castianeirinae of North and Central America (Araneae, Clubionidae). *Bulletin of the Museum of Comparative Zoology at Harvard College* 138: 163–325.
- World Spider Catalog (2016) World Spider Catalog, version 17.5. Natural History Museum Bern. <http://wsc.nmbe.ch> [accessed 20 January 2017]