

A new species of *Incestophantes* Tanasevitch, 1992 (Araneae, Linyphiidae) from Ukraine

Valery Gnelitsa

Sumy State Teacher's Training University, 87, Romenskaja street, Sumy 40002, Ukraine

urn:lsid:zoobank.org:author:C22EB46D-F6BA-48B0-9454-8AF0BE60F9D1

Corresponding author: Valery Gnelitsa (gnelitsa@mail.ru)

Academic editor: Pavel Stoev | Received 23 January 2009 | Accepted 12 April 2009 | Published 29 July 2009

urn:lsid:zoobank.org:pub:E938B9A8-019F-4FE5-9021-17DFD51A7959

Citation: Gnelitsa V (2009) A new species of *Incestophantes* Tanasevitch, 1992 (Araneae, Linyphiidae) from Ukraine. In: Stoev P, Dunlop J, Lazarov S (Eds) A life caught in a spider's web. Papers in arachnology in honour of Christo Deltchev. ZooKeys 16: 309-317. doi: 10.3897/zookeys.16.225

Abstract

A new spider species, *Incestophantes australis* sp. n. (Linyphiidae), found on the Crimean Peninsula, Ukraine is described and illustrated. It is morphologically closely related to *Incestophantes crucifer* (Menge, 1866), but is easily distinguished by its palpal conformation.

Keywords

Araneae, Linyphiidae, *Incestophantes australis* sp. n., *I. crucifer*, Ukraine

Introduction

The genus *Incestophantes* Tanasevitch, 1992 includes 23 species so far (Platnick 2008). Only *Incestophantes crucifer* (Menge, 1866) has been recorded from the forest zone and in the Carpathian region of the Ukraine (Gnelitsa 2000, 2002; Evtushenko 1993). It has also been mentioned from the Crimean Peninsula (Gnelitsa 2004). Comparison of the Crimean specimens of *Incestophantes crucifer* with those from the Ukraine revealed some morphological differences. It is now thought that the Crimean *Incestophantes* represents an undescribed species, closely related to *Incestophantes crucifer* (Menge, 1866). Therefore, this species is described here as *Incestophantes australis* sp. n. and comparative drawings and a brief re-description of *Incestophantes crucifer* are provided as well.

Material and methods

Specimens were collected using a hand-held suction sampler. Identification was made with a binocular microscope MBS-10; drawings were made using a camera lucida. The holotype male is deposited in the collection of the Zoological Department of Sumy State Teacher's Training University (SSTTU). All other specimens are deposited in author's private collection (VGC). Abbreviations of the names of palp and epigyne structures follow Hormiga (2000), Saaristo and Tanasevitch (1996), and Thaler et al. (1994): APc – apical part of paracymbium; E – embolus; i – keel of the lamella characteristic; Lch – lamella characteristic; LPc – lateral process of paracymbium; m – membrane; MPc – middle part of paracymbium; Pc – paracymbium; Ph – pit hook; R – radix; St – subtegulum; Su – suprattegulum; T – tegulum; TA – terminal apophysis; TA1 – first terminal apophysis; TA2 – second terminal apophysis; Ps – proscapus; DPs – distal part of scape. Other abbreviations in text: Fe – femur; Pt – patella; Mt – metatarsus; Ti – tibia; d – dorsal, p – prolateral, r – retrolateral position of leg spines. All measurements are in millimeters.

Taxonomic part

Incestophantes australis sp. n.

urn:lsid:zoobank.org:act:BEBFC5C6-21AC-4A08-BAD8-89E84238128D

Figs 1, 2a-c, 3a-d, 4a-d

Material. Holotype. Male, Ukraine, Crimea, **Yalta** Distr., Cape Martjan reserve, 44°30'N 34°15'E: *Quercus pubescens*, *Juniperus excelsa*, *Arbutus andrachne*, *Ruscus ponticus*, *Hedera taurica*, leaf litter, 23.IV.2002, V. Gnelitsa leg. (SSTTU). – *Paratypes* [all found in Crimea, Ukraine]: **Yalta** Distr., Shaitan Merdven rise, 44°25'N 33°51'E: *Quercus pubescens*, *Carpinus betulus*, *C. orientalis*, *Acer*, *Tilia*, *Euonimus verrucosa*, *Juniperus excelsa*, *Cotinus coggygia*, *Ruscus*, *Brachipodium*, *Hedera*, in grass, male, 12.X.2001; foot of Kilse-Burun Mt., 44°24'N 33°50'E, terraced slope, *Pinus pallasiana*, *Juniperus excelsa*, *Quercus pubescens*, *Carpinus orientalis*, *Hedera*, *Ruscus*, *Clematis*, with *Cotinus coggygia*, in needle litter and in grass, 8 females 4 males, 14.X.2001. **Alushta** Distr., Demerdji Mountain massif, 44°46'N 34°26'E, *Fagus* forest, in leaf litter, male, 17.IV.2002; SSE slope cereals and motley grass meadow, with sparse *Pyrus eleagnifolia*, *Fraxinus* and *Rosa*, in grass, male, 11.X.2002; meadow, in the grass along the stream, male, 10.X.2002. **Yalta**, Botkin path, E slope, 44°29'N 34°08'E, *Pinus*, *Quercus pubescens*, *Carpinus betulus*, *Swida*, *Ruscus*, *Crataegus*, needle litter, 2 females 1 male, 17.X.2002. **Pheodosia** Distr., Karadag reserve, 44°55'N 35°14'E, NNW slope of Tumanov ravine, *Quercus pubescens*, *Pinus pallasiana*, sparse *Fraxinus* with *Cornus mas*, in litter, 16 females 3 males, 25.IV.2003, 13.X.2003; Syvataya Mt. slope between North pass and Gyaur-Cheshme spring, *Quercus petraea*, *Acer*, *Paliurus spina-christi*, sparse *Fraxinus* with *Cornus mas*, in litter and in grass, male, 20.IV.2003; Syuryu-Kaya



Figure 1. Distribution of *Incestophantes* species in Ukraine: filled star – *I. australis*; filled circle – *I. crucifer*.

Mt. slope, sparse *Quercus pubescens*, *Fraxinus* with *Cornus mas*, in grass, 3 females 3 males, 11.X.2003, all leg. V. Gnelitsa (all preserved in the VGC).

Diagnosis. Male palp: Lamella characteristic 'Lch' beneath the distal spire with a wide, rounded keel 'i' (Fig. 2a,b). Shortened and widened apical part of paracymbium 'Apc' (Fig. 2a) distally truncate, cf. 'Apc' of *I. crucifer* which is elongate and gradually narrows towards the rounded end (Fig. 2d). Paracymbium with elongate lateral process

Table 1. Length of leg segments in male *Incestophantes australis*.

Legs	Femur	Patella	Tibia	Metatarsus	Tarsus
I	1.48	0.32	1.55	1.57	0.92
II	1.34	0.31	1.27	1.34	0.77
III	1.05	0.27	0.84	1.01	0.55
IV	1.41	0.29	1.23	1.40	0.71

Table 2. Length of leg segments in female *Incestophantes australis*.

Legs	Femur	Patella	Tibia	Metatarsus	Tarsus
I	1.43	0.33	1.44	1.41	0.90
II	1.27	0.32	1.16	1.18	0.76
III	1.02	0.28	0.80	0.90	0.55
IV	1.37	0.28	1.18	1.26	0.71

‘LPC’ (Fig. 2a); first terminal apophysis ‘TA1’ of the new species (Fig. 2b,c) narrowing towards the end, without parallel sides as in *I. crucifer* (Fig. 2e,f). Thick, long spine on palpal patella gradually tapers to a sharp tip, while this spine in *I. crucifer* distally widens and then tapers to a point.

Female distinguished by the narrower and elongated distal part of the scape ‘DPS’ (Fig. 3c). Because of considerable variation in the shape of the proscapus ‘Ps’ (Figs 3a, 4a-d) females are difficult to distinguish.

Description. Male. Total length 2.20; carapace: 1.12 long, 0.91 wide; yellow with thin dark border and median longitudinal stripe; sternum: 0.63 long, 0.62 wide, caudally extended between coxae IV, yellow-gray with slightly darker margin; posterior median eyes 0.6 diameter apart, anterior margin of chelicerae with two medium sized teeth, posterior margin with two tiny teeth at the base of the fang; stridulating files distinct. Abdomen yellow with dorsal pattern of two longitudinal rows of gradually diminishing grey blotches which are drawn together caudally. Legs yellow, femur, tibia and metatarsus all with dark ring, tibia darkened distally; spination: FeI – 1p, TiI – 2d:1p:1r, MtI – 1d; TiII – 2d:1r, MtII – 1d; TiIII – 2d and 2 spines (1p 1r) at the distal end, MtIII – 1d; TiIV – 2d, MtIV – 1d; position of metatarsal trichobothrium: I – 0.19, II – 0.19, III – 0.18.

Palp: see Fig. 2a-c.

Female. General appearance as in the male. Total length 2.75; carapace 1.06 long, 0.81 wide, yellow with dark gray border and median longitudinal stripe; sternum 0.63 long, 0.59 wide, caudally extended between coxae IV, yellow with slightly darker border; posterior median eyes 0.6 diameter apart; anterior margin of chelicerae with three medium sized teeth, posterior margin with four tiny teeth close together; stridulating files indistinct.

Epigyne: see Fig. 3a-c; vulva: see Fig. 3d.




Figure 2. Male palp. *Incestophantes australis* sp. n.: **a** lateral aspect **b** ventral aspect **c** mesal aspect. *I. crucifer*: **d** lateral aspect **e** ventral aspect **f** mesal aspect. Scale bars = 0.1 mm.




Figure 3. *Incestophantes australis* sp. n., epigyne: **a** ventral aspect **b** caudal aspect **c** dorsal aspect **d** vulva. *I. crucifer*, epigyne: **e** ventral aspect **f** caudal aspect **g** aspect **h** vulva. Scale bars = 0.1 mm.




Figure 4. *Incestophantes australis* sp. n., variation of the epigyne (ventral and caudal views) of specimens from a single locality. Scale bars = 0.1 mm.

Incestophantes crucifer (Menge, 1866)

Bathyphantes c. Menge, 1866: fig. 41.

Bolyphantes c.: Wiehle 1956: fig. 268-272; Miller 1971: plate XLIII, figs 7-8; Palmgren 1975: fig. 9.23-24; Pichka 1983: fig. 2.

Lepthyphantes c.: Simon 1884: figs 18-19; Thaler et al. 1994: fig. 30-39.

Incestophantes c.: Saaristo and Tanasevitch 2000: p. 257.

Figs 1, 2d-f, 3e-h

Material. Ukraine: Volyn region, **Shatsk** Distr., Shatsky Nature Park, 51°30'N 23°55'E, *Quercus* – *Pinus* forest, male, female, IV-X.1990, K. V. Evtushenko leg. (Evtushenko's private collection); Chernovtsy region, **Vyzhnitsa** Distr., Dolishny Shepot environs, 48°01'N 25°16'E, Chiuchelka Mt. slope, *Picea*, in needle litter and in moss on the ground, 3 females, 2.VII.2001; Sumy region, **Konotop** Distr., 2 km to the North of Kuzky village, 51°18'N 33°17'E, *Pinus*, sparse *Quercus*, *Betula*, *Sorbus* with *Frangula*, *Vaccinium myrtillus* and *Calluna vulgaris*, at the base of *Pine* trunk and in litter, 3 females 2 males, 22.V.2003, **Yampol** Distr., Prudishche environs, 51°57'N 33°43'E, *Pinus*, sparse *Sorbus*, at the base of *Pine* trunk, 6 females 1 male, 7.V.2000; **Seredinobuda** Distr., Staraya Guta environs, Desnyansko-Starogutsky Nature Park, 52°19'N 33°46'E, *Pinus*, sparse *Quercus*, *Sorbus* with *Frangula*, at the base of *Pine*

Table 3. Length of leg segments in male *Incestophantes crucifer*.

Legs	Femur	Patella	Tibia	Metatarsus	Tarsus
I	1.19	0.28	1.25	1.26	0.82
II	1.05	0.27	1.02	1.07	0.68
III	0.80	0.22	0.65	0.76	0.48
IV	1.04	0.24	0.98	1.09	0.63

Table 4. Length of leg segments in female *Incestophantes crucifer*.

Legs	Femur	Patella	Tibia	Metatarsus	Tarsus
I	1.05	0.26	0.99	0.97	0.67
II	0.91	0.24	0.81	0.81	0.57
III	0.70	0.22	0.53	0.60	0.39
IV	0.95	0.22	0.78	0.83	0.55

trunk, female, 22.VI.2000; moss, 6 females, 13.V.2001, 5 females, 15.V.2001; *Pinus*, *Betula*, sparse *Quercus*, *Sorbus* with *Frangula* and *Vaccinium myrtillus*, in litter, female, 16.V.2001; *Sphagnum-Eriophorum* bog, on the rising ground in the grass near the base of *Betula* trees, female, 8.V.1990, 2 females, 22.VI.2000, 2 females, 17.IX.2000, 5 females 1 male, 18.IX.2000, all collected by V. Gnelitsa (all preserved in the VGC).

Description. Male. Total length 1.69, carapace: 0.94 long, 0.74 wide, yellow with dark border and median longitudinal stripe; sternum: 0.56 long, 0.55 wide, caudally extended between coxae IV, yellow with slightly darker margin; posterior median eyes 0.6 diameter apart. Anterior margin of chelicerae at the base of the fang with five medium sized teeth and a small one remotely positioned, posterior margin with tiny tooth at the base of fang; stridulating files distinct. Abdomen with dorsal pattern of two longitudinal rows of dark patches on a light background. Legs yellow, femur, tibia and metatarsus all with dark ring, tibia darkened distally; spination: FeI – 1p, TiI – 2d:1p:1r, MtI – 1d; TiII – 2d:1r, MtII – 1d; TiIII – 2d and 2 spines (1p, 1r) at the distal end, MtIII – 1d; TiIV – 2d and 2 spines (1p, 1r) at the distal end, MtIV – 1d; position of metatarsal trichobothrium: I – 0.20, II – 0.19, III – 0.18.

Palp: see Fig. 2d-f.

Female. General appearance as in the male. Total length 2.23; carapace 0.78 long, 0.62 wide, yellow with dark gray border and median longitudinal stripe; sternum 0.49 long, 0.45 wide, caudally extended between coxae IV, yellow-gray with slightly darker border. Posterior median eyes 0.6 diameter apart. Anterior margin of chelicerae with three (two medium sized, one small) teeth, posterior margin with five tiny teeth close together; stridulating files as in male. Position of metatarsal trichobothrium: I – 0.25, II – 0.23, III – 0.20.

Epigyne: see Fig. 3e-g; vulva: see Fig. 3h.

Remarks. *Incestophantes crucifer* is known from Austria, Belarus, Belgium, Bulgaria, the Czech Republic, Estonia, Finland, France, Germany, Hungary, Italy, Liechtenstein, Norway, Poland, Romania, Russia, Slovakia, Sweden, Switzerland, Ukraine

and the former Yugoslavia (Van Helsdingen 2007). The spiders usually inhabit dry coniferous forests as well as bogs. Their webs are spanned in a slanting direction from tree trunks towards the litter.

References

- Evtushenko K (1993) To the study of spiders (Aranei) of the Volynskoe Polesye. Vestnik Zoologii, deposited in VINITI 06.01.93 № 27-B93: 1-13.
- Gnelitsa V (2000) The Linyphiid spiders of the Ivot river bottom land (the Sumy Area). Nature Reserves in Ukraine 6 (1-2): 77-80.
- Gnelitsa V (2002) The spiders of Linyphiidae family of the Desnansko–Starogutskiy Nature park. Nature Reserves in Ukraine 8 (2): 69-73.
- Gnelitsa V (2004) Preliminary data on the Linyphiidae spiders studying in Karadag Nature reserve. In: Karadagsky Prirodny zapovednik UNAN, Letopis prirody, Sonat, Simferopol, 135-138.
- Helsdingen PJ van (2007) Fauna Europaea Database <http://www.european-arachnology.org> [accessed 20.I.2009]
- Hormiga G (2000) Higher level phylogenetics of erigonine spiders (Araneae, Linyphiidae, Erigoninae). Smithsonian Contributions to Zoology 609: 1-160.
- Menge A (1866) Preussische Spinnen. Erste Abtheilung. Schriften der Naturforschenden Gesellschaft in Danzig (N.F.) 1: 1-152.
- Miller F (1971) Pavouci-Araneida. Klic zviřeny CSSR 4: 51-306.
- Palmgren P (1975) Die Spinnenfauna Finnlands und Ostfennoskandiens VI: Linyphiidae 1: Fauna fennica 28: 1-102.
- Pichka V (1983) New spider species of the USSR fauna. Vestnik Zoologii 3: 3-7.
- Platnick N (2008) The world spider catalog, version 9.0. American Museum of Natural History. <http://research.amnh.org/entomology/spiders/catalog/index.html>
- Saaristo M, Tanasevitch A (1996) Redelimitation of the subfamily Micronetinae Hull, 1920 and the genus *Lepthyphantes* Menge, 1866 with description of some new genera (Aranei, Linyphiidae). Berichte des naturwissenschaftlich-medizinischen Vereins in Innsbruck 83: 163-186.
- Saaristo M, Tanasevitch A (2000) Systematics of the *Bolyphantes-Poecilometes* genus-group of the subfamily Micronetinae Hull, 1920 (Arachnida: Araneae: Linyphiidae). Reichenbachia 33: 255-265.
- Simon E (1884) Les arachnides de France. Paris 5: 180-885.
- Thaler K, Helsdingen P, Deltsev C (1994) Vikariante Verbreitung im Artenkomplex von *Lepthyphantes annulatus* in Europa und ihre Deutung (Araneae, Linyphiidae). Zoologischer Anzeiger 232: 111-127.
- Wiehle H (1956) Spinnentiere oder Arachnoidea (Araneae). 28. Familie Linyphiidae-Baldachinspinnen. Tierwelt Deutschlands 44: i-viii, 1-337.