

# Species compositions of terrestrial isopods in public parks of a commuter town in Japan

Takahiro Nasu<sup>1</sup>, Kana Kitagawa<sup>1</sup>, Shigenori Karasawa<sup>2</sup>

**1** Fukuoka University of Education, 1-1 Akamabunkyo-machi, Munakata, Fukuoka 811-4192, Japan **2** Department of Life and Environmental Agricultural Sciences, Faculty of Agriculture, Tottori University, 4-101 Koyama-machi Minami, Tottori, Tottori 680-8553, Japan

Corresponding author: Shigenori Karasawa ([dojyoudoubutu@gmail.com](mailto:dojyoudoubutu@gmail.com))

---

Academic editor: K. Szlavecz | Received 28 October 2017 | Accepted 27 August 2018 | Published 3 December 2018

<http://zoobank.org/69BB2485-9129-42D9-8004-1ECD77F577A4>

---

**Citation:** Nasu T, Kitagawa K, Karasawa S (2018) Species compositions of terrestrial isopods in public parks of a commuter town in Japan In: Hornung E, Taiti S, Szlavecz K (Eds) Isopods in a Changing World. ZooKeys 801: 389–399. <https://doi.org/10.3897/zookeys.801.21875>

---

## Abstract

The species compositions of terrestrial isopod assemblages in 150 public parks in a commuter town in Japan were investigated. Eleven species of terrestrial isopods were present, of which seven were considered native or oriental species, and four were exotic species originally distributed in the Mediterranean and European regions. An exotic species *Armadillidium vulgare* Latreille, 1804 was found in all parks. Logistic model analysis indicated that the surrounding land use affected the distributions of three native species, *Burmoniscus kathmandius* (Schmalfuss, 1983), *Ligidium koreanum* Flasarova, 1972, and *Mongoloniscus koreanus* Verhoeff, 1930, indicating that landscape properties are important factors that limit the distributions of terrestrial isopods. The present study also showed that the public parks surrounded by forests or semi-natural environments in a commuter town provide habitats for native terrestrial isopods.

## Keywords

GIS, GLM, landscape environments, local environments, logistic distributions

## Introduction

Urbanization has rapidly spread throughout the world and has changed species compositions of regions through decreased diversity of native species (McKinney 2008) and increased numbers of exotics and/or generalist species (Niemelä 1999, Kotze et al. 2011), but the patterns of diversity changes are substantially different depending on biological group and climatic region (Faeth et al. 2011). Thus, understanding and protecting the biodiversity of urban regions is a major ecological concern. Many researchers have contributed to our knowledge of urban biodiversity in recent decades (e.g., McKinney 2008, Jones and Leather 2012, Nielsen et al. 2014), and some suggest that terrestrial isopods are the dominant macro-arthropods in soils of urban regions (Bolger et al. 2000, Smith et al. 2006, Magura et al. 2008a, b, Lee and Kwon 2015). Moreover, it has been observed that urban green spaces in parks provided important habitats that conserve biodiversity in urban environments (Kotze et al. 2011, Barratt et al. 2015). In Japan, many researchers have studied diversity of vegetation, birds, and insects in public parks (e.g., Ishii et al. 1991, Hata et al. 2003, Imai and Nakashizuka 2010, Hattori 2015). However, there has been no comprehensive study of the terrestrial isopods in public parks of Japan.


The Mediterranean region is considered a hotspot of terrestrial isopod diversity (Sfenthourakis and Taiti 2015) and some species that were originally distributed in the Mediterranean and European regions have become dispersed as exotic species throughout the world (e.g., Schmalfuss 2003, Cochard et al. 2010, Hornung et al. 2015). In Japan, seven species that are considered exotic have distribution areas with a potential for expansion (Nunomura 2007, Karasawa and Nakata 2018). These facts suggest that exotic species might already have occupied urban habitats and displaced native species in Japan.

The aims of this study were: 1) to describe the terrestrial isopod fauna in public parks of a commuter town in Japan; 2) to evaluate environments that limit species distributions; and 3) to determine whether public parks in the town provide habitats for native terrestrial isopods.

## Materials and methods

### Study area and sampling method

One hundred fifty public parks were selected in Munakata City, Fukuoka, Japan (Figure 1). Munakata City has been urbanized as a commuter town over several decades. The areas of the parks studied ranged from 74 m<sup>2</sup> to 17,089 m<sup>2</sup>. Two researchers walked around each park and searched for sites where terrestrial isopods were distributed, before collecting animals. To equalize the sampling effort among parks, the total time for animal collection was standardized as 7 min 30 s. All specimens found were collected, except for the highly abundant *Armadillidium vulgare* Latreille, 1804, for which only three or four voucher specimens were collected. Sampling was carried out



**Figure 1.** Location of Munakata city (red circle in the top right map) and map of the study area. Yellow circles represent sampling sites (parks). Blue denotes rice paddy; red denotes residential region; pink denotes grass; green denotes forest; the white line is Japan National Route 3. The map was created by modifying the high resolution land-use and land-cover map ver16.02 ([http://www.eorc.jaxa.jp/ALOS/lulc/jlulc\\_jpn.htm](http://www.eorc.jaxa.jp/ALOS/lulc/jlulc_jpn.htm)).

from 27 March to 27 August 2015. In addition, species lists were constructed for forests and a grassland area of the same city to determine the possible source of species in parks. To develop the species list for the forests, five sites (diameter ca. 5 m) were set up inside and at the edge of 18 forests (i.e., 10 sites per forest). In the survey, one researcher collected all terrestrial isopods for 10 min at each site. The species list for grassland was based on that of Tanaka and Karasawa (2018), which was carried out on a grass field in the same city.

Environmental variables were recorded at two levels: local and landscape scales. For local environments, the presence or absence of flower beds and woods in the parks were recorded during sampling. In addition, the percentage of the area of grass in parks was estimated using Google Map (<https://www.google.co.jp/maps>). For landscape environments, the areas of land use around the parks were measured using geographical information system (GIS) data. A raster data set for a high-resolution land-use and land-cover map (ver. 16.02) was obtained from the advanced land-observing satellite ([http://www.eorc.jaxa.jp/ALOS/lulc/jlulc\\_jpn.htm](http://www.eorc.jaxa.jp/ALOS/lulc/jlulc_jpn.htm); Hashimoto et al. 2014). For categorizing land use, grassland and crops were combined as ‘grass’, and four types of forest (broad-leaved, deciduous, evergreen and conifer) were combined as ‘forest’. Thus, four types of landscape environments were considered: ‘rice paddy’, ‘residential region’, ‘grass’, and ‘forest’. To evaluate suitable areas for analysis of landscape environments, buffer zones at five scales were created around each park (widths 50 m, 100 m, 200 m, 500 m, and 1000 m) and the areas of land use in each of the buffers were measured. GIS analyses were carried out using QGIS 2.14 (QGIS Development Team 2009).

### Statistical analyses

To evaluate the appropriate areas to use for the landscape environments, the Akaike information criteria (AIC) of the generalized linear models (GLMs) were compared at the five buffer widths for each species. The GLM was also used to evaluate relationships between the presence of 10 species and the species richness of native and exotic species (as response variables) with respect to environmental variables (as explanatory variables). Logistic (binary) and Poisson distributions were applied to the response variables of presence and species richness (of native and exotic species), respectively. Environmental variables were composed of three local variables (area percentage of grass, presence of flower beds and woods), and four landscape variables (areas of rice paddy, residential region, grass and forest). Using this procedure, perfect or quasi-perfect separations were found in the logistic models for some species. Thus, Firth’s bias-reduced logistic regression (Firth 1993; Heinze and Schemper 2002) was applied to the analysis of the presence of species. The proposed models were compared with the null model by the Likelihood Ratio Test. Significances of coefficient values in the proposed models were evaluated using the Chi-squared test (for presence of species) and the Wald test (for species richness). Multicollinearity among explanatory variables was assessed using variance inflation factors (VIF) for each model: variables with values > 10 were excluded. In addition, spatial autocorrelation of each species and species richness was evaluated by Moran’s *I*. To calculate the Moran’s *I*, distance weight was based on the distances among sampling sites: pairs of sites that are close together have higher values than pairs of sites that are far apart. All statistical analyses were carried out using the functions `glm`, `logistf`, `vif`, `lrtest`, and `Moran.I` in R version 3.4.1 (R Core Team 2017). *Armadillidium vulgare* was excluded from the latter analysis because this species was present in all parks (see Results section).

**Table 1.** Species, numbers of individuals collected and collection sites in the urban parks, grassland and forest in Munakata City, Japan.

Species	Urban park		Grassland <sup>a</sup> No. of ind.	Forest No. of ind.
	No. of ind.	Sites		
<b>Native species</b>				
<i>Agnara pannuosa</i>	18	9	a few <sup>b</sup>	1089
Armadillidae sp. 1 <sup>c</sup>	9	5	–	1955
Armadillidae sp. 2	–	–	–	60
Armadillidae sp. 3	–	–	–	1
<i>Burmoniscus dasystylus</i>	–	–	–	52
<i>Burmoniscus kathmandius</i>	17	5	–	–
<i>Exalloniscus cortii</i>	–	–	–	1
<i>Ligidium koreanum</i>	8	3	–	683
<i>Lucasioides</i> spp.	15	8	–	853
<i>Mongoloniscus koreanus</i>	172	36	3781	2
<i>Mongoloniscus vannamei</i>	110	36	a few	13
<i>Mongoloniscus</i> sp.	–	–	a few	–
<b>Exotic species</b>				
<i>Armadillidium vulgare</i> <sup>d</sup>	372	150	4526	2220
<i>Haplophthalmus danicus</i>	–	–	–	1
<i>Porcellio laevis</i>	60	34	–	–
<i>Porcellio scaber</i>	264	26	–	21
<i>Porcellionides pruinosus</i>	51	10	–	1

<sup>a</sup> The data from Tanaka and Karasawa (2018)


<sup>b</sup> This species was identified as Agnaridae in Tanaka and Karasawa (2018)

<sup>c</sup> This species was identified as *Spherillo* sp. in Karasawa et al. (2014)

<sup>d</sup> Three or four specimens only were collected from each park.

## Results

A total of 17 isopod species was recorded in the city, eleven of which were collected in the public parks. Five species, *A. vulgare*, *Haplophthalmus danicus* Budde-Lund, 1880, *Porcellio laevis* Latreille, 1804, *P. scaber* Latreille, 1804 and *Porcellionides pruinosus* (Brandt, 1833), are exotic species in Japan (Karasawa and Nakata 2018). *Armadillidium vulgare* was found in all parks and has also become the dominant species in the forests. Seven native species were collected in the parks but five of them were found in nine or fewer parks. In contrast, *Agnara pannuosa* (Nunomura, 1987), Armadillidae sp. 1, *Ligidium koreanum* Flasarova, 1972, *Lucasioides* spp. were abundant in the forests. *Mongoloniscus koreanus* Verhoeff, 1930, *M. vannamei* (Arcangeli, 1927), *P. laevis*, and *P. scaber* were widely distributed in the parks (> 25 parks) but their abundances were low in the forests. *Mongoloniscus koreanus* was dominant in grassland, as well as *A. vulgare* (Table 1). Overall, species richness in each park ranged from one to five (Figure 2). At least one exotic species was found in each park, but 80 parks lacked native species. The maximum species richness in a park was four for the native species and three for the exotic species.



**Figure 2.** Frequencies of species richness of native, exotic, and total species in parks

**Table 2.** Coefficients of explanatory variables and *p* values of the Likelihood ratio test.

Species	Local environments			Landscape environments			Intercept	Likelihood ratio test <i>p</i> -value	
	Grass	Flower bed	Wood	Residential	Rice paddy	Grass			Forest
<i>Agnara panmuosa</i>	0.181	1.493	-0.634	<b>-0.105</b>	-0.083	–	0.074	-2.098	0.177
Armadillidae sp. 1	1.756	-0.026	-1.340	–	-0.068	0.027	0.194	-3.114	0.399
<i>Burmoniscus kathmandius</i>	18.912	-0.472	0.745	<b>-0.047</b>	-0.036	–	-0.011	-18.440	<b>0.045</b>
<i>Ligidium koreanum</i>	-2.809	-1.331	-1.137	<b>-0.001</b>	-0.001	0.0001	–	4.447	<b>0.048</b>
<i>Lucasioides</i> spp.	-0.210	<b>-1.906</b>	0.148	–	0.001	0.001	-0.002	-2.262	<b>0.015</b>
<i>Mongoloniscus koreanus</i>	-0.035	-0.031	0.063	–	<b>0.009</b>	<b>0.012</b>	<b>0.011</b>	-3.367	<b>0.00001</b>
<i>Mongoloniscus vannamei</i>	0.893	1.103	-0.729	–	-0.001	0.001	-0.0002	-1.919	0.243
<i>Porcellio laevis</i>	-0.010	0.250	0.632	–	-0.010	-0.018	-0.036	-1.222	0.158
<i>Porcellio scaber</i>	1.200	0.452	0.398	–	-0.0004	<b>0.0000001</b>	-0.00002	-2.296	0.272
<i>Porcellionides pruinosus</i>	0.102	-0.636	0.961	-0.079	0.032	–	-0.150	-2.481	0.395
Number of species									
Native species	0.374	0.061	0.109	–	0.002	<b>0.006</b>	<b>0.004</b>	<b>-1.647</b>	<b>0.0002</b>
Exotic species	0.107	0.079	0.327	–	-0.0001	-0.001	-0.001	0.002	0.906

Bold represents *p* < 0.05 by the Chi-squared test.

Values of Moran's  $I$  revealed that distribution of *P. scaber* showed significant spatial autocorrelation ( $I = 0.027$ ,  $p < 0.05$ ), while other species and species richness showed no significant spatial autocorrelation. There were differences among species in the spatial scales having the lowest AIC values: *A. pannuosa*, 25 m; Armadillidae sp. 1, 25 m; *B. kathmandius* (Schmalfuss, 1983), 50 m; *L. koreanum*, 500 m; *Lucasioides* spp., 250 m; *M. koreanus*, 100 m; *M. vannamei*, 250 m; *P. laevis*, 50 m; *P. scaber*, 500 m; *P. pruinus*, 25 m (Supplementary material 1). In addition, the spatial scale of 100 m showed the lowest AIC for the species richness of native and exotic species. Coefficients of variables for the models fitted are described in Table 2. The likelihood ratio test significantly rejected the null model for *B. kathmandius*, *L. koreanum*, *Lucasioides* spp., and *M. koreanus*, and for the species richness of native species. Conversely, there was no significant difference between the proposal and null models for another species and species richness of exotic species, indicating that their distributions were not explained by the seven environmental variables. Local environment significantly affected the distribution of *Lucasioides* spp. only: the presence of flower beds may have a negative effect on the presence of *Lucasioides* spp. The distributions of four native species, *A. pannuosa*, *B. kathmandius*, *L. koreanum*, and *M. koreanus* were significantly affected by landscape properties, although there was no significant difference between the proposed and null models for *A. pannuosa*. *Agnara pannuosa*, *B. kathmandius*, and *L. koreanum* were negatively affected by the surrounding residential region, while natural or semi-natural environments (rice paddy, grass and forest) had positive effects on the distribution of *M. koreanus*. In addition, the model for the species richness of native species showed significant positive values for grass and forest areas, indicating that species richness of native species was high at sites surrounded by natural conditions. However, the model for *P. scaber* indicated a significant positive value for rice paddy only, although *P. scaber* showed spatial autocorrelation and the null model was not rejected.

## Discussion

Lee and Kwon (2015) reported that terrestrial isopods comprised approximately 90% of ground arthropods in urban parks of Osaka, a Japanese mega-city, but they did not identify them at species level. The present study found a total of eleven terrestrial isopod species in the public parks of a commuter town, four of which were exotic species. At least one terrestrial isopod species was found in the parks and the maximum species richness was five. These results imply that urban parks in Japan provide important habitats for terrestrial isopods, as observed in another country (Giurginca et al. 2017).

The species compositions in patchily distributed areas such as parks are known to be affected by local environments within the areas and by landscape environments around the parks (Jokimäki 1999, Germaine and Wakeling 2001, Smith et al. 2005). This study indicated that landscape environments play a more important role in determination of species distributions than do local environments; three native species

(*B. kathmandius*, *L. koreanum* and *M. koreanus*) preferred parks surrounded by forest or semi-natural environments (rice paddy and grass) to parks located in residential regions. *Ligidium koreanum* and *M. koreanus* have become abundant in forests and grassland within the city. In addition, *B. kathmandius* is widely distributed in East Asia and some Pacific Islands (Schmalfuss 1983, Taiti and Ferrara 1991, Karasawa 2016), and is usually found in grasslands of southern regions in Japan (SK, personal observations). Thus, these species are considered grass or forest specialists that may migrate to parks from surrounding semi-natural environments or forests. This finding suggests a hypothesis that the species richness of native species may increase in the parks surrounded by semi-natural and forest environments. The model analysis supported this hypothesis and these observations also indicate that an understanding of the distribution of terrestrial isopods requires evaluation of the surrounding landscape (Dauber et al. 2005).

Four exotic species with original distributions in the Mediterranean and European regions (Schmalfuss 2003, Cochard et al. 2010) were widely distributed in the public parks of the commuter town, but the present study failed to clarify the environmental factors that determined the distribution of exotic species; the null models were not rejected, and *P. scaber* also tended to be geographically concentrated. Three exotic species, *A. vulgare*, *P. scaber*, and *P. pruinosus*, are considered to be a habitat generalist, an urban specialist and a synanthropic species, respectively (Dangerfield and Telford 1990; Magura et al. 2008b), and they have been commonly found in urban regions (Vilisis et al. 2007, 2012, Vilisis and Hornung 2009, Giurginca et al. 2017). This may explain why human activity plays a more important role than environmental factors for their distributions in the city. Moreover, *A. vulgare* was distributed in all parks. This species is able to walk 24 m d<sup>-1</sup> and is a potentially able to walk 50 m h<sup>-1</sup> (Furukawa et al. 2017). This high dispersal ability could facilitate the distribution of *A. vulgare* via roads and residential regions and could be an important factor in the establishment and success of populations over a wide range. The establishment of populations of exotic species in public parks has the potential to be a source of exotic species in the future.

## Conclusions

Eleven terrestrial isopod species were found in the urban parks of a commuter town. Native species tended to be distributed in the parks located adjacent to natural environments, while their distributions had little relationship with the local environments within the parks. It is proposed that the location of parks is an important factor to consider in their design to protect the largest number of species of native terrestrial isopods.

## Acknowledgments

We thank members of the Laboratory of Animal Ecology at Fukuoka University of Education for assistance with field surveys, and Dr. Takahito Kuroki for advices on

GIS. We also thank the reviewers for their significant comments and efforts towards improving our manuscript. This work was supported by a Grant-in-Aid for Young Scientists (B) Grant Number 26830145. We thank Harry Taylor, PhD, from Edanz Group ([www.edanzediting.com/ac](http://www.edanzediting.com/ac)) for editing a draft of this manuscript.

## References

- Barratt BIP, Dickinson KJM, Freeman C, Porter S, Johnstone PD, Wing J, van Heezik Y (2015) Biodiversity of Coleoptera and other invertebrates in urban gardens: a case study in a New Zealand city. *Insect Conservation and Diversity* 8: 428–437. <https://doi.org/10.1111/icad.12120>
- Bolger DT, Suarez AV, Crooks KR, Morrison SA, Case TJ (2000) Arthropods in urban habitat fragments in southern California: area, age and edge effects. *Ecological Applications* 10: 1230–1248. <https://doi.org/10.2307/2641029>
- Cochard P-O, Vilisics F, Sechet E (2010) Alien terrestrial crustaceans (Isopods and Amphipods) Chapter 7.1. *BioRisk* 4: 81–96. <https://doi.org/10.3897/biorisk.4.54>
- Dangerfield JM, Telford SR (1990) Breeding phenology, variation in reproductive effort and offspring size in a tropical population of the woodlouse *Porcellionides pruinosus*. *Oecologia* 82: 251–258. <https://doi.org/10.1007/BF00323542>
- Dauber J, Purtauf T, Allspach A, Frisch J, Voigtländer K, Wolters V (2005) Local vs. landscape controls on diversity: a test using surface-dwelling soil macroinvertebrates of differing mobility. *Global Ecology and Biogeography* 14: 213–221. <https://doi.org/10.1111/j.1466-822X.2005.00150.x>
- Faeth SH, Bang C, Saari S (2011) Urban biodiversity: patterns and mechanisms. *Annals of the New York Academy of Sciences* 1223: 69–81. <https://doi.org/10.1111/j.1749-6632.2010.05925.x>
- Firth D (1993) Bias reduction of maximum likelihood estimates. *Biometrika* 80: 27–38. <https://doi.org/10.2307/2336755>
- Furukawa T, Mizushima M, Horinosono S, Karasawa S (2017) Estimation of the walking ability of an exotic terrestrial isopod *Armadillidium vulgare* Latreille by field and laboratory measurements. *Edaphologia* 101: 27–32.
- Germaine SS, Wakeling BF (2001) Lizard species distributions and habitat occupation along an urban gradient in Tucson, Arizona, USA. *Biological Conservation* 97: 229–237. [https://doi.org/10.1016/S0006-3207\(00\)00115-4](https://doi.org/10.1016/S0006-3207(00)00115-4)
- Giurginca A, Baba ŞC, Munteanu C-M (2017) New data on the Oniscidea, Diplopoda and Chilopoda from the urban parks of Bucharest. *North-Western Journal of Zoology* 13: 234–243.
- Hashimoto S, Tadono T, Onosato M, Hori M, Shiomi K (2014) A new method to derive precise land-use and land-cover maps using multi-temporal optical data. *The Remote Sensing Society of Japan* 34: 102–112. [In Japanese with English abstract]
- Hata H, Nakao S, Yabu S, Nakashima A, Yamada H (2003) Micro-habitat selection of grasshoppers in green space of park. *Journal of The Japanese Institute of Landscape Architecture* 66: 607–612. [In Japanese with English abstract]
- Hattori T (2015) Park biodiversity strategy. *Landscape Ecology and Management* 20: 37–40. [In Japanese]

- Heinze G, Schemper M (2002) A solution to the problem of separation in logistic regression. *Statistics in Medicine* 21: 2409–2419. <https://doi.org/10.1002/sim.1047>
- Hornung E, Szlavecz K, Dombos M (2015) Demography of some non-native isopods (Crustacea, Isopoda, Oniscidea) in a Mid-Atlantic forest, USA. *ZooKeys* 515: 127–143. <https://doi.org/10.3897/zookeys.515.9403>
- Imai H, Nakashizuka T (2010) Environmental factors affecting the composition and diversity of avian community in mid- to late breeding season in urban parks and green spaces. *Landscape and Urban Planning* 96: 183–194. <https://doi.org/10.1016/j.landurbplan.2010.03.006>
- Ishii M, Yamada M, Hirowatari T, Yasuda T (1991) Diversity of butterfly communities in urban parks in Osaka Prefecture. *Japanese Journal of Environmental Entomology and Zoology* 3: 183–195.
- Jokimäki J (1999) Occurrence of breeding bird species in urban parks: effects of park structure and broad-scale variables. *Urban Ecosystems* 3: 21–34. <https://doi.org/10.1023/A:1009505418327>
- Jones EL, Leather SR (2012) Invertebrates in urban areas: a review. *European Journal of Entomology* 109: 463–478. <https://doi.org/10.14411/eje.2012.060>
- Karasawa S (2016) Eleven nominal species of *Burmoniscus* are junior synonyms of *B. kathmandius* (Schmalfuss, 1983) (Crustacea, Isopoda, Oniscidea). *ZooKeys* 607: 1–24. <https://doi.org/10.3897/zookeys.607.8253>
- Karasawa S, Kanazawa Y, Kubota K (2014) Redefinitions of *Spherillo obscurus* (Budde-Lund, 1885) and *S. dorsalis* (Iwamoto, 1943) (Crustacea: Oniscidea: Armadillidae), with DNA markers for identification. *Edaphologia* 93: 11–27.
- Karasawa S, Nakata K (2018) Invasion stages and potential distributions of seven exotic terrestrial isopods in Japan. *BioRisk* 13: 53–76. <https://doi.org/10.3897/biorisk.13.23514>
- Kotze J, Venn S, Niemelä J, Spence JR (2011) Effects of Urbanization on the Ecology and Evolution of Arthropods. In: Niemelä J, Breuste JH, Elmqvist T, Guntenspergen G, James P, McIntyre NE (Eds) *Urban Ecology: Patterns, Processes, and Applications*. Oxford University Press, Oxford, 159–166. <https://doi.org/10.1093/acprof:oso/9780199563562.003.0019>
- Lee CM, Kwon TS (2015) Response of ground arthropods to effect of urbanization in southern Osaka, Japan. *Journal of Asia-Pacific Biodiversity* 8: 343–348. <https://doi.org/10.1016/j.japb.2015.10.007>
- Magura T, Tóthmérész B, Hornung E, Horváth R (2008a) Urbanization and ground-dwelling invertebrates. In: Wagner LN (Ed.) *Urbanization: 21<sup>st</sup> Century Issues and Challenges*. Nova Science Publishers, New York, 213–225.
- Magura T, Hornung E, Tóthmérész B (2008b) Abundance patterns of terrestrial isopods along an urbanization gradient. *Community Ecology* 9: 115–120. <https://doi.org/10.1556/ComEc.9.2008.1.13>
- McKinney ML (2008) Effects of urbanization on species richness: a review of plants and animals. *Urban Ecosystems* 11: 161–176. <https://doi.org/10.1007/s11252-007-0045-4>
- Nielsen AB, van den Bosch M, Maruthaveeran S, van den Bosch CK (2014) Species richness in urban parks and its drivers: a review of empirical evidence. *Urban Ecosystems* 17: 305–327. <https://doi.org/10.1007/s11252-013-0316-1>
- Niemelä J (1999) Is there a need for a theory of urban ecology? *Urban Ecosystems* 3: 57–65. <https://doi.org/10.1023/A:1009595932440>

- Nunomura N (2007) *Armadillidium vulgare* is exotic species. In: The Japanese Society of Soil Zoology (Ed) Practical guide to soil zoology: From sampling to data analysis. Tokai University Press, Kanagawa, 223–224. [In Japanese]
- QGIS Development Team (2009) QGIS Geographic Information System. Open Source Geospatial Foundation. <http://qgis.osgeo.org>
- R Core Team (2017) R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna. <https://www.R-project.org/>
- Schmalfuss H (1983) Terrestrial isopods from Nepal. *Senckenbergiana Biologica* 63: 373–392.
- Schmalfuss H (2003) World catalog of terrestrial isopods (Isopoda: Oniscidea). *Stuttgarter Beiträge zur Naturkunde, Serie A* 654: 1–341.
- Sfenthourakis S, Taiti S (2015) Patterns of taxonomic diversity among terrestrial isopods. *ZooKeys* 515: 13–25. <https://doi.org/10.3897/zookeys.515.9332>
- Smith RM, Gaston KJ, Warren PH, Thompson K (2005) Urban domestic gardens (V): relationships between landcover composition, housing and landscape. *Landscape Ecology* 20: 235–253. <https://doi.org/10.1007/s10980-004-3160-0>
- Smith RM, Warren PH, Thompson K, Gaston KJ (2006) Urban domestic gardens (VI): environmental correlates of invertebrate species richness. *Biodiversity and Conservation* 15: 2415–2438. <https://doi.org/10.1007/s10531-004-5014-0>
- Taiti S, Ferrara F (1991) Terrestrial isopods (Crustacea) from the Hawaiian Islands. *Bishop Museum Occasional Papers* 31: 202–227.
- Tanaka R, Karasawa S (2018) Life histories and soil water content preferences of sympatric exotic and native terrestrial isopods. *Entomological Science* 21: 198–209. <https://doi.org/10.1111/ens.12298>
- Vilisics F, Elek Z, Lövei GL, Hornung E (2007) Composition of terrestrial isopod assemblages along an urbanisation gradient in Denmark. *Pedobiologia* 51: 45–53. <https://doi.org/10.1016/j.pedobi.2006.12.004>
- Vilisics F, Hornung E (2009) Urban areas as hot-spots for introduced and shelters for native isopod species. *Urban Ecosystems* 12: 333–345. <https://doi.org/10.1007/s11252-009-0097-8>
- Vilisics F, Szekeres S, Hornung E (2012) Size dependent differences in litter consumption of isopods: preliminary results. *ZooKeys* 176: 247–259. <https://doi.org/10.3897/zookeys.176.2470>

## Supplementary material I

### AIC values of GLMs at the five spatial scales

Authors: Takahiro Nasu, Kana Kitagawa, Shigenori Karasawa

Data type: species data

Copyright notice: This dataset is made available under the Open Database License (<http://opendatacommons.org/licenses/odbl/1.0/>). The Open Database License (ODbL) is a license agreement intended to allow users to freely share, modify, and use this Dataset while maintaining this same freedom for others, provided that the original source and author(s) are credited.

Link: <https://doi.org/10.3897/zookeys.801.21875.suppl1>