

# Planthoppers of Delaware (Hemiptera, Fulgoroidea), excluding Delphacidae, with species incidence from adjacent States

Charles R. Bartlett<sup>1</sup>, Erin R. Adams<sup>1</sup>, Anthony T. Gonzon, Jr.<sup>2</sup>

**1** Department of Entomology and Wildlife Ecology, 250 Townsend Hall, University of Delaware, Newark, Delaware, 19717-1303. U.S.A. **2** DE Natural Heritage and Endangered Species Program, DE Division of Fish and Wildlife, DNREC 4876 Hay Point Landing Road, Smyrna, Delaware, 19977. U.S.A.

Corresponding author: Charles R. Bartlett (Bartlett@udel.edu)

---

Academic editor: Michael Wilson | Received 8 November 2010 | Accepted 29 January 2011 | Published 25 February 2011

**Citation:** Bartlett CR, Adams ER, Gonzon AT Jr. (2011) Planthoppers of Delaware (Hemiptera, Fulgoroidea), excluding Delphacidae, with species incidence from adjacent States. *ZooKeys* 83: 1–42. doi: 10.3897/zookeys.83.1176

---

## Abstract

The number of species of planthoppers (excluding Delphacidae) known from Delaware is updated from 7 (in 4 families) to 62 species (in 9 families). Specimen abundance is tallied by county and seasonally by two week intervals. The Chao1 abundance estimator suggests that the true fauna may be 74 species, although species incidence tallied from adjacent states (MD, NJ, PA and DC) suggests that a total fauna of approximately 100 species may be possible. An artificial key is presented to genus and select species with photos of most included taxa.

## Keywords

Auchenorrhyncha, Fulgoromorpha, Acanaloniidae, Achilidae, Caliscelidae, Cixiidae, Derbidae, Dictyopharidae, Flatidae, Issidae, species inventory, generic key

## Introduction

The distribution of planthoppers (Hemiptera: Auchenorrhyncha: Fulgoroidea) in the eastern United States was most recently summarized by Wilson and McPherson (1980a). Excluding the Delphacidae, Wilson and McPherson (1980a) reported only 4 planthopper species from Delaware, specifically *Acanalonia conica* (Say, 1830, Acanaloniidae), *Catonia cinctifrons* (Fitch, 1956, Achilidae), *Melanoliarius ecologus* (Caldwell,

1947; as *Oliarus*, Cixiidae), and *Metcalfa pruinosa* (Say, 1830, Flatidae). Additional species were later reported by Kramer in his revisions of the Cixiidae, specifically *Cixius nervosus* (Linnaeus, 1758) by Kramer (1981), *Bothriocera cognita* Caldwell, 1943, and *Bothriocera drakei* Metcalf, 1923, by Kramer (1983), bringing the reported fauna to 7 species in 4 families.

Our objectives were to provide an abundance-based list of planthopper species found in Delaware (excluding Delphacidae) established primarily on specimen records from the University of Delaware Insect Reference Collection (UDCC) in Newark, DE; provide a measure of completeness of this inventory using the Chao1 abundance-based diversity estimator (Chao 1984) and by comparison with incidence records from adjacent states (MD, NJ, PA and DC); and begin to assess their biology by providing preliminary information regarding the seasonality of the planthoppers of Delaware. We provide an artificial key to genus and select species to allow users to recognize planthopper species in the Mid-Atlantic States more easily.

## Methods

Planthopper specimens from Delaware, Maryland, New Jersey, and Pennsylvania in the UDCC were identified to species. Identification of some taxa requires dissection of male genitalia, in which case the abdomen was removed (sometimes after relaxing the specimen overnight in high humidity) and cleared for 24 hours in 15% potassium hydroxide (KOH), rinsed in water and transferred to glycerol for observation and manipulation (see, e.g., Wilson and McPherson 1980b, Bartlett and Deitz 2000). Species identification was made according to the following sources: Metcalf (1923, Derbidae except *Cedusa*, Dictyopharidae except *Scolops* and *Phylloscelis*, Flatidae), McAtee (1923, Derbidae: *Otiocerus*), Breakey (1928, Dictyopharidae: *Scolops*), Doering (1938, Issidae; 1939, Caliscelidae: *Bruchomorpha*; 1941, Caliscelidae: *Apheloneuma*), Beirne (1950, Achilidae: *Cixidia* [as *Epiptera*]), O'Brien (1971, Achilidae: Plectoderini), Kramer (1977, Cixiidae: *Oecleus*; 1979, Cixiidae: *Haplaxius* [as *Mynodus*]; 1981, Cixiidae: *Cixius*; 1983, Cixiidae: *Bothriocera*, *Pintalia*), Mead and Kramer (1982, Cixiidae: *Melanoliarus* [as *Oliarus*]), Flynn and Kramer (1983, Derbidae: *Cedusa*), Freund and Wilson (1995, Acanaloniidae), McPherson and Wilson (1995, Dictyopharidae: *Phylloscelis*). The specific identities of some taxa were confirmed by comparison with authoritatively determined specimens at the US Smithsonian Institution National Museum of Natural History (USNM), although in a few cases we examined types, or photographs of types (specifically the derbids *Otiocerus signoretii* Fitch, *Anotia burnetii* Fitch, and *Anotia robertsonii* Fitch from the USNM; and *Otiocerus stollii* Kirby and the purported type of *Anotia bonnetii* Kirby [but see discussion] from the Hope Entomological Collections Oxford University Museum of Natural History, OUMNH). Additional Kirby types were sought (from the British Museum, Manchester Museum, and Oxford), but are apparently missing. Females of some genera (e.g., Derbidae: *Cedusa* and many Cixiidae) cannot be identified to spe-

cies with confidence. These specimens were tallied at the generic level and included in the specimen counts, but not included in species counts or calculation of the Chao1 statistic (see below). The artificial key to genus and select species was constructed for all taxa not requiring dissection for identification. The key was developed by modification of keys within the above listed taxonomic references. Author and year for all species is provided in table 2.

Family-level nomenclature follows Emeljanov (1999) in recognizing Acanaloniidae and Caliscelidae as independent from Issidae. Keys to families of Fulgoroidea can be found in Wilson (2005). Generic nomenclature has been updated for Cixiidae following Emeljanov (2001) and Holzinger and colleagues (2002) and for Issidae by Gnezdilov (2004).

Incidence records were listed for Maryland, New Jersey, Pennsylvania, and the District of Columbia based on literature (see below) and specimen records. Specimen records were compiled both from the UDCC and USNM collections. Specimens from Delaware were totaled by county and collection date increment. For collection date tallies, each month was divided into two increments, "early" (the 1–15<sup>th</sup> of each month), and "late" (the 16<sup>th</sup>–end of month) dates. Specimens with incomplete date information were omitted from these counts (resulting in the number of specimens tallied for seasonal data for some species to be less than the number of specimens observed). Because some species were at times found in abundance, seasonality records were tallied in two ways; complete specimen counts, and observation records where each series (all specimens recorded from a particular location and date) was tallied as a single observation.

To help assess completeness of the inventory, literature records were compiled from published sources (viz. Wilson and McPherson 1980a, Kramer 1981, 1983; Mead and Kramer 1982, Flynn and Kramer 1983, and McPherson and Wilson 1995) into a species incidence table. Specimen incidence records were compiled with literature records, but independently annotated.

Photographs were taken using a Nikon SMZ-1500 Digital Imaging Workstation with Nikon DS-U1 digital Camera and NIS Elements Imaging software (version 3.0). Line drawings were made by Kimberley Shropshire (see acknowledgements) by tracing photographs and rendering detail freehand with reference to specimens.

Total planthopper species richness for Delaware was also evaluated using Chao's (1984) abundance based estimator of species richness calculated as  $S_{chao} = S_{obs} + F_1^2/2F_2$ , where  $S_{obs}$  = # observed species,  $F_1$  = # of species observed by exactly one specimen,  $F_2$  = # of species observed by exactly two specimens.

## Results

Among 1,734 specimens from Delaware we observed 62 planthopper species in 27 genera and 9 families (Table 1), including 55 new state records. Not surprisingly, specimen records were strongly biased (72% of observed specimens) toward New Castle

County where the main campus of University of Delaware is located. Some females in the genera *Bothriocera*, *Cixius*, *Haplaxius*, *Melanoliarius* (all Cixiidae) and *Cedusa* (Derbidae), representing 88 specimens, could not be definitively identified to species and these female specimens were subsequently excluded from the species tally and the calculation of the Chao1 statistic; however, one of the female *Bothriocera* specimens appears to represent an additional species. Specimens of *Omolicna* evidently represented 2 species, but we were unable to identify them or parse the species with confidence. For this reason, we have reported the specimens identified to the generic level and included them in the species count and calculations.

The most abundant species were *Melanoliarius placitus* (18% of observed specimens), *Aphelonema simplex* (10%), *Acanalonia conica* (9%), *Flatormenis chloris* (7%), and *Scolops sulcipes* (5%), collectively representing 49% of the specimens observed (Figure 1). However, for *Aphelonema simplex* there were only 5 collecting events, one of which comprised 70, and a second 69 specimens (out of 165 total observed specimens). In contrast, *Metcalfa pruinosa* (5%) and *Acanalonia bivittata* (3%) were both observed in many collecting events, but these frequently encountered species are readily recognized in the field and either avoided by collectors or not accessioned by the collection manager, and therefore are probably relatively underrepresented.

The Chao1 biodiversity estimator was calculated as 74.08 species, indicating that 12 additional planthopper species are predicted to occur. The incidence list for Delaware and adjacent states (Table 2) includes 112 taxa, of which 50 species were recorded from surrounding states with no Delaware records. In addition, 22 species from MD, 5 from NJ, 8 from PA, and 21 from DC are new state records.

The seasonality data suggests that the optimal time of year to find planthoppers in Delaware is between late June and early August (Table 1). It appears that most species have one generation per year, although the available data is sparse for some taxa. *Bruchomorpha oculata*, *Aphelonema simplex*, and *Cixius nervosus* may have two generations a year. It is evident from specimens collected in logs in March that *Apache degeerii* overwinters as adults (early record March 1: 9 specimens from 3 collection events), although the overwintering status of other taxa is not clear from this data. Records of cixiids from late April may indicate overwintering as immatures, as has been reported for cixiids in Germany (Nickel and Remane 2002).

Specimens reported incidentally by Zuefle (2006) and Zuefle and colleagues (2008) (Table 3) provide host data for 3 Delaware planthopper species. Zuefle (2006) sampled insect use of 45 woody plants that were: 1) native, 2) non-native with native US congeners, and 3) 'alien' plant species with no US congeners, using pesticide knock-down or vacuum sampling. Vouchers were reported in Zuefle (2006) as '*Oliarius sablensis*' were mostly *Melanoliarius ecologus* (32 of 35 dissected males were *M. ecologus* and the remaining 3 *M. sablensis*), so we here reported her cixiids as *Melanoliarius* spp. The hosts for the 3 *M. sablensis* specimens were *Rhododendron mucronatum*, *Cotoneaster lucidus*, and *Betula pendula*. Her results confirm a polyphagous host use for *Flatormenis chloris* and suggest that adult *Melanoliarius*, or at least *M. ecologus*, are polyphagous on woody plants.

**Table 1.** County and seasonality records for Delaware planthoppers. Number of observed specimens given for county records, with distribution of records over the year provided, including earliest and latest observation. For seasonality records, records were divided into early (day 1–15 of the month) and late (remainder of month) observations, and for each observation a specimen count is followed parenthetically by number of independent collecting events (see methods). Sum of seasonality records may be less than sum of specimen records as ambiguous date records were omitted from seasonality tally. Column totals below seasonal entry is a count of the number of species observed during that time interval.

	County records			March		April		May		June		July		August		September		October	
	New Castle	Kent	Sussex	Sum	Late date	Early date	Late	Early	Late	Early	Late	Late	Early	Late	Early	Late	Late	Early	Late
<b>Acanaloniidae</b>																			
<i>Acanalonia bivittata</i>	29	5	11	45	16-May	5-Oct			1(1)			5(1)	8(4)	6(5)	11(6)	9(7)	4(3)	1(1)	
<i>Acanalonia conica</i>	138	10	2	150	2-Jul	7-Oct						16(7)	25(17)	14(10)	24(15)	49(18)	22(8)	22(7)	2(2)
<b>Achilidae</b>																			
<i>Catonia carolina</i>	9		15	24	29-Jul	7-Oct							1(1)	4(1)	6(3)	10(1)	1(1)	1(1)	
<i>Catonia cinctifrons</i>	1			1	26-Jul								1(1)						
<i>Catonia nava</i>	10			10	3-Aug	27-Sep								1(1)				8(1)	
<i>Catonia picta</i>				1	7-Oct														1(1)
<i>Catonia pumila</i>				3	4-Aug	8-Sep								1(1)				2(1)	
<i>Cixidia fusca</i>				4	28-Jul	7-Oct							1(1)						3(1)
<i>Cixidia opaca</i>		1		1	28-Aug										1(1)				
<i>Cixidia variegata</i>				1	7-Oct														1(1)
<b>Caliscelidae</b>																			
<i>Aphelonema simplex</i>	15	141	9	165	30-May	3-Oct			8(1)				9(1)	4(1)	69(1)			70(1)	1(1)
<i>Bruchomorpha sp. n.</i>			7	7	29-Jun	21-Aug							3(2)	2(1)					
<i>Bruchomorpha oculata</i>	53		1	54	22-Jun	9-Oct						4(1)	6(4)		3(2)	15(3)		12(5)	12(6)





	County records			March		April		May		June		July		August		September		October			
	New Castle	Kent	Sussex	Sum	Early date	Late date	Early	Late	Early	Late	Early	Late	Early	Late	Early	Late	Early	Late	Early	Late	
					1	16	3	11	1	2	23	36	14-Jun	7-Oct	1	2	26-Jun	June	15-Jul	26-Jul	9-Aug
<i>Cedusa vulgaris</i>	1			1	3-Aug										1(1)						
<i>Cedusa</i> spp. Female	16			16	9-Jun	26-Jul				1(1)		3(1)	12(1)								
<i>Neocnethra heidemanni</i>	3			3	2-Sep	6-Sep										3(2)					
<i>Omolocna</i> spp.	11	2		23	14-Jun	7-Oct				1(1)		2(1)	1(1)	17(6)	4(3)				7(1)		
<i>Otiocerus coquebertii</i>	1			1	(22-26)-June	12-Jul					1(1)	1(1)									
<i>Otiocerus francilloni</i>				1	15-Jul							1(1)									
<i>Otiocerus reamurii</i>	2			2	26-Jul	9-Aug							1(1)	1(1)							
<i>Otiocerus wolffi</i>	7			5	(19-20)-July	7-Sep							2(2)	4(3)	6(2)						
<i>Patana vanduzeei</i>	2			2	3-Jul	4-Aug						1(1)	2(1)	1(1)							
<i>Sikaitana harri</i>		1		1	1-Jul							1(1)									
<b>Dictyopharidae</b>																					
<i>Rhynchomirina lingula</i>				14	14	21-Aug	1-Sep								10(2)	4(1)					
<i>Rhynchomirina microrhina</i>	16	1		18	27-Jul	13-Sep						3(2)	4(3)	11(7)							
<i>Scolops angustatus</i>	8			8	18-Jul	20-Jul							2(2)								
<i>Scolops perdis</i>				7	19-Aug	21-Aug									7(2)						
<i>Scolops pungens</i>				1	17-Jul								1(1)								
<i>Scolops sulcipes</i>	92	2		94	11-Jul	1-Oct						1(1)	50(14)	4(4)	12(4)	3(1)	11(4)	10(1)			

	County records			March		April		May		June		July		August		September		October				
	New Castle	Kent	Sussex	Sum	Early date	Late date	Early	Late	Early	Late	Early	Late	Early	Late	Early	Late	Early	Late				
					21-Oct	2-Jul	21-Oct	22-Jun	11-Oct	22-Jul	1-Oct	21-Oct	2-Jul	11-Oct	22-Jul	1-Oct	21-Oct	2-Jul	11-Oct	22-Jul	1-Oct	
<b>Flatidae</b>																						
<i>Flatormenis chloris</i>	85	18	9	112	2-Jul	21-Oct						2(1)	14(12)	16(11)	23(18)	19(12)	15(12)	8(5)	11(3)			
<i>Metacalfa pruinosa</i>	58	12	13	83	22-Jun	11-Oct				2(2)		6(6)	15(11)	20(14)	12(7)	14(10)	5(5)	4(3)				
<i>Ormenoides venusta</i>	18	2	1	21	22-Jul	1-Oct							2(2)	2(2)	8(3)	5(3)		2(1)				
<b>Fulgoridae</b>																						
<i>Cyrtoptus befragei</i>	4			4	4-Jun	18-Jul				2(2)	1(1)		1(1)									
<b>Issidae</b>																						
<i>Thionia bullata</i>	1	1		2	2-Aug	17-Oct								1(1)					1(1)			
<i>Thionia simplex</i>	14			14	4-Jul	16-Sep						2(2)	1(1)	1(1)	2(2)	7(5)	1(1)					
<b>Totals</b>	1253	209	272	1734			1	0	0	3	2	6	9	24	28	36	27	19	17	13	14	5




Figure 1. Rank abundance frequency distribution of planthopper species of Delaware. Number of specimens of each species found in Table 1.

**Table 2.** Planthopper incidence list for Delaware, Maryland New Jersey, Pennsylvania, and the District of Columbia. Specimen records are indicated by “S”, literature records by “L”, tentative or subsequent questioned records are annotated by “?”, and records reported as erroneous by “E”. Records from Wilson and McPherson (1980) except as noted.

Species	DE	MD	NJ	PA	DC	References and comments
<b>Acanaloniidae</b>						
<i>Acanalonia bivittata</i> (Say, 1825)	S	S,L	S,L	S,L	S,L	
<i>Acanalonia conica</i> (Say, 1830)	S,L	S,L	S,L	S,L	S	
<i>Acanalonia servillei</i> Spinola, 1839		S		S,L		<i>Acanalonia latifrons</i> (Walker, 1851) synonymized with <i>A. servillei</i> by Fennah 1971: 334-6.
<b>Achilidae</b>						
<i>Catonia carolina</i> (Metcalf, 1923)	S	S,L	S		S	
<i>Catonia cinctifrons</i> (Fitch, 1856)	S,L	S,L	S,L	S,L	S	
<i>Catonia lunata</i> Metcalf, 1923		S,L	S,L		S	
<i>Catonia nava</i> (Say, 1830)	S	S,L			S	
<i>Catonia picta</i> Van Duzee, 1908	S		S,L			
<i>Catonia pumila</i> Van Duzee, 1908	S	S,L	S,L	S,L	S	
<i>Cixidia fusca</i> (Walker, 1852)	S	S	S,L		S	
<i>Cixidia opaca</i> (Say, 1830)	S	L	L	S,L		
<i>Cixidia pallida</i> (Say, 1830)			L	L	S,L	
<i>Cixidia septentrionalis</i> (Provancher, 1889)			L	L	L	
<i>Cixidia variegata</i> (Van Duzee, 1908)	S	S	S,L		S	
<i>Synecoche dimidiata</i> (Van Duzee, 1910)		S,L	S,L	S,L		
<i>Synecoche grisea</i> (Van Duzee, 1908)		S,L		L		
<i>Synecoche impunctata</i> (Fitch, 1851)		S	L	L	S	
<b>Caliscelidae</b>						
<i>Aphelonema decorata</i> (Van Duzee, 1908)			L			
<i>Aphelonema histrionica</i> (Stål, 1864)		S				
<i>Aphelonema rugosa</i> (Ball, 1932)		L?				
<i>Aphelonema simplex</i> Uhler, 1876	S	S,L	S,L			
<i>Bruchomorpha dorsata</i> Fitch, 1856			S,L	L		
<i>Bruchomorpha jocosa</i> Stål, 1862			L		L	
<i>Bruchomorpha oculata</i> Newman, 1838	S	S,L	S,L	S,L	S	
<i>Bruchomorpha pallidipes</i> Stål, 1862	S	S	S		S,L	
<i>Bruchomorpha</i> sp. n.	S	S	S			
<i>Bruchomorpha tristis</i> Stål, 1862			L		L	
<i>Fitchiella robertsonii</i> (Fitch, 1856)		L		L		
<b>Cixiidae</b>						
<i>Bothriocera bicornis</i> (Fabricius, 1803)		E	E			Noted as error by Kramer 1983
<i>Bothriocera cognita</i> Caldwell, 1943	S,L		S,L	L		Kramer 1983
<i>Bothriocera drakei</i> Metcalf, 1923	S					
<i>Bothriocera maculata</i> Caldwell, 1943	S					

Species	DE	MD	NJ	PA	DC	References and comments
<i>Bothriocera signoreti</i> Stål, 1864		E				Noted as error by Kramer 1983
<i>Cixius angustatus</i> Caldwell, 1938	S	S				
<i>Cixius apicalis</i> Metcalf, 1923				L		Kramer 1981
<i>Cixius coloepeum</i> Fitch, 1856		S	L	L		
<i>Cixius misellus</i> Van Duzee, 1906			L	L		Kramer 1981 (PA record)
<i>Cixius nervosus</i> (Linnaeus, 1758)	S,L	S,L	S,L	S,L		Kramer 1981
<i>Cixius nike</i> Kramer, 1981				S,L		Kramer 1981
<i>Cixius pini</i> Fitch, 1851		S,L		S,L	S	Kramer 1981
<i>Cixius stigmatus</i> (Say, 1825)			L	L		
<i>Haplaxius enotatus</i> (Van Duzee, 1909)		S,L				
<i>Haplaxius ovatus</i> (Ball, 1933)	S	S,L	S,L			
<i>Haplaxius pictifrons</i> (Stål, 1862)	S	S,L	L	S,L	S	
<i>Haplaxius pusillus</i> (Van Duzee, 1909)			S,L			
<i>Haplaxius radicus</i> (Osborn, 1903)	S	S,L			S	
<i>Haplaxius wheeleri</i> (Wilson, 1996)			S,L	S		
<i>Melanoliarius chuliotus</i> (Ball, 1934)	S				L	
<i>Melanoliarius ecologus</i> Caldwell, 1947	S	S,L	S,L	S,L	S,L	Mead and Kramer 1982(MD/NJ/PA)
<i>Melanoliarius humilis</i> (Say, 1830)	S	S,L	L	S,L	L	Mead and Kramer 1982(NJ/PA)
<i>Melanoliarius montanus</i> (Metcalf, 1923)	S	S,L	S	S,L		Mead and Kramer 1982
<i>Melanoliarius placidus</i> Van Duzee, 1912	S	S,L	S,L	S	L	Mead and Kramer 1982
<i>Melanoliarius quinquelineatus</i> (Say, 1830)	S	S,L	S,L	S,L		Mead and Kramer 1982
<i>Melanoliarius sablensis</i> (Caldwell, 1951)	S	S,L	S,L	S,L	L	Mead and Kramer 1982(MD/NJ/PA)
<i>Melanoliarius near sablensis</i>	S	S		S		
<i>Melanoliarius spp. females</i>	S	S		S		
<i>Pentastiridius cinnamomeus</i> (Provancher, 1889)	S		L	L		Kramer 1981
<i>Oecleus borealis</i> Van Duzee, 1912		S,L	S,L	S,L	S	
<i>Oecleus productus</i> Metcalf, 1923	S	S,L				
<i>Pintalia delicata</i> (Fowler, 1904)		S,L				
<i>Pintalia vibex</i> Kramer, 1981	S	S,L				Kramer 1981
<b>Derbidae</b>						
<i>Anotia bonnetii</i> Kirby, 1821			L			
<i>Anotia burnettii</i> Fitch, 1856				L		
<i>Anotia fitchi</i> (Van Duzee, 1893)				L		
<i>Anotia kirkaldyi</i> Ball, 1902	S			S,L		
<i>Anotia robertsonii</i> Fitch, 1856	S		L	L		
<i>Anotia westwoodi</i> Fitch, 1856	S	S	L	S,L	S	
<i>Apache degeerii</i> (Kirby, 1821)	S	S,L	S,L	S,L		
<i>Cedusa carolinensis</i> Flynn & Kramer, 1983	S	S,L			S	Flynn and Kramer 1983
<i>Cedusa cedusa</i> McAtee, 1924		S,L			S	Flynn and Kramer 1983
<i>Cedusa chuluota</i> Ball, 1928			S,L			Flynn and Kramer 1983

Species	DE	MD	NJ	PA	DC	References and comments
<i>Cedusa edentula</i> (Van Duzee, 1912)		S	L		S	Flynn and Kramer 1983
<i>Cedusa gedusa</i> McAtee, 1924		S,L	S,L	S,L		Flynn and Kramer 1983
<i>Cedusa hedusa</i> McAtee, 1924		S,L				Flynn and Kramer 1983
<i>Cedusa incisa</i> (Metcalf, 1923)		S		S,L		Flynn and Kramer 1983
<i>Cedusa kedusa</i> McAtee, 1924	S	S,L	S,L		S	Flynn and Kramer 1983
<i>Cedusa maculata</i> (Van Duzee, 1912)		S,L		S,L		Flynn and Kramer 1983
<i>Cedusa mallochi</i> McAtee, 1924	S	L				Flynn and Kramer 1983
<i>Cedusa obscura</i> (Ball, 1902)		S,L	S,L		S	Flynn and Kramer 1983
<i>Cedusa redusa</i> McAtee, 1924	S	S,L				Flynn and Kramer 1983
<i>Cedusa shawi</i> Flynn & Kramer, 1983		S,L				Flynn and Kramer 1983
<i>Cedusa vulgaris</i> (Fitch, 1851)	S	S,L				Flynn and Kramer 1983
<i>Cedusa</i> spp. Females	S	S		S		
<i>Neocenchrea heidemanni</i> (Ball, 1902)	S	L	L		L	
<i>Omolicna uhleri</i> (Ball, 1902)		L	L		L	
<i>Otiocerus amyotii</i> Fitch, 1856			L	L	L	
<i>Otiocerus coquebertii</i> Kirby, 1821	S	L	L	S,L		
<i>Otiocerus francilloni</i> Kirby, 1821	S		L			
<i>Otiocerus reaumurii</i> Kirby, 1821	S					
<i>Otiocerus signoretii</i> Fitch, 1856		S	L	L		
<i>Otiocerus stollii</i> Kirby, 1821		S	L	L		
<i>Otiocerus wolfii</i> Kirby, 1821	S	S,L	L	S,L		
<i>Patara vanduzeei</i> Ball, 1902	S	S		L		
<i>Shellenius ballii</i> (McAtee, 1923)		S,L				
<i>Shellenius schellenbergii</i> (Kirby, 1821)			L			
<i>Sikaiana harti</i> (Metcalf, 1923)	S	S				
<b>Dictyopharidae</b>						
<i>Mitrops dioxys</i> (Walker, 1858)		L	L			
<i>Phylloscelis atra</i> Germar, 1839		L	L	S,L	L	
<i>Phylloscelis pallescens</i> Germar, 1839		L	L	L		
<i>Phylloscelis rubra</i> Ball, 1930			S,L			
<i>Rhynchomitra lingula</i> (Van Duzee, 1908)	S	S	S,L			
<i>Rhynchomitra microrrhina</i> (Walker, 1851)	S	S	L	S		
<i>Scolops angustatus</i> Uhler, 1929	S	S,L	S,L		L	
<i>Scolops grossus</i> Uhler, 1876			L?			Record probably in error.
<i>Scolops perdix</i> Uhler, 1900	S	L	S,L	S	L	
<i>Scolops pungens</i> (Germar, 1830)	S	L	S,L	L	L	
<i>Scolops sulcipes</i> (Say, 1825)	S	S,L	S,L	S,L	L	
<b>Flatidae</b>						
<i>Flatormenis chloris</i> (Melichar, 1902)	S	S,L	S,L	S,L	L	<i>Anormenis septentrionalis</i> auct. (nec. Spinola, 1839) synonymized with <i>A. chloris</i> by O'Brien 1985: 657-660, and transferred to <i>Flatormenis</i> by Medler 2003: 593.

Species	DE	MD	NJ	PA	DC	References and comments
<i>Cyarda melichari</i> Van Duzee, 1907					L	Species needs confirmation.
<i>Metcalfa pruinosa</i> (Say, 1830)	S,L	S,L	S,L	S,L	L	
<i>Ormenoides venusta</i> (Melichar, 1902)	S	S	S	S	S	
<b>Fulgoridae</b>						
<i>Cyrpoptus belfragei</i> Stål, 1869	S	L				
<i>Poblícia fuliginosa</i> (Olivier, 1791)		S				
<b>Issidae</b>						
<i>Exortus punctiferus</i> (Walker, 1851)			L			Originally reported by Smith (1890) as <i>Issus aciculatus</i> Uhler, 1876, possibly in error.
<i>Thionia bullata</i> (Say, 1830)	S	S	S,L	S,L	S,L	
<i>Thionia elliptica</i> (Germar, 1830)		S	L		L	
<i>Thionia simplex</i> (Germar, 1830)	S	S,L	S,L	S	L	
New records*	55	22	5	8	21	
Total species*	62	88	74	60	46	

\*Unidentified females and errors excluded, *Melanoliarus* near *sablensis* included with *M. sablensis*, 2 species of *Omolicna* counted for Delaware.

**Table 3.** Planthoppers reported by Zuefle (2006) collected in Delaware by host sampled 2004–2005. Host species were segregated into 3 categories; **1** Native woody plants **2** Non-native plants congeneric with US species; and **3** “Alien” woody plants - those with no US congeners. The *Melanoliarus* species were reported as ‘*Oliarius sablensis*’, but voucher specimens in the UDCC were found to be mostly *Melanoliarus ecologus* with a few *M. sablensis*.

Plant Family	Plant species	Planthopper species		
		<i>Melanoliarus</i> spp.	<i>Flatormenis chloris</i>	<i>Thionia simplex</i>
<b>Native</b>				
Aceraceae	<i>Acer rubrum</i>	0	0	0
Betulaceae	<i>Betula nigra</i>	0	2	0
Betulaceae	<i>Carpinus caroliniana</i>	2	0	0
Cornaceae	<i>Cornus florida</i>	0	0	0
Fagaceae	<i>Fagus grandifolia</i>	0	0	0
Hamamelidaceae	<i>Hamamelis virginiana</i>	0	0	0
Juglandaceae	<i>Juglans nigra</i>	4	0	0
Moraceae	<i>Morus rubra</i>	6	0	0
Rosaceae	<i>Prunus serotina</i>	0	2	0
Ericaceae	<i>Rhododendron periclymenoides</i>	7	0	0
Rosaceae	<i>Rosa carolina</i>	2	0	0

Plant Family	Plant species	Planthopper species		
		<i>Melanoliarus</i> spp.	<i>Flatormenis</i> <i>chloris</i>	<i>Thionia</i> <i>simplex</i>
Salicaceae	<i>Salix nigra</i>	1	1	1
Tiliaceae	<i>Tilia americana</i>	1	0	0
Ulmaceae	<i>Ulmus americana</i>	0	0	0
Caprifoliaceae	<i>Viburnum dentatum</i>	8	0	0
	<b>Subtotal</b>	31	5	1
<b>Non-native congeneric plants</b>				
Aceraceae	<i>Acer platanoides</i>	0	2	0
Betulaceae	<i>Betula pendula</i>	3	0	0
Betulaceae	<i>Carpinus betulus</i>	0	0	0
Cornaceae	<i>Cornus kousa</i>	2	0	0
Fagaceae	<i>Fagus sylvatica</i>	3	0	0
Hamamelidaceae	<i>Hamamelis mollis</i>	2	0	0
Juglandaceae	<i>Juglans regia</i>	2	0	0
Moraceae	<i>Morus alba</i>	0	0	0
Rosaceae	<i>Prunus serrulata</i>	1	1	0
Ericaceae	<i>Rhododendron mucronatum</i>	42	0	0
Rosaceae	<i>Rosa multiflora</i>	7	1	0
Salicaceae	<i>Salix babylonica</i>	0	0	0
Tiliaceae	<i>Tilia cordata</i>	2	0	0
Ulmaceae	<i>Ulmus parvifolia</i>	2	5	0
Caprifoliaceae	<i>Viburnum dilatatum</i>	14	2	0
	<b>Subtotal</b>	80	11	0
<b>Alien plants</b>				
Lardizabalaceae	<i>Akebia quinata</i>	9	0	0
Fabaceae	<i>Albizia julibrissin</i>	5	1	0
Rosaceae	<i>Cotoneaster lucidus</i>	16	2	0
Fabaceae	<i>Cytisus scoparius</i>	9	1	0
Oleaceae	<i>Forsythia suspensa</i>	10	0	0
Ginkgoaceae	<i>Ginkgo biloba</i>	1	0	0
Araliaceae	<i>Hedera helix</i>	6	0	0
Sapindaceae	<i>Koelreuteria paniculata</i>	1	0	0
Lythraceae	<i>Lagerstroemia indica</i>	5	0	0
Oleaceae	<i>Ligustrum vulgare</i>	6	1	0
Scrophulariaceae	<i>Paulownia tomentosa</i>	1	0	0
Rutaceae	<i>Phellodendron amurens</i>	0	0	0
Rutaceae	<i>Poncirus trifoliata</i>	1	0	0
Rosaceae	<i>Pyrus pashia</i>	4	0	0
Oleaceae	<i>Syringa vulgaris</i>	1	0	1
	<b>Subtotal</b>	75	5	1
	<b>Total</b>	<b>186</b>	<b>21</b>	<b>2</b>

## Systematics

Artificial key to genus and select planthopper species from Delaware and vicinity.

- 1 Hind tibiae with large movable spur at apex (Fig. 2A)..... **Delphacidae**  
 – Hind tibiae without movable spur at apex (e.g., Figs 2B–D)..... **2**  
 2 Second tarsomere of hind legs with row of apical spines (Fig. 2E) ..... **3**  
 – Second tarsomere of hind legs with one apical spine on each side (Fig. 2F) or spines absent..... **7**  
 3 Larger species, greater than 10 mm, with patterned forewings (Figs 3H, I); hindwings with numerous cross veins near apex and in anal area; uncommon in study area ..... **Fulgoridae, 71**  
 – Mostly smaller species, forewings variable; hindwings without cross veins near apex or in anal area..... **4**  
 4 Forewings overlapping posteriorly (Figs 4G–L, 5F–L, 6F–H), trailing margins angled; body flattened ..... **Achilidae, 13**  
 – Forewings not overlapping posteriorly; body variable..... **5**  
 5 Beak with apical segment subequal in length and width (except *Cedusa*); forewings often with tubercles on claval veins (Figs 8G, 9B); antennae may bear projections (Figs 10E, F) or subtended by a shelf-like structure (Figs 10A–D); median carina of frons often absent; parameres of male much longer than pygofer ..... **Derbidae (most), 41**  
 – Beak with apical segment longer than wide; forewings without tubercles on claval veins (or with tubercles on all veins); antennae never bearing projections or subtended by a shelf-like structure; median carina of frons present; parameres of male shorter than length of pygofer..... **6**  
 6 Frons with two or three median carinae and/or head with elongate anterior projection (Figs 13–14); median ocellus absent; wing vein tubercles usually absent ..... **Dictyopharidae, 60**  
 – Frons with one median carina; head not elongate; median ocellus usually present above frontoclypeal suture (Figs 6D–E, 7B, H); usually with tubercles on veins of wings ..... **Cixiidae, 35**  
 7 Forewings with tubercles on claval veins (e.g., Figs 8G, 9B), if tubercles present in claval area (Figs 3D–G) then forewings waxy with row of many small peripheral cells; beak with apical segment subequal in length and width; frons often compressed with median carina absent (Figs 10B, D, G); parameres much longer than pygofer..... **Derbidae (few), 41**  
 – Forewings without tubercles on claval veins (or with tubercles on all veins); beak with apical segment longer than wide; frons not compressed, median carina generally present (e.g., Figs 12, 15A–D); parameres shorter than length of pygofer ..... **8**  
 8 Forewings waxy, bearing tubercles between veins on clavus (Figs 3D–G) and with numerous costal crossveins..... **Flatidae, 68**



**Figure 2.** Hind legs of planthoppers. **A** Delphacidae, tibia with calcar **B** Acanaloniidae, tibia without spines **C** Caliscelidae, tibia with 1 spine **D** Issidae, tibia with 2 spines **E** Dictyopharidae, second tarsal segment with row of teeth **F** Acanaloniidae, second tarsal segment with pair of spines.


- Forewings not waxy, without tubercles on clavus; without numerous costal crossveins (e.g., Figs 3A–C, 15E–H) ..... **9**
- 9 Hind tibiae without lateral spines (Fig. 2B); forewings with reticulate venation, usually extending to apex of abdomen (even in brachypters); usually green (occasionally pink) (Figs 3A–C)..... **Acanaloniidae, 11**
- Hind tibiae with lateral spines (Figs 2C–D); forewing venation not reticulate (Figs 15E–H), brachypters may have forewings short (Fig. 11), exposing several segments in dorsal view; color not green, usually brown, black, or straw (pinkish in males of 1 species)..... **10**
- 10 Usually brachypterous with forewings shorter than abdomen (Fig. 11); frons with sublateral carinae bordering a large disc-like or elongate areolet, sublateral carinae of frons meeting ventrally (or nearly so) (Fig. 12); hind tibiae with single lateral spine (Fig. 2C)..... **Caliscelidae, 26**
- Forewings covering abdomen (both brachypters and macropters) (Figs 15E–H); frons with median carina, with or without sublateral carinae; if present, not meeting ventrally (Figs 15A–D); hind tibiae with two lateral spines (Fig. 2D)..... **Issidae, 72**

### **Acanaloniidae**

- 11 Body green (rarely pink) with conspicuous brownish to reddish marking along lateral portions of thoracic nota (Fig. 3B), continuing onto wings ..... ***Acanalonia bivittata***
- Body uniformly green (rarely pink) (Figs 3A, C); may have middorsal vitta on thorax ..... **12**
- 12 Head distinctly produced conically (Fig. 3A); without prominent median carina across vertex and thorax; abundant in Mid-Atlantic states..... ***Acanalonia conica***
- Head not produced conically (Fig. 3C); with prominent median carina across vertex and thorax; southeastern species occasional in Mid-Atlantic States ..... ***Acanalonia servillei***


### **Achilidae**

- 13 Head, including eyes, less than 2/3 as wide as pronotum (Figs 5F–J) (Myconini) ..... ***Cixidia, 14***
- Head including eyes at least 2/3 as wide as pronotum (Figs 4G–L, 6F–H) (Plectoderini) ..... **18**
- 14 Clypeus and upper half of frons dark brown or black, strongly contrasting with pale lower half of frons (Fig. 5B) ..... ***Cixidia opaca***
- Frons more uniformly colored, upper half not strongly contrasting (Figs 5A, C–E)..... **15**
- 15 Vertex short, projecting in front of eye for distance less than length of eye (Fig. 5J); frons distinctly and uniformly speckled (Fig. 5E)... ***Cixidia variegata***




**Figure 3.** Lateral habitus of Acanaloniidae, Flatidae, and Fulgoridae. **A** *Acanalonia conica* **B** *A. bivittata* **C** *A. servillei* **D** *Flatormenis chloris* **E** *Cyarda* sp. **F** *Metcalfa pruinosa* **G** *Ormenoides venusta* **H** *Cyrpoptus belfragei* **I** *Poblizia fuliginosa*.

- Vertex elongate, projecting in front of eye for distance equal to or greater to length of eye; frons more uniformly colored (Figs 5A, C)..... **16**
- 16 Frons and clypeus uniformly colored (Fig. 5D)..... *Cixidia septentrionalis*
- Clypeus distinctly darker than frons (Figs 5A, C)..... **17**
- 17 Vertex projected in front of eye for distance greater than eye length, vertex 1.3–1.5× as long as basal width (Fig. 5F); frons and clypeus about as dark as pronotum; forewings nearly uniform brown ..... *Cixidia fusca*
- Vertex projected in front of eye for distance about equal to eye length, vertex length about equal (1–0.95x) to basal width (Fig. 5H); frons and clypeus paler than pronotum; forewing variegated with grayish white ..... *Cixidia pallida*
- 18 Subcostal cell of forewing longer than 1/3 length of forewing, narrow throughout (Fig. 16B); medioventral lobe of male pygofer entire (Fig. 16G)..... *Synecdoche*, **19**
- Subcostal cell of forewing about 1/3 length of forewing, wider before its apex (Fig. 16A); medioventral lobe of male pygofer apically bifurcate (Fig. 16F)... *Catonia*, **21**
- 19 Frons entirely pale (Fig. 6B)..... *Synecdoche grisea*
- Frons with dark transverse bands or all dark (Figs 6A, C)..... **20**
- 20 Frons with dark bands (Fig. 6C) ..... *Synecdoche impunctata*
- Frons uniformly dark, contrasting with pale clypeus (Fig. 6A) ..... *Synecdoche dimidiata*



**Figure 4.** Habitus of *Catonia* (Achilidae) (A–F frons, G–K dorsal view). A, G *Catonia carolina* B, H *C. cinctifrons* C, I *C. lunata* D, J *C. nava* E, K *C. picta* F, L *C. pumila*.




**Figure 5.** Habitus of *Cixidia* (Achilidae) (A–E frons, F–J dorsal view). A, F *Cixidia fusca* B, G *C. opaca* C, H *C. pallida* D, I *C. septentrionalis* E, J *C. variegata*.


- 21 Upper dark band of frons mottled, distinctly paler than lower band (Fig. 4D); larger species usually more than 5.8 mm ..... *Catonia nava*
- Frons, if banded (Figs 4A–C, E), with upper dark band not mottled and not paler than lower, or frons not dark banded (Fig. 4F); size less than 6.2 mm ...  
..... 22
- 22 Frons with two very dark transverse bands (Figs 4B, E)..... 23
- Frons pale with pale bands, or uniformly pale (Figs 4A, C, F)..... 24
- 23 Lower dark band distinctly paler near frontoclypeal suture giving frons a tri-colored appearance (Fig. 4E); body often with orangish cast ... *Catonia picta*
- Lower dark band uniformly dark (Fig. 4B); body brown or grayish.....  
..... *Catonia cinctifrons*
- 24 Pale transverse marking at frontoclypeal suture not reaching lateral margin of frons (Figs 4A, C) ..... 25
- Frons uniformly colored or pale transverse marking at frontoclypeal suture extending to lateral margin of frons (Fig. 4F)..... *Catonia pumila*
- 25 Pale transverse marking at level of ocelli complete, reaching lateral margin of frons (Fig. 4C) ..... *Catonia lunata*
- Pale transverse marking at level of ocelli incomplete, not reaching lateral margin of frons (Fig. 4A) ..... *Catonia carolina*

**Caliscelidae**

- 26 Head produced into weevil-like snout (Figs 11E–H); usually black..... 30
- Head not produced (Fig. 11A–D); paler ..... *Aphelonema*, 27
- 27 Vertex very broad, width at least 5–6× median length (Figs 12A, D); frons greatly exposed above, fastigium rounded when viewed laterally; mostly straw to pink colored (Figs 11A, D), may have darker wings and abdomen..... 28
- Vertex longer, width 2–3× median length, frons not as exposed from above (Figs 12B–C); fastigium angled when viewed laterally; mostly black and pale colored (Figs 11B–C)..... 29
- 28 Head and thorax orange-tan, rest of dorsum blackish brown (Fig. 11A, especially in males); central frontal tablet of frons pointed below (Fig. 12A); found mostly in the southeast, reported from NJ ..... *Aphelonema decorata*
- Uniformly pale ochreous (females) to pink (most males) in color (Fig. 11D); central frontal tablet of frons almost circular (Fig. 12D).... *Aphelonema simplex*
- 29 When viewed from the side, fastigium of head produced forward, frons slanted; vertex somewhat triangular (Fig. 11B)..... *Aphelonema histrionica*
- When viewed from the side, fastigium not produced, frons not slanted; vertex broadly rounded anteriorly (Fig. 11C) ..... *Aphelonema rugosa*
- 30 Middle and front tibiae expanded ..... *Fitchiella robertsonii*
- Middle and front tibiae not expanded..... *Bruchomorpha*, 31
- 31 Dorsal light stripe broad and conspicuous, extending from near apex of face to apex of forewings or beyond ..... *Bruchomorpha sp. n.*
- Dorsal light stripe not broad and conspicuous, generally of lesser extent ... 32



**Figure 6.** Habitus of *Syneccoche* (Achilidae), *Haplaxius* and *Oecleus* (Cixiidae) (**A–F** frons, **F–J** dorsal view). **A, F** *Syneccoche dimidiata* **B, G** *S. grisea* **C, H** *S. impunctata* **D, I** *Haplaxius pictifrons* **E, J** *Oecleus borealis*.



**Figure 7.** Habitus of Cixiidae. **A** *Bothriocera cognita*, head, lateral view **B** same, frons **C** same, dorsal view **D** *Cixius pini*, dorsal view **E** *Melanoliarius placidus*, head, lateral view **F** same, dorsal view **G** *Pentastiridius cinnamomeus*, dorsal view **H** *Pintalia vibex*, frons **I** same, lateral view **J** same, dorsal view.

- 32 Nasal process distinctly pronounced, head concave ventrally in lateral view (Fig. 11F); in dorsal view extending anteriorly beyond eye for a distance equal or greater than length of eye.....***Bruchomorpha oculata***
- Nasal process less pronounced, head weakly convex ventrally; in dorsal view extending anteriorly beyond eye for a distance less than length of eye (Figs 11E, G–H).....**33**
- 33 Reddish-brown in color with a dark spot on clypeus..***Bruchomorpha jocosus***
- Uniformly black, usually with light stripe on vertex (sometimes reaching thorax) .....**34**
- 34 Legs pale (Fig 11G); small species, less than 2.6 mm.....
- .....***Bruchomorpha pallidipes***
- Legs dark (Fig 11H); large species, more than 2.6 mm.....***Bruchomorpha tristis***

**Cixiidae**

- 35 Antennae arising from elongated cup-like cavities anterior to eyes (Fig. 7A)..
- .....***Bothriocera***
- Antennae not within cup-like cavities, arising below eyes (Fig. 7E, 7I).....**36**
- 36 Hind tibiae without spines (similar to Fig. 2B) .....**37**
- Hind tibiae with one or more spines along axis before apex (similar to Figs 2C–D).....**38**
- 37 Mesonotum with 5 carinae; crown strongly narrowed (Fig. 6J) .....***Oecleus***
- Mesonotum with 3 carinae; crown slightly narrowed (Fig. 6I) ..... ***Haplaxius***
- 38 Mesonotum with 5 longitudinal carinae (although intermediate pair sometimes obsolete); posterior margin of crown angularly incised (Figs 7F–G) ...
- .....**39**
- Mesonotum with 3 carinae; posterior margin of crown quadrately or roundly incised (Figs 7D, J) .....**40**
- 39 Apex of basitarsus of hind leg with 12 teeth ..... ***Pentastiridius***
- Apex of basitarsus of hind leg with no more than 10 teeth ..... ***Melanoliarius***
- 40 Forewings roof-like in position with distal portions clearly separated (Fig. 7D); spines on hind tibiae conspicuous ..... ***Cixius***
- Forewings vertical in position with distal portions oppressed (Figs 7I–J); spines on hind tibiae inconspicuous..... ***Pintalia***


**Derbidae**

- 41 Clavus open (Figs 16C–D; combined anal veins reaching posterior cubitus and usually curved to follow wing margin); most taxa with head projecting well beyond eyes in lateral view (e.g., Figs 10E–F); frons very narrow (Fig. 10G); forewings twice as long as body or more, delicate appearing (Otiocerinae: Otiocerini and Sikaianini).....**42**
- Clavus closed (Fig. 16E; combined anal veins reaching wing margin within claval area); most taxa with head projecting only slightly beyond eyes



**Figure 8.** Lateral habitus of Derbidae I. **A** *Anotia bonnetii* **B** *A. kirkaldyi* **C** *A. robertsonii* **D** *A. westwoodi* **E** *Apache degeerii* **F** *Neocenchrea beidemanni* **G** *Patara vanduzeei* **H** *Sikaiana harti* **I** *Anotia fitchi* **J** *Anotia westwoodi*, head lateral view; **K** *Sayiana sayi*, head lateral view.


- (Figs 10A, C); frons usually not as narrow (Figs 10B, D) (except *Patara*, see Fig. 8G); forewings not as long, most taxa less delicate (Otiocerinae: Patarini; Cedusinae; and Derbinae: Cenchreini) .....57
- 42 Antennae with 2 or 3 conspicuous appendages (Figs 10E–F) ..... 43
- Antennae lacking appendages (Figs 8J, 10G) ..... 51
- 43 General color uniformly rose or reddish (Fig. 8E); head in lateral view with vertex distinctly concave in apical third and apex pointed (Fig. 10F); dorsal margin of wings in repose sharply angled upward in apical third; forewings with dusky spots in cells.....*Apache degeerii*
- General color white or yellow (e.g., Figs 9D–J), although red markings may be present; head in lateral view with vertex rounded (Fig. 9I, J, 16H, I), or nearly flat (Fig. 10E); dorsal margin of wings straight or curved slightly upward ..... 44
- 44 In lateral view, demarcation between vertex and frons obtusely angular (Fig. 10E) ..... *Otiocerus*, 46
- In lateral view, demarcation between vertex and frons smoothly rounded (Figs 9I, J; 16H–I) ..... *Shellenius*, 45
- 45 Head in lateral view 1.5× as long as broad (Figs 9J, 16I); forewing brownish apically in trailing portion of wing; red markings reduced or absent .....  
..... *Shellenius schellenbergii*



**Figure 9.** Lateral habitus of Derbidae II. **A** *Cedusa* sp. **B** *Omolicna* sp. **C** *Sayiana sayi* **D** *Otiocerus coquebertii* **E** *O. francilloni* **F** *O. reaumurii* **G** *O. stollii* **H** *O. wolfii* **I** *Shellenius balli* **J** *S. schellenbergii*.

- Head in lateral view 2.0× as long as broad (Figs 9I, 16H); forewings with very pale brown markings widely distributed; with red markings on head and wing ..... *Shellenius balli*
- 46 Wings with conspicuous round dusky spots in cells (Figs 9E, F, H) ..... 47
- Wings without conspicuous round dusky spots in cells (Figs 9D, G) ..... 50
- 47 Apical margin of forewings with a row of spots in the cells (Figs 9E, H) ... 48
- Spots not in row within apical cells (Fig. 9F) ..... 49
- 48 Apex of head with a black line laterally followed by a broader red line (Fig. 9H); forewings with spots throughout ..... *Otiocerus wolfii*
- Apex of head without a black line laterally (Fig. 9E); forewings with spots mostly in proximal half ..... *Otiocerus francilloni*
- 49 Forewings with a large black spot on the sutural margin (in the clavus) and four smaller ones in a square, including 1 in costal cell.... *Otiocerus signoretii*
- Forewings with spots arranged differently from above (Fig. 9F) ..... *Otiocerus reaumurii*

- 50 Color of the wings dark, without distinct band (Fig. 9G) ... ***Otiocerus stollii***  
 – Color of the wings pale with distinct reddish forked band (Fig. 9D).....  
 ..... ***Otiocerus coquebertii***
- 51 In lateral view, head projecting in front of eyes for a distance of less than half  
 width of eyes; forewings with scattered spots..... ***Sikaiana harti***  
 – In lateral view, head projecting in front of eyes for a distance subequal to  
 width of eyes (Fig. 8J); color mostly following veins ..... ***Anotia*, 52**
- 52 Costa narrow; forewings with veins not crowded together to give appearance  
 of a stigma (Figs 8A–D); some or most veins of forewings with smoky bor-  
 ders..... **53**  
 – Costa broader; Sc and R vein tips crowded together to give appearance of a stig-  
 ma (Fig. 8I); forewings more extensively marked with fuscous ..... ***Anotia fitchi***
- 53 First 3 segments of abdomen with middorsal black stripe .... ***Anotia burnetii***  
 – Abdomen without middorsal black stripe ..... **54**
- 54 Forewings mostly pale with a few fuscous marked crossveins (Fig. 8C); apex  
 of forewing without dark round spots ..... ***Anotia robertsonii***  
 – Forewings more extensively marked; most veins with smoky borders (Figs 8B,  
 D); apex of forewing often with dark round spots..... **55**
- 55 Head with a single marking, below antennae; apical border of forewings with  
 four dark round spots in the cells (Fig. 8A) ..... ***Anotia bonnetii***  
 – Head with dark or red markings above and below antennae; apical border of  
 forewings usually without round spots in the cells..... **56**
- 56 At least some veins dark in color (Fig. 8B) ..... ***Anotia kirkaldyi***  
 – All veins pale (Fig. 8D) ..... ***Anotia westwoodi***
- 57 Antennae terete, subtended by flattened subantennal process from gena or  
 anterior portion of lateral margin of pronotum (Figs 10A–D), often strongly  
 modified into a reversed “c” (in lateral view) directly behind antennae, or  
 strongly keeled; face not strongly compressed, frons evident; clavus at least  
 half as long as whole forewing (Derbinae: Cenchreini, and Cedusinae)..... **58**  
 – Second segment of antennae flattened (more evident in males than females),  
 antennae not subtended by process; lateral margin of pronotum not strongly  
 modified; face strongly compressed, frons keel-like (similar to Fig. 10G); clavus  
 less than half as long as whole forewing (Fig. 9C) (*Otiocerinae*: Patarini).....  
 ..... ***Patara vanduzeei***
- 58 Subantennal process large, extending from gena, completely subtending an-  
 tennae as a shelf (Fig. 10A); reduced (or absent) sensory pits on head and  
 wings; color uniform, near black or deep grey (Fig. 9A), infrequently near  
 white with yellowish brown patches (Cedusinae) ..... ***Cedusa***  
 – Subantennal process extending from pronotum, smaller (Fig. 10C); lateral  
 carinae of vertex and second claval vein with sensory pits; color usually orange  
 to pale (Figs 8F, 9B) (Derbinae: Cenchreini)..... **59**
- 59 Media with more than two branches, connected to cubitus by crossvein; size  
 less than 6 mm, usually distinctly orangish (Fig. 9B)..... ***Omolicna***




**Figure 10.** Heads of Derbidae. **A** *Cedusa* sp. lateral view **B** *Cedusa* sp., frontal view **C** *Omolicna* sp., lateral view **D** *Omolicna* sp., frontal view **E** *Otiocerus wolffi*, lateral view **F** *Apache degeerii*, lateral view **G** *Anotia robertsonii*, frontal view.

- Media and cubitus each with two branches, not connected by crossveins; size over 7 mm; color orangish white (Fig. 8F) ..... *Neocenchrea heidemanni*


**Dictyopharidae**

- 60 Head projected in front of eyes (Figs 13, 14G, H); front femora not foliaceous..... **63**
- Head not projected in front of eyes (Figs 14A–F); front femora foliaceous....  
..... *Phylloscelis*, **61**
- 61 Eight or fewer longitudinal veins on the forewing; color either uniformly black to dark brown in dorsal view or yellowish body with reddish-brown forewings with prominent yellow wing veins (Figs 14C, F); carinae of frons indistinct ..... *Phylloscelis atra*
- With more than 8 longitudinal veins; color not as above; carinae of frons distinct (Figs 14A–B, D–E) ..... **62**
- 62 Veins concolorous with forewings; body black to light reddish brown (Fig. 14D)..... *Phylloscelis rubra*




**Figure 11.** Lateral view of Caliscelidae. **A** *Aphelonema decorata* **B** *A. bistrionica* **C** *A. rugosa* **D** *A. simplex* **E** *Bruchromorpha jocosa* **F** *B. oculata* **G** *B. pallidipes* **H** *B. tristis*.


- Veins of forewings dark mottled with pale; body light grey-brown (Fig. 14E)..... *Phylloscelis pallescens*
- 63 Forewings clear, macropterous; head projection anterior to eyes subequal in width to vertex; body green (Figs 14G–H)..... *Rhynchomitra*, 64
- Forewings patterned, usually brachypterous; head projection anterior to eyes narrower than vertex; body brownish (Fig. 13)..... *Scolops*, 65



**Figure 12.** Frontal view of Caliscelidae. **A** *Aphelonema decorata* **B** *A. histrionica* **C** *A. rugosa* **D** *A. simplex* **E** *Bruchomorpha jocosa* **F** *B. oculata* **G** *B. pallidipes* **H** *B. tristis*.




**Figure 13.** Habitus of *Scolops* (Dictyopharidae). **A**, **E** *Scolops angustatus* **B**, **F** *S. perdix* **C**, **G** *S. pungens* **D**, **H** *S. sulcipes*.



**Figure 14.** *Phylloscelis* and *Rhynchomitra* (Dictyopharidae) (**A–C, G** Dorsal view habitus, **D–F** frontal view, **H** Dorsal view, head and thorax). **A, D** *Phylloscelis rubra* **B, E** *P. pallelescens* **C, F** *P. atra* **G** *Rhynchomitra microrrhina* **H** *R. lingula*.

- 64 Head projection long (Fig. 14G), in dorsal view narrowing anterior to eyes, projected in front of eyes greater than width of vertex; upcurved in lateral view ..... *Rhynchomitra microrrhina*
- Head projection short (Fig. 14H), in dorsal view rather quadrate, projected in front of eyes for distance about width of vertex; not distinctly upcurved in lateral view ..... *Rhynchomitra lingula*
- 65 Costal cell of forewing with costal vein and membrane white (Fig. 13A) ..... *Scolops angustatus*
- Costal cell of forewing with costal vein variegated (Figs 13B–D) ..... **66**
- 66 Forewings reticulate over apical half (especially brachypters), veins margined with dark (Figs 13D, H) ..... *Scolops sulcipes*
- Forewings not reticulate over apical half (Figs B–C, F–G) ..... **67**
- 67 Pronotum and usually vertex with dark markings (Fig. 13G); body with grayish cast ..... *Scolops perdix*
- Pronotum and vertex without dark markings (Fig. 13F); body with brownish cast ..... *Scolops pungens*



**Figure 15.** Issidae (A–D frontal view E–H Lateral view). A, E *Exortus punctiferus* B, F *Thionia bullata* C, G *T. elliptica* D, H *T. simplex*.

**Flatidae**


- 68 Wings much longer than wide, distinctly narrowing caudally to caudal apex (Fig. 3E); brown ..... *Cyarda*
- Wings slightly longer than wide, truncate to broadly rounded caudally (Figs 3D, F–G); green or grey ..... **69**
- 69 Body grey to blackish (Fig. 3F); forewings with single row of marginal cells along apical and trailing margin (set off by a submarginal vein) ..... *Metcalfa pruinosa*
- Body green (Figs 3D, G); forewings with one or two rows of marginal cells.... ..... **70**
- 70 Frons broader than long; forewings with two rows of marginal cells along apical and trailing margin (set off by two submarginal veins) (Fig. 3D); wings usually rather truncate apically..... *Flatormenis chloris*
- Frons longer than broad; forewings with one row of marginal cells (Fig. 3G); wings usually rounded apically (forewings often with orangish cast along apices) ..... *Ormenoides venusta*

**Fulgoridae**

- 71 Forewings and much of body nearly black (Fig. 3I); caudal abdominal tergites red; head in lateral view with frons at acute angle from vertex; flange of head behind eye small ..... *Poblicia fuliginosa*
- Forewings and body mottled (Fig. 3H), predominately reddish brown; abdomen not red; head in lateral view with frons at sharp angle from vertex; flange of head behind eye distinct ..... *Cyrpoptus belfragei*

**Issidae**

- 72 Hind wings absent or rudimentary; smaller insects, less than 4.5 mm (Figs 15A, E); southeastern species, reported from NJ, possibly in error..... *Exortus punctiferus*
- Hind wings present, entire, with strongly marked notches at the joints of the folds, anal area large; larger insects varying from 5.5 to 8.0 mm (Figs 15B–D, F–H) ..... *Thionia*, **73**
- 73 Uniformly colored, lacking proximal bulla (Fig. 15H); carinae of face weak (Fig. 15D)..... *Thionia simplex*
- Body patterned, wings with proximal bulla (Figs 15F–G); carinae of face conspicuous (Figs 15B–C) ..... **74**
- 74 Vertex broader than long; distinctly concave in frontal view with lateral margins elevated (Fig. 15B)..... *Thionia elliptica*
- Vertex longer than broad, slightly concave in frontal view, lateral margins not strongly elevated (Fig. 15C) ..... *Thionia bullata*



**Figure 16.** Line drawings of Achilidae and Derbidae (**A–E** Forewing, head left, costal margin top **F–G** Male genitalia, ventral view **H–I** *Shellenius* spp., head, left lateral view). **A** *Catonia pumila*; **B** *Synecdoche rubella* (Van Duzee, 1910); **C** *Anotia fitchi*; **D** *A. robertsonii*; **E** *Cedusa* sp.; **F** *Catonia picta*; **G** *Synecdoche dimidiata*; **H** *Shellenius balli*; **I** *Shellenius schellenbergii* (**A–B** Redrawn from O’Brien, 1971; **E** redrawn from Metcalf, 1923).

## Discussion

### Biodiversity

This survey brings the known diversity of Delaware planthoppers (excluding Delphacidae) from 7 to 62, plus provides new state records for MD (22), NJ (5), PA (8) and DC (21) providing species counts for those states as 88, 74, 60 and 46 respectively (Table 2). The Chao1 estimator suggests an additional 12 species may be found in the state. State-level incidence records of 112 species (Table 2) provides some basis for speculation of which species might be missing from the current inventory, and might be interpreted to suggest that the true diversity of planthoppers in Delaware may be closer to 100 species. A better understanding of the habits and finer-scale distribution pat-

terns would be desirable in order to construct a candidate list of species not yet detected in the Delaware fauna. However, some species detected were not previously known from the region (viz. *Aphelonema histrionica*, *Bothriocera drakei*, *B. maculata*, *Cixius angustatus*, *Sikaiana harti*, *Poblicia fuliginosa*, and *Otiocerus reaumurii*), suggesting that the compiled species list may yet be substantially incomplete for the combined states.

In addition to the planthopper fauna reported here, a preliminary inventory of the delphacids of Delaware suggests at least 54 species in the state, although additional taxa are likely to be found before the completion of that inventory.

## Taxonomy

**Cixiidae:** A number of specimens presented taxonomic difficulties. In the Cixiidae, specimens that appeared close to *Melanoliarus sablensis* differed from that depicted by Mead and Kramer (1982: 474) by having an additional ventral process on the aedeagus and a differing arrangement (size and orientation) of the other ventral processes. Similar specimens were observed in the Great Smoky Mountains National Park (Gonzon et al. 2007). In addition to the odd specimens, a specimen much more similar to that depicted by Mead and Kramer (1982) was found. While the possibility that these specimens represent an undescribed species should be investigated, we feel it is likely that they simply represent a variant of the more conventional form, and we have treated them as the same species with respect to biodiversity estimation calculations. Also, a group of females of *Melanoliarus* with uniformly dark wings were separated from others because they appear to represent a species not found among the males; they were excluded from the species counts.

Emeljanov (2001) moved several Nearctic Pentastirini from *Melanoliarus* to *Pentastiridius* and *Reptalus*. *Pentastiridius* can be separated from the other two genera by having 12 teeth at the apex of the basitarsus, versus 10 or fewer in *Melanoliarus* and *Reptalus*; however, the features of *Melanoliarus* have not been investigated relative to *Reptalus* and diagnostic features separating these genera have not been defined. It is probable that *Melanoliarus* as currently defined is not monophyletic.

**Achilidae:** Species of *Cixidia* were identified primarily using features described by Beirne (1950), whose key emphasized color, particularly that of the face. He admitted that there was “some variation” (Beirne 1950: 186) within taxa, and key color features were often relativistic, making species difficult to distinguish without access to authoritatively identified specimens, particularly in the context of this study *C. fusca*, *C. pallida*, *C. variegata*, and *C. septentrionalis*. Unfortunately, Beirne (1950) did not describe sufficient structural features to assist in doubtful cases. A revision of *Cixidia* would be desirable to address ambiguities, and to describe potential new species from the southwestern US.

**Dictyopharidae:** The only member of *Phylloscelis* collected by the authors (or the senior author’s students) was *Phylloscelis rubra* in New Jersey on cranberry (*Vaccinium macrocarpon* Aiton). This genus is a good example of a taxon that is likely to be in

Delaware, but has not yet been found. While there are only 4 species in the genus, and 3 in the study area (Figure 14), the species are best confirmed by genitalic features as presented in McPherson and Wilson (1995).

**Derbidae:** A number of taxonomic issues were found among the Derbidae, including problems separating species in the genera *Omolicna*, *Cedusa* and two genera of Otiocerinae (*Anotia* and *Otiocerus*). Specimens of *Omolicna* (Derbidae) could not be definitively identified to species despite there being only 4 described North American species, and only 3 of these eastern - *O. fulva* (Van Duzee, 1909), *O. mcateei* (Dozier, 1928), and *O. uhleri* (Ball, 1902). While literature records suggest that *O. uhleri* (Ball, 1902) should be the only northern species, it was evident from the genitalia of Delaware specimens that at least 2 species are present. Because the original descriptions are incomplete, and at times conflicting with subsequent authors, we were unable to determine which of the specimens were *O. uhleri*, and whether the remainder were *O. mcateei*, *O. fulva* or undescribed.

The derbid genus *Cedusa* is diverse and its members require examination of male genitalia for identification, and even then considerable study is required. Two species within this genus were found to differ from the descriptions provided by Flynn and Kramer (1983). *Cedusa kedusa* bears a large bifid process on both the left and right sides of the aedeagus. For the horizontal ramus of the bifid process on the left side, Flynn and Kramer (1983: 235) state that the apex may be "...occasionally trifurcate and dentate antepically with the number of teeth varying from none to four...". Most of the observed specimens in this study had 4–6 teeth, but otherwise agreed with the description of this species. For *Cedusa cedusa*, a feature in the key (couplet 72) states that this species has the "paramere with inner ventral margin truncately incised in basal portion" (Flynn and Kramer 1983: 135); but for most of our specimens, this feature was rounded or acute. Variations (in this feature and/or details of the processes of the aedeagus) contrast to Flynn and Kramer's (1983: 228) comment that "all specimens [of *C. cedusa*] seen are similar to the illustration", and have led us to consider our specimens as 'near *cedusa*' until further evaluation of the variation in this species can be made.

Species in the Otiocerinae tended to be problematic, particularly since most taxa are rare in collections. It is also a problem that otiocerines have been described primarily based on superficial color features whose diagnostic value has not been verified by reference to genitalic features. While attempting to verify our species concepts, we solicited photographs or examined type specimens of select otiocerines. We found that many of the Fitch types (deposited at the USNM) are in poor shape and greatly faded. It is likely that some of the Kirby collection had been lost (see Horn and Kahle, 1935), and 6 of 8 otiocerine Kirby types could not be located at this time (specifically *Otiocerus schellenbergii*, *O. reaumurii*, *O. degeerii*, *O. abbotii*, *O. coquebertii* and *Anotia bonnetii* [but see below]). It is clear that both *Anotia* and *Otiocerus* are in need of revision. The revision should reference genitalic features to verify species identities, provide a critical reexamination of geographic records, and (as needed) designate neotypes for the apparently missing Kirby types, although Kirby (1821) generally provided adequate descriptions. Also, based on Kirby's (1821) description, it is possible that the

*balli* of McAtee (1923) is the same as Kirby's *schellenbergii*. While McAtee (1923) and Metcalf (1923) may have misapprehended these species, we have retained their view of these taxa until definitive evidence (esp. Kirby's *schellenbergii* type) can be found.

Ten species of *Anotia* are reported from the United States (including species formerly in *Amalopota* Van Duzee, 1889, subsumed under *Anotia* by Fennah, 1951: 152). Of the 10 species, *A. caliginosa* Ball, 1937, and *A. lineata* Ball, 1937, are southwestern species (recorded from Arizona) and *A. mcateei* (Dozier, 1928), reported from Illinois and Mississippi, does not occur in the study area. Of the remainder, 5 (*A. burnetii*, *A. bonnetii*, *A. kirkaldyi*, *A. robertsoni*, and *A. westwoodi*) are similar in appearance in having white wings whose veins are variably bordered with dark. It is not clear how much intraspecific variation would be expected in features of wing color or pattern, and such patterns were difficult to interpret in the greatly faded Fitch type specimens (we examined types of *Anotia robertsonii* and *A. burnetii*). *Anotia kirkaldyi* and *A. westwoodi* share with *A. bonnetii* the presence of dark spots in the apical cells of the forewing, although they may be more prominent in the latter species. *Anotia kirkaldyi* and *A. westwoodi* can be separated with difficulty based on the presence of darkened wing veins in the former species, but these taxa are otherwise very similar and may not be distinct. *Anotia robertsonii* is similar to *A. burnetii* in possessing less extensive wing markings than *A. kirkaldyi*, *A. westwoodi*, and *A. bonnetii*; and in possessing dark markings on the dorsum of the abdomen, although in *A. burnetii* the markings are confined to the middorsum of segments 1–3 and in *A. robertsonii* the entire dorsum of subsequent terga (5–7 or 8).

The type specimen of *Anotia bonnetii* (the type species of the genus) was also sought, along with types of other otiocerines described by Kirby (1821). Kirby (1821) specified that he had a single *A. bonnetii* specimen, which he described and illustrated. The specimen photographed as the type of *A. bonnetii* (at OUMNH) is pinned and spread, missing the abdomen, both wings on the left side, and the head anterior to the eyes; but it was clear that the specimen was not the one used to describe *A. bonnetii*. We feel the type has been mislabelled, and this specimen is actually the type of *Otiocerus francilloni*. The specimen could readily have been mislabeled when the Oxford Museum type collection was evacuated to the cellar underneath the Ashmolean Museum during World War II. Kirby (1821: 17), reports black spots and bands (“elytris nigro punctatis et fasciatis”) for *O. francilloni*, with the black band interrupted, which is consistent with this specimen.

Nine species of *Otiocerus* are reported from the north of Mexico; two species, *O. abbotii* Kirby, 1821, and *O. kirbyii* Fitch, 1851; are not reported from the study area (but see below). We examined the types of *O. signoretii* and *O. stollii* to help confirm features attributed to these species. The type specimen of *O. signoretii*, at the USNM, is in rather poor condition, faded, and partially enmeshed in mycelium, but shows the pattern of spots described by Fitch (1856: 394) that was used in subsequent keys to the genus (“...four dots... placed at the angles of an imaginary square...”). Fitch (1856) also reports “...a broad dusky cloud-like stripe from the base to the middle of the inner margin, and extending thence obliquely across to the outer margin at its tip, and send-

ing a very broad branch to the tip of the inner margin...". In the type specimen, these markings are very faint. The type specimen of *Otiocerus stollii* Kirby (at OUMNH) consists of only of one front and one hind wing (evidently of the right side), but the forewing was consistent with our understanding of that species.

McAtee (1923: 47) noted within his key that *O. reaumurii*, *O. wolfii*, and *O. signoretii* "may be one species". While we are confident that *O. wolfii* is distinct from the other taxa, *O. reaumurii* and *O. signoretii* are very similar. Because the type specimen of *O. signoretii* is greatly faded, we attempted to diagnose this species from *O. reaumurii* by the distribution of dark spots on the wing, in particular the presence of a spot in the costal cell of *O. signoretii*. From the available material, these species appear to differ externally mainly in the spot organization. McAtee (1923: 46–47) noted that between the two species, the vitta of *O. reaumurii* was broader and 'percurrent', and the vitta of *O. signoretii* was 'forked at apex of clavus', but we have been unable to verify these features. These species are both similar to *O. stollii* except for more extensive dark markings of *O. stollii*. Interestingly, all observed specimens of *O. reaumurii* and *O. signoretii* were female, and all observed *O. stollii* were male, possibly suggesting that all these species are part of a single sexually dimorphic species. However, we did not observe a sufficient number of specimens to exclude the possibility that this sex ratio was obtained by chance alone. Also, Fitch reported the type of *O. signoretii* to be a male, but the condition of the type specimen makes this difficult to confirm.

A single specimen of *Otiocerus* from Maryland was not clearly associated with any of the described species. The specimen is uniformly pale, head without markings, forewings without spots and with a very faint band. A similar specimen was found among undetermined Derbidae at the USNM. It is possible that this specimen is *Otiocerus kirbyii*, but we were unable to confirm this identification.

**Flatidae:** The genus *Cyarda* is under revision by S. Wilson (S. Wilson, pers. comm.). Species in this genus are largely Caribbean. Four *Cyarda* have been reported from the United States: *Cyarda acuminipennis* (Spinola, 1839), *C. melichari* Van Duzee, 1907, *C. sordida* Fennah, 1965 (= *C. sp. nr. acutissima* Metcalf & Bruner, 1948; see Fennah, 1965: 115) and *C. walkeri* Metcalf, 1923. However, Fennah (1965: 112) noted that for *C. walkeri* it "...must be assumed that this species occurs only in Jamaica". Metcalf (1923) reported *C. acuminipennis* from the eastern US, and later from Florida by Metcalf and Bruner (1948), but occurrence of this species in the US has not been subsequently substantiated (e.g., by Fennah 1965). Of the remaining species, *C. sordida* is reported only from Florida (Fennah 1965) and *C. melichari* is widely reported in the eastern United States (including the District of Columbia), but its genitalic features have not been compared to the other US species, so it cannot be assumed that *Cyarda* found outside of Florida (including the D.C. record) are *C. melichari* as has apparently been previously assumed. The image used here (Figure 3E) is from an undetermined specimen from Ft. Lauderdale, FL.

**Caliscelidae:** Specimens reported as *Bruchomorpha* sp. n. were collected at Phillips Landing, Sussex Co., DE (on 3 dates) as well as single specimens from Medford, NJ and Baltimore, MD. Superficially, these specimens are similar to *Bruchomorpha dor-*

*sata*, which has been reported in the Mid-Atlantic region by Dozier (1928), Doering (1939), and Wilson and McPherson (1980a); however, the snout is longer than that described by Doering (1939) for *B. dorsata*, and females are larger than the reported size range for this species. The specimens are also superficially similar to *Bruchomorpha beameri* Doering, 1939; a Midwestern species, but the dimensions of the snout and coloration of the legs do not match. Unlike both *B. beameri* and *B. dorsata*, the aedeagus bears no dorsally directed process, and the ventral process is strongly retrose curved. We conclude this taxon to be an undescribed species, which will be described after further review of *Bruchomorpha* species,

### Seasonality

Seasonality data were compiled from available Delaware specimens as a way to begin to understand the life history of local planthopper taxa. From the available seasonality information, it appears that all non-delphacid planthoppers have a single generation a year in Delaware, with the possible exceptions of *Bruchomorpha oculata*, *Aphelonema simplex*, and *Cixius nervosus*. This would be in general agreement with Nickel and Remane (2002) who report that all non-delphacid planthoppers in Germany have a single generation a year. From these data it is evident that *Apache degeerii* overwinters as an adult, and based on April records that at least *Bothriocera cognita* and *Melanoliarius placitus* may overwinter as nymphs. Of the remaining species little can be determined concerning overwintering stage. Nickel (2003) reports that 18.6% of Fulgoromorpha (including Delphacidae) in Germany overwinter as eggs, 61.4% as nymphs, and 12.4% as adults, with the remainder unclear. Published literature reports that *Flatormenis chloris*, *Metcalfa pruinosa*, *Ormenoides venusta*, *Acanalonia bivittata*, *A. conica*, *Thionia elliptica* and *Phylloscelis pallescens* overwinter as eggs (Wilson and McPherson 1981a, b; Wilson and Wheeler 1987, McPherson and Wilson 1996). Nickel and Remane (2002) report for the German fauna that all cixiids and achilids overwinter as nymphs.

A large number of *Melanoliarius placitus* were collected in early July of 2002 by the senior author and several students. The series was collected at mercury vapor lights (many specimens landed on trees near the lights instead of at the lights). Interestingly, this time period fell between the last quarter (July 2, 2002) and the New Moon (July 10 2002), which is similar to observations made by Bartlett and colleagues (2008) concerning Membracidae, where large numbers were collected at lights at times near a new moon.

### Conclusion

While the planthoppers of the eastern United States may be characterized as relatively well known from a taxonomic perspective, their faunistics and ecology remain

poorly understood. Although Delaware is near the two largest insect collections in the US (the USNM and the American Museum of Natural History, both of which employ hemipterists), it is a testament to our inchoate understanding of US planthopper faunistics that this study has increased our known Delaware fauna by over 700%. The diversity of planthopper species in Delaware is expected to be relatively modest relative to other states because it is small and physiographically rather uniform, and because planthopper diversity tends to generally increase inversely with latitude (and within North America, is greatest overall in the southwest). Here we also report totals of 88 species for Maryland, 74 for New Jersey, 60 for Pennsylvania, and 46 for the District of Columbia based on a compilation of literature records and available specimens. The only other state with a modern, relatively complete, survey of its planthopper fauna is Illinois (Wilson and McPherson 1980b), which reported 150 species, of which 66 were delphacids. In comparison, the total North American planthopper fauna appears to be 12 families, 165 genera and 935 species, of which 61 genera and 338 species are delphacids, and approximately 2/3 of all US planthopper species are western (unpublished data from species checklist compiled by S. W. Wilson, L. B. O'Brien, and C. R. Bartlett). Clearly our understanding of the faunistics of US planthoppers is limited, and our appreciation of planthopper ecology remains in its infancy. Further regional investigations would be helpful in gaining a more complete understanding of the US planthopper fauna.

## **Acknowledgements**

We are indebted to Kimberley Shropshire (University of Delaware) for the line art and most of the photography used in this work. We are grateful to Lois O'Brien for advice, support, specimens, and many helpful comments on early versions of this text; Stuart McKamey (USNM) for the loan of specimens, photographs of types, and assistance at the USNM collection; Zoë Simmons (Hope Entomological Collections, Oxford University Museum of Natural History) for locating and photographing some of the Kirby derbid types; Dmitri Logunov (Manchester Museum, University of Manchester, UK) and Mick Webb (British Natural History Museum, London) for seeking Kirby types in their collection. We are particularly grateful for past and present students whose diligent collecting has provided much of the basis for this work, especially Lawrence Barringer, Leo Donovall, Christopher Heckscher, Ashley Kennedy, Nate Nazdrowicz, Rob Snyder, and Katie Weglarz. We also thank an anonymous reviewer for helpful comments. This project was supported by the USDA Agriculture and Food Research Initiative Competitive Grants Program Grant No. 2009-55605-05006 from the National Institute of Food and Agriculture and Hatch Project W-2185 Biological Control in Pest Management Systems of Plants. Additional support was provided by the University of Delaware Department of Entomology and Wildlife Ecology.

## References

- Bartlett CR, Deitz LL (2000) *Revision of the New World delphacid planthopper genus Pissonotus (Hemiptera: Fulgoroidea)*. Thomas Say Publications in Entomology: Monographs, Lanham, Maryland, 234pp.
- Bartlett CR, Deitz LL, Rothschild MJ, Wallace MS (2008) Treehopper diversity (Hemiptera: Membracidae) of Little Orleans, Allegany Co., Maryland. *Proceedings of the Washington Entomological Society* 110: 130–143.
- Beirne BP (1950) The Canadian species of *Epiptera* (Homoptera: Achilidae). *Canadian Entomologist* 82: 186–190.
- Breakey EP (1928) The genus *Scolops* (Homoptera, Fulgoridae). *University of Kansas Science Bulletin* 18(6): 417–455.
- Chao A (1984) Non-parametric estimation of the number of classes in a population. *Scandinavian Journal of Statistics* 11: 265–270.
- Doering KC (1938) A contribution to the taxonomy of the subfamily Issinae in America north of Mexico (Fulgoroidea: Homoptera). Part II. *University of Kansas Science Bulletin* 25(20): 447–575.
- Doering KC (1939) A contribution to the taxonomy of the subfamily Issinae in America north of Mexico (Fulgoroidea: Homoptera). Part III. *University of Kansas Science Bulletin* 26(2): 83–167.
- Doering KC (1941) A contribution to the taxonomy of the subfamily Issinae in America north of Mexico (Fulgoroidea: Homoptera). Part IV. *University of Kansas Science Bulletin* 27(10): 185–233.
- Dozier HL (1928, dated 1926) The Fulgoridae or planthoppers of Mississippi, including those of possible occurrence. *Technical Bulletin of the Mississippi Agricultural Experiment Station* 14: 1–152.
- Emeljanov AF (1999) Notes on delimitation of families of the Issidae group with description of a new species of Caliscelidae belonging to a new genus and tribe (Homoptera, Fulgoroidea). *Zoosystematica Rossica* 8: 61–72.
- Emeljanov AF (2001) The generic position of some Nearctic Pentastirini (Homoptera: Fulgoroidea: Cixiidae). *Zoosystematica Rossica* 9: 122.
- Fennah RG (1952) On the generic classification of Derbidae (Homoptera: Fulgoroidea) with descriptions of new Neotropical species. *Transactions of the Royal Entomological Society of London* 103(4): 109–170.
- Fennah RG (1965) New Species of Fulgoroidea (Homoptera) from the West Indies. *Transactions of the Royal Entomological Society, London* 117(4): 95–126.
- Fitch A (1856) Third report on noxious and other insects of the State of New York. *Transactions of the New York State Agricultural Society* 16: 315–490.
- Flynn JE, Kramer JP (1983) Taxonomic study of the planthopper genus *Cedusa* in the Americas (Homoptera: Fulgoroidea: Derbidae). *Entomography* 2: 121–260.
- Freund RL, Wilson SW (1995) The planthopper genus *Acanalonia* in the United States (Homoptera: Issidae): male and female genitalic morphology. *Insecta Mundi* 9(3–4): 195–215.

- Gnezdilov VM (2004) Two new genera of the family Issidae (Homoptera: Cicadina: Fulgoroidea) from North America. *Russian Entomological Journal* 13 (1–2): 1–2.
- Gonzon AT, Bartlett CR, Bowman JL (2007) (Dated 2006) Planthopper (Hemiptera: Fulgoroidea) diversity in the Great Smoky Mountains National Park. *Transactions of the American Entomological Society* 132: 243–260.
- Holzinger WE, Kammerlander I, Emeljanov AF (2002) The family Cixiidae Spinola 1839 (Hemiptera: Fulgoromorpha) - a review. In: Holzinger W (Ed) *Zikaden: Leafhoppers, Planthoppers, and Cicadas (Insecta: Hemiptera: Auchenorrhyncha)*. *Denisia* 4: 113–138.
- Horn W, Kahle I (1935) *Über entomologische Sammlungen (Ein Beitrag zur Geschichte der Entomo-Museologie)*. *Entomologische Beihefte aus Berlin-Dahlem, Teil 2*: 1–160.
- Kirby W (1821) The characters of *Otiocerus* and *Anotia*, two new genera of Hemipterous insects belonging to the family of Cicadiadae [sic]: with a description of several species. *Transactions of the Linnaean Society of London Zoology* 13: 12–23.
- Kramer JP (1977) Taxonomic study of the planthopper genus *Oecleus* in the United States (Homoptera: Fulgoroidea: Cixiidae). *Transactions of the American Entomological Society* 103: 379–449.
- Kramer JP (1979) Taxonomic study of the planthopper genus *Myndus* in the Americas (Homoptera: Fulgoroidea: Cixiidae). *Transactions of the American Entomological Society* 105: 301–389.
- Kramer JP (1983) Taxonomic study of the planthopper family Cixiidae in the United States (Homoptera: Fulgoroidea). *Transactions of the American Entomological Society* 109: 1–57.
- McAtee WL (1923) A new species of *Otiocerus* (Homoptera; Fulgoridae). *Proceedings of the Biological Society of Washington* 36: 45–48.
- McPherson KR, Wilson SW (1995) The planthopper genus *Phylloscelis* in the United States (Homoptera: Dictyopharidae). *Insecta Mundi* 9: 177–188.
- McPherson KR, Wilson SW (1996) Life history and descriptions of immatures of the dictyopharid planthopper *Phylloscelis pallelescens* (Homoptera: Fulgoroidea). *Journal of the New York Entomological Society* 103: 170–179.
- Mead FW, Kramer JP (1982) Taxonomic study of the planthopper genus *Oliarus* in the United States (Homoptera: Fulgoroidea: Cixiidae). *Transactions of the American Entomological Society* 107: 381–569.
- Medler JT (2003) Types of Flatidae. XXIV. Type designations and taxonomic notes on species in the Natural History Museum of Geneva (Homoptera, Auchenorrhyncha, Fulgoroidea). *Revue Suisse de Zoologie* 110: 591–597.
- Metcalf ZP (1923) A key to the Fulgoridae of Eastern North America with descriptions of new species. *Journal of the Elisha Mitchell Scientific Society* 38: 139–230, plate(s): 38–70.
- Metcalf ZP, Bruner SC (1948) Cuban Flatidae with new species from adjacent regions. *Annals of the Entomological Society of America* 41: 63–118.
- Nickel H (2003) *The leafhoppers and planthoppers of Germany (Hemiptera, Auchenorrhyncha): patterns and strategies in a highly diverse group of phytophagous insects*. Pensoft Publishers, Sofia-Moscow, 460 pp.

- Nickel H, Remane R (2002) Check list of the planthoppers and leafhoppers of Germany, with notes on food plants, diet width, life cycles, geographic range and conservation status (Hemiptera, Fulgoromorpha and Cicadomorpha). *Artenliste der Zikaden Deutschlands* 5: 27–64 [English reprint of *Beiträge zur Zikadenkunde* 5: 27–64],
- O'Brien LB (1971) Systematics of the tribe Plectoderini (Insecta, Fulgoroidea, Achilidae) in America North of Mexico. *University of California Publications in Entomology* 64: 1–79.
- O'Brien LB (1985) New synonymies and combinations in New World Fulgoroidea (Achilidae, Delphacidae, Flatidae, Fulgoridae: Homoptera). *Annals of the Entomological Society of America* 78: 657–662.
- Smith JB (1890) Sub-order Homoptera. In: *Catalogue of insects found in New Jersey. Geological Survey of New Jersey. Final Report of the State Geologist, volume 2.* John L. Murphy, Trenton, New Jersey, 436–447.
- Wilson SW (2005) Keys to the Families of Fulgoromorpha with Emphasis on Planthoppers of Potential Economic Importance in the Southeastern United States (Hemiptera: Auchenorrhyncha). *Florida Entomologist* 88: 464–481.
- Wilson SW, Wheeler AG Jr (1987) Life history of *Thionia elliptica* (Homoptera: Fulgoroidea) with description of a new *Thionia* species from Texas. *Journal of the New York Entomological Society* 95: 440–451.
- Wilson SW, McPherson JE (1980a) The Distribution of the Fulgoroidea of the Eastern United States (Homoptera). *Transactions of the Illinois State Academy of Science* 73(4): 7–20.
- Wilson SW, McPherson JE (1980b) Keys to the planthoppers, or Fulgoroidea of Illinois (Homoptera). *Transactions of the Illinois State Academy of Science* 73(2): 1–61.
- Wilson SW, McPherson JE (1981a) Life histories of *Acanalonia bivittata* and *A. conica* with descriptions of immature stages. *Annals of the Entomological Society of America* 74: 289–298.
- Wilson SW, McPherson JE (1981b) Life histories of *Anormenis septentrionalis*, *Metcalfa pruniosa*, and *Ormenooides venusta* with descriptions of immature stages. *Annals of the Entomological Society of America* 74: 299–311.
- Zuefle ME (2006) The impact of non-native woody plants on the native herbivorous insect community of northern Delaware. M.S. Thesis, University of Delaware, Department of Entomology and Applied Ecology, 66 pp.
- Zuefle ME, Brown WP, Tallamy DW (2008) Effects of non-native plants on the native insect community of Delaware. *Biological Invasions* 10: 1159–1169.