

# *Acanthodasys caribbeanensis* sp. n., a new species of Thaumastodermatidae (Gastrotricha, Macrodasysida) from Belize and Panama

Rick Hochberg<sup>†</sup>, Sarah Atherton<sup>‡</sup>

University of Massachusetts Lowell, One University Avenue, Lowell, MA 01854, 01.978.934.2885

<sup>†</sup> [urn:lsid:zoobank.org:author:8C5BB2F6-35A3-4A4D-9F67-C2A21A5DADE1](https://zoobank.org/urn:lsid:zoobank.org:author:8C5BB2F6-35A3-4A4D-9F67-C2A21A5DADE1)

<sup>‡</sup> [urn:lsid:zoobank.org:author:1F597997-CD78-4F36-A82B-977B14DCAA6C](https://zoobank.org/urn:lsid:zoobank.org:author:1F597997-CD78-4F36-A82B-977B14DCAA6C)

Corresponding author: Rick Hochberg ([rick\\_hochberg@uml.edu](mailto:rick_hochberg@uml.edu))

---

Academic editor: Antonio Todaro | Received 21 July 2010 | Accepted 30 August 2010 | Published 13 October 2010

[urn:lsid:zoobank.org:pub:E249DC0B-8B28-4484-97F3-6E7A8F354B3C](https://zoobank.org/urn:lsid:zoobank.org:pub:E249DC0B-8B28-4484-97F3-6E7A8F354B3C)

---

**Citation:** Hochberg R, Atherton S (2010) *Acanthodasys caribbeanensis* sp. n., a new species of Thaumastodermatidae (Gastrotricha, Macrodasysida) from Belize and Panama. ZooKeys 61: 1–10. doi: 10.3897/zookeys.61.552

---

## Abstract

We describe one new species of *Acanthodasys* (Gastrotricha, Macrodasysida, Thaumastodermatidae) collected from sublittoral sites around Carrie Bow Cay, Belize and Isla Colón in the Bocas del Toro archipelago, Panama. Though eight species of *Acanthodasys* are currently recognized, no species has yet been reported from the Caribbean. *Acanthodasys caribbeanensis* sp. n. is characterized by the lack of lateral adhesive tubes, the presence of ventrolateral adhesive tubes, and with cuticular armature in the form of both spineless and spined scales. The spineless scales are not elliptical as in other species of *Acanthodasys*, but are instead variable in shape and closely resemble the spineless scales of species of *Diplodasys*. Spined scales bear uniancres up to 50 µm long and are the largest reported in the genus. Uniancres are arranged dorsally around the mouth rim and distributed in five distinguishable columns. Adult size varies from 325–625 µm long.

## Keywords

Meiofauna, Caribbean, Panama, Belize, gastrotrich, taxonomy

## Introduction

The phylum Gastrotricha contains some of the smallest (0.1 mm–3 mm) benthic marine invertebrates. In general, marine gastrotrichs are restricted to the interstitial pore spaces between sand grains, making them overlooked in most tropical biodiversity investigations. Knowledge of gastrotrich biodiversity in the Tropical Northwestern Atlantic (aka wider Caribbean) is almost entirely limited to species lists or descriptions of novel taxa from only a handful of sites in the five Caribbean ecoregions: the Gulf of Mexico (Todaro 1994; Todaro et al. 1995), South Florida ecoregion (Thane-Fenchel 1970; Schoepfer-Sterrer 1974; Decho et al. 1985; Evans and Hummon 1991; Evans 1994), Bahamian ecoregion (Renaud-Debyser 1963), Lesser Antilles ecoregion (Kisielewski 1984; Hummon 2010) and Central Caribbean ecoregion including Colombia (Hummon 1974) and Panama (Hochberg 2008). Additional unpublished records of marine gastrotrich diversity and biogeography can be found in Hummon (2009).

Here, we document a new species of *Acanthodasya* (Macrodasyida, Thaumastodermatidae) from two Central Caribbean islands, Carrie Bow Cay on the Belizean barrier reef complex and Isla Colón in the Bocas del Toro archipelago, Panama. Investigations on both islands were conducted as part of a Smithsonian sponsored biodiversity project to barcode the local marine meiofauna.

## Methods

Gastrotrichs were collected from a variety of sublittoral sites around Carrie Bow Cay, Belize (Jan. 13–28, 2010) and Bocas del Toro, Panama (June 6–19, 2010). Research was performed at the Smithsonian field station on Carrie Bow Cay, Belize and Smithsonian Tropical Research Institute, Bocas del Toro, Panama, respectively. Sediments were collected by SCUBA in bags and buckets and brought back to the field stations for subsampling. Extraction of gastrotrichs was performed with the following protocol: 1) approximately 100 cm<sup>3</sup> of sediment was combined with ca. 900 cm<sup>3</sup> of 7.5% MgCl<sub>2</sub> in a 1 L Erlenmeyer flask and allowed to rest for 15 mins; 2) the flask was gently shaken and the supernatant was decanted over a 48 µm mesh; and 3) the mesh was gently washed with seawater into a petri dish. Specimens were sorted with a Leica EZ4 stereomicroscope, transferred to a glass slide, and viewed with a Zeiss A1 compound microscope equipped with DIC (differential interference contrast). Light micrographs were captured with a Sony Handycam digital camera. Measurements of individual specimens were performed with an ocular micrometer. Lengths and positions of organ systems are described in terms of percentage body units, where total body length from anterior (U00) to posterior (U100) is 100 units.

Specimens were prepared for museum archival using the following protocol: fix in 2.5% glutaraldehyde in 0.1M phosphate buffer saline (PBS; pH 7.4) for at least one week; rinse with PBS for 1 hr; expose to 1% OsO<sub>4</sub> in 0.1M PBS for 1 min (to increase contrast); rinse in PBS for 15 min; dehydrate through an ethanol series;

transferr to propylene oxide for 30 min; and embed in epon on a glass microscope slide (coverslipped and placed in an oven at 60°C for 24 hrs). Type specimens are deposited in the National Museum of Natural History, Smithsonian Institution, Washington, DC.

Abbreviations: PIJ, pharyngeointestinal junction; TbA, anterior adhesive tubes below mouth rim; TbL, lateral adhesive tubes; TbP, posterior adhesive tubes on caudal pedicles; TbVL, ventrolateral adhesive tubes.

## Results

**Order Macrodasysida Remane, 1925 [Rao & Clausen, 1970]**

**Family Thaumastodermatidae Remane, 1927**

**Subfamily Diplodasyinae Ruppert, 1978**

**Genus *Acanthodasys* Remane, 1927**

***Acanthodasys caribbeanensis* sp. n.**

urn:lsid:zoobank.org:act:2ED007EB-D7F8-4E0D-9DC4-37BAFA49B247

Figs 1–3

**Type locality.** Station CBC10.19, Carrie Bow Cay Reef, coarse sand patches on ridge, 15 m depth, 16°48.127N, 88°04.607W. Sediments collected by Cheryl Thacker on January 23, 2010.


**Holotype.** Adult specimen, reproductively mature, 625 µm long, resin preparation: specimen in dorsoventral orientation. USNM #1145897

**Other localities.** Station BRS2010-104, Wild Cane Rock, Bocas del Toro, Panama, 14 m depth, medium coarse sand plain, 9°21.016N, 82°10.335W. Sediment collected by Daniel Gouge on June 8, 2010.


**Paratypes.** Cat. No. USNM 1145898, Adult specimen, minimum 425 µm long (curled), resin preparation: specimen on side. Cat. No. USNM 1145899, Adult specimen, minimum 450 µm long (curled), resin preparation: specimen in dorsoventral orientation.

**Material examined.** Ten specimens. Three prepared for archival.

**Diagnosis.** *Acanthodasys* with an adult body length from 375 µm to 625 µm long. Maximum body width at mouth/PIJ/midpoint of body is 30/62/100 µm. Pharynx to 238 µm long. Cuticle of spineless and spined scales. Spineless scales of various shape. Spined scales bear uniancres to 50 µm long, distributed as one dorsal column flanked by two dorsolateral columns and two lateral columns. Uniancres also extend dorsally across mouth rim. Ten epidermal glands per side. Up to nine TbA per side inserting directly on body surface at mouth rim. TbL absent. At least 43 TbVL per side beginning at PIJ and extending onto the outer edges of the caudal lobes. Two TbP insert terminally on caudal lobes, two insert medially. Hermaphroditic, with paired testes and single glandular caudal organ. Rosette organ on left dorsolateral side of body; single ovary present.




**Figure 1.** *Acanthodasya caribbeanensis* sp. n. from Carrie Bow Cay, Belize **A** Dorsal view, transmitted light **B** Closeup of anterior end, dorsal view, DIC optics **C** Closeup of posterior end, dorsal view, DIC optics. Abbreviations: cu caudal uniancre, du dorsal uniancre, dlu dorsolateral uniancre, eg egg, lu lateral uniancre, ou oral uniancre, TbP posterior adhesive tubes.



**Figure 2.** *Acanthodasys caribbeanensis* sp. n. **A** Habitus, drawn in dorsal view. **B** Habitus, drawn in ventral view. **C** Closeup of head region showing the distribution of TbA **D** Examples of different shapes of uniancres on the body **E** Examples of different scale shapes on dorsal and ventral sides. Abbreviations: co caudal organ, cu caudal uniancres, dlu dorsolateral uniancre, du dorsal uniancre, ep epidermal gland, lu lateral uniancre, ou oral uniancre, ro rosette organ, TbA anterior adhesive tube, TbP posterior adhesive tubes, TbVI ventrolateral adhesive tubes, te testis.

**Etymology.** Named after the region of origin, referring to the Caribbean (Latin adjectival ending: *ensis*; *caribbeanensis*).

**Description.** The description is based on the holotype (adult, 625 µm long), with ranges given from specimens measured in vivo. Body strap-shaped and 425–625 µm long (Fig. 1A). Mouth region narrow; body inflates at U04, narrows again at U08, and widens toward the trunk. Widths of mouth at U04/pharynx region at U08/PIJ at U36/trunk at U50/caudal base at U92 are 30/67/105/125/20 µm. Pharynx 238



**Figure 3.** *Acanthodasys caribbeanensis* sp. n. from Carrie Bow Cay, Belize (A, B) and Bocas del Toro, Panama (C) **A** Closeup of dorsal cuticle showing dorsal uniancres, dorsolateral uniancres and spineless dorsal scales **B** Closeup of dorsal uniancres **C** Paratype (Cat. No. USNM 1145899) of Panamanian specimen showing the rosette organ (gland). Abbreviations: du dorsal uniancres, ep epidermal gland, lu lateral uniancre, ro rosette organ, TbVL ventrolateral adhesive tubes.

µm long with pharyngeal pores at U32. Dorsal rim of oral hood, ca. 10 µm wide, free of cuticular spines and fringed with numerous mobile cilia and few stiff sensory hairs to 10 µm long. Epidermal glands to 18 µm diameter and to 10 per side. Ventral locomotory cilia present as a complete field beginning at ca. U08 and extending to the caudum (Fig. 2B).

**Cuticular armature.** Cuticular spines evident at low magnification and arranged around the periphery of the head, along the trunk and on the caudal lobes (Figs 1, 2). Spined- and spineless scales present. Approximately 10–12 oral uniancres line the dorsal periphery of the head behind the “naked” region of the oral hood (Fig. 1B). Uniancres slightly staggered in position; uniancres increase in size from medial position (10  $\mu\text{m}$  long) to a lateral position along head (U06, 23  $\mu\text{m}$  long). Uniancres in lateral and dorsolateral positions continue along the trunk as individual columns. The trunk contains a total of five columns of uniancres: one dorsal, two dorsolateral, and two lateral columns (Figs 1A, 2A). Sixteen uniancres in dorsal column begin at U08 and extend to U80, increasing in size from 20–50  $\mu\text{m}$  long down the length of the body, and with a slight bend or posterior curvature at the apex (Fig. 3B). Dorsolateral columns of ca. 14–16 spines that increase in length along the trunk, from 17–35  $\mu\text{m}$  long and with a slight posterior curvature or distinguishable bend (Fig. 2D); column ends at U85. Lateral columns of 23–26 spines that increase in length along the trunk, from 20–35  $\mu\text{m}$ , and mostly with a straight shape but oriented in a slight posterior direction; column ends at U87. Five small uniancres present on each caudal lobe, 8–20  $\mu\text{m}$  long, in somewhat dorsolateral position (Fig. 1C). Cuticle between uniancres present as spineless scales. Scales of various shape; most dorsal scales have an obtuse triangular shape with the apex pointed posteriorly; some ventral scales same shape as dorsal scales; others of varying shapes (Figs 1B, 2E, 3A). Center of each scale with a triangular or eye-shaped depression. Scales ca. 15  $\mu\text{m}$  wide and arranged in 5 columns on dorsal surface; ventral surface also with scales but the number of columns was undetermined.

**Adhesive tubes.** Anterior adhesive tubes (TbA), 8 per side to 12  $\mu\text{m}$  long, distributed along the mouth margin in a single row (Fig. 2B,C). TbL are absent. TbVL arranged in bilateral columns, to 23  $\mu\text{m}$  long, begin at PIJ and extend posteriorly on to the lateral edge of the caudal lobes; tubes on caudal lobe edges each ca. 15  $\mu\text{m}$  long (Fig. 2B). TbP distributed on caudal lobes as two terminal tubes to 16  $\mu\text{m}$  long and two medial tubes on the inner edge of the caudal lobes to 12  $\mu\text{m}$  long.

**Digestive tract.** Mouth terminal to 30  $\mu\text{m}$  wide. Pharynx to 238  $\mu\text{m}$  long with pharyngeal pores at base around U32. Intestine narrow and tapering toward posterior end. Anus at U88.

**Reproductive system.** Hermaphroditic, with paired testes at PIJ. Vasa deferentia extend posteriorly, but their point of termination was not observed. Glandular caudal organ, ca. 30  $\mu\text{m}$  long, at U90 (Fig. 2A). Rosette organ, ca. 27  $\mu\text{m}$  diameter, present at U42 on left, dorsolateral surface. Glands of the rosette organ stain intensely with  $\text{OsO}_4$  (Fig. 3C). Single egg present, ca. 57  $\mu\text{m}$  x 100  $\mu\text{m}$  (Fig. 1C).

## Taxonomic remarks

At present, there are eight valid species of *Acanthodasys* described globally: *Acanthodasys aculeatus* Remane, 1927; *A. algarvensis* Hummon, 2008 (see Hummon and Todaro 2010); *A. arcassonensis* Kisiulewski, 1987; *A. carolinensis* Hummon, 2008; *A. fibro-*

*sus* Clausen, 2004; *A. flabellicaudus* Hummon & Todaro, 2009; *A. lineatus* Clausen, 2000; and *A. silvulus* Evans, 1992. Five species mentioned by Ruppert (1978) are not described and considered nomina nuda (Kisielewski 1987; Hummon 2008). Among described species, *A. aculeatus*, the type species, has the widest geographic distribution with conspecifics reported from the coasts of Europe (Kisielewski 1987) and the Mediterranean and Black Seas (reviewed in Todaro et al. 2003) to as far south as the coastlines of India (Naidu and Rao 2004), the Maldive Islands (Gerlack 1961), and the Atlantic coast of Florida, USA (Evans 1992). Hummon (2009) has reported *A. aculeatus* from a variety of other locales. However, to date, no species of *Acanthodasys* are reported from the Caribbean.

Among the eight described species, only *A. aculeatus* Remane, 1927 and *A. arcassonensis* Kisielewski, 1987 possess spineless scales distributed among their uniancres as found in *A. caribbeanensis* sp. n. However, the new species differs from both described species in several significant ways: the head profile differs between *A. caribbeanensis* sp. n. and *A. arcassonensis*; the uniancres are arranged in distinguishable columns in the new species (Figs 1, 2); the uniancres are larger in *A. caribbeanensis* sp. nov (up to 50  $\mu\text{m}$  long) compared to all other species (e.g. *A. arcassonensis* to 11  $\mu\text{m}$  long; *A. fiborosos* to 20  $\mu\text{m}$ ); the spineless scales are not elliptical in the new species but instead variously shaped; and the new species possesses adhesive tubes in a ventrolateral (TbVL) as opposed to a lateral (TbL) position. This latter characteristic is noteworthy because the arrangement of TbVL of *A. caribbeanensis* sp. n. is more similar to the arrangement observed in species of *Diplodasys*, the sister taxon of *Acanthodasys* within the Diplodasyinae (Ruppert 1978), than to other species of *Acanthodasys* (e.g., compare to *A. flabellicaudus* Hummon & Todaro, 2009 and *D. sanctimariae* Hummon & Todaro, 2009).

Interestingly, there are several aspects of the cuticle of *A. caribbeanensis* sp. n. that appear unique and worth further mentioning. First, the uniancres of the new species are the largest described spines (see Fig 1A; up to 50  $\mu\text{m}$  long on an adult specimen 625  $\mu\text{m}$  long) in the genus and the largest spines known from any species of Macrodasysida. Second, the spineless scales of *A. caribbeanensis* sp. n. are more similar in appearance to the spineless scales of species of *Diplodasys* than they are to the scales described for other species of *Acanthodasys*. For example, the scales of *D. minor* Remane, 1936 have a similar shape to those of *A. caribbeanensis* sp. n., and several species have scales with a wide depressed region in their center like the scales of *A. caribbeanensis* (e.g., see *D. ankei* ssp. *pacifica* Schmidt, 1974; comparison of species' scales in Kisielewski 1987). Note, however, that the scales of *A. caribbeanensis* sp. n. appear to lack any obvious ribbing or texture (Fig. 3A) that often characterizes the scales of species of *Diplodasys*.

Another peculiar feature of *Acanthodasys caribbeanensis* sp. n. is the relatively wide size distribution of reproductive adults. We found specimens ranging from 325  $\mu\text{m}$  long in Panama to 625  $\mu\text{m}$  long in Belize, and all specimens showed clear evidence of reproductive maturity i.e., paired testes and/or a solitary rosette organ were present (Fig. 3C). According to Ruppert (1978), the rosette organ is a glandular ring around the female gonopore, but whether it is present in specimens prior to the maturation

of female gametes is not mentioned. According to our surveys, specimens of *A. caribbeanensis* sp. n. may have a rosette organ but no obvious ova or testes. As evidence, we have prepared a paratype (Cat. No. USNM 1145898) of a smaller adult specimen, ca. 400  $\mu\text{m}$  long (slightly curled), with a clear rosette organ but without evidence of eggs or testes. Unfortunately, the rosette organ can be difficult to see in live specimens, but becomes apparent if animals are fixed briefly in  $\text{OsO}_4$  (see methods); the  $\text{OsO}_4$  greatly enhances the contrast of many tissues including the epidermal glands (see Fig. 3C).

## Acknowledgements

We thank Dr. M. Antonio Todaro and two anonymous reviewers for their helpful comments on this manuscript. The first author thanks Dr. Jon Norenburg (Smithsonian Institution) for inviting him to participate in the sampling conducted on Carrie Bow Cay, Belize. Both authors also thank Dr. Norenburg and Dr. Rachel Collins (Smithsonian Tropical Research Institute) for their invitation to participate in the Meiofauna Workshop at STRI. Thanks are also extended to Daniel Gouge, Cheryl Thacker and Katrine Woorsae for their assistance with SCUBA collection of samples. This material is based upon work supported by the National Science Foundation under Grant No. DEB 0918499. This research received partial support from a Marine Science Network award to Jon Norenburg and an award from the Encyclopedia of Life Biodiversity Synthesis Group to Rachel Collin and Jon Norenburg.

## References

- Clausen C (2000) Gastrotricha Macrodasysida from the Tromso region, northern Norway. *Sarsia* 85: 357–384.
- Clausen C (2004) A new species of *Acanthodasys* (Gastrotricha: Thaumastodermatidae) from the west coast of Norway. *Sarsia* 89: 137–141.
- Decho AW, Hummon WD, Fleeger JW (1985) Meiofauna – Sediment interactions around tropical seagrass sediments using factor analysis. *Journal of Marine Research* 43: 237–255.
- Evans WA (1992) Five new species of marine Gastrotricha from the Atlantic coast of Florida. *Bulletin of Marine Science* 51: 315–328.
- Evans WA (1994) Morphological variability in warm-temperate and subtropical populations of *Macrodasys* (Gastrotricha: Macrodasysida: Macrodasysidae) with description of seven new species. *Proceedings of the Biological Society of Washington* 107: 239–255.
- Evans WA, Hummon WD (1991) A new genus and species of Gastrotricha from the Atlantic coast of Florida, U.S.A. *Transactions of the American Microscopical Society* 110: 321–327.
- Gerlach SA (1961) Über Gastrotrichen aus dem Meeressand der Malediven (Indischer Ozean). *Zoologischer Anzeiger* 167: 471–475.
- Hochberg R (2008) Gastrotricha of Bocas del Toro, Panama: A preliminary report. *Meiofauna Marina* 16: 101–107.

- Hummon WD (1974) Intertidal marine Gastrotricha from Colombia. *Bulletin of Marine Science* 24: 396–408.
- Hummon WD (2008) Gastrotricha of the North Atlantic Ocean: 1. Twenty four new and two redescribed species of Macrodsyida. *Meiofauna Marina* 16: 117–174.
- Hummon WD (2009) Global database for marine Gastrotricha (Taxonomic, Geographic, Bibliographic and Video). [<http://132.235.243.28>] or [<http://hummon-nas.biosci.ohiou.edu>].
- Hummon WD (2010) Marine Gastrotricha of the Caribbean Sea: A review and new descriptions. *Bulletin of Marine Science* 86: 661–708.
- Hummon WD, Todaro MA (2007) A new species of Xenotrichulidae (Gastrotricha) from southern and southeastern USA. *Cahiers de Biologie Marine* 48: 297–302.
- Hummon WD, Todaro MA (2009) Italian marine Gastrotricha: VI. Seven new species of Macrodsyida. *Zootaxa* 2278: 47–68.
- Hummon WD, Todaro MA (2010) Analytic taxonomy and notes on marine, brackish-water and estuarine Gastrotricha. *Zootaxa* 2392: 1–32.
- Kisielewski J (1984) *Xenotrichula guadelupense* sp. n., a new marine gastrotrich from Guadeloupe. *Bulletin du Museum D'Histoire Naturelle, Paris* 6A(1): 37–40.
- Kisielewski J (1987) New records of marine Gastrotricha from the French coasts of Manche and Atlantic I. Macrodsyida, with descriptions of seven new species. *Bulletin du Museum D'Histoire Naturelle, Paris* 9: 837–877.
- Naidu KV, Rao GC (2004) *The Fauna of India and the Adjacent Countries—Gastrotricha*. Director, Zoological Survey of India, Kolkata, 169 pp.
- Remane A (1927) Neue Gastrotricha Macrodsyioidea. *Zoologische Jahrbuecher Abteilung fuer Systematik Oekologie und Geographie der Tiere* 54: 203–242.
- Remane A (1936) Gastrotricha. In: Bronns HG (Ed) *Klassen Ordnungen des Tierreichs, Band 4, Abteilung II, Buch I, Teil 2, Lieferungen 1–2*. Akademie Verlagsgesellschaft, Berlin.
- Renaud-Debyser J (1963) Recherches écologiques sur la faune interstitielle des sables: Bassin d'Arcachon et île de Bimini, Bahamas. *Vie Milieu Supplement* 15 :1–157.
- Ruppert EE (1978) The reproductive system of gastrotrichs. III. Genital organs of Thaumastodermatinae subfam. n. and Diplodasyinae subfam. n. with discussion of reproduction in Macrodsyida. *Zoologica Scripta* 7: 93–114.
- Schmidt P (1974) Interstitielle Fauna von Galapagos. IV. Gastrotricha. *Mikrofauna des Meeresbodens* 26: 1–76.
- Schoepfer-Sterrer C (1974) Five new species of *Urodasy*s and remarks on the terminology of the genital organs in Macrodsyids (Gastrotricha). *Cahiers de Biologie Marine* 15: 229–259.
- Thane-Fenchel A (1970) Interstitial gastrotrichs in some south Florida beaches. *Ophelia* 7: 113–137.
- Todaro MA (1994) *Chaetonotus triacanthus* and *Heteroxenotrichula texana*, two new chaetonotid gastrotrichs from the Gulf of Mexico. *Transactions of the American Microscopical Society* 113: 15–21.
- Todaro MA, Matinato L, Balsamo M, Tongiorgi P (2003) Faunistics and zoogeographical overview of the Mediterranean and Black Sea marine Gastrotricha. *Biogeographia* 24: 131–160.
- Todaro MA, Fleegeer JW, Hummon WD (1995) Marine gastrotrichs from the sand beaches of the northern Gulf of Mexico: Species list and distribution. *Hydrobiologia* 310: 107–117.