

# ***Apoxyrina hirtuosa* (Wiedemann, 1821) comb. n., lectotype designation, redescription and identification key to species of *Apoxyrina* Schiner, 1866 (Asilidae, Laphriinae)**

Edgar Alvim<sup>1,2</sup>, Rosaly Ale-Rocha<sup>1</sup>, Freddy Bravo<sup>3</sup>

**1** Instituto Nacional de Pesquisas da Amazônia (INPA), Coordenação de Pesquisas em Entomologia, Av. André Araújo, 2936, Petrópolis, CEP 69011-970, Manaus, Amazonas, Brasil **2** Fellowship CNPq **3** Departamento de Ciências Biológicas, Universidade Estadual de Feira de Santana, Departamento de Ciências Biológicas, Av. Transnordestina S/N, Bairro Novo Horizonte 44036-900, Feira de Santana, BA, Brasil

Corresponding author: Edgar Alvim (edgar.alvim@gmail.com)

---

Academic editor: Martin Hauser | Received 11 July 2011 | Accepted 1 August 2011 | Published 26 August 2011

---

**Citation:** Alvim E, Ale-Rocha R, Bravo F (2011) *Apoxyrina hirtuosa* (Wiedemann, 1821) comb. n., lectotype designation, redescription and identification key to species of *Apoxyrina* Schiner, 1866 (Asilidae, Laphriinae). ZooKeys 125: 51–57. doi: 10.3897/zookeys.125.1790

---

## **Abstract**

The type specimens of *Neodiognites hirtuosus* (Wiedemann, 1821), two males and one female, deposited at the Museum für Naturkunde der Humboldt-Universität (ZMHB), Berlin were examined. The specimens show the diagnostic characters of *Apoxyrina* Schiner, 1866: face strongly pronounced, swollen and curved hind tibia, and terminalia with epandrium large and hypandrium short and obtuse. A new combination is suggested, *Apoxyrina hirtuosa* (Wiedemann, 1821), and lectotype and paralectotypes are designated. The species is redescribed, the male terminalia is described and illustrated for the first time, and an identification key to *Apoxyrina* is presented.

## **Keywords**

Neotropical, taxonomy, Laphystiini

## **Introduction**

*Dasypogon hirtuosus* Wiedemann, 1821, a species classified in the subfamily Dasypogoninae (Papavero 2009), is known from Brazil, with no additional details of the collection locality, was described based on two males and one female without a holotype

designation (i.e., syntypes) (Wiedemann 1821). Carrera (1949) transferred *D. hirtuosus* to *Laustauroides* Carrera, 1949 (Dasypogoninae) based on specimens collected in Brazil, however without examination of the type material. Subsequently, Artigas and Papavero (1988) synonymized *Laustauroides* with *Neodiogmites* Carrera, 1949, another genus of Dasypogoninae, and therefore the current name of the species is *Neodiogmites hirtuosus*.

During a revision of the genus *Neodiogmites*, the three syntypes of *N. hirtuosus* were examined. It was apparent that this species is not a Dasypogoninae and consequently, is not a member of *Neodiogmites*. Here, we propose a new combination for this species, with a lectotype designation.

## Material and methods

The specimens studied are deposited in the Museum für Naturkunde der Humboldt-Universität (ZMHB), Berlin, Germany. The material includes three syntypes, two males and one female; and their labels read only “Brazil”. To observe the terminalia, it was detached from the abdomen and cleared in cold 10% KOH, followed by neutralization in acetic acid, dehydration in ethanol, and washing in distilled water. The dissected terminalia was placed in glycerin in a microvial pinned with the respective specimen. Morphological terms follow Cumming and Wood (2009).

## Taxonomy

### *Apoxyria hirtuosa* (Wiedemann, 1821) **comb. n.**

[http://species-id.net/wiki/Apoxyria\\_hirtuosa](http://species-id.net/wiki/Apoxyria_hirtuosa)


Figs 1–7

*Dasypogon hirtuosus* Wiedemann 1821: 227; Wiedemann 1828: 402 (redescription); Walker 1854: 443 (check list); Schiner 1866: 679 (check list); Williston 1891: 67 (catalogue); Kertész 1909: 128 (catalogue);

*Lastauroides hirtuosus*; Carrera 1949: 97, Fig. 27; 1958a: 146; Carrera and Vulcano 1961: 69 (prey); Carrera and Papavero 1962: 53 (check list); Hull 1962: 241, Figs 545, 1074, 1083 (check list); Martin and Papavero 1970: 29 (catalogue).

*Neodiogmites hirtuosus*; Artigas and Papavero 1988: 213 (key), 151, Fig. 157; Papavero 2009: 1 (catalogue); Geller-Grimm 2011 (online catalogue).

**Type-material examined.** Lectotype male (ZMHB), present designation, labeled: “Brazil\ ?, [without date] V. Olfers coll.” A red label written “Lectotype” was added. Specimen in reasonable condition, head slightly dusty, flagellum and middle leg lost, right wing mounted on permanent slides, abdomen dissected and placed in a micro-vial with glycerin, pinned together with the specimen. Paralectotypes: 1 male and 1 female, same locality as lectotype. Paralectotype male (ZMHB) in reasonable condition, head with a




**Figures 1–4.** *Apoxyria hirtuosa* (Wiedemann, 1821) comb. n. **1** General lateral view of lectotype, male **2** General lateral view of paralectotype female **3** Frontal view of head of lectotype **4** Wing of lectotype. Scale = 1mm.

little dust, lacking flagellum, mesonotum broken posteriorly, abdomen cracked between the second and third segment. Paralectotype female (ZMHB) in good condition, but the mesonotum is perforated posteriorly and the left flagellum is lost.

Lectotype male: Measurements: 12.5 mm (body length excluding antennae); 9.0 mm (wings).

**Diagnosis.** Face pronounced, covered by yellow pruinescence; dorsocentral setae of the same length as the scutum setae, however black; scutellum covered by short yellow setae, with several yellow long and slender apical scutellar setae; wings with  $r_1$  open.

**Redescription: Lectotype male.** Head (Fig. 3): face black, covered by yellow pruinescence that is denser on the sides, pronounced, not ending abruptly on upper part, occupying 2/3 of face; mystax black and yellow with some yellow setae between the antenna and facial swelling; frons black with sparse yellow pruinescence, yellow setae below and beside ocellar tubercle; vertex black; ocellar tubercle with several yellow setae; orbital setae yellow; postocular setae black with thin yellow setae between them; occiput black with yellow pruinescence, with black setae and yellow lower setae; proboscis black,



**Figures 5–7.** Terminalia of *Apoxyria hirtuosa* (Wiedemann, 1821) comb. n., lectotype male. **5** Dorsal **6** Ventral **7** Lateral. Scale = 1mm. Abbreviations: apc proc goncxl-apical process gonocoxal; cerc- cercus; epand- epandrium; epand arm- epandrial arm; 1° proc goncxl-first gonocoxal process; goncx-gonocoxito; gonst-gonostylus; hypd-hypandrium; proc cerc- process cercal; ph-phallus; 2° proc goncxl- second process gonocoxal.

apex obtuse, with short yellow setae ventrally; palpus black, longer than half length of proboscis, yellow setae basally and black on the remainder; antenna black, scape and pedicel almost the same length with black setae, scape with some basal yellow setae.

Thorax (Fig 1): black; second cervical sclerite black with sparse yellow pruinescence and black setae; anteprenotum with black setae; postpronotum covered by yellow pruinescence and yellow setae laterally, some black setae mixed; postpronotal lobe

with black setae anteriorly and yellow posteriorly; proepisternum and proepimeron with yellow pruinescence and yellow setae; scutum covered by short yellow setae; dorsocentral setae of the same length as the scutum setae, however black; two notopleural setae; three to four supra-alar setae and two postalar setae, all black; scutellum covered by short yellow setae, with several yellow long and slender apical scutellar setae; mesopleura covered by yellow pruinescence, except anteriorly on the anepisternum and katapisternum; anepisternum with yellow setae and some black setae posteriorly; katapisternum with yellow setae; katatergite with brownish setae.

Legs (Fig. 1): shiny black; coxae with yellowish pruinescence and yellow setae; femora covered by short yellow setae with some black setae on the dorsum; fore and hind tibiae covered by yellow setae, longer on ventral and posteroventral margin, and with dense short yellow setae on ventral margin which extend onto first tarsomere, these setae are also present on the hind tibiae, but are denser than on the other legs; fore tibia with anterodorsal, posterodorsal, and posteroventral rows of stout black setae; mid tibia with dorsal, posterodorsal, posteroventral, and ventral rows of stout black setae; hind tibia covered by yellow and black setae of different lengths, with anterodorsal, posterodorsal, and posteroventral rows of stout black setae and dense yellow setae on ventral and posteroventral margins. Tarsi covered by short yellow setae and stout black setae; claws black; pulvilli yellowish.

Wing (Fig. 4): membrane slightly infusate; veins brown; alula reduced, but a small lobe is still distinct;  $r_1$  open, apex of  $R_{2+3}$  arching sharply anteriorly in  $90^\circ$  angle,  $R_4$  strongly sinuate,  $R_4$  and  $R_5$  diverge from each other at the wing margin,  $r_5$  open,  $m_3$  closed and petiolate; cell cup closed; haltere yellow.

Abdomen (Fig 1): black, covered by short yellow setae, longer on the sides of anterior three tergites; posterior margin of tergites 6 and 7 reddish. Terminalia (Figs 5–7): terminalia reddish with black setae; epandrium long, with deep, rounded sulcus on apical fourth, forming long arms laterally; subepandrial sclerite with medial evagination forming laterally arms with rounded apex; cercus with two projections dorsally; gonocoxite with a claw-shaped projection on the apex and with two expansions, the second expansion with shell-shaped apex; gonostylus with rounded apex; hypandrium short and boomerang-shaped.

Female (Fig. 2): flagellum longer than scape and pedicel combined, with and a spine on the apex, black covered by brown pruinescence denser in the base; postpronotal lobe with black setae anteriorly; scutellum with some short black setae anteriorly; legs and abdomen with fewer setae than in male; terminalia pale brown.

**Distribution:** Brazil.

**Discussion.** After the original description of *Dasypogon hirtuosus* by Wiedemann (1821), the type specimens have not been examined until now. All the transfers to other genera were based on other specimens collected in Brazil (Carrera 1949; Artigas and Papavero 1988). The morphological study of the syntypes of this species revealed the lack of a spine on the prothoracic tibia, one of the most important characters to identify members of Dasypogoninae (Hull 1962; Papavero 1973; Dikow 2009). Moreover, the specimens studied have important characters that places them in the Laphriinae:

the male with only six abdominal tergites visible dorsally, as defined by Hull (1962), wings whit  $r_1$  open, apex of  $R_{2+3}$  arching sharply anteriorly in  $90^\circ$  angle and  $R_4$  strongly sinuate (Dikow 2009).

Among the genera of Laphriinae, the specimens show the diagnostic characters of *Apoxyria*: numerous long and slender apical scutellar setae, the face strongly pronounced, swollen hind femora, swollen and curved hind tibia, and terminalia with a large epandrium and short and obtuse hypandrium.

At the moment, the genus *Apoxyria* is known only from Brazil, now with three species: *Apoxyria apicata* Schiner, 1866, *Apoxyria americana* Carrera, 1955 and *Apoxyria hirtuosa*. There are insufficient data to determine their distributions in detail, because few specimens of this genus have been collected.

### Key to species of *Apoxyria* Schiner, 1866.

- 1  $R_{2+3}$  ends in  $R_1$  (not reaching C), cell  $r1$  close, petiolate; male with posterior margins of tergites 6–7 black (Brazil) ..... ***A. apicata* Schiner, 1866**
- $R_{2+3}$  ends in C, cell  $r1$  open (Fig. 4); male with posterior margins of tergites 6–7 either reddish or yellow ..... **2**
- 2 Mesonotum covered by short yellow setae (Fig 1); male with posterior margins of tergites 6–7 reddish; epandrial arms long and slender (Fig. 5) (Brazil) ..... ***A. hirtuosa* (Wiedemann, 1821) comb. n.**
- Mesonotum covered by short black setae; male with posterior margins of tergites 6–7 yellow; epandrial arms short and thick (Fig. 66 from Artigas et al. 1997) (Brazil: Goiás and Santa Catarina) .... ***A. americana* Carrera, 1955**

### Acknowledgments

Our thanks to Dr. Joachim Ziegler, curator of ZMHB, for the loan of specimens; to the Proequipamento CAPES-DCEN project of INPA, for the use of Automontage facilities. LDW and RAR are grateful to the Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq) for the support provided and the fellowship granted.

### References

- Artigas JN, Papavero N (1988) The American genera of Asilidae (Diptera): Keys for identification with an atlas of female spermathecae and other morphological details. II. Key to the genera of Dasypogoninae Macquart, with descriptions of new genera and species and new synonymies. *Gayana Zoologica* 52(3/4): 199–260.
- Artigas JN, Papavero N, Costa, CA (1997) The American genera of Asilidae (Diptera): Keys for identification with an atlas of female spermathecae and other morphological details.



- VIII. Subfamily Laphystiinae G.H. Hardy, with descriptions of five new genera and species and a catalogue of the Neotropical species. *Arquivos de Zoologia do Estado de São Paulo* 34(1): 1–55.
- Carrera M (1949) Contribuição ao conhecimento dos Asilidae neotropicais (Diptera). I. Sôbre as espécies brasileiras com esporão na tibia. *Arquivos de Zoologia do Estado de São Paulo* 7: 1–148.
- Carrera M (1958) Dipteros de Boracéia I. Asilidae. *Arquivos de Zoologia do Estado de São Paulo* 13(12): 141–154.
- Carrera M, Vulcano MA (1961) Relação de alguns Asilidae (Diptera) e suas presas. (IV). *Revista brasileira de Entomologia* 10: 67–80.
- Carrera M, Papavero N (1962) Saropogonini neotropicais (Diptera, Asilidae, Dasypogoninae). *Studia Entomologica* 5: 39–64.
- Cumming JM, Wood DM (2009) Adult morphology and terminology. In: Brown BV, Bor-kent A, Cumming JM, Wood DM, Woodley NE, Zumbado MA (Eds) *Manual of Central American Diptera*, Volume 1, NRC Research Press, Ottawa, Ontario, 950 pp.
- Dikow T (2009) Phylogeny of Asilidae inferred from morphological characters of imagines (Insecta: Diptera: Brachycera: Asiloidea). *Bulletin of the American Museum of Natural History* 319: 1–175. doi: 10.1206/603.1
- Kertész K (1909) *Catalogus dipterorum hucusque descriptorum* (IV). *Oncodidae, Nemestrinidae, Mydidae, Asilidae* 4: 348.
- Geller-Grimm F (2010) Catalog of species. Available from: <http://www.geller-grimm.de/catalog/species.htm> [accessed 25, March, 2011]
- Hull FM (1962) Robber Flies of the World: The Genera of the Family Asilidae. *Bulletin of the United States National Museum* 224(43): 1–906. doi: 10.5479/si.03629236.224
- Martin CH, Papavero N (1970) Family Asilidae. In: Papavero N (Ed) *A catalogue of the Diptera of the Americas south of the United States*. Museu de Zoologia, Universidade de São Paulo 35(b): 1–139.
- Papavero N (1973) Studies of Asilidae (Diptera) systematics and evolution. I. A preliminary classification in subfamilies. *Arquivos de Zoologia do Estado de São Paulo*, 23(3): 217–274.
- Papavero N (2009) Catalogue of Neotropical Diptera. Asilidae. *Neotropical Diptera* 17: 1–179.
- Schiner JR (1866) Die Wiedemannischen Asiliden, interpretiert und in die seither errichteten neuen Gattungen eingereiht. *Verhandlungen der zoologisch-botanischen Gesellschaft in Wien*, 16: 649–722.
- Walker F (1854) List of the specimens of dipterous insects in the collection of the British Museum (3): 507–775.
- Wiedemann CRW (1821) *Diptera exotica*. Part 1 Kiliae 244, 179–242.
- Williston SW (1891) Catalogue of the described species of South American Asilidae. *Transaction of the American Entomological Society* 18: 67–91.