

A new species of *Schrankia* Hübner, 1825 from China (Lepidoptera, Erebidae, Hypenodinae)

Oleg Pekarsky^{1,†}

1 H-1068 Budapest, Felsőerdősor u. 16-18, Hungary

† <urn:lsid:zoobank.org:pub:77E41AD1-6CAA-4BCC-ABC3-98967AFF1E22>

Corresponding author: Oleg Pekarsky (opbp@t-online.hu)

Academic editor: D. Lafontaine | Received 15 August 2012 | Accepted 6 November 2012 | Published 15 November 2012

<urn:lsid:zoobank.org:author:40DC027F-FCF3-4953-AC60-C071E814A768>

Citation: Pekarsky O (2012) A new species of *Schrankia* Hübner, 1825 from China (Lepidoptera, Erebidae, Hypenodinae). ZooKeys 242: 43–50. doi: [10.3897/zookeys.242.3856](https://doi.org/10.3897/zookeys.242.3856)

Abstract

A new species of the genus *Schrankia* Hübner, 1825, *S. pelicano* sp. n. is described from Sichuan, China. A diagnostic comparison is made with *Schrankia taenialis* (Hübner, [1809]) and *Schrankia separatalis* (Herz, 1904); illustrations of the male holotype and its genitalia are provided. A checklist of the genus with synonyms is given.

Keywords

Lepidoptera, Erebidae, *Schrankia*, new species, China

Introduction

Schrankia Hübner, 1825 is a widespread genus of Erebidae, being distributed in the Northern hemisphere, the Oriental and Australasiatic regions. It is represented by three species in Europe (Fibiger et al. 2010, Sinev 2008), three species in Africa and a single species on the Seychelles (Hacker 2004), twelve species in eastern and south-eastern Asia (Kononenko et al. 1998, Kononenko and Han 2007, Inoue 1979, Sugi 1982, Holloway 2008, Galsworthy 1997, Kendrick 2002). A considerable number of species live in Australasiatic region (Holloway 1977, Robinson 1975, Evenhuis 2007); three

species in Central America one of which, *S. macula* (Druce, 1891), also inhabits North America (Lafontaine and Schmidt 2010); there are only two valid taxa remaining in Hawaii after the remarkable revisionary work of Medeiros et al. (2009). The European taxon, *S. intermedialis* Reid, 1972 is treated by Fibiger et al. (2010) as an interspecific hybrid of *S. costaestrigalis* and *S. taenialis*. The genus has been reported twice from Hong Kong (Galsworthy 1997, Kendrick 2002) but has not been mentioned in any Chinese literature (Chen et al. 1991, Chen 1999, Li-zhong 2005); this is the first record of the genus from south-west China.

Systematic part

Genus *Schrankia* Hübner, 1825

<http://species-id.net/wiki/Schrankia>

Figs 3, 8

Type-species. *Pyralis taenialis* Hübner, [1809]

Remarks. The genus *Schrankia* is characterized by the slender body and narrow, weakly sclerotized wings, light-brown ground color of the forewing and abdomen, long, straight labial palps, which are three to four times as long as the diameter of the eye (Zimmerman 1958) and the absence of ocelli. In the male genitalia (Fig. 8), uncus long, slightly curved; valva elongated with acute apex, bearing three well-developed processes in the middle; juxta X-shaped, composed of two well-sclerotized, bent bars; aedeagus thin, elongated, slightly curved with club-like caecum. The externally often confusingly similar species of the genus *Hypenodes* (type-species *H. humidalis* Doubleday, 1850) have a smaller and thinner body, narrow wings, grey or brownish forewing ground color, upcurved labial palps; ocelli also are absent. The configuration of the male genitalia is very uniform throughout the genus, having a very simple, long, narrow valva with small and thin processes at the base, and a short, wide aedeagus with a tapered caecum.


Schrankia pelicano sp. n.

urn:lsid:zoobank.org:act:033BC935-14EC-4A82-B753-15C530773B3D

http://species-id.net/wiki/Schrankia_pelicano

Figs 1, 2, 5–7


Type material. Holotype male. China, Sichuan, N29°52.808', E102°50.240', near Ying Jing, bamboo forest, 700 m, 4.IV.2011, leg. Floriani; slide No.: OP1429m (coll. O. Pekarsky, deposited in the HNHM Budapest). **Paratypes.** China, Sichuan: 4 ♂♂, with same data as the holotype (coll. A. Saldaitis, Vilnius); 1 ♂ (coll. O. Pekarsky); 1 ♂ (coll. W. Speidel, Munich).



Figures 1–4. Adults. **1–2** *Schrankia pelicano* sp. n. **1** holotype male, China **2** paratype male, China **3** *S. taenialis* male, Hungary **4** *S. separatalis* male, Korea.

Etymology. The species name refers to the resemblance of the opened male genitalia to a pelican.

Diagnosis. The new species possesses a number of the diagnostic characters of the genus *Schrankia* (e.g., absence of ocelli; long, straight palpi; wide valval base; presence of medial complex of processes; thin, elongated aedeagus). The autapomorphic features of *S. pelicano* are the very long palpi, being much longer than in other species of *Schrankia*, the biarticulate uncus, the extremely large sacculus, which is almost equal in size and similar in shape to the distal half of the valva. Considering the diagnostic characters mentioned previously, the new species is placed into *Schrankia*, but its generic position could change through a much needed revision of the entire genus. Among *Schrankia*, the new species is closest to the Oriental group of species known from Korea and Japan, *S. separatalis*, *S. dimorpha*, *S. kogii*, *S. masuii* and *S. seinoi*. This species-group is characterized by the flat, thin, weakly sclerotized (almost transparent) distal half of valva, which has a rounded apex, wider and more strongly sclerotized haunch-like valval base, the well-developed apical saccular projection, and the less broadened caecum. The numerous autapomorphic features of the male genitalia of the new species (Figs 5–7) make it difficult to determine its closest relative within *Schrankia*. The comparison is provided here with *S. taenialis* (Fig. 8), the type-species of the genus and *S. separatalis* (Fig. 9), the most similar representative of the Oriental group of species. *S. pelicano* is similar externally to *S. taenialis* and *S. separatalis* (Fig. 4) but has longer labial palpi, narrower, more elongated forewing with a less acute apex and straight,



Figures 5–9. Male genitalia. **5–7** *Schrankia pelicanus* sp. n., holotype, China, slide No. OP1429m **5** clasping apparatus and aedeagus with vesica inverted **6** juxta (medially enlarged) **7** 8th abdominal segment **8** *S. taenialis* Hungary, slide No. 10059 RL **9** *S. separatalis* Korea, slide V. Kononenko.

almost parallel crosslines without undulations or dentations. The male genitalia of *S. pelicanus* differ from those of the related species by the long and narrow, subapically constricted valva, very large sacculus, digitiform ampulla, and basally clavate uncus

with a long, narrow, acute distal part. *S. taenialis* (Fig. 8) has a narrow valva with an acute apex, a large and broad ampulla and a long uncus. The genitalia of *S. separatalis* have, in comparison with *S. pellicano*, short and wide valva with small sacculus and much thicker, continuously curved uncus having half-cylindrical cross-section (Fig. 9).

Description. Male (Figs 1, 2). Wingspan 16–17 mm, length of forewing 8 mm. Head, thorax and abdomen ochreous grey; ocelli absent; tongue well developed; palpi very long (almost 5 times longer than diameter of eye), straight, 2nd segment (with appressed scales) about 2.8 times longer than 3rd segment, which is porrect; antenna with fine ciliation ventrally along full length, basal joint without pectination. Forewing elongate, narrow with acute apex, outer margin with rounded termen; ground color grey brown; costa straight with five milk-white patches; basal line barely visible, semi-circular; antemedial line black, straight, oblique, curved upward near costa; postmedial line black, straight, oblique, extending from 2/3 from base on hind margin of wing to 9/10 from base on costal margin, edged on outer side by narrow yellow postmedial fascia; subterminal line faint, pale, irregular, parallel to outer margin of wing; terminal line black, most prominent between wing veins; cilia yellow at base with dark medial line, grey distally. Hindwing ochreous grey, discal spot grey, terminal line black; cilia pale yellow at base, grey distally. Abdomen slender, long. Female unknown.

Male genitalia (Figs 5–7). Uncus biarticulate, consisting of clavate and setose main part and long, narrow, subapically curved, bill-like extension with pointed tip; tegumen narrow, as long as vinculum; scaphium well sclerotized, distally dilated; subscaphium membranous; juxta large, triangular, wider at base, weakly sclerotized (almost transparent), with two narrow, strongly sclerotized lateral plates with serrated inner edges (Fig. 6); vinculum strong, cup shaped. Valva elongated, conspicuously constricted subapically; cucullus almost rounded with acute tip; corona absent; sacculus very large, elongated, with very long, heavily sclerotized, distally dilated saccular extension, almost as long as distal half of valve; clavus unspecialised; clasper/ampulla complex large, sclerotized, setose, cuneate with broad base and finely pointed apex. Aedeagus tubular, thin and gracile, finely undulate. Vesica relatively short (shorter than aedeagus), thinly tubular (as broad as average width of aedeagus), with fine granulose scobination throughout.

Distribution. The species is known only from the type-locality, South-West China, Province Sichuan.

Checklist of *Schrankia*

Europe

S. taenialis (Hübner, [1809]) TL: Europe

syn. *albistrigatis* Haworth, 1809 [TL: Britain]

S. costaestrigalis (Stephens, 1834) TL: Wittlesea Mere, Swaffham, Norfolk

syn. *exsularis* Meyrick, 1888 TL: New Zealand, Taranaki

syn. *triangulalis* Hudson, 1923

- syn. *costistrigalis* Dannehl, 1925 [TL: Italy]
 syn. *lugubralis* Dannehl, 1925 TL: South Tirol, Italy
 syn. *monotona* Lempke, 1949 [TL: Holland]
 syn. *unicolor* Lempke, 1949 [TL: Holland]
 syn. *virgata* Lempke, 1966 [TL: Holland]
 syn. *hartigi* Berio, 1991 TL: Sardinia
S. balneorum (Alphéraky, 1880) TL: N. Caucasus
 syn. *bosporella* Budashkin & Klyuchko, 1990 TL: Crimea

Africa

- S. solitaria* Fletcher, 1961 TL: Ruwenzori, Mahoma River [Uganda]
S. microscopica (Berio, 1962) TL: Aldabra Islands [Seychelles]
S. namibiensis Hacker, 2004 TL: Namibia, Brandberg, Am Königstein
S. scopariooides Hacker, 2004 TL: Namibia, Brandberg, Hungarob-valley

Asiatic region

- S. obstructalis* (Walker, [1866]) TL: Sarawak [Malaysia]
S. croceipicta (Hampson, 1893) TL: Ceylon, Pundaloya
 syn. *croceipicta aegrota* Berio, 1962; 179, TL: Seychelles, Mahe B., Vallon
S. aurantilineata (Hampson, 1896) TL: Ceylon
S. separatalis (Herz, 1904) TL: Korea
 syn. *squalida* Wileman & South, 1917; 28, TL: Japan
S. dimorpha Inoue, 1979 TL: Kagawa Pref., Shinoe, Fudodaki [Japan]
S. kogii Inoue, 1979 TL: Hokkaido, Shintoku, Kuttari
S. masuii Inoue, 1979 TL: Kagawa Pref., Shinoe, Oyashiki [Japan]
S. seinoi Inoue, 1979 TL: Amami-Oshima Is., Sumiyoson [Japan]
S. bilineata Galsworthy, 1997 TL: Hong Kong
S. pelicano sp. n. TL: China, Sichuan
S. bruntoni Holloway, 2008 TL: Ulu Temburon, Brunei
S. dusunorum Holloway, 2008 TL: Sabah, Ulu Dusun, 30mls W of Sandakan [Malaysia]
S. spiralaedeagus Holloway, 2008 TL: Sarawak, Gunong Mulu Nat. Park [Malaysia]

Australasiatic region

- S. calligrapha* Snellen, 1880 TL: New Hebrides, Aneityum, Red Crest, 3 km NE of Anelgauhat
S. taona (Tams, 1935) TL: Samoa, Savaii
S. capnophanes (Turner, 1939) TL: Tasmania, Mt. Wellington
S. dochmographa Fletcher, 1957 TL: Solomon Is., Rennell I., Hutuna
S. furoroa Robinson, 1975 TL: Fiji, Rotuma, Furoroa
S. vitiensis Robinson, 1975 TL: [Fiji]
S. boisea Holloway, 1977 TL: New Caledonia, Port Boise
S. cheesmaniae Holloway, 1977 TL: New Hebrides, Aneityum, Red Crest, 3 km NE of Anelgauhat

- S. daviesi* Holloway, 1977 TL: Norfolk Is., N. Mission Road
S. erromanga Holloway, 1977 TL: New Hebrides, Erromango I., Nouankao Camp
S. karkara Holloway, 1977 TL: New Guinea, Karkar I., Dampier I.
S. nokowula Holloway, 1977 TL: New Hebrides, Sanot, Mt. Tabwemasana, Nokowula
S. nouankaoa Holloway, 1977 TL: New Hebrides, Erromango I., Nouankao Camp
S. tabwemasana Holloway, 1977 TL: New Hebrides, Santo, Mt. Tabwemasana, Nokowula
S. tamasi Holloway, 1977 TL: Samoa, Upolu I., Mt. Vaea

Neotropical region

- S. macula* (Druce, 1891) TL: Panama, Chiriquí
S. flualis (Schaus, 1916) TL: Panama, Trinidad River
S. musalis (Schaus, 1916) TL: Panama, Trinidad River

Oceanian region

- S. altivolans* (Butler, 1880) TL: Hawaii, Mauna Loa
syn. *simplex* (Butler, 1881) TL: Hawaii
syn. *oxygramma* (Meyrick, 1899) TL: Kaua, Kaholuamano [Hawaii]
syn. *sarothrura* (Meyrick, 1899) TL: Hawaii, Ola
syn. *arrhecta* (Meyrick, 1904) TL: Hawaii, Mt. Waimea
S. howarthi Davis & Medeiros, 2009 TL: Hawaii

Acknowledgements

I would like to express my sincere thanks to Aidas Saldaitis for access to his collection; to László Ronkay for his critical comments on the manuscript, to Gábor Ronkay for the image of *Srankia taenialis*, Jürgen Schmidl for help with finding literature and to Vladimir Kononenko for the images of *Schrankia separatalis*, help with finding literature, valuable suggestions and critical comments on the manuscript.

References

- Chen YX (1999) Lepidoptera, Noctuidae. In: Zhu HF et al. (Ed) *Fauna Sinica Insect*, Vol. 16. Science Press, Beijing, 1596 pp.
Chen YX, Wang BH, Lin DW (1991) The Noctuids Fauna of Xizang. Henan Scientific and Technical Publishing House, Zhengzhou, 409 pp.
Evenhuis NL (2007) Checklist of Fijian Lepidoptera. Bishop Museum Technical Report 38(13) 53 pp. <http://hbs.bishopmuseum.org/fiji/checklists/lepidoptera.html>
Fibiger M, Ronkay L, Yela J, Zilli A (2010) Rivulinae, Boletobiinae, Hypenodinae, Araeopteroninae, Eublemininae, Herminiinae, Hypeninae, Euteliinae and Micronoctuidae in-

- cluding Supplement to Volumes 1–11. Noctuidae Europaea, volume 12. Entomological press, Sorø, 451pp.
- Galsworthy AC (1997) New and revised species of macrolepidoptera from Hong Kong. Memoirs of the Hong Kong Natural History Society. 21: 127–150, 25 figures, 1 plate.
- Hacker H (2004) Nolidae and Noctuidae (excluding Catocalinae: Audeini and Tachosini) (Lepidoptera). In: May EW (Ed) The Lepidoptera of the Brandberg Massif in Namibia Part 1. Esperiana Meimor 1, Delta Druck u. Verlag, Schwanfeld, 261–304.
- Holloway JD (1977) The Lepidoptera of Norfolk Island, their biogeography and ecology. Series Entomologica 13. W. Junk, The Hague: 292 pp.
- Holloway JD (2008) The Moths of Borneo. Vol 17. Family Noctuidae, subfamilies Rivulinae, Phytometrinae, Herminiinae, Hypeninae and Hypenodinae. Malayan Nature Journal 60(1-4): 1–268.
- Inoue H (1979) Genera *Schränkia* Hübner and *Luceria* Walker from Japan (Lepidoptera: Noctuidae) Tinea 10 (30): 299–308.
- Kendrick RC (2002) Moths (Insecta: Lepidoptera) of Hong Kong. University of Hong Kong, Hong Kong 660 pp. <http://www.libsearch.com/visit/463010>
- Kononenko VS, Ahn SB, Ronkay L (1998) Illustrated catalogue of Noctuidae in Korea (Lepidoptera). In: Insects of Korea [3], 509 pp.
- Kononenko VS, Han HL (2007) Atlas genitalia of the Noctuidae in Korea (Lepidoptera). In: Park KT (Ed) Insects of Korea [11]. 464 pp.
- Lafontaine JD, Schmidt BC (2010) Annotated check list of the Noctuoidea (Insecta, Lepidoptera) of North America north of Mexico. ZooKeys 40: 1–239. doi: 10.3897/zookeys.40.414
- Li-zhong H (2005) List of Chinese Insects. Vol. III. Lepidoptera (Sun Yat-Sen) University Press, Guangzhou, 595 pp.
- Medeiros MJ, Davis D, Howarth FG, Gillespie R (2009) Evolution of cave living in Hawaiian *Schränkia* (Lepidoptera: Noctuidae) with description of a remarkable new cave species. Zoological Journal of the Linnean Society, 2009, 156,:114–139. doi: 10.1111/j.1096-3642.2008.00477.x
- Robinson GS (1975) Macrolepidoptera of Fiji and Rotuma: a taxonomic and biogeographic study. E. W. Classey Ltd., 362 pp.
- Sinev SY (Ed) (2008) Catalogue of the Lepidoptera of Russia. KMK Scientific Press, St. Petersburg. 424 pp.
- Sugi S (1982) Noctuidae (except Herminiinae). In Inoue H, Sugi S, Kuroko H, Moriuti S, Kawabe A. Moths of Japan, Vol. 1, 669–913 pp, Vol. 2, 334–405 pp, Kodansha, Tokyo.
- Zimmerman EC (1958) Insects of Hawaii, Vol. 7: Macrolepidoptera. University of Hawaii Press, Honolulu, 542 pp.