ZooKeys 326: 55–67 (2013) doi: 10.3897/zookeys.326.5871 www.zookeys.org

RESEARCH ARTICLE



Revision of the genera *Hovadelium* Ardoin and *Mimolaena* Ardoin (Coleoptera, Tenebrionidae, Laenini) from Madagascar, with remarks on tribal assignment¹

Wolfgang Schawaller^{1,†}

Staatliches Museum für Naturkunde, Rosenstein 1, D-70191 Stuttgart, Germany

+ http://zoobank.org/D3B396E3-69A8-405D-8500-0504698C1DAA

Corresponding author: Wolfgang Schawaller (wolfgang.schawaller@smns-bw.de)

Academic editor: Patrice Bouchard Received 28 June 2012 Accepted 20 August 2013	Published 26 August 2013

Citation: Schawaller W (2013) Revision of the genera *Hovadelium* Ardoin and *Mimolaena* Ardoin (Coleoptera, Tenebrionidae, Laenini) from Madagascar, with remarks on tribal assignment. ZooKeys 326: 55–67. doi: 10.3897/zook eys.326.5871

Abstract

The genera *Hovadelium* Ardoin, 1961 and *Mimolaena* Ardoin, 1961, endemic in Madagascar, are revised and assigned to the tribe Laenini Seidlitz, 1896 (subfamily Lagrinae Latreille, 1825). New species: *Hovadelium ardoini* **sp. n.**, *Hovadelium bremeri* **sp. n.** and *Mimolaena janaki* **sp. n.** An identification key is compiled for all taxa. Distribution of *Hovadelium* (5 species) and *Mimolaena* (3 species) is mapped. The congeners might be indicator species for the highly endangered mature forests in Madagascar.

Keywords

Coleoptera, Tenebrionidae, Lagriinae, Laenini, Hovadelium, Mimolaena, taxonomy, new species, Madagascar

Introduction

Ardoin (1961) described the genera *Hovadelium* Ardoin, 1961 (type species *H. discoidale* Ardoin, 1961) and *Mimolaena* Ardoin, 1961 (type species *M. pauliani* Ardoin, 1961), endemic to Madagascar, and placed them into the tenebrionid tribe Adeliini. Subsequently, additional species were described by Ardoin (1976) and Ferrer (1998).

¹ Contributions to Tenenbrionidae no. 113. For no. 112 see: Annales Zoologici 63, 2013.

Copyright Wolfgang Schawaller. This is an open access article distributed under the terms of the Creative Commons Attribution License 3.0 (CC-BY), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

In the revision of the tribe Adeliini, Matthews (1998) mentioned, that the tribal assignment of the Malagasy genera *Hovadelium* and *Mimolaena* either to Adeliini or to Laenini remains doubtful. Both genera are placed now herein finally into Laenini because of the lack of defensive glands. Different genera of this tribe are also known from South Africa (Endrödy-Younga and Schawaller 2002, Ferrer 2005).

So far, all descriptions were based only on single specimens. Recently, a huge number of newly collected specimens, mainly of *Hovadelium*, were handed over to the author for examination by Prof. Dr. H. J. Bremer (Osnabrück, Germany). This material, including three so far undescribed species, is represented herein, together with reexamination of the previously described taxa.

As other members of Laenini, all species are wingless and have restricted distributional patterns (Map see Fig. 1). So far, all records originate from the southeastern part of the island, additional taxa might be present in the northeastern part. Living in litter of the broadleaved evergreen forests, the congeners can be considered as indicator species for these mature and highly endangered forests in Madagascar.

Depositories

CRFL	Collection René Fouquè, Liberec, Czech Republic
HNHM	Hungarian Natural History Museum, Budapest, Hungary
MNHN	Muséum national d'Histoire naturelle, Paris, France
MZUF	Museo Zoologico de "La Specola", Firenze, Italy
NMPC	National Museum, Prague, Czech Republic
SMNS	Staatliches Museum für Naturkunde, Stuttgart, Germany
TMSA	Ditsong National Museum of Natural History, Pretoria, South Africa
ZSM	Zoologische Staatssammlung, München, Germany

Tribal assignment

Some specimens of *Hovadelium ardoini* sp. n. were sent to E. Matthews (Adelaide) for a personal dissection of the female genital tract. The examination showed, that defensive glands are completely absent in all the dissected specimens, which is characteristic for Laenini and the only difference to Adeliini.

Matthews (in litteris): "I have dissected the females and can't see any trace of defensive glands, certainly not the long ones between segments 8 and 9 which are found in all Adeliini. There are no 7/8 glands either, although finding those in Laenini would not be surprising since they are characteristic of many Lagriinae. Stridulatory files (plectron) are absent. There are also no vaginal sclerites, such as the ones I found in one *Laena* (and most Adeliini). The spermatheca consists of three short wide tubules on the side of the vagina, similar to those of a species of *Laena* (Matthews 1998: fig. 57) which however has two long narrow tubules. Too few Laenini have been dissected for us to know the



Figure 1. Records of Laenini in Madagascar, vegetation map (modified from NationMaster.com). *Hovadelium discoidale* could not be mapped (known only from "Madagascar" without detailed locality).

significance of these details, but the general configuration of the female system is typical of Laenini/Adeliini. The aedeagus of the male is of the usual simple type."

Ovipositor and female genital tract (Fig. 15): Paraproct and coxite subequal in length, coxite lobes 3 and 4 fused, digitiform, gonostyles terminal in position, coxite baculi transverse, paraproct baculi longitudinal, spiculum gastrale a slender rod without terminal fork. No bursa copulatrix, three short and wide spermathecal tubules attached to side of vagina, vaginal sclerites absent, long slender spermathecal accessory gland attached to anterior end of vagina.

The genus Hovadelium

Hovadelium aeneum Ardoin, 1961

http://species-id.net/wiki/Hovadelium_aeneum Figs 2, 10

Reexamined type-material. C Madagascar, Plateau Soaindrana, Andringitra-Ambalavao, 2090 m, 16.I.1958, leg. R. Paulian, $\stackrel{\circ}{\rightarrow}$ holotype MNHN.

New material. C Madagascar, Andringitra, Andohariana, 2000–2100 m, mission C.N.R.S., 1 ex. MNHN (det. Ardoin). – E Madagascar, Massiv Ambondrombe, Ikoka, 1300–1400 m, 12.–13.III.1996, leg. J. Janák & P. Moravec, 11 ex. SMNS, 5 ex. ZSM. – E Madagascar, Massiv Ambondrombe, 1300–1400 m, 14.III.1996, leg. J. Janák & P. Moravec, 11 ex. SMNS, 5 ex. TMSA, 5 ex. ZSM. – E Madagascar, Massiv Ambondrombe, cote 1579, 1500–1600 m, 15.–18.III.1996, leg. J. Janák & P. Moravec, 11 ex. SMNS, 5 ex. NMPC, 5 ex. ZSM. – E Madagascar, Massiv Ambondrombe, 1700 m, 17.III.1996, leg. J. Janák & P. Moravec, 1 ex. SMNS.

Diagnostic characters. Body length 2.7–4.0 mm (the holotype has a length of 4.0 mm and not of 5.0 mm as given by Ardoin 1961). Pronotum subquadrate, widest before the middle, with rounded anterior and posterior angles, surface slightly convex, with fine and sparse punctation, between punctation slightly shagreened. Elytra with punctural rows in striae, intervals convex, slightly shagreened and with an irregular row of tubercles, interval 7 at base near shoulders with a longer seta. Aedeagus see Fig. 10.

Hovadelium ardoini sp. n.

http://zoobank.org/9E7A178D-AC88-4A2A-A947-5A05D8931C50 http://species-id.net/wiki/Hovadelium_ardoini Figs 3, 11

Type specimens. Holotype male: E Madagascar, Fianaratsoa Prov., Ambohimahamasoa, 1300–1400 m, 21.–23.III.1996, leg. J. Janák & P. Moravec, SMNS. – Paratypes: Same data as holotype, 20 ex. SMNS, 5 ex. HNHM, 5 ex. TMSA, 5 ex. ZSM. – E Madagascar, Ranomafana NP, Sahavondrona, 1150–1250 m, 3.–4.II.1995, leg. J. Janák, 13 ex. SMNS, 5 ex. NMPC, 5 ex. ZSM. – E Madagascar, Ranomafana NP, Vohiparara, 1100–1200 m, 21.–24.I.1993, leg. J. Janák, 2 ex. SMNS. – E Madagascar, Fianaratsoa Prov., Ranomafana, 29.XI.–2.XII.1995, leg. I. Jeniš, 1 ex. SMNS. – E Madagascar, Maromiza, Andasibe (Périnet), 1000–1200 m, 9.II.1993, leg. J. Janák, 1 ex. SMNS. – E Madagascar, Maromiza, Andasibe (Périnet), 930–1000 m, 7.–10.I.1995, leg. J. Janák, 5 ex. SMNS. – E Madagascar, Maromiza, Andasibe (Périnet), 7.XI.1998, leg. R. Müller, 1 ex. TMSA.

Diagnosis. *H. ardoini* sp. n. is similar to *H. discoidale* Ardoin, 1961, but lacks the striking deep groove ventral of the eyes. Both can be separated also by the shape of the pronotum widest behind the middle in *H. discoidale* (Fig. 5), but widest in the middle in *H. ardoini* sp. n. (Fig. 3), and by the anterior angles of the pronotum, which are



Figures 2–5. Dorsal view of *Hovadelium* species. 2 *H. aeneum*, holotype MNHM 3 *H. ardoini* sp. n., paratype SMNS 4 *H. bremeri* sp. n., holotype SMNS 5 *H. discoidale*, holotype MNHN. – Scale line 2 mm.

distinctly marked in *H. discoidale*, and completely rounded in *H. ardoini* sp. n. The elytral punctural rows are identical in both species, but the disc of the elytra is flattened in *H. discoidale*, whereas in *H. ardoini* sp. n. the elytra are more convex. See also under *Hovadelium bremeri* sp. n. and key below.

Description. Body length 3.3–4.7 mm, unicoloured dark brown. Eyes flat, not prominent; without a deep groove ventral of the eyes. Head with deep clypeal suture and two pairs of long setae as characteristic for the genus; frons shining and without punctures. Shape of the antennomeres see Fig. 3. Pronotum subquadrate, widest in the middle, anterior and posterior angles completely rounded, anterior and posterior margin finely bordered, lateral margins with broader border, anterior margin not excavated; surface slightly convex, with fine and sparse punctation, punctures only weakly impressed, surface between punctation shining and only slightly shagreened; propleura shining, without punctation. Elytra with nine punctural rows in distinct striae, these punctures small and elongate, only slightly broader than striae, without setae; intervals convex, shining and without punctures nor tubercles, interval 7 at base near shoulders with a longer seta. Ventrites shining, ventrites 1–4 in the middle with a pair of longer setae, last ventrite in both sexes unbordered. Femora and tibiae in both sexes without teeth or other modifications. In males protarsi only slightly dilatated, without other external differences. Aedeagus see Fig. 11.

Etymology. Named in honour of Jean Paul Ardoin (1918–1978), former pharmacist in Arcachon (France), author of the Malagasy genera of Laenini and specialist of other tenebrionids from Africa and Madagascar.

Hovadelium bremeri sp. n.

http://zoobank.org/1F59CF9F-4AF5-4510-9915-F1BB12C6CFAA http://species-id.net/wiki/Hovadelium_bremeri Figs 4, 12

Type specimens. Holotype male: E Madagascar, 30 km ESE Betroka, Vohitrosa Forest, 1400–1500 m, 17.–18.XII.1998, leg. J. Janák, SMNS. – Paratypes: Same data as holotype, 7 ex. SMNS, 2 ex. ZSM.

Diagnosis. *Hovadelium bremeri* sp. n. and *Hovadelium ardoini* sp. n. are similar, both share the general body shape, the lacking of a groove ventral of the eyes, the shining surface of pronotum and elytra, the elytral interval 7 at base near shoulders with a longer seta, and the lacking tubercles on the elytral intervals. In *H. bremeri* sp. n., the body length is somewhat shorter in the average, the pronotum is narrower towards base, the pronotal punctation is larger and denser, the anterior pronotal margin is unbordered in the middle, the punctures of the elytral rows are larger, and the apicale of the aedeagus is shorter. See also under *Hovadelium ardoini* sp. n. and key below.

Description. Body length 2.8–3.5 mm, unicoloured dark brown. Eyes flat, not prominent; without a deep groove ventral of the eyes. Head with deep clypeal suture and two pairs of long setae as characteristic for the genus; frons shining and without punctures. Shape of the antennomeres see Fig. 4. Pronotum subquadrate, widest in the

middle, anterior and posterior angles completely rounded, anterior margin unbordered in the middle, posterior margin finely bordered, lateral margins with broader border, anterior margin not excavated; surface slightly convex, with irregular larger, but not confluent punctation, punctures only weakly impressed, surface between punctation shining; propleura shining, without punctation. Elytra with nine punctural rows in weak striae, these punctures large and broader than striae, without setae; intervals convex, shining and without punctures nor tubercles, interval 7 at base near shoulders with a longer seta. Ventrites shining, ventrites 1–4 in the middle with a pair of longer setae, last ventrite in both sexes unbordered. Femora and tibiae in both sexes without teeth or other modifications. In males protarsi only slightly dilatated, without other

Etymology. Named in honour of Prof. Dr. H. J. Bremer (Osnabrück, Germany), who provided me with most of the newly collected specimens, and allowed to keep the larger part in SMNS.

Hovadelium discoidale Ardoin, 1961

external differences. Aedeagus see Fig. 12.

http://species-id.net/wiki/Hovadelium_discoidale Fig. 5

Reexamined type-material. "Madagascar" (without detailed data), collection Oberthuer, male holotype MNHN.

Remarks. The type specimen has a quite unique character, namely the head with a deep groove ventral of the eyes (Fig. 5). Ardoin (1961) assumed that this might be a sexualdimorph character of males within the genus. However, in all the plenty herein presented males and females of other species of *Hovadelium*, such a groove is not present. Thus, this structure (of unknown biological function) is considered as not generic but just as specific for *H. discoidale*. Unfortunately, an exact type locality is unknown.

Diagnostic characters. Body length 4.2 mm (not 5.0 mm as given by Ardoin 1961). Head with a deep groove ventral of the eyes (Fig. 5). Pronotum subquadrate, widest somewhat behind the middle, with marked anterior and rounded posterior angles, surface slightly convex, with fine and sparse punctation, between punctation slightly shagreened. Elytra with punctural rows in striae, intervals convex, shining and without tubercles, interval 7 at base near shoulders with a longer seta. Aedeagus not examined herein (because of the fragility of the type).

Hovadelium elongatum Ardoin, 1976 http://species-id.net/wiki/Hovadelium_elongatum Figs 6, 13

Reexamined type-material. None, not found in MNHN. Type locality. SE Madagascar, Plateau Andohahelo, SE Trafonaomby, 1770–1950 m.



Figures 6–9. Dorsal view of *Hovadelium* and *Mimolaena* species. 6 *H. elongatum*, non-type SMNS 7 *M. clarissae*, holotype MZUF 8 *M. janaki* sp. n., holotype SMNS. 9 *M. pauliani*, holotype MNHN. – Scale line 2 mm.

New material. SE Madagascar, 3 km NW Fort Dauphin, Pic St. Louis, 150–250 m, 19.II.2004, leg. P. Bulirsch, 6 ex. CRFL, 2 ex. SMNS. – E Madagascar, 38 km ESE Betroka, Kalambatrita Forest, 3 km SSE Ambaro, 1400 m, 29.XII.1998, leg. J. Janák, 2 ex. SMNS.

Remarks. I hope not to fail in assigning the newly collected specimens to this species, described upon a single female. Distinct differences between the description and the new specimens do not exist, and the larger part of the new material originates from the surroundings of the type locality nearby Fort Dauphin.

Diagnostic characters. Body length 4.0–6.0 mm (holotype 5.0 mm). Pronotum cordiform, anterior margin regularly excavated, widest before the middle, with rounded anterior and posterior angles, surface slightly convex, with fine and sparse punctation, between punctation distinctly shagreened. Elytra with punctural rows in striae, intervals convex, slightly shagreened and somewhat uneven ("petit granules peu saillants" in description), interval 7 at base near shoulders with a longer seta. Aedeagus see Fig. 13.

The genus Mimolaena

Mimolaena clarissae Ferrer, 1998

http://species-id.net/wiki/Mimolaena_clarissae Fig. 7

Reexamined type-material. E Madagascar, Ambila Lemaitso (labelled as Ambila La Maintso), V.1990, leg. C. Raharimina, female holotype MZUF.

Diagnostic characters. Body length 4.4 mm. Pronotum subquadrate, with slightly prominent anterior and with rounded posterior angles, surface slightly convex and with rough and partly confluent punctures. Elytra with rough irregular punctation without any separation in rows and intervals. Aedeagus unknown, only \mathcal{Q} holotype known.

Mimolaena janaki sp. n.

http://zoobank.org/CB2004A1-7F5A-423C-99C7-AF4BAC9E1EE3 http://species-id.net/wiki/Mimolaena_janaki Figs 8, 14

Type specimens. Holotype male: E Madagascar, Ranomafana NP, Vohiparara, 1100–1200 m, 21.–24.I.1993, leg. J. Janák, SMNS. – Paratypes: Same data as holotype, 3 ex. SMNS, 2 ex. ZSM. – E Madagascar, Massiv Ambondrombe, 1600–1700 m, 17.III.1996, leg. J. Janák & P. Moravec, 1 \bigcirc SMNS. – E Madagascar, Massiv Ambondrombe, 1500–1600 m, 15.–18.III.1996, leg. J. Janák & P. Moravec, 1 \bigcirc ZSM. – E Madagascar, Massiv Ambondrombe, 1300–1400 m, 14.III.1996, leg. J. Janák & P. Moravec, 1 \bigcirc SMNS.



Figures 10–14. Aedeagus of *Hovadelium* and *Mimolaena* species. 10 *H. aeneum*, non-type SMNS 11 *H. ardoini* sp. n., holotype SMNS 12 *H. bremeri* sp. n., holotype SMNS 13 *H. elongatum*, non-type SMNS 14 *M. janaki* sp. n., holotype SMNS. – Scale line 0.5 mm.

Diagnosis. To be recognized by the shape of the pronotum with spine-like posterior angles, by scattered and fine punctation of the pronotum, by only six elytral rows of punctures extinguished in the posterior and external part of elytra, and by the shape of the aedeagus. The two other known species of *Mimolaena* possess nearly rounded posterior angles of the pronotum, and the punctation on the pronotum is either fine and the elytra bear punctural rows (*M. pauliani* Ardoin, 1961), or the punctation on the pronotum is rough and dense and the elytra bear an dense irregular punctation not separated in rows and intervals (*M. clarissae* Ferrer, 1998). See also key below.

Description. Body length 3.5–4.7 mm, unicoloured dark brown. Eyes (Fig. 8) not reduced, slightly prominent. Shape of the antennomeres see Fig. 8. Shape of pronotum see Fig. 8, disc with a few scattered punctures, most punctures bearing a longer erect seta; surface without any impressions, surface shining, lateral margins bordered, basal margin bordered and not bent downwards, posterior angles prominent spine-like, propleura unpunctured. Elytra (Fig. 8) with only six punctural rows without striae, these rows extinguishing in posterior and external part of elytra, punctures of rows of similar size as pronotal punctures, punctures of the elytral rows without setae,



Figure 15. Ovipositor and female genital tract of *Hovadelium ardoini* sp. n. (drawing by Eric Matthews). – Scale line 1 mm.

a few additional punctures apart from the rows laterally and distally on the elytra bear a longer erect seta, intervals shining without any punctures and setation, intervals flat. Ventrites shining, in males with fine punctation and short setation, in females nearly unpunctured and without setation, last ventrite in both sexes unbordered. Femora and tibiae in both sexes without teeth or other modifications. Aedeagus see Fig. 14.

Etymology. Named in honor of J. Janák, one of the collectors of the type series and of other Malagasy Laenini.

Mimolaena pauliani Ardoin, 1961

http://species-id.net/wiki/Mimolaena_pauliani Fig. 9

Reexamined type-material. C Madagascar, Plateau Soaindrana, Andringitra-Ambalavao, 2090 m, 16.I.1958, leg. R. Paulian, male holotype and 1 male paratype MNHN.

Diagnostic characters. Body length 4.4–4.8 mm. Pronotum subquadrate, with rounded anterior and posterior angles, surface flat and with sparse and fine punctation. Elytra with distinct punctural rows without striae, size of the punctures diminishing

laterally and apically. Aedeagus not examined herein (because of the fragility of the type). It is said (but not figured) in the description, that the apicale is short and acute at the tip, and the basale is long and bent.

Identification key of Laenini from Madagascar

Base of elytra excavated for pronotum, humeral angle protruding; head be-1 tween eyes and clypeus each with a pair of long tactile setae; elytral interval 7 at base near shoulders with a long tactile seta (genus *Hovadelium*)2 Base of elytra not excavated, humeral angle rounded; head with irregular short setation, without pairs of long setae; elytral interval 7 at base near shoulders without a long tactile seta (genus *Mimolaena*)......6 2 3 Pronotum cordiform, surface of head and pronotum distinctly mat and sha-Pronotum subquadrate, surface of head and pronotum shining or at most weakly shagreened......4 Pronotum widest before the middle, elytral intervals with an irregular row of 4 5 Lateral margin of pronotum regularly rounded towards anterior and posterior angles, anterior margin of pronotum completely bordered, pronotal punctation fine, punctures of elytral rows small, elongate and not distinctly Lateral margin of pronotum narrower towards base, anterior margin of pronotum unbordered in the middle, pronotal punctation larger and denser, punctures of elytral intervals larger and broader than weak striae - Fig. 4 6 Posterior angles of pronotum prominent spine-like – Fig. 8.... M. janaki sp. n. Posterior angles of pronotum rounded......7 7 Pronotum with fine and separate punctation, elytra with punctural rows -Pronotum with rough and partly confluent punctation, elytra with rough irregular punctation without any separation in rows and intervals – Fig. 7

Acknowledgements

I cordially thank Prof. Dr. Hans J. Bremer (Osnabrück), who provided me with most of the newly collected specimens already many years ago, and allowed to keep the larger part in SMNS. Dr. Luboš Purchart (Brno) kindly searched in Paris in the course of his visit successfully for most of the types of Ardoin. For the loan of material from the collections under their care I thank Dr. Luca Bartolozzi (Firenze), René Fouquè (Liberec), Antoine Mantilleri (Paris) and Ruth Müller (Pretoria). Dr. Eric Matthews (Adelaide) dissected some female genitalia, provided a figure, recognised the assignment to the tribe Laenini and generously allowed me to publish the results herein. The photographs were taken by Johannes Reibnitz (Stuttgart) with a Leica DFC320 digital camera on a Leica MZ16 APO microscope and subsequently processed by him with Auto-Montage (Syncroscopy) software. Two referees, unknown to me, improved the manuscript by their comments and corrections.

References

- Ardoin P (1961) Deux nouveaux genres d'Adeliini Malgaches. Bulletin de l'Académie Malgache 37(1959): 31–38.
- Ardoin P (1976) Les Tenebrionidae des Chaines Anosyennes Madagascar sud-est (1) (Coleoptera). Annales de la Société entomologique de France 12: 3–29.
- Endrödy-Younga S, Schawaller W (2002) First record of Laenini (Coleoptera: Tenebrionidae) from South Africa, with descriptions of two new genera and several new species. Annals of the Transvaal Museum 39: 9–21.
- Ferrer J (1998) Contribution à la connaissance des Tenebrionidae de Madagascar (Insecta, Coleoptera). Entomofauna, Zeitschrift für Entomologie 19: 353–404.
- Ferrer J (2005) Descriptión de una nueva especies de *Enicmosoma* Gebien de Angola y de un nuevo género y especie de Transvaal (Coleoptera, Tenebrionidae, Lagriinae). Nouvelle Revue d'Entomologie (NS) 21 (2004): 199–203.
- Matthews E G (1998) Classification, phylogeny and biogeography of the genera of Adeliini (Coleoptera: Tenebrionidae). Invertebrate Taxonomy 12: 685–824. doi: 10.1071/IT97008