

A new species of *Lycodon* Boie, 1826 (*Serpentes, Colubridae*) from northern Vietnam

Helen Y. Janssen^{1,2}, Cuong T. Pham³, Hanh Thi Ngo⁴, Minh Duc Le^{5,6,7},
Truong Q. Nguyen^{3,8}, Thomas Ziegler^{1,2}

1 AG Zoologischer Garten Köln, Riehler Strasse 173, D-50735 Cologne, Germany **2** Institute of Zoology, University of Cologne, Zülpicher Strasse 47b, D-50674 Cologne, Germany **3** Institute of Ecology and Biological Resources, Vietnam Academy of Science and Technology, 18 Hoang Quoc Viet, Cau Giay, Hanoi, Vietnam **4** Faculty of Biology, Hanoi University of Science, Vietnam National University, 334 Nguyen Trai Road, Hanoi, Vietnam **5** Faculty of Environmental Sciences, Hanoi University of Science, Vietnam National University, 334 Nguyen Trai Road, Hanoi, Vietnam **6** Central Institute for Natural Resources and Environmental Studies, Hanoi National University, 19 Le Thanh Tong, Hanoi, Vietnam **7** Department of Herpetology, American Museum of Natural History, Central Park West at 79th Street, New York, New York 10024, USA **8** Graduate University of Science and Technology, Vietnam Academy of Science and Technology, 18 Hoang Quoc Viet, Cau Giay, Hanoi, Vietnam

Corresponding author: Truong Q. Nguyen (nqt2@yahoo.com); Thomas Ziegler (ziegler@koelnerzoo.de)

Academic editor: Robert Jadin | Received 5 May 2019 | Accepted 25 July 2019 | Published 10 September 2019

<http://zoobank.org/F54D525B-03CD-4640-A750-1EEE13D71B06>

Citation: Janssen HY, Pham CT, Ngo HT, Le MD, Nguyen TQ, Ziegler T (2019) A new species of *Lycodon* Boie, 1826 (*Serpentes, Colubridae*) from northern Vietnam. ZooKeys 875: 1–29. <https://doi.org/10.3897/zookeys.875.35933>

Abstract

A new species of the genus *Lycodon* is described from Cao Bang Province, Vietnam, based on three individuals with distinct differences in morphology and molecular data. The new species is differentiated from its congeners by a combination of the following characters: dorsal scales in 17–17–15 rows, smooth throughout; supralabials usually eight (rarely nine); infralabials ten; one elongated loreal on each side, in contact with the eye; precloacal plate single; ventral scales 212–218 (plus one or two preventral scales); subcaudals 90 or 91; maxillary teeth 13 or 14; dorsal surface of body with 28 or 29 light body bands; dorsal surface of tail with 13 cream bands, forming a distinct blotch in the vertebral region. Based on phy-

logenetic analyses of mitochondrial cytochrome *b* sequence data, the new species is recovered as the sister species to a clade containing *L. multizonatus* and *L. liuchengchaoi* with strong support from the Bayesian analysis. The new species is at least 7.5% divergent from other species within this clade in uncorrected pairwise distance calculated using a fragment of more than 1000 bp of the mitochondrial cytochrome *b*. This discovery increases the number of *Lycodon* species known from Vietnam to 16.

Keywords

Cao Bang Province, *Lycodon pictus* sp. nov., morphology, phylogeny, taxonomy

Introduction

The genus *Lycodon* Boie, 1827 is one of the most diverse genera of colubrid snakes, with 61 currently recognised species (Uetz et al. 2019, Luu et al. 2019). Recent phylogenetic studies showed that the genera *Dinodon*, *Dryocalamus* and *Lepturophis* nested within *Lycodon* and suggested to place them into the genus *Lycodon* sensu lato (Guo et al. 2013; Siler et al. 2013; Figueroa et al. 2016). The members of this genus have a broad distribution from eastern Iran to southern China and Japan, southward to the Philippines as well as the Indo-Australian Archipelago (Lanza 1999; Siler et al. 2013; Neang et al. 2014). Six species of *Lycodon* have been described in the last five years, namely *L. zoosvictoriae* Neang, Hartmann, Hun, Souter & Furey, 2014 from Cambodia; *L. caverniculus* Grismer, Quah, Anuar, Muin, Wood & Nor, 2014 from Malaysia; *L. sidiki* Wostl, Hamidy, Kurniawan & Smith, 2017 from Indonesia; *L. banksi* Luu, Bonkowski, Nguyen, Le, Calame & Ziegler, 2018 from Laos, *L. namdongensis* Luu, Ziegler, Ha, Le & Hoang, 2019 from Vietnam and *L. gibsonae* Vogel & David, 2019 from Thailand. From Vietnam, fifteen species of *Lycodon* have been reported to date, comprising *L. capucinus* (Boie, 1827), *L. cardamomensis* (Daltry & Wüster, 2002), *L. davisonii* (Blanford, 1878), *L. fasciatus* (Anderson, 1879), *L. flavozonatus* (Pope, 1928), *L. futsingensis* (Pope, 1928), *L. laoensis* Günther, 1864, *L. meridionalis* (Bourret, 1935), *L. namdongensis* Luu, Ziegler, Ha, Le & Hoang. 2019, *L. paucifasciatus* Rendahl in Smith, 1943, *L. rosozonatus* (Hu & Zhao, 1972), *L. rufozonatus* Cantor, 1842, *L. ruhstrati abditus* Vogel, David, Pauwels, Sumontha, Norval, Hendrix, Vu & Ziegler, 2009, *L. septentrionalis* (Günther, 1875), and *L. subcinctus* Boie, 1827 (Uetz et al. 2019, Luu et al. 2019).

Our recent field surveys in the Ha Lang and Trung Khanh districts, Cao Bang Province, northern Vietnam, revealed a snake population that was referable to the genus *Lycodon* based on the following characters: nostril enlarged; robustly arched upper maxillary bone with an inward curve in the anterior part; anterior and posterior maxillary teeth interrupted by a diastema; dorsal scales smooth or weakly keeled, in 17 rows anteriorly and at midbody, and posteriorly 15 rows (Lanza 1999; Grismer et al. 2014). However, the series of three individuals from Cao Bang were morphologically distinct from other named species. These morphological results were further corroborated by the analysis of a fragment of the mitochondrial cytochrome *b* gene, and so herein we describe the population from Cao Bang Province, northern Vietnam, as a new *Lycodon* species.

Materials and methods

Sampling

The field surveys were led by TQN in October 2011 and from April to May 2012. The collected specimens were euthanised with ethyl-acetate, fixed in approximately 85% ethanol for 10 hours, and subsequently transferred to 70% ethanol for permanent storage. Liver tissue samples were preserved separately in 95% ethanol. The specimens were deposited in the collections of the Institute of Ecology and Biological Resources (IEBR), Hanoi, Vietnam and of the Zoologisches Forschungsmuseum Alexander Koenig (ZFMK), Bonn, Germany.

Morphological analysis

Identification of sex was performed by dissection (inspection of gonads and presence of hemipenes). Maxillary teeth were counted by dissecting the right maxilla for teeth / sockets. Scalation and maxillary teeth number were examined with a binocular dissecting microscope. Measurements were taken following Ziegler et al. (2018) with a measuring tape to the nearest 1 mm.

Abbreviations of morphological characters are as follows:

SVL	Snout-vent length (from tip of snout to vent);
TaL	tail length;
TaL / TL	ratio of tail length / total length;
TL	total length;
DSR	dorsal scale rows number at one head length posterior to the head – number of dorsal scale rows at midbody – number of dorsal scale rows at one head length anterior to the vent;
SL	supralabials (counted on upper lips);
SL / orbit	number of supralabials entering orbit;
IL	infralabials (counted on lower lips);
Lor	loreals;
Lor / eye	loreal scale touching the eye (yes or no);
PreOc	preoculars;
PostOc	postoculars;
Atem	number of anterior temporals;
PTem	number of posterior temporals;
BodySc	scalation of the body (keeled or smooth);
PreVen	number of preventral scales;
Ven	number of ventral scales;
SubC	number of subcaudal scales;
Prec	precloacal (or cloacal) plate (single or divided);
Teeth max	number of maxillary teeth / alveoli.

Scale counts were taken following Vogel et al. (2009). Ventral scales (Ven) were counted according to Dowling (1951). Bilateral scale counts were given as left / right.

Comparisons were mainly based on the data provided by Boulenger (1893), Pope (1928), Smith (1943), Leviton (1965), Ota and Ross (1994), Manthey and Grossmann (1997), Lanza (1999), Vogel et al. (2009), Vogel and David (2010) and Neang et al. (2014), with additional references provided in the comparisons and legends of the tables. Additionally, studied specimens are listed in the Appendix 1.

Molecular data and phylogenetic analyses

Representative taxa of the genus *Lycodon* were included in the study. Sequences of the species were downloaded from GenBank. Two samples of the population from Cao Bang Province (ZFMK 93746, ZFMK 93747) were incorporated in the analysis. *Boiga cynodon* (Boie 1827) and *Dipsadoboa flavigularis broadleyi* (Broadley & Stevens, 1971) were used as outgroups based on Figueiroa et al. (2016).

DNA was extracted using DNeasy Blood and Tissue kit (Qiagen, Germany) following the manufacturer's instructions. A fragment of the mitochondrial cytochrome *b* gene was amplified using the primer pair L14910 (5'-GACCTGTGATMTGAAAACCAY-CGTTGT-3') and H16064 (5'-CTTGGTTACAAGAACAAATGCTTA-3'; Burbrink et al. 2000). Extracted DNA was amplified using HotStarTaq Mastermix (Qiagen, Germany) with 21 µl volume consisting of 10 µl of Mastermix, 5 µl of water, 2 µl of each primer at 10 pmol/ml and 2 µl of DNA. PCR conditions were 95 °C for 15 minutes to activate the Taq, with 40 cycles of 95 °C for 30 s, 45 °C for 45 s, 72 °C for 60 s, and a final extension at 72 °C for six minutes. The mitochondrial cytochrome *b* gene was utilised in this study because it has been widely used in previous molecular analyses of *Lycodon* (e.g., Guo et al. 2013, Siler et al. 2013), and has been shown to be informative in revealing new species of *Lycodon* (e.g., Grismer et al. 2014, Luu et al. 2018, 2019).

PCR products were visualised using gel electrophoresis through a 2% low melting-point agarose gel stained with ethidium bromide. Successful amplifications were purified to eliminate PCR components using GeneJET™ PCR Purification kit (ThermoFisher Scientific, Lithuania). Purified PCR products were sent to FirstBase (Malaysia) for sequencing.

The obtained sequences were aligned in ClustalX 1.8.3 (Thompson et al. 1997) using the default settings. Data were analysed using maximum parsimony (MP) as implemented in PAUP*4.0b10 (Swofford 2001) and Bayesian inference (BI) as implemented in MrBayes v3.2 (Ronquist et al. 2012). Settings for these analyses followed Le et al. (2006), except that the number of generations in the Bayesian analysis was increased to 1×10^7 . For the maximum likelihood (ML) analysis, we used IQ-TREE v.1.6.7.1 (Nguyen et al. 2015) with a single model and 10,000 ultrafast bootstrap replications. For ML and BI, the optimal model for nucleotide evolution was set to TrN+I+G by Modeltest v3.7 (Posada and Crandall 1998). For BI, the analysis was conducted with

a random starting tree and run for 10^7 generations. Four Markov chains, one cold and three heated (utilising default heating values), were sampled every 1000 generations. Log-likelihood scores of sample points were plotted against generation time to detect stationarity of the Markov chains. The burn-in value was set to 26 in the BI analysis, as $-\ln L$ scores reached stationarity after 26,000 generations in both runs. Two independent analyses were run simultaneously. Nodal support was evaluated using Bootstrap replication (BP) as estimated in PAUP*4.0b10 and IQ-TREE v1.6.7.1 and posterior probability (PP) in MrBayes v3.2. BP ≥ 70 and PP $\geq 95\%$ are regarded as strong support for a clade. Uncorrected pairwise distances (p) were calculated in PAUP*4.0b10.

Results

Molecular data and phylogenetic analyses

The final matrix consisted of 1011 bp aligned characters and the alignment contained no gaps. In total, 404 characters were found to be parsimony informative. MP analysis resulted in five most parsimonious trees having 1662 steps (CI = 0.41, RI = 0.72). Our tree topologies are very similar to those recovered by Guo et al. (2015) and Luu et al. (2018). The new species was recovered to be the sister species to a clade containing *L. multizonatus* + *L. liuchengchaoi*, with strong support in BI (PP = 96), but weak support in MP and ML (BP_{MP} = 56, BP_{ML} = 69) (Fig. 1). The new species has an uncorrected p -distance of at least 7.5% and 8.1% from *Lycodon liuchengchaoi* Zhang, Jang, Vogel & Rao, 2011 and *L. multizonatus* Zhao & Jiang, 1981, respectively.

Lycodon pictus sp. nov.

<http://zoobank.org/FEA7DFD1-BF41-4608-A477-93861FF13AD4>

Figs 2–6

Holotype. IEBR 4166 (field number CB 2012.97), adult male, collected on 18 April 2012 by TQN et al. (altitude 701 m a.s.l.), Trung Khanh District, Cao Bang Province.

Paratypes. ZFMK 93747, juvenile, collected on 15 October 2011 by TQN et al. (altitude 588 m a.s.l.), Ha Lang District, Cao Bang Province; ZFMK 93746, adult female, collected on 10 April 2012 by TQN et al., Ha Lang District, Cao Bang Province.

Diagnosis. *Lycodon pictus* sp. nov. can be differentiated from its congeners by the following morphological characters: dorsal scales in 17–17–15 rows, all smooth; supralabials usually eight (rarely nine); infralabials ten; one elongated loreal on each side, in contact with the eye; precloacal plate single; ventral scales 212–218 (plus one or two preventral scales); subcaudals 90 or 91; a total length of 597+ mm in males and 543 mm in females; tail / total length ratio 0.211–0.215; maxillary teeth 13 or 14; dorsal surface of body with 28 or 29 light body bands; dorsal surface of tail with 13 cream




Figure 1. Bayesian cladogram based on the partial mitochondrial cytochrome *b* gene of snakes belonging to the genus *Lycodon*. Numbers above and below branches are bootstrap values of MP/ML analyses ($\geq 50\%$) and Bayesian posterior probabilities (BC), respectively.

bands forming a distinct blotch in the vertebral region; ventral surface of body and tail mostly cream with the dark body bands in part extending towards the venter, sometimes forming complete dark bands around the body.

Description of the holotype. Head elongate, moderately distinct from the neck, rather flattened, longer than wide, narrow anteriorly; nostril lateral, located in the middle of the nasal; eye large, pupils vertically elliptic; rostral triangular, much broader than high, hardly visible from above; nasal divided into two scales by a vertical ridge along posterior edge of nostril; two internasals, anteriorly rounded, slightly wider than high, bordered by two large, pentagonal prefrontals posteriorly; frontal single, enlarged, pentagonal to hexagonal, narrowed posteriorly; parietals longer than wide, in contact with each other medially, with upper anterior and posterior temporals, paraparietal laterally and four nuchal scales posteriorly; paraparietals elongated, anterior part widened; loreal 1/1, elongate, not entering orbit; supralabials 8/8, first and second in contact with nasal, third to fifth entering orbit, sixth largest; infralabials 10/10, first pair in broad contact with each other, first to fifth in contact with anterior pair of chin shields; anterior and posterior pairs of chin shields elongate, of the same size and shape, second pair not meeting in midline; preocular 1/1; postoculars 2/2, lowermost smaller,



Figure 2. Holotype of *Lycodon pictus* sp. nov. (IEBR 4166) in life. Photograph T. Lehmann.

bordering anterior temporals; anterior temporals 2/2, posterior temporals 3/3, upper ones thinner than lower ones. Left maxilla arched, with an angular apex, distinctly bent inwards anteriorly. A total of 13 maxillary teeth or teeth alveola, with the following formula: five small anterior teeth, with the last two ones being somewhat enlarged + two strongly enlarged teeth, thick, and not much curved + a wide gap, somewhat wider than the length of the largest teeth + four small teeth + a small gap + two enlarged posterior teeth.

Body elongate, SVL 488 mm; TaL > 109 mm (tail tip lost); preventral 1, ventrals 212, from behind neck region distinctly notched laterally; subcaudals > 54 (tail tip lost), paired; precloacal plate single; DSR 17-17-15, all smooth; the vertebral scales not enlarged; DSR reduction from 17 to 15 at the position of ventral 150.

Coloration in preservative. Head, neck, and dorsal surface of body brownish black; light body bands beginning after 1.5 times the head length behind the head, in total 29 transverse light bands on body and at least nine light bands on tail; the first four body bands yellowish cream, and distinctly widened towards the venter, increased in size posteriorly; a dark mottling in the vertebrate region more prominent posteriorly; the subsequent light body bands with two distinct indentations on each side, fused in the middle in the last third of the body. In dorsal view the light bands forming a distinct blotch in the vertebral region, with a dark centre and a lighter frame; laterally, the middle part of the light bands forming blotches, but wider and with an extended



Figure 3. Head views of *Lycodon pictus* sp. nov. (IEBR 4166) in preservative (scale bar refers to head in dorsal view). Photographs T. Ziegler (upper), C.T. Pham (lower).

dark centre, fused laterally in the last third part of the body; the lower and widest part of the light body bands with a dark small blotch in the centre in the anterior part of body; the light bands on the tail with a blotch like pattern in the vertebral region, but



Figure 4. Female paratype of *Lycodon pictus* sp. nov. (ZFMK 93746) in life. Photographs T.Q. Nguyen.

less pronounced than that on body, and one light blotch at the lateral side of tail, widened towards the venter, with a dark centre; ventral surface of head and neck yellowish cream, belly cream and greyish cream in the last third part of body and on lower tail



Figure 5. Juvenile paratype of *Lycodon pictus* sp. nov. (ZFMK 93747) in preservative (upper, dorsal view; lower, ventral view). Photographs T. Ziegler.

surface; the dark dorsal bands (28 on body and at least nine on tail) in part extending towards the venter (most prominent in the anterior five dark body bands), not forming complete dark bands around the body, but complete on the tail; lateral side of the head dark above and light below, with the lighter pattern beginning in the supralabial region; tip of lower jaw and infralabial region in part greyish; dorsal surface of the head and upper head sides a bit paler than the remaining head dorsum.

Hemipenis. Hemipenes elongated, not fully everted, not turgid. Truncus without spines. Spine ornamentation starting at truncus region with somewhat enlarged, medium sized spines. Apex with microspines. Sulcus stretches in the middle to apex. Apex not fully everted, ending somewhat widened with an oblique opening, with microspines inside, pointing to the not fully everted condition of the outer genital organ.

Variations. In the juvenile ZFMK 93747, the number of supralabials on the left side is nine, with fourth to sixth entering the orbit. The loreal does not touch the eye on the right side. The lower anterior temporal scale is not touching the postocular scale on the left side. In general, the coloration is more intense in the juvenile. The creamy pattern on the posterior third of the body sides is connected by a horizontal cream-colored stripe. It has a yellowish cream band on the head that reaches from SL 5 behind the jaws and distinctly lightens the posterior half of the head but does not touch the frontal. In the juvenile, the banded pattern is more simple, consisting of dark bands



Figure 6. Dorsal head and neck pattern of the juvenile paratype of *Lycodon pictus* sp. nov. (ZFMK 93747) in preservative. Photograph T. Ziegler.

which narrow towards the venter and light bands which widen towards the venter and bear a dark pattern and a more or less distinct dark blotch at the lower side (see Fig. 5).

In the female ZFMK 93746, the lower anterior temporal scale is not touching the postocular scale on the right side. For measurements and scalation data of the examined specimens see Table 1.

Dentition. Female ZFMK 93746 and juvenile ZFMK 93747: Left maxilla arched, with an angular apex, distinctly bent inwards anteriorly. A total of 13 (in female) or 14 (in juvenile) maxillary teeth or teeth alveola, with the following formula: five small anterior teeth, with the last two ones being somewhat enlarged + two strongly enlarged teeth, thick, and not much curved + a wide gap, somewhat wider than the length of the largest teeth + four small teeth + a small gap + two enlarged posterior teeth in the female and three posterior teeth in the juvenile, with the anterior two ones enlarged.

Comparisons. In our phylogenetic analysis, *Lycodon pictus* sp. nov. is most closely related to *L. liuchengchaoi* and *L. multizonatus*. From *L. liuchengchaoi*, the new species differs in terms of body scalation (all smooth in the new species vs. feebly keeled in several median rows in *L. liuchengchaoi*), head scalation (ten infralabials vs. 7–9) and dentition (13 or 14 maxillary teeth vs. 8 or 9). In addition, the new species differs from the latter in having 28 or 29 cream body bands (vs. 40 yellow rings on the body in *L. liuchengchaoi*) (Zhang et al. 2015).

Table 1. Sex, measurements (in mm), scalation data, and coloration pattern of *Lycodon pictus* sp. nov. For abbreviations see Materials and methods. Key: asterisk (*) lower Atem not touching PostOc; plus sign (+) tail tip lost.

	Holotype IEBR 4166	Paratype ZFMK 93746	Paratype ZFMK 93747
Sex	male	female	juvenile
TL	597	543	237
SVL	488	426	187
TaL	109+	117	50
TaL/TL	—	0.215	0.211
Teeth max	13	13	14
SL	8/8	8/8	9/8
SL/orbit	3–5	3–5	4–6/3–5
IL	10/10	10/10	10/10
PreOc	1/1	1/1	1/1
PostOc	2/2	2/2	2/2
Lor	1/1	1/1	1/1
Atem	2/2	2'2	2/2'
PTem	3/3	3/3	3/3
DSR	17-17-15	17-17-15	17-17-15
PreVen	1	2	1
Ven	212	216	218
Prec	single	single	single
Subc	54+	91	90
BodySc	smooth	smooth	smooth
Dark bands on body	28	29	28
Light bands on body	29	29	28
Dark bands on tail	9	13	13
Light bands on tail	9+	13	13

The new species differs from *L. multizonatus* by having more maxillary teeth (13 or 14 vs. 10 or 11 in *L. multizonatus*), more infralabials (10 vs. 8) and a single precloacal plate (vs. divided). In addition, the new species differs from the latter in terms of body scalation (minimum 212 ventrals and minimum 90 subcaudals vs. 190–195 ventrals and 68–75 subcaudals in *L. multizonatus*). Furthermore, *L. pictus* sp. nov. has fewer light body bands (28 or 29 vs. 55–73 in *L. multizonatus*) (Lei et al. 2014).

From its Vietnamese congeners, the new species can be differentiated as follows: *Lycodon pictus* sp. nov. differs from *L. capucinus* in having a single precloacal plate (vs. divided), a loreal touching the eye (vs. not in contact with the eye), in having more ventrals (minimum 212 vs. 182–211) and more subcaudals (90 or 91 vs. 59–74), and in terms of dorsal pattern (banded vs. reticulated) (Luu et al. 2019).

Lycodon pictus sp. nov. differs from *L. cardamomensis* in terms of dorsal scalation (17-17-15 smooth DSR vs. 19-17-15 weakly keeled DSR), in having a loreal in contact with the eye (vs. separated) and in having 28 or 29 light body bands (vs. 12–14 pinkish orange body bands) (Daltry and Wüster 2002, Do et al. 2017).

Lycodon pictus sp. nov. differs from *L. davisonii* in having 17 midbody dorsal scale rows (vs. 13 midbody dorsal scale rows), fewer ventral scales (maximum 218 vs. 235–265), more infralabials (10 vs. 8) and the absence of preocular (vs. present). In addition, the new species differs from the latter in having a different dorsal pattern

(28 or 29 cream bands on body vs. 36 white rings on the body) (Blanford 1878, Boulenger 1893).

Lycodon pictus sp. nov. differs from *L. fasciatus* in having smooth dorsal scales (vs. keeled) and more maxillary teeth (13 or 14 vs. 11). Additionally, the colour pattern of *Lycodon pictus* sp. nov. differs in being dark brownish black with light body bands turning into a marbling posteriorly, whereas *L. fasciatus* is black or purplish black above with yellowish cross-bars of irregular outline and has a dark median stippling (Pope 1928, Smith 1943). Werner (1922) described *Dinodon yunnanensis* from Yunnan Fu, now Kunming, Yunnan Province, southwestern China. This species was synonymized with *Lycodon fasciatus* by Pope (1935: 188), but according to Vogel and David (2010), this taxon might be a distinct species (see also Vogel and David 2019). *Lycodon pictus* sp. nov. differs from *Dinodon yunnanensis* Werner, 1922 in having more ventrals (minimum 212 vs. 193), more subcaudals (90 or 91 vs. 66), more infralabials (10 vs. 9) and more light body bands (28 or 29 vs. 23) (Werner 1922, Vogel and David 2010, Vogel and David 2019).

Lycodon pictus sp. nov. differs from *L. flavozonatus* in terms of dorsal scalation (smooth vs. keeled), in having more subcaudals (90 or 91 vs. 80–88), the loreal in contact with the eye in *Lycodon pictus* sp. nov. (vs. separated in *L. flavozonatus*) and in coloration pattern (brownish black with 28 or 29 cream body bands and 9–13 light bands on the tail vs. black with 68 yellow body bands and 21 on the tail) (Pope 1928, Vogt in Pope 1928).

Lycodon pictus sp. nov. differs from *L. futsingensis* in having more ventrals (minimum 212 vs. 193–208) and more subcaudals (minimum 90 vs. 72–87). Additionally, the loreal does not enter the orbit in *L. futsingensis*, whereas it enters the orbit in *Lycodon pictus* sp. nov. (Vogel et al. 2009).

Lycodon pictus sp. nov. differs from *L. laoensis* in having a single precloacal plate (vs. divided), more ventral scales (minimum 212 vs. 163–192), more subcaudal scales (minimum 90 vs. 60–76), an elongated loreal scale in contact with the orbit (vs. separated) and cream body bands (vs. yellow) (Grismer et al. 2014, Neang et al. 2014).

Lycodon pictus sp. nov. differs from *L. meridionalis* in having smooth dorsals (vs. feebly keeled in 10–12 median rows), a lower ventral scale count (maximum 218 vs. 227–240) and fewer subcaudals (maximum 91 vs. 96–106). In addition, the new species differs in having cream body bands (vs. yellow thin crossbars) (Gawor et al. 2016 and examined ZFMK specimens, see Appendix 1).

Lycodon pictus sp. nov. differs from *L. namdongensis* in having more subcaudals (90 or 91 vs. 85) and the loreal in contact with the eye (vs. separated from the eye in *L. namdongensis*). The new species also differs in coloration pattern (brownish black with 28 or 29 light bands on the body vs. grey with 23 cream cross rings on the body in *L. namdongensis*), and in having irregular bands turning into a marbling posteriorly (vs. clearly demarcated cross bands on the body) (Luu et al. 2019).

Lycodon pictus sp. nov. differs from *L. paucifasciatus* in terms of dorsal scalation (17–17–15 smooth DSR vs. 19–(19–17)–15 DSR, the upper one or two plus vertebral

row distinctly keeled) and fewer ventral scales (maximum 218 vs. 219–222). In addition, the new species has a loreal entering the eye (vs. separated) and 28 or 29 light body bands (vs. 14–25 beige or dirty cream body bands) (Vogel et al. 2009).

Lycodon pictus sp. nov. differs from *L. rosozonatus* in having 17–17–15 smooth DSR (vs. 19–19–15(17) keeled DSR), fewer ventral scales (maximum 218 vs. 221–234) and a loreal in contact with the eye (vs. separated). In addition, the new species has cream body bands (vs. pinkish red) (Hu et al. 1975, Neang et al. 2014).

Lycodon pictus sp. nov. differs from *L. ruhstrati abditus* in having smooth dorsals (vs. 7–8 dorsal scale (including vertebral) rows keeled), an elongated loreal in contact with the eye (vs. separated), smooth dorsal scales (vs. feebly keeled in the posterior body part), and in coloration pattern (28 or 29 cream body bands vs. 44–52 light red body bands) (Zhao 2006, Luu et al. 2018).

Lycodon pictus sp. nov. differs from *L. ruhstrati abditus* in having smooth dorsals (vs. 7–8 dorsal scale (including vertebral) rows keeled), an elongated loreal in contact with the eye (vs. separated), and in having irregular bands turning into a marbling posteriorly (vs. clearly demarcated cross bands on the body) (Vogel et al. 2009).

Lycodon pictus sp. nov. differs from *L. septentrionalis* by its smooth dorsal scales (vs. 7–9 median rows feebly keeled), 10 infralabials (vs. 7 or 8), and the loreal entering the orbit (vs. separated in *L. septentrionalis*). In addition, the new species differs in having cream irregular bands on a brown body (vs. white narrow bands on a black body forming complete annuli) (Günther 1875, Boulenger 1893, Neang et al. 2014).

Lycodon pictus sp. nov. differs from *L. subcinctus* in having 10 infralabials (vs. 8 or 9), one preocular (vs. preocular absent), smooth dorsal scales (vs. feebly keeled) and 28 or 29 cream bands on the body and 9–13 on the tail (vs. 9–15 bands on the body and none on the tail) (Boulenger 1893, Neang et al. 2014).

Lycodon pictus sp. nov. differs from *L. ophiophagus*, a species from southern Thailand but with similar scalation, in having a loreal entering the eye (vs. separated) and in dorsal colour pattern (28 or 29 light bands on a brown body vs. 20 or 21 white bands on a dark body) as well as and in having irregular bands turning into a marbling posteriorly (vs. clearly demarcated cross bands on the body) (Vogel et al. 2009).

For additional measurements, dentition, and scalation data see Tables 2–8.

Distribution. *Lycodon pictus* sp. nov. is currently known only from Ha Lang and Trung Khanh districts, Cao Bang Province, northern Vietnam (Fig. 7).

Etymology. The name of the species *pictus* means painted or decorated in Latin and refers to its unique dorsal colour pattern.

Natural history. *Lycodon pictus* sp. nov. seems to be closely associated with karst environment. Specimens were found at night between 19:00 and 23:00, on forest paths or on the ground near cave entrances. The surrounding habitat was secondary karst forest, consisting of medium and small hardwood trees mixed with shrubs and vines. Air temperature was 23.4–29.6°C and humidity was 66–79%. Other reptiles were also found at the site, including *Acanthosaura lepidogaster* (Cuvier, 1829), *Gekko adleri* Nguyen, Wang, Yang, Lehmann, Le, Ziegler & Bonkowski, 2013, *Goniurosaurus lului* Grismer, Viets & Boyle, 1999, *Lycodon futsingensis* (Pope, 1928), and *Protobothrops trungkhanhensis* Orlov, Ryabov & Nguyen, 2009 (Fig. 8).



Figure 7. Map showing the type locality of *Lycodon pictus* sp. nov. in Cao Bang Province.



Figure 8. Habitat of *Lycodon pictus* sp. nov.: the female paratype (ZFMK 93746) was found in Ha Lang District, Cao Bang Province.

Table 2. Measurements (in mm), dentition, and scalation data of *Lycodon* species from Vietnam. Data taken from Blanford (1878), Pope (1928), Smith (1943), Boulenger (1893), Dalmat and Wüster (2002), Jackson and Fritts (2004), Vogel et al. (2009), Neang et al. (2014), Do et al. (2016), and Luu et al. (2019); distinguishing characters are marked in bold. For abbreviations see Materials and methods.

	<i>Lycodon pictus</i> sp. nov.	<i>L. capricornis</i>	<i>L. cardamomensis</i>	<i>L. davisonii</i>	<i>L. fasciatus</i>	<i>L. flavozonatus</i>	<i>L. fusingensis</i>	<i>L. laoensis</i>
TL	597	816	896	920 ♂	894	1440 ♂, 1210 ♀	850 ♂, 773 ♀	470
Teeth max	13 or 14	15	10-12	8	11	13	12-15	
SL	8	9 or 10	8	7	8	8	8 (7)	9 or 10
SL/orbit	3-5	3-5	3-4	3-5	3-5	3-5 (4-5; 2-4; 4-6)	3-5	3-5
IL	10	9 or 10	10	8	9 (8, 10)	10	10 (9, 11)	10
PreOc	1	1	1	0	1	1	1	1
PostOc	2	2	2 or 3	1 or 2	2	2	2 (3)	2 (3)
Lor	1	1	1	1	1	1	1	1
Lor/eye	yes	no	yes	yes	no	no	no	
Atem	2	2	2	1 or 2	2	2	2 (1)	2
pTem	3	3	2 or 3	2	2	2 or 3	3 (2)	3
DSR	17-17-15	17-17-15	19-17-15	13	17-17-15	17-17-15	17-17 (16)-15	17-17-15
Ven	212-218	182-211	215-228	233-265	182-225	211-221 ♂, 212-218 ♀	193-204 ♂, 198-208 ♀	163-192
Prec	single	divided	single	single	single	divided/ single	single	divided
Subc	59-74	87-93	90-108	65-94	81-88 ♂, 80-84 ♀	72-87 ♂, 78-85 ♀	60-76	smooth
BodySc	smooth	weakly keeled	weakly keeled	smooth	keeled	7 rows feebly keeled at midbody	smooth	smooth

Table 3. Measurements (in mm), dentition, and scalation data of *Lycodon* species from Vietnam (continuation of Table 2). Data from Günther (1875), Boulenger (1893), Hu et al. (1975), Zhao (2006), Vogel et al. (2009), Guo et al. (2013), Neang et al. (2014), Gavoty et al. (2016), Luu et al. (2018), Luu et al. (2019), and based on examined specimens from ZFMK (see Appendix 1). Key distinguishing characters are marked in bold; * Luu et al. (2019) mention 240 ventrals as maximum for *L. rufozonatus*; however, that is a transcription error from Bourret (1935) who gave 204 as the maximum number of ventrals. For abbreviations see Materials and methods.

	<i>L. meridionalis</i>	<i>L. nandongensis</i>	<i>L. paucifasciatus</i>	<i>L. rufozonatus</i>	<i>L. rubrivittatus</i>	<i>L. rubriflavus</i>	<i>L. septentrionalis</i>	<i>L. subcinctus</i>
TL	1139	723	763	1060	1234	964	1163	1000
Teeth max	11	12	11 or 12	12 or 13	11–13	11–13	7	8–14
SL	8	8	8	8	8	8	8	8
SL/orbit	3–5	3–5	3–5	3–5	3–5	3–5	3–5	3–5 (3–6)
IL	10	10	10	10 (9)	10 (9,11)	7 or 8	7 or 8	8 or 9
PreOc	1	2/1	1	1	1	1	1	0
PostOc	2	3	2	2	2	2	2	2 or 3
Lor	1	1	1	1	1	1	1	1
Lor/eye	no	no	no	no	no	no	no	yes
Atem	2	2	2	2	2	2 (1)	2	1
PTem	3	2	3	3	3	3 (2)	3	2
DSR	17–17–15	17–17–15	19–(19–17)–15	19–19–15(17)	(19–17)–17–15	17–17–15	17–17–15	17–17–15
Ven	227–240	218	219–222	221–234	184–225*	197–229	202–224	192–230
Prec	single	single	single	single	single	single	single	divided (rarely single)
Subc	96–106	85	90–92	53–98	90–103	83–104	83–104	60–91
BodySc	dorsals feebly keeled in 10–12 median rows, smooth in outer rows	smooth	upper 1 or 2 dorsal scale row(s) plus vertebral row distinctly keeled	weakly keeled in the posterior body part	feebly keeled in the posterior body part	smooth (DSR 1–6), distinctly keeled (DSR 7–8 and vertebral row)	7 or 9 median rows feebly keeled	feebly keeled

Table 4. Measurements (in mm), dentition and scalation data of *Lycodon* species from China, Laos and Cambodia. Data taken from Zhao and Jiang (1981), Bouenger (1893), Vogel and David (2010) Vogel and Luu (2011), Zhang et al. (2011), Lei et al. (2012), Lei et al. (2014), Neang et al. (2015), Ganesh and Vogel (2018), and Luu et al. (2018). Key: plus sign (+) tail incomplete; distinguishing characters are marked in bold. For abbreviations see Materials and methods.

	<i>L. aulicus</i>	<i>L. banksi</i>	<i>L. davidi</i>	<i>L. gongshan</i>	<i>L. liuchengchaoi</i>	<i>L. multizonatus</i>	<i>L. symapior</i>	<i>L. zoosvictoriae</i>
TL	719	465 +	389.5	963 ♂	676	505	487	520.7
Teeth max			11		8 or 9	10 or 11	10	9
SL	9 (8 or 10)	8	8	7 or 8	8 (rarely 7)	8	8	8
SL/orbit	3–5	3–5	3–5	3–5	3–5	3–5	3–5	3–5/4–5
IL	10 or 11	10	10	8	8 (7,9)	8 (7)	8	10
PreOc	1	1	1	1	1	0 or 1	1	1 or 2
PostOc	2	2	2	2	2	2	2	2
Lor	1	1	1	1	1	1	1	1
Lor/eye	no	yes	no	yes	yes	yes	no	no
Atem	2	2	2	2	1–3	2 (1)	2	2
PTem	3	3	2 or 3	2 or 3	1–3	3 (2)	2	2
DSR	17–17.15	17–17.15	17–17–15	17–17–15	17–17–15	15 or 17–17–15	17–17–15	17–17–15
Ven	180–215	241	224 ♂	210–216 ♂, 215 ♀	190–228	190–195	201–203 ♀	213 ♀
Prec	single	single	single	single	divided	divided	single	single
Subc	57–78	26 +	99 ♂	95 or 96 ♂, 92 ♀	68–77	68–75	68 or 69 ♀	85 ♀
BodySc	smooth and glossy	smooth (six central DSR of posterior 1/3 feebly keeled)	middorsal scale rows slightly keeled, outermost rows entirely smooth	upper dorsal rows 6–12 and vertebral row keeled	feebly keeled in median rows	smooth	6–7 upper rows and vertebral row feebly keeled	weakly keeled

Table 5. Measurements (in mm), dentition, and scalation data of *Lycodon* species from Thailand and Myanmar. Data taken from Günther (1864), Boulenger (1893), Boulenger (1900), Smith (1943), Lanza (1999), Slowinski et al. (2001), Daltry and Wüster (2002), Vogel et al. (2009), Bahuguna and Bhata (2010), Grismer et al. (2014), Luu et al. (2018), and Vogel and David (2019); distinguishing characters are marked in bold. For abbreviations see Materials and methods.

	<i>L. albofuscus</i>	<i>L. butleri</i>	<i>L. effraensis</i>	<i>L. gibsonae</i>	<i>L. gracilis</i>	<i>L. jara</i>	<i>L. kundui</i>	<i>L. ophiophagus</i>	<i>L. subannulatus</i>	<i>L. zawi</i>
TL	1480	876	700	906	533	535 ♂, 550 ♀	909	684	480	
Teeth max	12			13	9		11-13	8 or 10	12	
SL	8	8 or 9	9	8	8	8 or 9	7	8	7	8 or 9
SL/orbit	3-5	3-5	3-5	3-4 and 3-5	3-4	3-5	3-4	3-5	3-4	3-5
IL	9 or 10	10 or 11	10				10	8		9 or 10
PreOc	1	1	1	1	2	1		1	1	1
PostOc	2	2	2-3	2	2	2	2	2	2	2 (1)
Lor	1	1	0	1	1 (united with lower PreOc)	1	1	1	1	1
Lor/eye	no	yes	no Lor	yes	yes	no	no	no	yes	no
Atem	2	2	2	2	2	1 or 2	1	2	2	2 (3)
pTem	2	2	2 or 3	3	3	2 or 3	2	3	2	3 (4)
DSR	17	19 (17 in Boulenger 1900)	17	17-17-15	15	17-17-15	15-15-15	17-17-15	15-15-15	17-17-15
Ven	241	220-227	215-228	223-226 ♂	234	167-188	186	211-212	225-244	179-194 ♂, 207 ♀
Prec			single	single	single	divided	divided (entire in Lanza 1990)	single	single	divided
Subc	155-208	81-96	72-99	91-92 ♂	81-83	52-74	70	87-90	93-111	45-75 ♂
BodySc	keeled	keeled	smooth	upper 3 or 4 rows keeled	keeled	smooth	smooth	keeled	smooth	smooth

Table 6. Measurements (in mm), dentition and scalation data of *Lycodon* species from India. Data taken from Boulenger (1893), Wall (1906), Smith (1943), Taylor (1950), Captain (1999), Vijayakumar and David (2005), Mukherjee and Bhupathy (2007), Misri et al. (2007), Vogel and Lou (2011), Vogel and Harikrishnan (2013), Ganesh and Vogel (2018), and Melvinselvan et al. (2018); distinguishing characters are marked in bold. For abbreviations see Materials and methods.

	<i>L. anamallensis</i>	<i>L. flavicollis</i>	<i>L. flavomaculatus</i>	<i>L. gammiei</i>	<i>L. hypsinthoides</i>	<i>L. macconnelli</i>	<i>L. nymphpha</i>	<i>L. siamensis</i>	<i>L. tenuaria</i>	<i>L. travancoricus</i>
TL	522	543	520	1150	717♂, 563♀	365	574	432	790	600♂, 625♀
Teeth max						8-10				
SL	9	9	9	8(7,9)	9	8(7)	7(8 or 6)	9		9
SL/orbit	3-5	3-5	3-5	3-5(3-4)	3-5	3-5	3-4	3-5		3-5
IL	10 or 11	11	10		10	8		11		
PreOc	1	1	1	1	1	1	1 or 2	1	1	
PostOc	2 or 3	2	2	2(1)	2	2	2	2	2	2
Lor	1/1 (except Holotype 2/2)	1	1	1	1	1 (0 in Wall 1906)	1	1	1	1
Lor/eye		no	no	no	no	no	yes	no	no	
Atem	2	2(3)	2(1)	2 or irregular	2	1 or 2	2	2 rarely 1		2 or 3
P'Tem	3+4	3 (rarely 2)	3 (rarely 2)	2 or irregular	3	2 or 3	2 or 3	3 rarely 2		3
DSR	17-17.15	17-17.15	17-17.15	17-17(19)-15	17-17.15	17-17.15	13-13.13	17-17.15	?	17-17.15
Ven	174-186♂, 186-204♀	210-224	165-183	205-220	188-202♂, 199-210♀	163-187	200-243	153-178	218-237	176-206
Prec	divided (except holotype)	divided	divided	single	divided	divided	divided	divided	divided	single
Subc	63-73♂, 60-74♀	65-72	53-63	98-111	68-75♂, 61-68♀	48-56	65-88	42-66	61-102	64-76
BodySc	smooth	smooth with single apical pit	smooth	9 dorsal rows keeled, 5 rows at each side smooth	smooth	smooth	keeled	smooth	smooth	

Table 7. Measurements (in mm), dentition and scutation data of *Lycodon* species from the Philippines. Data taken from Griffin (1909), Taylor (1922), Leviton (1965), ORa and Ross (1994), Lanza (1999), Gaulke (2002), and Gaulke et al (2003); distinguishing characters are marked in bold. For abbreviations see Materials and methods.

Table 8. Measurements (in mm), dentition and scelation data of *Lycodon* species from Sri Lanka, Malaysia, Japan and Indonesia. Data taken from Boulenger (1893), Stejneger (1907), Smith (1943), Vogel et al. (2009), Grismer et al. (2014), and Wostl et al. (2017); distinguishing characters are marked in bold. For abbreviations see Materials and methods.

	<i>L. carinatus</i>	<i>L. c垣erniculus</i>	<i>L. multifasciatus</i>	<i>L. orientalis</i>	<i>L. semicarinatus</i>	<i>L. sidiki</i>	<i>L. stormi</i>	<i>L. tristrigatus</i>
TL	730	508.2	700	660	1100	715	597	360
Teeth max				10 or 11		7		8 or 10
SL	8 or 9	9 or 10		8	8	8	8	7
SL/orbit	3-5	4-6	3-5	3-5	3-5	3-5	3-4	3-4
IL		10 or 11			10/9			
PreOc	1	1	0		1	0	1	0
PostOc	2	2	2	2	2	2	2	2
Lor	1	1	1	1	1	1	1	1
Lor/eye	no	yes	no	yes	no	no	no	yes
Atem	2	3 (2)		2	2	2	1	2
PTem	2 or 3	3 or 4		3	3	2	3	2 or 3
DSR	17 or 19-19-17	17-17-15	17-17-?	17	17	17-17-15	19	15
Ven	185-202	245 ♂, 232 ♀	232-237 ♂, 229-235 ♀	200-208	211-234	195	217	224
Prec	single	single		divided	single	divided	single	single
Subc	51-64	113 ♂, 92 ♀	115-119 ♂, 106-117 ♀	68-74	65-105	85	75	86
BodySc	strongly keeled	the 8 medial rows weakly keeled	keeled	scales with a very faint keel along their anterior half	scales keeled along anterior half (4 outer rows smooth, other with a feeble though faint keel along their anterior half)	keeled	smooth	keeled

Discussion

Our phylogenetic analyses reveal *Lycodon pictus* sp. nov. to be the sister taxon to a clade containing *L. multizonatus* and *L. liuchengchaoi* from China, but only with strong statistical support in the BI. The new species differed from the latter by at least 7.5% in uncorrected pairwise sequence distance. There has been some taxonomic confusion in the genus *Lycodon*. Two of the *L. liuchengchaoi* sequences (KC733201, KC733202) in the phylogenetic tree had previously been identified as *L. fasciatus*, but the phylogenetic analysis by Guo et al. (2015) correctly assigned them to *L. liuchengchaoi*. *Lycodon "flavozonatus"*, on the other hand, appears to be paraphyletic with *Lycodon "meridionalis"* (MH669271, MH669268). Moreover, the *Lycodon subcinctus* species group is likely to contain cryptic diversity. In terms of uncorrected pairwise genetic distance of populations within this species group, two samples (GenBank numbers KX822581 and KX822582) are approximately 9.1–9.2% divergent from KC733203 and 6.3–6.5% from KC010384 and KC010385. The latter two clades differ by approximately 8.0% from each other. These issues need to be further investigated in future studies.

This new discovery increases the number of *Lycodon* known from Vietnam to 16, of which nine are confined to karst formations, underlining the importance of this habitat in promoting reptile speciation (Luu et al. 2018). Although Vietnam is located in the region with one of the most extensive limestone outcrops in the world (Day and Urich 2000) many of the areas are still poorly surveyed, and likely contain a high level of cryptic diversity. Recent studies show that this habitat harbours a significant portion of endemic diversity in the region and should be protected from anthropogenic threats (Clement et al. 2006, Nguyen et al. 2015, Luu et al. 2016).

Acknowledgements

We are grateful to the directorates of the Forest Protection Department of Cao Bang Province, Forest Protection Units of Ha Lang and Trung Khanh districts for support of our field work and issuing relevant permits. Thanks to D.T. Le, H.T. An (Hanoi), S. Herbst, T. Lehmann, and M. Bonkowski (Cologne) for their assistance, E. Sterling (New York) and K. Koy (Berkeley) for providing the map. We thank J. Wang from the State Key Laboratory of Biocontrol / The Museum of Biology, School of Life Sciences, Sun Yat-sen University, Guangzhou for kindly providing the translation of the original description of *Lycodon multizonatus*. Many thanks also to B.L. Stuart (Raleigh) and T.T. Nguyen (Hanoi) for helping to improve a previous version of the manuscript. For the fruitful cooperation within joint research projects we cordially thank S.V. Nguyen (IEBR, Hanoi), as well as T. Pagel and C. Landsberg (Cologne Zoo). Cologne Zoo is partner of the World Association of Zoos and Aquariums (WAZA): Conservation Project 07011 (Herpetodiversity Research). This research is supported by the National Foundation for Science and Technology Development (NAFOSTED, Grant No. 106.05-2017.329). Field work in Cao Bang was partially supported by the Nagao Natural Environment Foundation (Japan).

References

- Anderson J (1879) Reptilia and amphibia. In: Anatomical and Zoological Researches: Comprising an Account of the Zoological Results of the Two Expeditions to Western Yunnan in 1868 and 1875. Bernard Quarich, London1: 705–860, 969–975. <https://doi.org/10.5962/bhl.title.55401> [index 1878]
- Bahuguna B, Bhuta PT (2010) Extension of the range and redescription of *Lycodon jara* (Shaw) (Reptilia: Serpentes: Colubridae). Records of the Zoological Survey of India 110(4): 37–39.
- Blanford WT (1878) Notes on some Reptilia from the Himalayas and Burma. Journal of the Asiatic Society of Bengal (2) 47: 125–131.
- Boie F (1827) Bemerkungen über Merrem's Versuch eines Systems der Amphibien, 1. Lief-erung: Ophidier. Isis van Oken 20: 508–566.
- Boulenger GA (1893) Catalogue of the Snakes in the British Museum (Natural History). Vol I. Trustees of the British Museum, London, 448 pp. [28 pls]
- Boulenger GA (1900) Description of a new snake from the Perak Hills. Journal of the Bombay Natural History Society 13: 336.
- Bourret R (1935) Notes herpétologiques sur l'Indochine française. IX. Les Serpents de Chapa. Bulletin général de l'Instruction publique 14(1934–1935): 7 (Mars): 239–251 [Reprint: 5–17].
- Burbrink FT, Lawson R, Slowinski JB (2000) Mitochondrial DNA phylogeography of the polytypic North American rat snake (*Elaphe obsoleta*): a critique of the subspecies concept. Evolution 54: 2107–2118. <https://doi.org/10.1111/j.0014-3820.2000.tb01253.x>
- Cantor T (1842) General features of Chusan, with remarks on the flora and fauna of that island [part 1]. Annals & Magazine of Natural History 9(1): 265–278. <https://doi.org/10.1080/03745484209445337>
- Captain A (1999) On the identification of *Lycodon flavomaculatus* Wall 1907. Journal of the Bombay Natural History Society 96: 323–327.
- Clements R, Sodhi NS, Schilthuizen M, Ng PKL (2006) Limestone karsts of Southeast Asia: imperilled arks of biodiversity. BioScience 56: 733–742. [https://doi.org/10.1641/0006-3568\(2006\)56\[733:LKOSAI\]2.0.CO;2](https://doi.org/10.1641/0006-3568(2006)56[733:LKOSAI]2.0.CO;2)
- Cuvier GJLNFD (1829) Le Règne Animal Distribué, d'après son Organisation, pour servir de base à l'Histoire naturelle des Animaux et d'introduction à l'Anatomie Comparé. Nouvelle Edition [second edition]. Vol. 2. Les Reptiles. Déterville, Paris, 1–406.
- Daltry JC, Wüster W (2002) A new species of wolf snake (Serpentes: Colubridae: *Lycodon*) from the Cardamom Mountains, southwestern Cambodia. Herpetologica 58(4): 498–504. [https://doi.org/10.1655/0018-0831\(2002\)058\[0498:ansows\]2.0.co;2](https://doi.org/10.1655/0018-0831(2002)058[0498:ansows]2.0.co;2)
- Day M, Urich P (2000) An assessment of protected karst landscapes in Southeast Asia. Cave and Karst Science 27: 61–70.
- Do DT, Ngo CD, Ziegler T, Nguyen TQ (2017) First record of *Lycodon cardamomensis* Daltry & Wüster, 2002, (Squamata: Colubridae) from Vietnam. Russian Journal of Herpetology 24(2): 167–170.
- Dowling HG (1951) A proposed standard system of counting ventral in snakes. Journal of Herpetology 1: 97–99. <https://doi.org/10.2307/1437542>

- Figueroa A, McKelvy AD, Grismer LL, Bell CD, Lailvaux SP (2016) A species-level phylogeny of extant snakes with description of a new colubrid subfamily and genus. PLoS ONE 11: e0161070. <https://doi.org/10.1371/journal.pone.0161070>
- Ganesh SR, Vogel G (2018) Taxonomic reassessment of the Common Indian Wolf Snakes *Lycodon aulicus* (Linnaeus, 1758) complex (Squamata: Serpentes: Colubridae). Bonn Zoological Bulletin 67(1): 25–36.
- Gawor A, Pham CT, Nguyen TQ, Nguyen TT, Schmitz A, Ziegler T (2016) The herpetofauna of the Bai Tu Long National Park, northeastern Vietnam. Salamandra 52(1): 23–41.
- Gaulke M (2002) A new species of *Lycodon* from Panay Island, Philippines (Reptilia, Serpentes, Colubridae). Spixiana 25: 85–92.
- Gaulke M, Demegillo A, Reiter J, Tacud B (2003) Additions to the Herpetofauna of Panay Island, Philippines. Salamandra 39 (2) 111–122.
- Griffin LE (1909) A list of snakes found in Palawan. Philippine Journal of Science 4: 595–601.
- Grismer LL, Viets BE, Boyle LJ (1999) Two new continental species of *Goniurosaurus* (Squamata: Eublepharidae) with a phylogeny and evolutionary classification of the genus. Journal of Herpetology 33(3): 382–393. <https://doi.org/10.2307/1565635>
- Grismer LL, Quah ESH, Anuar S, Muin MA, Wood Jr PL, Aziza S (2014) A diminutive new species of cave-dwelling Wolf Snake (Colubridae: *Lycodon* Boie, 1826) from Peninsular Malaysia. Zootaxa 3815: 51–67. <https://doi.org/10.11646/zootaxa.3815.1.3>
- Guo P, Zhang L, Liu Q, Li C, Pyron RA, Jiang K, Burbrink FT (2013) *Lycodon* and *Dinodon*: one genus or two? Evidence from molecular phylogenetics and morphological comparisons. Molecular Phylogenetics and Evolution 68: 144–149. <https://doi.org/10.1016/j.ympev.2013.03.008>
- Guo P, Zhong G, Liu Q, Zhu F, Li F, Wang P, Xiao R, Fang M, Fu X (2015) Phylogenetic position of *Lycodon gongshan* Vogel & Luo, 2011, a snake endemic to China. Amphibia-Reptilia 36: 165–169. <https://doi.org/10.1163/15685381-00002985>
- Günther ACLG (1864) The reptiles of British India. Ray Society, London, 452 pp. [pls 1–26]
- Günther ACLG (1875) Second report on collection of Indian reptiles obtained by the British Museum. Proceedings of the Zoological Society of London 1875 (March): 224–234. [pls 30–34]
- Hu S, Djao [Zhao] E, Huang Z (1975) Three new species of reptiles from Hainan Island, Guangdong Province (in Chinese with English abstract). Acta Zoologica Sinica, Peking 21(4): 379–384. [English translation by Koshikawa A (1982) Smithsonian Herpetological Information Service, Washington 53: 1–9] <https://doi.org/10.5479/si.23317515.53.1>
- Jackson K, Fritts TH (2004) Dentitional specialisations for durophagy in the Common Wolf snake, *Lycodon aulicus capucinus*. Amphibia-Reptilia 25: 247–254. <https://doi.org/10.1163/1568538041975134>
- Lanza B (1999) A new species of *Lycodon* from the Philippines, with a key to the genus (Reptilia Serpentes Colubridae). Tropical Zoology 12: 89–104. <https://doi.org/10.1080/03946975.1999.10539380>
- Le M, Raxworthy CJ, McCord WP, Mertz L (2006) A molecular phylogeny of tortoises (Turtles: Testudinidae) based on mitochondrial and nuclear genes. Molecular Phylogenetics and Evolution 40: 517–531. <https://doi.org/10.1016/j.ympev.2006.03.003>

- Lei J, Sun X, Jiang K, Vogel G, Booth DT, Ding L (2014) Multilocus Phylogeny of *Lycodon* and the Taxonomic Revision of *Oligodon multizonatum*. Asian Herpetological Research 5: 26–37. <https://doi.org/10.3724/SP.J.1245.2014.00026>
- Leviton AE (1965) Contributions to a review of Philippine snakes, VIII The snakes of the genus *Lycodon* H. Boie. The Philippine Journal of Science 94(1): 117–140.
- Luu VQ, Bonkowski M, Nguyen TQ, Le MD, Schneider N, Ngo HT, Ziegler T (2016) Evolution in karst massifs: Cryptic diversity among bent-toed geckos along the Truong Son Range with descriptions of three new species and one new country record from Laos. Zootaxa 4107: 101–140. <http://doi.org/10.11646/zootaxa.4107.2.1>
- Luu VQ, Bonkowski M, Nguyen TQ, Le MD, Calame T, Ziegler T (2018) A new species of *Lycodon* Boie, 1826 (Serpentes: Colubridae) from central Laos. Revue suisse de Zoologie 125(2): 263–276. <https://doi.org/10.11646/zootaxa.4586.2.3>
- Luu VQ, Ziegler T, Ha NV, Le MD, Hoang TT (2019) A new species of *Lycodon* Boie, 1826 (Serpentes: Colubridae) from Thanh Hoa Province, Vietnam. Zootaxa 4586(2): 261–277. <https://doi.org/10.11646/zootaxa.4586.2.3>
- Maki M (1931) Monograph of the Snakes of Japan. Dai-ichi Shobo, Tokyo (1)7: 240 pp. [expanded edition 1933]
- Manthey U, Grossman W (1997) Amphibien & Reptilien Südostasiens. Natur und Tier-Verlag, Münster, 512 pp.
- Melvinselvan G, Nibedita D, Murali G (2018) Observations on the reproduction and feeding habit of a rare colubrid, Indian bridal Snake *Lycodon nympha* Daudin 1803 (Serpentes: Colubridae) from Southern India. Captive & Field Herpetology 2(1): 16–22.
- Mistry V, Vogel G, Tillack F (2007) Rediscovery of *Dinodon gammieei* (Blanford 1878) (Serpentes, Colubridae), with description of its validity. Hamadryad 31(2): 265–273.
- Mukherjee D, Bhupathy S (2007) A new species of wolf snake (Serpentes: Colubridae: *Lycodon*) from Anaikatti Hills, Western Ghats, Tamil Nadu, India. Russian Journal of Herpetology 14(1): 21–26.
- Müller L (1924) Über neue oder seltene Mittel- und Südamerikanische Amphibien und Reptilien. Mitteilungen aus dem Zoologischen Museum in Berlin 11(1): 75–93 [1923].
- Neang T, Hartmann T, Hun S, Souter NJ, Furey NM (2014) A new species of wolf snake (Colubridae: *Lycodon* Fitzinger, 1826) from Phnom Samkos Wildlife Sanctuary, Cardamom Mountains, southwest Cambodia. Zootaxa 3814(1): 068–080. <https://doi.org/10.11646/zootaxa.3814.1.3>
- Nguyen TQ, Le MD, Pham AV, Ngo HN, Hoang CV, Pham CT, Ziegler T (2015) Two new species of *Cyrtodactylus* (Squamata: Gekkonidae) from the karst forest of Hoa Binh Province, Vietnam. Zootaxa 3985: 375–390. <https://doi.org/10.11646/zootaxa.4162.2.4>
- Nguyen TQ, Wang YY, Yang JH, Lehmann T, Le MD, Ziegler T, Bonkowski M (2013) A new species of the *Gekko japonicus* group (Squamata: Sauria: Gekkonidae) from the border region between China and Vietnam. Zootaxa 3652(5): 501–518. <https://doi.org/10.11646/zootaxa.3895.1.4>
- Nguyen LT, Schmidt HA, von Haeseler A, Bui MQ (2015) IQ-TREE: A fast and effective stochastic algorithm for estimating maximum likelihood phylogenies. Molecular Biology and Evolution 32: 268–274. <https://doi.org/10.1093/molbev/msu300>

- Orlov NL, Ryabov SA, Nguyen TT (2009) Two New Species of Genera *Protobothrops* Hoge et Romano-Hoge, 1983 and *Viridovipera* Malhotra et Thorpe, 2004 (Ophidia: Viperidae: Crotalinae) from Karst Region in Northeastern Vietnam. Part I. Description of a New Species of *Protobothrops* Genus. Russian Journal of Herpetology 16(1): 69–82.
- Ota H, Ross CA (1994) Four new species of *Lycodon* (Serpentes: Colubridae) from the northern Philippines. Copeia 1994(1): 159–174. <https://doi.org/10.2307/1446682>
- Pauwels OSG, David P, Chan-ard T (2006) *Dryocalamus Davisonii tungsongensis* Nutphand, 1986 and *Lycodon suratensis* Nutphand, 1986 (Serpentes: Colubridae): translation of their original description and taxonomic Status. Hamadryad 30(1, 2): 114–120.
- Pope CH (1928) Seven new reptiles from Fukien Province, China. American Museum Novitates 320: 1–6.
- Pope CH (1928) Four new snakes and a new lizard from South China. American Museum Novitates 325: 1–4.
- Posada D, Crandall KA (1998) MODELTEST: testing the model of DNA substitution. Bioinformatics 14: 817–818. <https://doi.org/10.1093/bioinformatics/14.9.817>
- Ronquist F, Teslenko M, van der Mark P, Ayres DL, Darling A, Höhna S, Larget B, Liu L, Suchard MA, Huelsenbeck JP (2012) MrBayes 3.2: efficient Bayesian phylogenetic inference and model choice across a large model space. Systematic Biology 61: 539–542. <https://doi.org/10.1093/sysbio/sys029>
- Siler CD, Oliveros CH, Santanen A, Brown RM (2013) Multilocus phylogeny reveals unexpected diversification patterns in Asian Wolf Snakes (Genus *Lycodon*). Zoologica Scripta 42: 262–277. <https://doi.org/10.1111/zsc.12007>
- Slowinski JB, Pawar SS, Win H, Thin T, Gyi SW, Oo SL, Tun H (2001) A new *Lycodon* (Serpentes: Colubridae) from Northeast India and Myanmar (Burma). Proceedings of the California Academy of Science 52: 397–405. <https://www.mapress.com/j/zt/article/view/zootaxa.4276.4.6>
- Smith MA (1943) The fauna of British India, Ceylon and Burma, including the whole of the Indo-chinese subregion. Reptilia and Amphibia. Vol. III. Serpentes. Taylor & Francis, London, 583 pp.
- Stejneger L (1907) Herpetology of Japan and adjacent territory. Bulletin of the United States National Museum 58: 1–577. <https://doi.org/10.5479/si.03629236.58.i>
- Swofford DL (2001) PAUP*. Phylogenetic Analysis Using Parsimony (* and Other Methods), version 4. Sinauer Associates, Sunderland, Massachusetts. <https://doi.org/10.1111/j.0014-3820.2002.tb00191.x>
- Taylor EH (1922) The snakes of the Philippine Islands. Manila (Bureau of Printing or Science), Monograph 16: 1–312. <https://doi.org/10.5962/bhl.title.55346>
- Taylor EH (1950) The snakes of Ceylon. University of Kansas Science Bulletin 33(14): 519–603. <https://doi.org/10.5962/bhl.part.16131>
- Thompson JD, Gibson TJ, Plewniak F, Jeanmougin F, Higgins DG (1997) The ClustalX windows interface: Flexible strategies for multiple sequence alignment aided by quality analysis tools. Nucleic Acids Research 25: 4876–4882. <https://doi.org/10.1093/nar/25.24.4876>
- Uetz P, Freed P, Hošek J [Eds] (2019) The Reptile Database. <http://www.reptile-database.org> [accessed August 16, 2019]

- Vijayakumar SP, David P (2005) Taxonomy, natural history, and distribution of the snakes of the Nicobar Islands (India), based on new materials and with an emphasis on endemic species. *Russian Journal of Herpetology* 13(1): 11 – 40.
- Vogel G, David P (2010) A new species of the genus *Lycodon* (Boie, 1826) from Yunnan Province, China (Serpentes: Colubridae). *Bonn Zoological Bulletin* 57(2): 289–296.
- Vogel G, David P (2019) A new species of the *Lycodon fasciatus* complex from the Khorat Plateau, eastern Thailand (Reptiles, Squamata, Colubridae). *Zootaxa* 4577 (3): 515–528. <https://doi.org/10.11646/zootaxa.4577.3.6>
- Vogel G, David P, Pauwels OSG, Sumontha M, Norval G, Hendrix R, Vu NT, Ziegler T (2009) A revision of *Lycodon ruhstrati* (Fischer 1886) *auctorum* (Squamata Colubridae), with the description of a new species from Thailand and a new subspecies from the Asian mainland. *Tropical Zoology* 22: 131–182.
- Vogel G, Harikrishnan S (2013) Revalidation of *Lycodon hypsirhinoides* (Theobald, 1868) from Andamanislands (Squamata: Serpentes: Colubridae). *Taprobanica* 5(1): 19–31. <https://doi.org/10.4038/tapro.v5i1.5657>
- Vogel G, Luo J (2011) A new species of the genus *Lycodon* (Boie, 1826) from the southwestern mountains of China (Squamata: Colubridae). *Zootaxa* 2807: 29–40. <https://doi.org/10.11646/zootaxa.2807.1.2>
- Vogel G, Nguyen TQ, Kingsada P, Ziegler T (2012) A new species of the genus *Lycodon* (Boie, 1826) from Laos (Squamata: Serpentes: Colubridae). *North-Western Journal of Zoology* 8(2): 344–352. <http://biozoojournals.3x.ro/nwjz/index.html>
- Wall F (1906) A new Himalayan snake (*Lycodon mackinnoni*). *Journal of the Bombay Natural History Society* 17: 29–30.
- Werner F (1922) Neue Reptilien aus Süd-China, gesammelt von Dr. H. Handel-Mazzetti. *Anzeiger Österreichische Akademie der Wissenschaften. Wien, ser. Mathematik-Naturwissenschaften*, Vienna 59(24–25): 220–222.
- Wostl E, Hamidy A, Kurniawan N, Smith EN (2017) A new species of Wolf Snake of the genus *Lycodon* H. Boie in Fitzinger (Squamata: Colubridae) from the Aceh Province of northern Sumatra, Indonesia. *Zootaxa* 4276(4): 539–553. <https://doi.org/10.11646/zootaxa.4276.4.6>
- Zhang J, Jiang K, Vogel G, Rao D (2011) A new species of the genus *Lycodon* (Squamata, Colubridae) from Sichuan Province, China. 2982: 59–68.
- Zhang L, Peng LF, Yu L, Wang ZP, Huang LQ, Huang S (2015) New Record of *Lycodon liuchengchaoi* in Anhui. *Zoological Research* 36(1): 178–180.
- Zhao E (2006) Zhongguo Shelei [Snakes of China]. I. Anhui Science and Technology Publishing House, Hefei, 372 pp. [in Chinese].
- Zhao E, Jiang J (1981) Studies on amphibians and reptiles of Mt. Gongga Shan, Sichuan, China. I. A new species and a new subspecies of snake from Sichuan (in Chinese). *Acta Herpetologica Sinica*, Chengdu, [old ser.] 5(7): 53–58.
- Ziegler T, Ngo HN, Pham AV, Nguyen TT, Le MD, Nguyen TQ (2018) A new species of *Parafimbrrios* from northern Vietnam (Squamata: Xenodermatidae). *Zootaxa* 4527(2): 269–276. <https://doi.org/10.11646/zootaxa.4527.2.7>

Appendix I

Comparative specimens examined

- Lycodon fasciatus*. Vietnam: Quang Binh Province (ZFMK 86448)
Lycodon fasciatus. Vietnam: Quang Binh Province (ZFMK 86449)
Lycodon fasciatus. Vietnam: Quang Binh Province (ZFMK 86450)
Lycodon futsingensis. Vietnam: Cao Bang (IEBR 4165)
Lycodon futsingensis. Vietnam: Cao Bang (IEBR 4170)
Lycodon futsingensis. Vietnam: Vinh Phuc (ZFMK 89385)
Lycodon laoensis. Cambodia: Phnom Penh (ZFMK 54886)
Lycodon laoensis. Vietnam: Dong Nai (ZFMK 88928)
Lycodon meridionalis. Vietnam: Quang Ninh (ZFMK 95193)
Lycodon meridionalis. Vietnam: Hai Phong (ZFMK 94906)
Lycodon meridionalis. Vietnam: Bac Giang (ZFMK 89389)
Lycodon meridionalis. Vietnam: Vinh Phuc (ZFMK 89225)
Lycodon paucifasciatus. Vietnam: Quang Binh (ZFMK 86452)
Lycodon paucifasciatus. Vietnam: Quang Binh (ZFMK 80661)
Lycodon paucifasciatus. Vietnam: Quang Binh (ZFMK 80662)
Lycodon subcinctus. Indonesia: Bali (ZFMK 95499)
Lycodon subcinctus. Vietnam: Dong Nai (ZFMK 91899)