

# Italian Dermestidae: notes on some species and an updated checklist (Coleoptera)

Gianluca Nardi<sup>1</sup>, Jiří Háva<sup>2</sup>

**1** MiPAAF, Corpo Forestale dello Stato, Centro Nazionale per lo Studio e la Conservazione della Biodiversità Forestale “Bosco Fontana” di Verona, Sede di Bosco Fontana, Strada Mantova 29, I-46045 Marmirolo (MN), Italy **2** Department of Forest Protection and Entomology, Faculty of Forestry and Wood Sciences, Czech University of Life Sciences, Kamýcká 1176, CZ-165 21, Prague 6 - Suchdol, Czech Republic

Corresponding author: Gianluca Nardi ([l\\_nardi@hotmail.com](mailto:l_nardi@hotmail.com))

---

Academic editor: C. Majka | Received 20 July 2013 | Accepted 29 November 2013 | Published 6 December 2013

**Citation:** Nardi G, Háva J (2013) Italian Dermestidae: notes on some species and an updated checklist (Coleoptera). ZooKeys 360: 45–81. doi: 10.3897/zookeys.360.6023

---

## Abstract

An up-to-date checklist of the Italian Dermestidae is provided. The presence of 95 species in Italy is confirmed, while further 5 species (*Dermestes (Dermestes) vorax* Motschulsky, 1860, *Thorictus pilosus* Peyron, 1857, *T. wasmanni* Reitter, 1895, *Attagenus (Attagenus) simonis* Reitter, 1881 and *Globicornis (G.) breviclavus* (Reitter, 1878)) and 1 subspecies (*A. (A.) tigrinus pulcher* Faldermann, 1835) are excluded from the Italian fauna.

*Attagenus (Attagenus) calabricus* Reitter, 1881 and *A. (A.) lobatus* Rosenhauer, 1856 are for the first time recorded from Abruzzi and Tuscany respectively; *A. (A.) silvaticus* Zhantiev, 1976 is recorded for the first time from mainland Italy (Apulia); *Anthrenus (Anthrenus) angustefasciatus* Ganglbauer, 1904 is new to northern Italy (Friuli-Venezia Giulia), central Italy (Tuscany), Apulia and Basilicata; *A. (A.) munroi* Hinton, 1943 is new to central Italy (Elba Island); *A. (A.) delicatus* Kiesenwetter, 1851 is for the first time recorded from Apulia; *Globicornis (Globicornis) fasciata* (Fairmaire & Brisout de Barnevile, 1859) is new to southern Italy (Basilicata); *G. (Hadrotoma) sulcata* (C.N.F. Brisout de Barnevile, 1866) is for the first time recorded from central Italy (Abruzzi), Campania and Sicily, while *Trogoderma inclusum* LeConte, 1854 is new to Apulia.

Seven species (*Dermestes (Dermestes) peruvianus* Laporte de Castelnau, 1840, *D. (Dermestinus) carnivorus* Fabricius, 1775, *D. (Dermestinus) hankae* Háva, 1999, *D. (Dermestinus) intermedius intermedius* Kalík, 1951, *D. (Dermestinus) szekessyi* Kalík, 1950, *Anthrenus (Anthrenops) coloratus* Reitter, 1881 and *Trogoderma angustum* (Solier, 1849)) recently recorded from Italy (without further details) are discussed.

The lectotype and a paralectotype are designated for *Attagenus (A.) calabricus* Reitter, 1881 from Calabria.

*Attagenus pellio* (Linnaeus, 1758) var. *pilosissimus* Roubal, 1932 is removed from synonymy with *A. (A.) pellio* and recognized as a valid species (**stat. prom.**); it is known from Lombardy, Apulia and Calabria.

### Keywords

Dermestidae, lectotype designation, new status, taxonomy, distribution, new records, chorotypes, checklist, Italy

## Introduction

The main purpose of the present paper is to enhance knowledge of the Dermestidae of Italy. It is chiefly based on material collected by the staff of the Centro Nazionale per lo Studio e la Conservazione della Biodiversità Forestale “Bosco Fontana” di Verona (Italy), during entomological surveys in various Italian Protected Areas (e.g. Cerretti et al. 2003, Mason et al. 2006, Nardi and Vomero 2007, Cerretti et al. 2009, Nardi et al. 2011). Moreover, this paper gives an opportunity to provide a new checklist of the Dermestidae of Italy: it updates the nomenclature, systematics and the faunistic of the previous (Audisio et al. 1995) chiefly according to recent capital works (Háva 2007, Zhantiev 2011a, Háva 2011a).

## Material and methods

This paper is organized in three sections: “Notes on some species”, “Literature records” and “A new checklist of the Dermestidae of Italy”.

The classification of the subfamilies, tribes, genera and subgenera follows that proposed by Háva (2004, 2007); species are listed alphabetically according to the nomenclature of Háva (2007), Kadej et al. (2007), Háva (2008), Zhantiev (2009) and Háva (2011a).

The first section is based chiefly on study of new material. With very few exceptions, identifications were made by J. Háva.

The second section summarizes the literature data updating the Italian distribution of the species provided by Audisio et al. (1995) or by Háva and Nardi (2011).

In both sections, nomenclatural combinations and possible synonyms (listed chronologically-alphabetically) found in the literature that includes Italian records are provided below the valid name of each species or subspecies.

The first section concerns 25 species. As far as possible, the following data are provided for each record: region, province, general area and/or commune, locality of collection, altitude, UTM coordinates, date of collection, collector/s, collecting method, possible research project, number of specimens, and, in parentheses, abbreviation of depository. The labels of the examined specimens are generally written in Italian; hereunder, the regions and the collecting methods were translated into English. The label

data of type specimens are cited verbatim between quotation marks, a forward slash separating the different lines on a label, and a comma separating subsequent labels. Comments and interpretations are given in square brackets.

The mainland Italian regions are listed from north to south, and from west to east, all toponyms are listed alphabetically. Chorotypes were assigned according to Vigna Taglianti et al. (1993, 1999) and based on the distributions provided by Háva (2003, 2007, 2011b), Zhantiev (2009, 2011a, 2011b) and Háva et al. (2013), unless otherwise stated.

In the second section, the literature records updating the checklist of Dermestidae of Italy provided by Audisio et al. (1995) or by Háva and Nardi (2011) are detailed. These new faunistic data are based on papers subsequent to these two papers but also include some previously overlooked publications. These records concern 14 species. The data on distribution are listed according to the geographic divisions by Audisio et al. (1995): northern Italy (N), peninsular Italy (S), Sardinia and small circum-Sardinian islands (Sa) and Sicily and small circum-Sicilian islands (Si).

Finally, a new checklist of the Italian Dermestidae is provided (Table 1). The total number of species for each geographic division includes only those recorded with certain, so generic records for “Italy” and records followed by a question mark are excluded.

Acronyms for chorotypes follow Vigna Taglianti et al. (1993, 1999), while those for the geographic divisions of Italy follow Audisio et al. (1995); both are included in the foreword of the checklist.

The following abbreviations are used in the text: dint. = dintorni = environs; env. = environs; ex = specimen/s; FA = F. Angelini leg.; FEI = Forum Entomologi Italiani (<http://www.entomologiitaliani.net>); Fraz. = Frazione = hamlet; GC = G. Scaglioni leg.; GNP = Gargano National Park; leg. = legit or legerunt; PC = P. Cornacchia leg.; Prog. = Progetto = Project; prov. = province; sn = sweep net.

## Acronyms of specimen depositories

<b>AHG</b>	Andreas Herrmann private collection, Stade, Germany
<b>CNBFVR</b>	Centro Nazionale per lo Studio e la Conservazione della Biodiversità Forestale “Bosco Fontana” di Verona, Sede di Bosco Fontana. Marmirolo (Mantua), Italy
<b>GNAC</b>	Gianluca Nardi private collection, Cisterna di Latina (Latina), Italy
<b>HNHM</b>	Hungarian Natural History Museum, Budapest, Hungary
<b>JHAC</b>	Private Entomological Laboratory and Collection, Jiří Háva, Prague-west, Czech Republic
<b>MCZR</b>	Museo Civico di Zoologia, Rome, Italy
<b>MZUF</b>	Museo di Storia Naturale, Sezione di Zoologia, Università di Firenze, Florence, Italy
<b>PCOP</b>	Paolo Cornacchia private collection, Porto Mantovano (Mantua), Italy

## Notes on some species

### *Dermestes (Dermestes) peruvianus* Laporte de Castelnau, 1840

[http://species-id.net/wiki/Dermestes\\_peruvianus](http://species-id.net/wiki/Dermestes_peruvianus)

*Dermestes peruvianus* Laport.: Bertolini 1904: 58.

*Dermestes peruvianus* Casteln.: Luigioni 1929: 1024; Porta 1929: 301.

*Dermestes (Dermestes) peruvianus* Castelnau, 1840: Plata-Negrache 1972: 150; Zhantiev 2011a.

*Dermestes (Dermestes) peruvianus* Laporte de Castelnau, 1840: Audisio et al. 1995: 12, 16; Háva 2007: 300.

*Dermestes peruvianus* Laporte de Castelnau: Pollini 1998: 909.

**Material examined.** [Latium: Roma province,] Roma env., [no other data,] 1 ex (JHAC).

**Chorotype.** Subcosmopolitan (Háva 2007). This species is also recorded from Portugal (Grosso-Silva and Serrano 2000) and Switzerland (Kenis 2005, Zhantiev 2011a). Both countries must be added to the European distribution summarized by Háva (2007), while the former country must also be added to those provided by Zhantiev (2011a).

**Italian distribution.** Italy? (Bertolini 1904, Luigioni 1929, Porta 1929). Italian mainland? (Audisio et al. 1995). Italy (without further details) (Plata-Negrache 1972, Háva 2007). Italian mainland (Zhantiev 2011a).

**Remarks.** Species described from Peru (cf. Háva and Kalík 2005); the first precise record from Italy is shown above.

Domenichini et al. (1993: 272) wrote that this species is *common in food factories* and was reared experimentally in an insectarium in Piacenza University (northern Italy), but the place of origin of this captive population is not mentioned. This species, and *D. (Dermostinus) carnivorus* Fabricius, 1775 (see below), are also considered in an Italian manual of applied entomology (Pollini 1998), but no records from Italy are listed therein.

### *Dermestes (Dermostinus) carnivorus* Fabricius, 1775

[http://species-id.net/wiki/Dermestes\\_carnivorus](http://species-id.net/wiki/Dermestes_carnivorus)

*Dermestes carnivorus* Fabricius: Pollini 1998: 906.

*Dermestes (Dermostinus) carnivorus* Fabricius, 1775: Háva 2007: 300; Zhantiev 2011a.

**Material examined.** Tuscany: Firenze [= Florence], V.1948, [no collector,] 1 ♀ (JHAC).

**Chorotype.** Cosmopolitan (cf. Háva 2007).

**Italian distribution.** Italy (without further details) (Háva 2007). Italian mainland (Zhantiev 2011a).

**Remarks.** Pollini (1998: 906) wrote about this species that *si è acclimatato in Europa e in India [= it is naturalized in Europe and India]*, but no Italian locality was mentioned. The above generic Italian record by Háva (2007), was based on the above specimen.

***Dermestes (Dermestes) vorax* Motschulsky, 1860**

[http://species-id.net/wiki/Dermestes\\_vorax](http://species-id.net/wiki/Dermestes_vorax)

*Dermestes vorax* Motschulsky, 1860: Denux and Zagatti 2010: 357.

*Dermestes (Dermestes) vorax* Motschulsky, 1860: Háva and Nardi 2011: 408.

**Material examined.** None.

**Chorotype.** Asiatic (Háva 2003, 2007). This species was introduced to the coastal Croatia (Depoli 1928, Luigioni 1929, Porta 1934, all as *D. vorax* Motsch.), but this record was later ignored (Mroczkowski 1968, Háva 2003, 2007, Zhantiev 2011a).

**Italian distribution.** Italy (without further details) (Denux and Zagatti 2010).

**Remarks.** Denux and Zagatti (2010) wrote that an Italian record of this species was provided by Lohse (1979) and Háva (2003), but in these papers this species is not recorded from this country. Moreover, a recent record from “Italy” (<http://www.dermestidae.wz.cz/main.html>; <http://www.dermestidae.com/index.html>) is erroneous since concerns a dark specimen of *D. (Dermestes) lardarius* Linnaeus, 1758 (Háva, unpublished data), so it was ignored by Háva and Nardi (2011).

***Dermestes (Dermostinus) hankae* Háva, 1999**

[http://species-id.net/wiki/Dermestes\\_hankae](http://species-id.net/wiki/Dermestes_hankae)

*Dermestes (Dermostinus) hankae* Háva, 1999: Háva 2007: 301.

*Dermestes (Dermostinus) hankai* [sic!] Háva, 1999: Zhantiev 2011a.

**Material examined.** Italy mer. [= southern Italy], [no other data], 1 ♂ (JHAC).

**Chorotype.** W-Mediterranean: Spain, France, Italy and Algeria (Háva 2007, Prieto and Háva 2013).

**Italian distribution.** Italy (without further details) (Háva 2007). Italian mainland (Zhantiev 2011a).

**Remarks.** Species described from southern France (Háva 1999: 143); the generic record for Italy (Háva 2007) was based on the above specimen.

The specific name (*hankae*) is correct (ICZN 1999, art. 31.1.2), so the name *hankai* is only a subsequent incorrect spelling.

This species is very similar to *D. (Dermostinus) pardalis* Billberg, 1808 (Háva 1999, Herrmann and Bahillo de la Puebla 2003, Prieto and Háva 2013); the re-examination of specimens attributed to it could provide additional records of *D. (Dermostinus) hankae*. In Italy, *D. (Dermostinus) pardalis* is recorded from Sardinia (cf. Háva and Nardi

2011), generically from mainland Italy (Zhantiev 2011a), and from a non-specified locality of the northern mainland (Giachino 1982: 360, as *D. thoracicus* Dejean[, 1821 (nomen nudum)]), but this record, based on very old specimens, should be confirmed.

***Dermestes (Dermestinus) intermedius intermedius* Kalík, 1951**

[http://species-id.net/wiki/Dermestes\\_intermedius\\_intermedius](http://species-id.net/wiki/Dermestes_intermedius_intermedius)

*Dermestes (Dermestinus) intermedius intermedius* Kalík, 1951: Háva 2003: 18; Háva 2007: 301.

*Dermestes (Dermestinus) intermedius* Kalík, 1951: Zhantiev 2011a.

**Material examined.** None.

**Chorotype.** Turano-European reaching only as far as (*D. (D.) intermedius iranicus* Háva & Kalík, 1999) Iran, Iraq and Syria eastward (cf. Háva 2007, Háva et al. 2011).

**Italian distribution.** Italy (without further details) (Háva 2003, 2007). Italian mainland (Zhantiev 2011a).

**Remarks.** The above generic Italian records by Háva (2003, 2007) were based on a personal communication of V. Kalík, who identified an Italian specimen; unfortunately, no further details are available (Háva, unpublished data).

***Dermestes (Dermestinus) szekessyi* Kalík, 1950**

[http://species-id.net/wiki/Dermestes\\_szekessyi](http://species-id.net/wiki/Dermestes_szekessyi)

*Dermestes (Dermestinus) szekessyi* Kalík, 1950: Háva 2007: 302; Zhantiev 2011a.

**Material examined.** None.

**Chorotype.** Sibero-European (cf. Háva 2003, 2007, Háva et al. 2011, Zhantiev 2011a).

**Italian distribution.** Italy (without further details) (Háva 2007). Italian mainland (Zhantiev 2011a).

**Remarks.** The above generic Italian record of Háva (2007) was based on a personal communication of V. Kalík, who identified an Italian specimen; unfortunately, no further details are available (Háva, unpublished data).

***Dermestes (Montandonia) olivieri* Lepesme, 1939**

[http://species-id.net/wiki/Dermestes\\_olivieri](http://species-id.net/wiki/Dermestes_olivieri)

*Dermestes ater* Oliv. [= Olivier, 1790]: Ghiliani 1842: 33; Ghiliani 1847: 93; Reiche 1860: 720; Ghiliani 1887: 310; Bertolini 1890: 196; Iehl 1909: 26; Luigio-

ni 1929: 537; Porta 1929: 301; Gridelli 1949: 176; Horion 1955: 197; Faggioli 1956: 174; Gulli 1961: 4.

*Dermestes ater* Ol.: Villa and Villa 1844: 38 [444]; Rottenberg 1870: 238; Steck 1887: 181; Ragusa 1892: 203; Bertolini 1904: 58; Vitale 1904: 75; Holdhaus 1911: 445; Panganetti-Hummel 1918: 87; Zangheri 1969: 1334.

*Dermestes ater* Olivier: Giachino 1982: 361.

*Dermestes olivieri* Lep.: Angelini 1986: 74; Angelini and Montemurro 1986: 573; Angelini 1987: 38; Angelini 1991: 206.

*Dermestes (Dermestes) olivieri* Lepesme, 1939: Audisio et al. 1995: 12; Bordoni et al. 2006: 60; FEI 2011.

*Dermestes ater* Degeer, 1774 [sic!]: Sapuppo 2002: 189.

*Dermestes (Montandonia) olivieri* Lepesme, 1939: Háva and Nardi 2011: 416.

**Material examined.** Apulia: Foggia prov., [GNP] Mattinata, strada per M.te Sacro [= road to Mount Sacro], Masseria Vaira dint., 613 m, 33°T 584160 4821825, 24.V.2010, I. Toni leg., sn on *Ferula communis* [(Apiaceae)], Prog. Foresta Umbra, 1 ex (CNB-FVR); Taranto prov., Ginosa Marina, VIII.1985, P. Abbazzi leg., 2 ex (MZUF). Tuscany: Arezzo [prov.], Cavriglia, [Fraz.] Montegonzi, 16.V.1991, Lisa leg., 1 ex (MZUF); [Arezzo prov.] Ris. Nat. [= Riserva Naturale = Nature Reserve] “Valle Inferno-Bandella”, Casa Giardino, 8.V.1998, L. Bartolozzi, B. Cecchi A. & Sforzi leg., 1 ex (MZUF); [Firenze prov.] Firenze [= Florence], V.1896, ex-coll. Beccari, 1 ex (MZUF); Firenze prov., Greve [in Chianti], [Fraz.] Strada in Chianti, 20.I.1957, A. Bandinelli leg., 1 ex (MZUF); [Grosseto prov.], Parco Naturale della [= Natural Park of] Maremma, Mti [= Monti = Mountains] dell’Uccellina, S. Rabano, X.1988, L. Bartolozzi leg., 4 ex (MZUF).

**Chorotype.** Turano-European with extension to Tunisia (cf. Háva 2007, Háva and Herrmann 2009, 2011, Háva et al. 2011, Háva and Nardi 2011).

**Italian distribution.** All of Italy, Sicily and Sardinia (cf. Bertolini 1904, Luigioni 1929, Horion 1955, Audisio et al. 1995, Háva and Nardi 2011). These generic distributional data are most likely correct; nevertheless, precise records are known only from the following regions: Trentino (Bertolini 1890), Lombardy (Villa and Villa 1844, Giachino 1982), Piedmont (Ghiliani 1847, 1887), Aosta Valley (Iehl 1909), Romagna (Zangheri 1969), Tuscany (Bordoni et al. 2006), Latium (Horion 1955, FEI 2011), Abruzzi (Horion 1955), Apulia (Holdhaus 1911, Panganetti-Hummel 1918, Gridelli 1949, Horion 1955, Faggioli 1956, Angelini 1987), Basilicata (Angelini 1986, Angelini and Montemurro 1986), Calabria (Angelini 1991), Sicily (Ghiliani 1842, Rottenberg 1870, Steck 1887, Ragusa 1892, Vitale 1904, Luigioni 1929, Horion 1955, Audisio et al. 1995), and Sardinia (cf. Háva and Nardi 2011).

**Remarks.** New collecting sites from Tuscany and Apulia. In Tuscany, the species was formally recorded from a sole locality (Bordoni et al. 2006); it is also for the first time recorded here from a nature reserve, of which the beetle-fauna is well known (cf. Zinetti and Terzani 2009). The species was already known from Apulia, from various localities, including the Gargano National Park (see the references above).

Sapuppo (2002) recorded *Dermestes ater* DeGeer, 1774 from two Sicilian localities based on specimens collected by A. Gulli and identified by A. Porta. The data from a locality are exactly the same as those of “*Dermestes ater* Oliv.” provided by Gulli (1941), and thus, both sites are attributed here to *D. (Montandonia) olivieri*; the homonymy (*D. ater* DeGeer and *D. ater* Oliver) was very probably the cause of the error of Sapuppo (2002). *Dermestes (Dermestes) ater* DeGeer is a cosmopolitan species, but in Italy it is known from some northern regions only (Luigioni 1929: 537, as *D. (Dermestes) cadaverinus* Fabr. [= Fabricius, 1775], Minelli and Negrisolo 1993: 92, as *D. ater* Degeer, Audisio et al. 1995: 12, as *D. (Dermestes) ater* Degeer).

### ***Thorictus pilosus* Peyron, 1857**

[http://species-id.net/wiki/Thorictus\\_pilosus](http://species-id.net/wiki/Thorictus_pilosus)

*Thorictus pilosus* Peyron, 1857: Audisio et al. 1995: 12.

#### **Material examined.** None.

**Chorotype.** Turanian-E-Mediterranean: Cyprus, Egypt, Greece, Iraq, Israel, Lebanon, Libya, Syria, Tunisia, Turkey and Uzbekistan (John 1963, Löbl 2007).

**Italian distribution.** “S?” [= peninsular Italy?] (Audisio et al. 1995).

**Remarks.** The above doubtful record for peninsular Italy has already been ignored by other authors (Löbl 2007, Zhantiev 2011a); this species is excluded from Italy here, since when considering its geographical distribution, its occurrence in this country is improbable. A doubtful record from Corsica (cf. Bertolini 1874: 96, as *T. piliger* Schaum[, 1858], Luigioni 1929: 1022, as *T. pilosus* Pehr., Porta 1929: 204, as *T. pilosus* Peyron) has also never been confirmed (Löbl 2007, Zhantiev 2011a), and must be referred to *T. grandicollis grandicollis* Germar, 1842 (Háva, unpublished data).

### ***Thorictus wasmanni* Reitter, 1895**

[http://species-id.net/wiki/Thorictus\\_wasmanni](http://species-id.net/wiki/Thorictus_wasmanni)

*Thorictus wasmanni* Reitter, 1895: Audisio et al. 1995: 12.

#### **Material examined.** None.

**Chorotype.** Turanian: endemic of Uzbekistan (John 1963, Löbl 2007).

**Italian distribution.** “Si?” [= Sicily?] (Audisio et al. 1995).

**Remarks.** The above doubtful Sicilian record has already been ignored by other authors (Löbl 2007, Zhantiev 2011a); this species must be excluded from the Italian fauna, since its occurrence in Sicily is incompatible with its geonemy. It is known from the type specimens only (Zhantiev 2011b).

***Attagenus (Attagenus) calabricus* Reitter, 1881**

[http://species-id.net/wiki/Attagenus\\_calabricus](http://species-id.net/wiki/Attagenus_calabricus)

*Attagenus calabricus* Reitter, 1881: 37.

*Attagenus (Lanorus) calabricus* Reitt.: Reitter 1883: 89; Reitter 1891: 337; Ragusa 1892: 204; Reitter 1906: 378; Dalla Torre 1911: 52; Winkler 1926: 676; Luigioni 1929: 538; Porta 1929: 302.

*Megatoma calabrica* Rtrr.: Reitter 1887: 48.

*Attagenus calabricus* Reitt.: Bertolini 1904: 58; Panganetti-Hummel 1918: 87; Schatzmayr 1941: 73; Angelini 1986: 74; Angelini and Montemurro 1986: 573; Angelini 1987: 38; Angelini 1991: 206.

*Attagenus calabricus* Rtt.: Liebmann 1962: 4.

*Attagenus calabricus* Reitter, 1881: Mroczkowski 1968: 82; Sparacio 1997: 73.

*Attagenus (Lanorus) calabricus* Reitter, 1881: Audisio et al. 1995: 13.

*Attagenus (Attagenus) calabricus* Reitter, 1881: Háva 2003: 59; Háva 2007: 308; Zhantiev 2011a.

**Material examined.** Abruzzi: Pescara prov., Maiella, Valle Orfento, Caramanico, 23.VI.1988, G. Osella leg., 1 ex (JHAC). Apulia: Brindisi prov., Francavilla Fontana, 30.V.1998, C. Esposito leg., 1 ex (GNAC); [Foggia prov., GNP] Mattinata, 22.VI.1991, on flowers on the beach, M. Mantic leg., 4 ex (JHAC); [Foggia prov., GNP] Gargano, Mattinata, 8–10.V.1995, Z. Svec leg., 1 ex (JHAC); [Foggia prov., GNP, Vico del Gargano, Fraz.] San Menaio, V.1996, M. Fikacek leg., 2 ex (JHAC); Taranto prov., Circ. [= Circum] Mar Piccolo, 24.V.1992, Montemurro leg., 1 ex (JHAC); Taranto prov., Grottaglie, VI.1369 [sic!], FA, 1 ex (MCSV). Basilicata: [Matera prov.,] Marina di Nova Siri, 4.VI.2008, W. Apfel leg., 1 ex (JHAC); [Matera prov.,] Matera, 4.VI.1989, FA, 1 ex (AHG); [Matera prov.,] Matera, 350 m, 4.VI.1989, FA, 11 ex (MCSV); Potenza prov., Francavilla sul Sinni, 320 m, 5.VI.1988, querceta [= oak wood], FA, 1 ex (MCSV); [Potenza or Matera prov.,] Pollino, 19.VI.1994, FA, 1 ex (AHG). Calabria: “Calabria / Baudi [leg.]” [handwritten by unknown], “Holotypus 1881 / Attagenus / Calabricus / Reitter” [red-framed; handwritten by unknown], “Calabricus m. / Calab. Baudi [leg.]” [handwritten probably by Reitter], “coll. Reitter” [printed], “Lectotype / Attagenus calabricus Reitter / J. Háva des. 2012”, 1 ex (HNHM); “Calabria / Baudi [leg.]” [handwritten by unknown], “Paratypus 1881 / Attagenus / Calabricus / Reitter” [red-framed; handwritten by unknown], “coll. Reitter [printed]”, “Paralectotype / Attagenus calabricus Reitter / J. Háva des. 2012”, 1 ex (HNHM); Calabria [without further data], 1 ex (JHAC); [Cosenza prov., Cassano all’Ionio, Fraz.] Marina di Sibari env., 3 m, sandy dunes, 14.6.2009, P. Kresl leg., 6 ex (JHAC); [Cosenza prov.,] Roseto Capo Spulico, 27.V.–3.VI.2006, V. Hron leg., 1 ex (JHAC); [Cosenza prov.,] Sila, Mte. [= Mount] Oliveto, 2.VII.1929, C. Confalonieri leg., 1 ex (JHAC); [Cosenza prov.,] Sila, Sud [= South of] S. Giovanni [in] Fiore, 1100 m, 4.VII.[19]87, FA, 2 ex (MCSV); Reggio Calabria prov., Aspromonte. [= Aspromonte], str. [= strada = road] Cimina–Zomaro, 600–800 m, 6.VI.1994, FA, 6 ex (JHAC).

**Chorotype.** Apenninic endemic (Háva 2007).

**Italian distribution.** Campania (Luigioni 1929), Apulia (Panganetti-Hummel 1918, Luigioni 1929, Angelini 1987), Basilicata (Angelini 1986, Angelini and Montemurro 1986), Calabria (Bertolini 1904, Luigioni 1929, Porta 1929, Schatzmayr 1941, Angelini 1991) and Sicily (Ragusa 1892, Bertolini 1904, Winkler 1926, Luigioni 1929, Porta 1929, Liebmann 1962, Mroczkowski 1968, Audisio et al. 1995, Sparacio 1997, Háva 2003, 2007, Zhantiev 2011a).

The generic records for Italy (Reitter 1883, 1891, 1906, Dalla Torre 1911, Winkler 1926, Mroczkowski 1968, Háva 2003, 2007), peninsular Italy (Audisio et al. 1995), southern Italy (Sparacio 1997) and mainland Italy (Zhantiev 2011a) refer to the above listed regions.

**Remarks.** First record from the Abruzzi region; this new site (cf. Osella 1988) is at the northernmost limit of the species distribution, while the above record from the Brindisi province is at the easternmost limit. The southernmost site is the Pantelleria Island in the Sicilian Channel (Liebmann 1962).

Reitter (1881) described this species without giving any locality of collecting (see also Reitter 1887), but based on the specific name, the type locality is apparently in the Calabria region (southern peninsular Italy). The species was described based on an unspecified number of specimens without any type designation (Reitter 1881). The two type specimens above are thus syntypes and are designated here as lectotype and paralectotype, respectively; in this way “Calabria” becomes the type locality of the species (ICZN 1991, art. 76.2).

### *Attagenus (Attagenus) lobatus* Rosenhauer, 1856

[http://species-id.net/wiki/Attagenus\\_lobatus](http://species-id.net/wiki/Attagenus_lobatus)

*Attagenus (Attagenus) lobatus* Rosenhauer, 1856: Háva and Nardi 2011: 420.

**Material examined.** Tuscany: Firenze [= Florence], 28.IX.1988, L. Chelazzi leg., 1 ♂ (MZUF).

**Chorotype.** Centralasiatic-Mediterranean; this species was introduced to the USA (cf. Háva and Nardi 2011; Háva et al. 2013).

**Italian distribution.** Latium, Campania, Apulia, Calabria, Sicily and Sardinia (cf. Háva and Nardi 2011).

**Remarks.** First record for Tuscany. The new locality is the northernmost in Italy (cf. Háva and Nardi 2011).

### *Attagenus (Attagenus) pilosissimus* Roubal, 1932 stat. prom.

*Attagenus pellio* L. var. *pilosissimus* Roubal, 1932: 66.

*Attagenus pellio* Lin. v. *pilosissimus* Roubal: Porta 1934: 171.

*Attagenus pellio* (Linnaeus, 1758) var. *pilosissimus* Roubal, 1932: Mroczkowski 1968: 89.

*Attagenus pellio* var. *pilosissimus* Roub.: Angelini 1986: 74.

*Attagenus pellio* var. *pilosissimus* Roubal: Angelini 1991: 206.

*Attagenus* (*Attagenus*) *pellio* (Linné, 1758) = *pilosissimus* Roubal, 1932: Audisio et al. 1995: 13.

*Attagenus* (*Attagenus*) *pellio* (Linnaeus, 1758) = *Attagenus pellio* var. *pilosissimus* Roubal, 1932: Háva 2003: 66.

*Attagenus* (*Attagenus*) *pellio* (Linnaeus, 1758) = *pilosissimus* Roubal, 1932: Háva 2007: 57.

*Attagenus* (*Attagenus*) *pellio* (Linnaeus, 1758) = *Attagenus pellio* var. *pilosissimus* Roubal, 1932: Háva and Nardi 2007: 122.

**Material examined.** Apulia: Bari prov., Gioia del Colle, 22.IV.1993, De Marzo leg., 1 ♂ (JHAC); Foggia prov., [GNP] Foresta Umbra, Monte Sant'Angelo, caserme forestali [= barracks of the Italian National Forestry Service], 792 m, 33T 582306 4630267, 27.IV.2010, PC, death on windowsill, Prog. Foresta Umbra, 1 ♀ (JHAC). Lombardy: Bergamo prov., Oltre il Colle, 7.VII.2005, FA, 1 ♂ (JHAC).

**Chorotype.** Alpino-Apenninic endemic (cf. Háva and Nardi 2007).

**Italian distribution.** Apulia and Calabria (Roubal 1932, Porta 1934, Angelini 1986, 1991, Háva and Nardi 2007).

**Remarks.** *Attagenus pellio* var. *pilosissimus* from Apulia was listed as a synonym of *A. (A.) pellio* by Audisio et al. (1995), because Mroczkowski (1968) listed *pilosissimus* as a “variety”. This means that this name was implicitly considered as a junior synonym as Mroczkowski also treated subspecific taxa in his paper. This opinion was confirmed by Háva and Nardi (2007: 122) through a study of the holotype, a female specimen which is very worn. A photo of this holotype is published at <http://www.dermestidae.com/Attagenuspelliotipilosissimus.html>. Nevertheless, based on the study of the above new material, this variety is resurrected as a valid species (stat. prom.). *Attagenus* (*Attagenus*) *pilosissimus* and *A. (A.) pellio* show differences in the shape of the antennal club (Figs 1–2) and in the male genitalia (Figs 4–5): the median lobe of the aedeagus of the former species (Fig. 5) is longer and narrower, and parameres are parallel for most of their length. Moreover, *A. (A.) pilosissimus* also differs by the pubescence of the abdomen (Fig. 4) and pronotum (Fig. 3) and by the presence of only one small circular spot with whitish pubescence on each elytron (Fig. 3). *Attagenus* (*A.*) *pilosissimus* is known only from mainland Italy (Lombardy, Apulia and Calabria), where, in sympatry, *A. (A.) pellio* (Linnaeus, 1758) also occurs (cf. Holdhaus 1911, as *A. pellio* L., Háva and Nardi 2007).

### *Attagenus* (*Attagenus*) *silvaticus* Zhantiev, 1976

[http://species-id.net/wiki/Attagenus\\_silvaticus](http://species-id.net/wiki/Attagenus_silvaticus)

*Attagenus* (*Attagenus*) *silvaticus* Zhantiev, 1976: Háva and Nardi 2007: 122.

*Attagenus silvaticus* Zhantiev, 1976: Zhantiev 2011a.

**Material examined.** Apulia: Foggia prov., [GNP] Foresta Umbra, Vico del Gargano, Bosco Sfilzi, 391 m, 33T 585633 4634653, 23.V.2010, D. Birtele & I. Toni leg., window flight trap, Prog. Foresta Umbra, 1 ♀ (CNBFVR).

**Chorotype.** European (Czech Republic, Hungary, northwestern Russia, Sicily, Slovakia, Ukraine) with extension eastward to “Caucasus”, Anatolia, Iran and West Siberia (Háva 2007, Háva and Nardi 2007, Háva et al. 2010, as *A. silvaticus*, Zhantiev 2011a).

**Italian distribution.** Sicily (Háva and Nardi 2007, Zhantiev 2011a).

**Remarks.** First record for mainland Italy. The above specimen was intercepted in a beech wood (D. Birtele, pers. comm., 2012).

### *Attagenus (Attagenus) simonis* Reitter, 1881

[http://species-id.net/wiki/Attagenus\\_simonis](http://species-id.net/wiki/Attagenus_simonis)

*Attagenus tigrinus* F. [= (Fabricius, 1792)] v. *Simoni* [sic!] Reit.: Bertolini 1904: 58.

*Attagenus bifasciatus* Oliv. [= (Olivier, 1790)] v. *Simoni* [sic!] Reitt.: Luigioni 1923b: 4.

*Attagenus (Lanorus) bifasciatus* Oliv. a *Simoni* [sic!] Reitt.: Luigioni 1929: 538; Porta 1929: 302.

*Attagenus bifasciatus* (Olivier, 1790) var. *simoni* [sic!] Reitter, 1881: Sapuppo 2002: 187–188.

*Attagenus (Attagenus) simonis* Reitter, 1881: Háva and Nardi 2007: 418–419.

**Material examined.** None.

**Chorotype.** E-Mediterranean: Egypt, Israel, Jordan and Syria (Háva 2007).

**Italian distribution.** Latium (Luigioni 1929), Campania (Bertolini 1904, Luigioni 1929, Porta 1929), Capri Island (Luigioni 1923b, 1929, Porta 1929) and Sicily (Luigioni 1929, Sapuppo 2002).

**Remarks.** This species was recently excluded from the Italian fauna (Háva and Nardi 2007, 2011); also according to Mroczkowski (1968: 81, as *A. bifasciatus* var. *simonis*), it occurs only in some East-Mediterranean countries. Nevertheless, an escatomediterranean distributional pattern (Zilli 2000) could explain the occurrence of this species in Italy.

Gulli (1961: 4, as *A. bifasciatus* Oliv. var. *Rossii* Ganglb. [= Ganglbauer, 1904] and *A. b.* var. *Simoni* [sic!]) recorded Sicilian specimens of *A. (A.) rossii* (identified by A. Porta) with a colour similar to *A. (A.) simonis*; these specimens are also listed by Sapuppo (2002), but unfortunately their re-examination has so far been impossible.

### *Attagenus (Attagenus) tigrinus pulcher* Faldermann, 1835


[http://species-id.net/wiki/Attagenus\\_tigrinus\\_pulcher](http://species-id.net/wiki/Attagenus_tigrinus_pulcher)

*Attagenus pulcher* Fald.: Bargagli 1872: 102, Bertolini 1874: 99.

*Attagenus tigrinus* F. [= (Fabricius, 1792)] var. *pulcher* Falder.: Bertolini 1904: 58.

**Material examined.** None.

**Chorotype.** Mediterranean (cf. Háva and Nardi 2011); *A. (A.) tigrinus pulcher* is a Caucasian endemic (Zhantiev 2009).



**Figures 1–6.** *Attagenus (Attagenus)* spp., males **1** Antennal club of *A. (A.) pilosissimus* from Italy (Apulia, Bari prov., Gioia del Colle) **2** Antennal club of *A. (A.) pellio* from Czech Republic (Central Bohemia, Prague) **3** Habitus of *A. (A.) pilosissimus* from Italy (Apulia, Bari prov., Gioia del Colle) (length 4.5 mm) **4** Abdomen in ventral view of *A. (A.) pilosissimus* from Italy (Apulia, Bari prov., Gioia del Colle) **5** Aedeagus (setation omitted) of *A. (A.) pilosissimus* from Italy (Lombardy, Bergamo prov., Oltre il Colle) (length 0.5 mm) **6** Aedeagus (setation omitted) of *A. (A.) pellio* from Czech Republic (Central Bohemia, Prague) (length 0.5 mm) (photos and drawings by J. Háva).

**Italian distribution.** Sardinia (Bargagli 1872, Bertolini 1874, 1904).

**Remarks.** This subspecies is excluded here from the Italian fauna. Zhantiev (2009) recognized this taxon as a valid subspecies. This paper was overlooked by Háva and Nardi (2011), so these authors, according to Háva (2007), listed it as a synonym of *A. (A.) bifasciatus* (A.G. Olivier, 1790). Nevertheless, the status of this subspecies needs to be revised (Háva, unpublished data).

***Anthrenus (Anthrenops) coloratus* Reitter, 1881**

[http://species-id.net/wiki/Anthrenus\\_coloratus](http://species-id.net/wiki/Anthrenus_coloratus)

*Anthrenus (Anthrenops) coloratus* Reitter, 1881: Háva 2007: 311.

*Anthrenus (Florilinus) coloratus* Reitter, 1881: Zhantiev 2011a.

**Material examined.** Apulia: [Foggia prov., GNP] Gargano, V.1991, J. Háva leg., 12 ex (JHAC).

**Chorotype.** Turano-Europeo-Mediterranean with extension to Canary Islands, Tajikistan, “India” and Sudan; this species was introduced to and is widespread in the USA (Beal 1998, as *A. coloratus*, Háva 2003, 2007).

**Italian distribution.** Italy (Háva 2007). Italian mainland (Zhantiev 2011a).

**Remarks.** The generic Italian record by Háva (2007) was based on the above specimens.

***Anthrenus (Anthrenus) angustefasciatus* Ganglbauer, 1904**

[http://species-id.net/wiki/Anthrenus\\_angustefasciatus](http://species-id.net/wiki/Anthrenus_angustefasciatus)

*Anthrenus (Anthrenus) pimpinellae* Fabr. [= (Fabricius, 1775)] a. *angustefasciatus* Ganglb.: Luigioni 1929: 540.

*Anthrenus pimpinellae* Fabr. ab. *angustefasciatus* Ganglb.: Porta 1934: 171.

*Anthrenus (Anthrenus) angustefasciatus* Ganglbauer, 1904: Háva 2003: 79; Háva 2007: 311; Kadej et al. 2007: 728; Háva 2011b: 78; Zhantiev 2011a; Háva et al. 2013: 137.

**Material examined.** Friuli-Venezia Giulia: Trieste prov., Carso, [Sgonico, Fraz.] Gabrovizza S. Primo, 12.VI.1988, sn, PC, 1 ♀ (PCOP). Tuscany: Grosseto prov., Pitigliano, 23.I.[20]02, L. Saltini leg., 1 ex (GNAC). Apulia: [Brindisi prov.,] Brindisi env., 18.V.1997, J. Hrdlicka leg., 4 ex (JHAC). Basilicata: Potenza prov., [Pollino National Park,] Terranova di Pollino, 900 m, 27–30.VI.2005, GC, sn, 1 ♀ (PCOP).

**Chorotype.** European with extension southward to a part of the Maghreb: Algeria, Bosnia and Herzegovina, Croatia, Czech Republic, France mainland, Italy, Morocco, Portugal, Spain, Switzerland, Yugoslavia and European Turkey (cf. Háva 2011b, Zhantiev 2011a, Háva et al. 2013).

The records for Corsica (Luigioni 1929, Porta 1934) were later ignored (Sainte-Claire Deville 1937, Mroczkowski 1968, as *A. (A.) pimpinellae* var. *angustefasciatus*, Zhantiev 2011a, Háva et al. 2013).

**Italian distribution.** Calabria (Luigioni 1929, Porta 1934), Sardinia (Háva et al. 2013). Italy (without further details) (Háva 2003, 2007, Kadej et al. 2007, Háva 2011b). Italian mainland (Zhantiev 2011a).

**Remarks.** Ganglbauer (1904: 42) described *A. pimpinellae* var. *angustefasciatus* from “Dalmatien [= Dalmatia]” that was only recently raised to species level (Háva 2003: 79); its diagnostic features were provided by different authors (Kadej 2005, Kadej et al. 2007, Háva 2011b, Háva and Zahradník 2011).

First records for northern Italy (Friuli-Venezia Giulia), for central Italy (Tuscania), for Apulia and Basilicata; the former record is expected since this species occurs in neighbouring foreign regions (Depoli 1928, as *A. pimpinellae* f. *angustefasciatus* Ganglb., Novak 1952, as *A. pimpinellae* a. *angustefasciatus* Gglb.). The generic records for Italy (Háva 2003, 2007, Kadej et al. 2007, Háva 2011b) and, probably also those generic for mainland Italy (Zhantiev 2011a), refer to previous literature records and, in part, to the above material.

### *Anthrenus (Anthrenus) delicatus* Kiesenwetter, 1851

[http://species-id.net/wiki/Anthrenus\\_delicatus](http://species-id.net/wiki/Anthrenus_delicatus)

*Anthrenus pimpinellae* Fabr. [= (Fabricius, 1775)] var. *delicatulus* [sic!] Kiesw.: Ghiliani 1887: 311.

*Anthrenus pimpinellae* F. var. *delicatus* Kiesw.: Steck 1887: 181; Ragusa 1892: 206.

*Anthrenus pimpinellae* Fabr. Var. *delicatulus* [sic!] Kiesw.: Baudi 1890: 102.

*Anthrenus pimpinellae* F. [var.] *Isabellinae* Muls. [= Mulsant & Rey, 1868]: Ragusa 1892: 206.

*Anthrenus pimpinellae* F. var. *delicatus* K.: Bertolini 1904: 59.

*Anthrenus pimpinellae* F. ab. *delicatus* Kiesw.: Luigioni 1923a: 135.

*Anthrenus (Anthrenus) pimpinellae* F. a. *delicatus* Kiesw.: = *Isabellinae* Muls. Rey: Luigioni 1929: 540.

*Anthrenus (Anthrenus) pimpinellae* Fabr. V. *delicatus* Kiesw.: Porta 1929: 306.

*Anthrenus pimpinellae* v. *delicatus* Kiesw.: Liebmann 1962: 4.

*Anthrenus (Anthrenus) delicatus* Kiesenwetter, 1851: Audisio et al. 1995: 14; Kadej 2005: 728; Lo Cascio et al. 2006: 322; Háva and Nardi 2011: 423; Zhantiev 2011a.

*Anthrenus (Anthrenus) delicatus* Kiesenwetter, 1852 [sic!]: Háva 2007: 312.

**Material examined.** Apulia: Foggia prov., [GNP] Promontorio del Gargano, Vieste, SP 89 [= Strada Provinciale n. 89 = Provincial Road n. 89], Casa Cupari dint., 354 m, 33T 5920001 4630150, 29.V.2010, abandoned quarry, sn on grasses, PC, Prog. Foresta Umbra, 6 ex (CNBFVR; JHAC); Foggia prov., [GNP] Promontorio del Gargano, Peschici, Lampia del Principe dint., 161 m, 33T 587282 4639346, 28.IV.2010, maquis sn, PC, Prog. Foresta Umbra, 1 ♀ (CNBFVR). Latium: Roma prov., M.ti [= Monti = Mounts] Simbruini, Jenne dint., M.te [= Monte = Mount] Porcaro, 800 m, 31.V. 2007, sn, PC GC, 1 ex (PCOP). Sicily: Palermo prov., Corleone, Bosco della Ficuzza, 700 m, 1.VI. 2008, sn near a watering trough, PC GC, 1 ex (PCOP).

**Chorotype.** Mediterranean, reaching eastward as far as Caucasus and Iran (cf. Kadej 2005, Háva and Nardi 2011, Zhantiev 2011b). This species is also recorded from Cyprus (Zhantiev 2011a, Alziar and Lemaire 2012: 67, as *A. delicatus*), and this country must be added to the distribution summarized by Háva and Nardi (2011).

**Italian distribution.** Trentino, Venezia Giulia (Luigioni 1929), Piedmont (Ghiliani 1887, Baudi 1890, Bertolini 1904, Luigioni 1929), Lombardy (Háva and Nardi

2004), Liguria (Luigioni 1929), Latium, Campania (Luigioni 1923a, Porta 1929), Sicily (Ragusa 1892, Bertolini 1904, Luigioni 1929, Porta 1929, Liebmann 1962, Audisio et al. 1995, Lo Cascio et al. 2006) and Sardinia (Bertolini 1904, Luigioni 1929, Porta 1929, Audisio et al. 1995, Háva and Nardi 2011, Zhantiev 2011a).

Moreover, Luigioni (1929) recorded this species generically for central and southern Italy; generic records for northern and peninsular Italy (Audisio et al. 1995) and for “Italy” (Kadej 2005, Háva 2007) refer to the above regions, the same is probably valid for a generic record for mainland Italy (Zhantiev 2011a).

**Remarks.** First detailed records from Apulia. About Sicily, Ragusa (1892) wrote *Ne posseggo vari esemplari [= I have several specimens]* without giving the collecting sites, so the sole known precise Sicilian localities were the Pantelleria and Lipari Islands (Liebmann 1962, Lo Cascio et al. 2006); the above new site is located in one of the most important forested areas of western Sicily, a zone of great value for environmental conservation (cf. Sabella and Sparacio 2004, Mason et al. 2006, La Mantia et al. 2010, Falci et al. 2012). Luigioni (1923a) also recorded *A. (A.) delicatus* from the environs of Rome; this record was overlooked by Nardi (1997). As stated by Audisio et al. (1995: 16), the Italian distribution of this species should be verified, because it was considered for a long time as a variety or aberration of *A. (A.) pimpinellae*. Moreover, the colour of elytral fasciae in four of the above Apulian specimens are visually very similar to those of *A. (A.) mroczkowskii* Kalík, 1954, nevertheless their identification was established by the examination of the structure of antennae, 9th sternite and male genitalia (cf. Kadej 2005, Kadej et al. 2007). In this framework, it will be better to revise the specimens of *A. (A.) mroczkowskii* recorded so far from Italy (Angelini 1986, as *A. pimpinellae mroczkowskii*, Angelini and Montemurro 1986, as *A. pimpinellae mroczkowskii*). The distribution of *A. (A.) mroczkowskii* also includes the following countries: Albania, Bosnia and Herzegovina, Bulgaria, Corsica, Crete, Croatia, Greece, Slovenia, European Turkey and Algeria (introduced) (Háva 2007, as *A. (A.) pimpinellae mroczkowskii*, Kadej et al. 2007).

### *Anthrenus (Anthrenus) munroi* Hinton, 1943

[http://species-id.net/wiki/Anthrenus\\_munroi](http://species-id.net/wiki/Anthrenus_munroi)

*Anthrenus munroi* Hint.: Angelini 1986: 73; Angelini and Montemurro 1986: 573; Angelini 1987: 37.

*Anthrenus (Anthrenus) munroi* Hinton, 1943: Háva and Nardi 2011: 425.

**Material examined.** Tuscany: [Livorno prov.], Is. [= Isola = Island] d’Elba, dint. Marina di Campo, 15–17.VI.1999, Abbazzi, Bartolozzi, Lo Cascio & Sforzi leg., 1 ♂ (MZUF).

**Chorotype.** Mediterranean (eastwards as far as Ukraine, Turkey, Syria, Lebanon, Jordan, and Israel), with extension to Hungary and Portugal (cf. Kadej et al. 2007, Háva and Nardi 2011).

**Italian distribution.** Apulia, Basilicata, Calabria (Angelini 1986, Angelini and Montemurro 1986, Angelini 1987) and Sardinia (Háva and Nardi 2011).

**Remarks.** First record for central Italy; this new site is the northernmost in Italy. The species also occurs in the facing Corsica (cf. Háva and Nardi 2011).

*Anthrenus (A.) pimpinellae pimpinellae* (Fabricius, 1775) and *A. (Nathrenus) verbasci* (Linnaeus, 1767) were the sole congeneric species previously recorded from Elba Island (Holdhaus 1923, as *A. pimpinellae* F. and *A. verbasci* L.; Luigioni 1929, Porta 1929, both as *A. (A.) pimpinellae* Fabr. and *A. (N.) verbasci* Lin.)

#### *Anthrenus (Anthrenus) pimpinellae pimpinellae* (Fabricius, 1775)

[http://species-id.net/wiki/Anthrenus\\_pimpinellae\\_pimpinellae](http://species-id.net/wiki/Anthrenus_pimpinellae_pimpinellae)

*Anthrenus pimpinellae* F.: Ragusa 1892: 206; Bertolini 1904: 59; Holdhaus 1911: 445; Panganetti-Hummel 1918: 87; Holdhaus 1923: 101; Luigioni 1923b: 4; Marcuzzi 1985: 8.

*Anthrenus (Anthrenus) pimpinellae* Fabr.: Luigioni 1929: 540; Porta 1929: 306.

*Anthrenus (Anthrenus) pimpinellae pimpinellae* (Fabricius, 1775): Háva and Nardi 2004: 120; Háva and Nardi 2011: 426.

*Anthrenus (Anthrenus) pimpinellae* (Fabricius, 1775) s.l.: Lo Cascio et al. 2006: 322.

**Material examined.** Apulia: Foggia prov., [GNP] Promontorio del Gargano, Vieste, SP 89 [= Strada Provinciale n. 89 = Provincial Road n. 89], Casa Cupari dint., 354 m, 33T 5920001 4630150, abandoned quarry, sn on grasses, 29.V.2010, PC, Prog. Foresta Umbra, 4 ex (CNBFVR).

**Chorotype.** Cosmopolitan (cf. Háva and Nardi 2011).

**Italian distribution.** All of Italy and Sicily (Bertolini 1904, Luigioni 1929, Porta 1929; Háva and Nardi 2011). This species is also recorded from some minor islands: Elba (Holdhaus 1923), Capri (Luigioni 1923b, 1929, Porta 1929), Lampedusa (Ragusa 1892, Luigioni 1929) and Vulcano (Lo Cascio et al. 2006).

**Remarks.** Species common and widespread in Italy but from the Gargano National Park (cf. Angelini 1987) was recorded only by Holdhaus (1911); it is also known from other Apulian localities (Panganetti-Hummel 1918, Marcuzzi 1985).

#### *Anthrenus (Nathrenus) signatus* Erichson, 1846

[http://species-id.net/wiki/Anthrenus\\_signatus](http://species-id.net/wiki/Anthrenus_signatus)

*Anthrenus (Nathrenus) signatus* Erichson, 1846: Háva and Nardi 2011: 428.

**Material examined.** Basilicata: Potenza prov., Pignola, Ris. [= Riserva = Reserve] WWF, L. [= Lago = Lake] Pignola, 700 m, 21.VI.1992, FA, 1 ♂ (JHAC).

**Chorotype.** European (cf. Háva and Nardi 2011).

**Italian distribution.** South Tyrol, Venetia, Venezia Giulia, Latium?, Apulia, Basilicata and Sicily (cf. Háva and Nardi 2011).

**Remarks.** This species was recorded in Basilicata region from the Pollino National Park (Angelini 1986) only; the above specimen comes from a nature reserve, from which a sole congeneric species, *A. (N.) verbasci*, was known so far (Angelini 1996a, 1998, both as *A. verbasci*).

*Anthrenus signatus* ab. *bielawskii* Mroczkowski, 1958 from Bulgaria (Mroczkowski 1958: 6, 1968: 145) was recently ignored (Háva 2007, 2008, 2011a), since according to the ICZN (1999, art. 45.5 and 45.6.2), it is an infrasubspecific and unavailable name.

### ***Globicornis (Globicornis) breviclavis* (Reitter, 1878)**

[http://species-id.net/wiki/Globicornis\\_breviclavis](http://species-id.net/wiki/Globicornis_breviclavis)

*Hadrotoma breviclavis* Reitter: Baudi 1890: 101.

*Hadrotoma breviclavis* Reitt.: Bertolini 1904: 59.

*Globicornis (Globicornis) breviclavis* Reitt.: Ganglbauer 1904: 31; Reitter 1906: 379;

Dalla Torre 1911: 63; Winkler 1926: 679; Luigioni 1929: 539; Porta 1929: 304.

*Globicornis (Globicornis) breviclavis* (Reitter, 1878): Audisio et al. 1995: 13, 16.

### **Material examined.** None.

**Chorotype.** Endemic to the Near East: Armenia, Bulgaria, “Caucasus”, Georgia, European Turkey, Ukraine, and Southern European Russia (Adygeya, Krasnodar kraj) (Háva 2007, Háva and Legalov 2010, Zhantiev 2011a).

**Italian distribution.** Piedmont: Graian and Pennine Alps (Baudi 1890, Luigioni 1929, Porta 1929); Piedmont (without further details) (Bertolini 1904, Reitter 1906); Piedmont? (Dalla Torre 1911, Winkler 1926, Audisio et al. 1995). Liguria: Ligurian Apennines (Luigioni 1929); Liguria? (Audisio et al. 1995).

**Remarks.** This species described from Caucasus (Reitter 1878, as *Hadrotoma breviclavis*) is excluded here from the Italian fauna. Ganglbauer (1904) listed the paper by Baudi (1890), while Dalla Torre (1911) referred to Ganglbauer (1904). The Italian records were ignored by Mroczkowski (1968: 114), who recorded this species only from the Caucasus. No specimens of this species are present in the P. Luigioni collection (MCZR) (A. Zilli, pers. comm. 2013). The above Italian records probably concern the similar *G. (G.) luckowi* Herrmann, Háva & Kadej, 2011 which is known only from southern Switzerland, Piedmont and Liguria (Herrmann et al. 2011, Háva et al. 2013). Nevertheless, as discussed for *Attagenus (A.) simonis*, an escatomediterranean distributional pattern (Zilli 2000) could explain the occurrence of *G. (G.) breviclavis* in Italy.

**Globicornis (Globicornis) fasciata (Fairmaire & Brisout de Barneville, 1859)**  
[http://species-id.net/wiki/Globicornis\\_fasciata](http://species-id.net/wiki/Globicornis_fasciata)

*Hadrotoma fasciata* Fairm.: Bertolini 1904: 59; Luigioni 1923a: 135.  
*Globicornis fasciata* Fair.: Vitale 1904: 75.  
*Globicornis (Globicornis) fasciata* Frm. [18]59: Winkler 1926: 679.  
*Globicornis (Globicornis) fasciata* Fairm.: Luigioni 1929: 539; Porta 1929: 304.  
*Globicornis fasciatus* [sic!] Fairm.: Horion 1955: 208.  
*Globicornis (Globicornis) fasciata* (Fairmaire, 1859): Mroczkowski 1968: 114; Audisio et al. 1995: 13, 16; Zhantiev 2011a.  
*Globicornis (Globicornis) fasciata* (Fairmaire & Brisout, 1859): Háva 2003: 109; Háva and Nardi 2011: 429.  
*Globicornis (Globicornis) fasciata* (Fairmaire & Brisout de Barneville, 1859): Háva 2007: 315.

**Material examined.** Basilicata: Potenza prov., Parco Naz. del Pollino [= Pollino National Park], Casa del Conte dint., tra [= between] Acqua Tremola e [= and] Piani di S. Francesco, 1100–1500 m, 9.VII.2002, sn, GC, 3 ex (PCOP).

**Chorotype.** European (Belgium, Denmark, France, Germany, Italy, Spain, Sweden, Switzerland) with extension southward to Tunisia (cf. Háva et al. 2010, Háva and Nardi 2011).

**Italian distribution.** Piedmont (Porta 1929, Horion 1955), Tuscany (Luigioni 1929, Horion 1955), Latium (Luigioni 1923a, 1929, Porta 1929, Horion 1955), Sicily (Vitale 1904) and Sardinia (Bertolini 1904, Winkler 1926, Luigioni 1929, Porta 1929, Horion 1955, Mroczkowski 1968, Audisio et al. 1995, Háva 2003, 2007, Zhantiev 2011a).

The records for northern and peninsular Italy (Audisio et al. 1995) as those generic for Italy (Háva 2003) refers to above regions; the same is probably valid for a generic record for mainland Italy (Zhantiev 2011a).

**Remarks.** First record from southern Italy. Five species of this genus are now recorded from the Pollino National Park (cf. Angelini 1986).

The records for Piedmont (Porta 1929, Horion 1955) and those generic for northern Italy (Audisio et al. 1995: 13) refers to the Maritime Alps (Audisio et al. 1995: 16). The species also occurs in a neighbouring French department (Sainte-Claire Deville 1937: 252, as *G. fasciata* Fairm.). The record from Sicily (Vitale 1904) was omitted by subsequent authors (cf. Luigioni 1929, Audisio et al. 1995, Zhantiev 2011a), and thus, it should be confirmed.

In the literature (see also above), the authorship of this species was often attributed only to Fairmaire.

***Globicornis (Hadrotoma) sulcata* (C.N.F. Brisout de Barneville, 1866)**  
[http://species-id.net/wiki/Globicornis\\_sulcata](http://species-id.net/wiki/Globicornis_sulcata)

*Globicornis ? sulcata* (C. Brisout de Barneville, 1866): Angelini 1986: 74.

*Globicornis (Hadrotoma) ? sulcata* (C. Brisout de Barneville, 1866): Audisio et al. 1995: 16.

*Globicornis (Hadrotoma) sulcata* (C. Brisout de Barneville, 1866): Audisio et al. 1995: 13.

**Material examined.** Abruzzi: Parco Naz. [= National Park], L'Aquila prov., Pescasseroli, 1300 m, 1.IV.2003, FA, 1 ♀ (JHAC). Basilicata: Potenza prov., Moliterno, 500 m, 11.V.2003, FA, 1 ♀ (JHAC); Potenza prov., Ruoti, 29.IX.1996, *Abies–Quercus*, FA, 1 ♂ (JHAC). Campania: Avellino prov., Cilento, Novi Velia, M. [= Mount] Sacro, 1900 m, 7.XII.1996, *Fagus*, FA, 1 ♂ (JHAC). Sicily: Palermo prov., Ficuzza, 700 m, 1–4.V.2000, bosco leccio [= ilex wood], FA, 1 ♂ (JHAC); Palermo prov., Bosco della Ficuzza, 27.V.2006, M. Sarovec leg., 1 ♂, 4 ♀♀ (JHAC).

**Chorotype.** W-Mediterranean: France, Italy and Spain.

**Italian distribution.** Basilicata (Angelini 1987, Audisio et al. 1995).

**Remarks.** First records from central Italy (Abruzzi), Campania and Sicily. In the Sicilian site, it was also collected *Anthrenus (Anthrenus) delicatus* (see above).

***Trogoderma angustum* (Solier, 1849)**  
[http://species-id.net/wiki/Trogoderma\\_angustum](http://species-id.net/wiki/Trogoderma_angustum)

*Trogoderma angustum* (Solier, 1849): Háva 2007: 319; Zhantiev 2011a.

**Material examined.** None.

**Chorotype.** Subcosmopolitan (cf. Háva 2007). This species is also recorded from Switzerland (Kenis 2005); this country must be added to the European distributions recently summarized by Háva (2007) and by Zhantiev (2011a).

**Italian distribution.** Italy (without further details) (Háva 2007). Italian mainland (Zhantiev 2011a).

**Remarks.** The above generic Italian record by Háva (2007) was based on a personal communication from V. Kalík, who identified Italian material; unfortunately, no further details are available (Háva, unpublished data).

***Trogoderma inclusum* LeConte, 1854**  
[http://species-id.net/wiki/Trogoderma\\_inclusum](http://species-id.net/wiki/Trogoderma_inclusum)

*Trogoderma meridionalis* Kraatz, 1858: 146.

*Trogoderma meridionalis* Kraatz: Mulsant and Rey 1868: 132.

*Trogoderma meridionale* [sic!] Kraatz: Bertolini 1874: 99.

- Trogoderma versicolor* Creutz. [= (Creutzer, 1799)] var. *meridionale* [sic!] Kr.: Ragusa 1892: 205; Porta 1929: 305.
- Trogoderma versicolor* Creut. v. *meridionale* [sic!] K.: Bertolini 1904: 59.
- Trogoderma versicolor* Creutz. a. *meridionale* [sic!] Kr.: Luigioni 1929: 540.
- Trogoderma versicolor meridionale* [sic!] Kraatz, 1858; Mroczkowski 1968: 107.
- Trogoderma inclusum* Le Conte: Dal Monte 1972: 102.
- Trogoderma ? inclusum* Lec.: Angelini and Montemurro 1986: 573.
- Trogoderma inclusum*: Nardi R. et al. 1993: 623, fig. 9.
- Trogoderma [?] inclusum* J.L. Leconte, 1854: Audisio et al. 1995: 13, 16.
- Trogoderma inclusum* LeConte: Nicoli Aldini 1999: 8.
- Trogoderma inclusum* LeConte, 1854: Háva 2007: 319; Háva and Nardi 2011: 432; Zhantiev 2011a.

**Material examined.** Apulia: Lecce prov., Salento, Torre S. Giovanni dint., 1–7. VII.2002, PC GC, sn, 1 ex (PCOP). Venetia: Verona prov., Verona, [Fraz.] Cancelllo, [about 500 m], 22.VI.1986, sn, PC, 1 ex (PCOP).

**Chorotype.** Cosmopolitan (cf. Háva and Herrmann 2011, Háva and Nardi 2011, Háva et al. 2011).

**Italian distribution.** Lombardy (Nicoli Aldini 1999, Háva and Nardi 2011), Liguria (Luigioni 1929), Emilia-Romagna (Nicoli Aldini 1999), Latium (Nardi R. et al. 1993, Nicoli Aldini 1999), Basilicata [?] (Angelini and Montemurro 1986, Audisio et al. 1995), Sicily (Kraatz 1858, Mulsant and Rey 1868, Ragusa 1892, Bertolini 1904, Porta 1929, Mroczkowski 1968) and Sardinia (cf. Háva and Nardi 2011).

The first Italian record of this species was provided by Kraatz (1858) who recorded it from Sicily (without a precise locality), while some authors (Bertolini 1874, Dal Monte 1972, Háva 2007) and Zhantiev (2011a) recorded it from Italy and mainland Italy, respectively, without further details. Moreover, Nicoli Aldini (1999: 8) also recorded it from *alcune città del Nord-Italia* [= some towns of Northern Italy] but without mentioning the regions.

**Remarks.** In Italy, this species and *T. versicolor* (Creutzer, 1799) have often been confused (cf. Nicoli Aldini 2003b, Háva and Nardi 2011); old Italian records in the literature must be thus considered with caution. *Trogoderma inclusum* is recorded here for the first time from Apulia and Venetia. This species is a stored product pest; it also develops in nests of social aculeate Hymenoptera, and is a predator of egg masses of *Lymantria dispar* (Linnaeus, 1758) (Lepidoptera, Erebidae, Lymantriinae) (cf. Háva and Nardi 2011).

## Literature records

The old Sicilian records of *Attagenus (A.) trifasciatus* (Fabricius, 1787) and *Trogoderma glabrum* (Herbst, 1783) (Ghiliani 1842) were ignored (or overlooked?) by subsequent authors and are thus considered as doubtful here (Table 1).

*Thorictus mauritanicus* Lucas, 1846 and *T. stricticollis* Kraatz, 1859 were both recorded from “Italy” by Háva (2003, 2007) and from “Italian mainland” by Zhantiev (2011a). Háva’s (2003, 2007) records of *T. mauritanicus* refer only to Sardinia and Sicily, while those of *T. stricticollis* (Háva 2003, 2007) refer only to Sardinia (Háva, unpublished data). In this framework, the records of both species for mainland Italy (Zhantiev 2011a) are considered erroneous.

*Thorictus sicilianus* John, 1965 (= *sicilianus* John, 1966) from Sicily and Morocco (John 1965, 1966, Audisio et al. 1995, Löbl 2007) was recently recognized as a junior synonym of *T. grandicollis grandicollis* Germar, 1842 (Háva 2013); *Attagenus picipennis* Pic, 1894 from Sicily (Audisio et al. 1995) is a junior synonym of *A. (Attagenus) brunneus* Faldermann, 1835 (Háva 2007: 57); *Anthrenus (A.) scrophulariae suecius* Palm, 1940 that was recorded from southern Italy and Sicily (Audisio et al. 1995), was recently listed by Háva (2007: 313) as a junior synonym of *A. (A.) scrophulariae scrophulariae* (Linnaeus, 1758), but for an error this new synonym was not formally established (cf. Háva 2007: 57). These three taxa are thus excluded from the new checklist.

***Dermestes (Dermostinus) gyllenhalii gyllenhalii*** Laporte de Castelnau, 1840  
*Dermestes ? atomarius* Erichson, 1846: Sapuppo 2002: 189.  
Si? (Sapuppo 2002).

***Dermestes (Dermostinus) laniarius laniarius*** Illiger, 1801  
*Dermestes laniarius* Illig.: Gulli 1961: 4.  
*Dermestes laniarius* Illiger, 1801: Sapuppo 2002: 190.  
Si (Gulli 1961, Sapuppo 2002).

***Thorictus grandicollis grandicollis*** Germar, 1842  
*Thorictus grandicollis* Germ.: Cerruti 1954: 111.  
N (Cerruti 1954).

***Attagenus (Attagenus) cyphonoides*** Reitter, 1881  
*Attagenus alfierii* Pic[, 1910]: Trematerra 1987: 16.  
*Attagenus (Attagenus) cyphonoides* Reitter, 1881: Háva and Nardi 2004: 121, Háva 2007: 308.  
*Attagenus (Attagenus) cyphonoides* Reitter, 1881: Zhantiev 2011a.  
S (Háva and Nardi 2004). Italy (Trematerra 1987, Háva 2007). Italian mainland (Zhantiev 2011a).

***Attagenus (Attagenus) fasciatus*** (Thunberg, 1795)  
*Attagenus fasciatus* (Thunberg, 1795): Nicoli Aldini 2003a 155–156, Nicoli Aldini 2003b: 123–125, Nicoli Aldini 2004: 983, 985, Zhantiev 2011a.  
*Attagenus (Attagenus) fasciatus* (Thunberg, 1795): Háva and Nardi 2004: 121, Háva 2007: 308.

N (Nicoli Aldini 2003a, 2003b, 2004), S (Háva and Nardi 2004). Italy (Háva 2007). Italian mainland (Zhantiev 2011a).

***Attagenus (Attagenus) pantherinus* (Ahrens, 1814)**

*Attagenus pantherinus*: Henry 1938: 66.

*Attagenus (Attagenus) pantherinus* (Ahrens, 1814): Háva 2007: 309, Háva and Nardi 2007: 121.

*Attagenus pantherinus* (Ahrens, 1814): Zhantiev 2011a.

N (Henry 1938, Háva and Nardi 2007), Sa (Zhantiev 2011a). Italy (Háva 2007).

***Attagenus (Attagenus) rossii* Ganglbauer, 1904**

*Dermestes bifasciatus* Rossi, 1794: 79.

*Attagenus bifasciatus* Rossi: Rottenberg 1870: 238, Bertolini 1874: 99, Steck 1887: 181, Bertolini 1904: 58.

*Attagenus bifasciatus*: Failla-Tedaldi 1887: 158.

*Attagenus (Lanorus) bifasciatus* Rossi: Ragusa 1892: 204.

*Attagenus (Lanorus) bifasciatus* Oliv. [= (A.G. Olivier, 1790)] a. *Rossii* Ganglb.: Lui-gioni 1929: 538.

*Attagenus (Lanorus) bifasciatus* Oliv. v. *Rossii* Ganglb.: Porta 1929: 302.

*Attagenus bifasciatus* Oliv. var. *Rossii* Ganglb.: Gulli 1961: 4.

*Attagenus bifasciatus* (Olivier, 1790) var. *Rossii* Ganglbauer, 1904: Sapuppo 2002: 187.

*Attagenus (Attagenus) rossii* Ganglbauer, 1904: Háva 2007: 309, Háva and Nardi 2007: 123, Háva and Nardi 2011: 422.

*Attagenus rossii* Ganglbauer, 1904: Zhantiev 2011a.

*Attagenus rossii* Ganglbauer, 1804 [sic!]: FEI 2012.

S (Rossi 1794, Luigioni 1929, Porta 1929, Háva and Nardi 2007, 2011), Si (Rotten-berg 1870, Bertolini 1874, Failla-Tedaldi 1887, Steck 1887, Ragusa 1892, Ber-tolini 1904, Luigioni 1929, Porta 1929, Gulli 1961, Sapuppo 2002, Háva and Nardi 2007, Zhantiev 2011a, FEI 2012), Sa (cf. Háva and Nardi 2011). Italy (Háva 2007). Italian mainland (Zhantiev 2011a).

***Attagenus (Attagenus) trifasciatus* (Fabricius, 1787)**

*Attagenus trifasciatus* Fabr.: Ghiliani 1842: 33.

*Attagenus ? trifasciatus* (Fabricius, 1787): Sapuppo 2002: 189.

Si? (Ghiliani 1842, Sapuppo 2002).

***Anthrenocerus australis* (Hope, 1843)**

*Anthrenocerus australis* (Hope, 1843): Herrmann and Háva 2007: 5, Háva 2007: 315, Zhantiev 2011a.

N (Herrmann and Háva 2007). Italy (Háva 2007). Italian mainland (Zhantiev 2011a).

***Globicornis (Globicornis) luckowi*** Herrmann, Háva & Kadej, 2011

*Globicornis (Globicornis) luckowi* Herrmann, Háva & Kadej, 2011: Háva et al. 2013: 138.  
N (Háva et al. 2013).

***Globicornis (Hadrotoma) corticalis*** (Eichhoff, 1863)

*Globicornis corticalis* (Eichhoff, 1863): Biscaccianti et al. 2008: 50.  
S (Biscaccianti et al. 2008).

***Reesa vespulae*** (Milliron, 1939)

*Reesa vespulae* (Milliron, 1939): Nicoli Aldini 2003a: 156, 159, Nicoli Aldini 2003b:  
123, 127, Nicoli Aldini 2004: 983, 987.  
N (Nicoli Aldini 2003a, 2003b, 2004).

***Trogoderma glabrum*** (Herbst, 1783)

*Trogoderma elongatulum* Fabr. [= (Fabricius, 1801)]: Ghiliani 1842: 33.  
Si? (Ghiliani 1842).

***Trogoderma variabile*** Ballion, 1878

*Trogoderma variabile* Ballion, 1878: Nicoli Aldini 1998: 115, Háva 2003: 141, Háva  
2007: 320.  
*Trogoderma variabile* Ballion: Nicoli Aldini 2003a: 156, Nicoli Aldini 2003b: 126–127.  
N (Nicoli Aldini 1998, 2003a, 2003b), Sa (Zhantiev 2011a). Italy (Háva 2003, 2007).  
Italian mainland (Zhantiev 2011a).

## A new checklist of the Dermestidae of Italy

The present-day dermestid fauna of Italy is summarized in the checklist below (Table 1).

The following abbreviations are used: AFR = Afrotropical; ALAP = Alpino-Apenininc endemic; ALPC = Central-Alpine endemic; APPE = Apenninic endemic; AUST = Australian; CAE = Centralasiatic-European; CEM = Centralasiatic-Europo-Mediterranean; CEU = Central European; COS = Cosmopolitan; EME = E-Mediterranean; EUM = Europeo-Mediterranean; EUR = European; IT = Italy and/or Italian mainland without further details; (i) = introduced to Europe; MED = Mediterranean; N = northern Italy (Trentino-Alto Adige, Venetia, Friuli-Venezia Giulia, Lombardy, Piedmont, Aosta Valley, Liguria and Emilia-Romagna); NAF = N-African; OLA = Holarctic; PAL = Palearctic; S = peninsular Italy (the remainig of continental Italy); Sa = Sardinia and small circum-Sardinian islands; SCO = Subcosmopolitan; SEU = S-European; Si = Sicily and small circum-Sicilian islands; SICI = Sicilian endemic; SIE = Siberio-European; TEM = Turano-Europo-Mediterranean; TUE = Turano-European; TUM = Turano-Mediterranean; WME = W-Mediterranean.

**Table I.** Checklist of the Dermestidae of Italy.

Taxa	Distribution			Chorotype			
<b>Dermestinae Latreille, 1804</b>							
<b>Dermestini Latreille, 1804</b>							
<b>Genus <i>Dermestes</i> Linnaeus, 1758</b>							
<b>Subgenus <i>Dermestes</i> Linnaeus, 1758</b>							
1. <i>ater</i> DeGeer, 1774	N			COS			
2. <i>bicolor</i> Fabricius, 1781				PAL			
<i>b. bicolor</i> Fabricius, 1781	N	S		Sa			
3. <i>haemorrhoidalis</i> Küster, 1852	N			COS			
4. <i>hispanicus</i> Kalík, 1952			Sa	WME			
5. <i>lardarius</i> Linnaeus, 1758	N	S	Si	Sa			
6. <i>peruvianus</i> Laporte de Castelnau, 1840	N?	S		COS			
<b>Subgenus <i>Dermestinus</i> Zhantiev, 1967</b>							
7. <i>aurichalceus</i> Küster, 1846	N	S?		Sa			
8. <i>carnivorus</i> Fabricius, 1775		S		COS			
9. <i>erichsoni</i> Ganglbauer, 1904	N	S	Si	Sa			
10. <i>frischii</i> Kugelann, 1792	N	S	Si	Sa			
11. <i>gyllenhalii</i> Laporte de Castelnau, 1840				EUR			
<i>g. gyllenhalii</i> Laporte de Castelnau, 1840	N	S	Si?				
12. <i>hankae</i> Háva, 1999		S		WME			
13. <i>intermedius</i> Kalík, 1951				EUR			
<i>i. intermedius</i> Kalík, 1951	IT	IT					
14. <i>laniarius</i> Illiger, 1801				PAL			
<i>l. laniarius</i> Illiger, 1801	N	S	Si	Sa			
15. <i>maculatus</i> DeGeer, 1774	N	S	Si	Sa			
16. <i>murinus</i> Linnaeus, 1758				PAL			
<i>m. murinus</i> Linnaeus, 1758	N	S	Si	Sa			
17. <i>mustelinus</i> Erichson, 1846	N	S	Si	Sa			
18. <i>pardalis</i> Billberg, 1808	N?	IT		Sa			
19. <i>sardous</i> Küster, 1846				MED			
<i>s. sardous</i> Küster, 1846	N	S		Sa			
20. <i>szekessyi</i> Kalík, 1950	IT	IT		SIE			
21. <i>undulatus</i> Brahm, 1790	N	S	Si	OLA			
<b>Subgenus <i>Montandonia</i> Jacquet, 1886</b>							
22. <i>fuliginosus</i> P. Rossi, 1792	N	S	Si	EUR			
23. <i>hirticollis</i> Fabricius, 1792			Sa	WME			
24. <i>olivieri</i> Lepesme, 1939	N	S	Si	TEM			
<b>Thorictinae Agassiz, 1846</b>							
<b>Thaumaphrastini Anderson, 1949</b>							
<b>Genus <i>Thorictodes</i> Reitter, 1875</b>							
25. <i>heydeni</i> Reitter, 1875	N	S		COS			
<b>Thorictini Agassiz, 1846</b>							
<b>Genus <i>Thorictus</i> Germar, 1834</b>							
26. <i>grandicollis</i> Germar, 1842				TUM			
<i>g. grandicollis</i> Germar, 1842	N	S	Si	Sa			
27. <i>mauritanicus</i> Lucas, 1846			Si	Sa			
28. <i>stricticollis</i> Kraatz, 1859				MED			
<i>s. stricticollis</i> Kraatz, 1859				Sa			

Taxa	Distribution				Chorotype
	N	S	Si	Sa	
29. <i>studti</i> John, 1971			Si		SICI
<b>Orphilinae LeConte, 1900</b>					
<b>Genus <i>Orphilus</i> Erichson, 1848</b>					
30. <i>niger</i> (P. Rossi, 1790)	N	S	Si	Sa	CEM
<b>Trinodinae T.L. Casey, 1900</b>					
<b>Thylodriini Semenov, 1909</b>					
<b>Genus <i>Thylodrias</i> Motschulsky, 1839</b>					
31. <i>contractus</i> Motschulsky, 1839	N				COS
<b>Trinodini T.L. Casey, 1900</b>					
<b>Genus <i>Trinodes</i> Dejean, 1821</b>					
32. <i>hirtus</i> (Fabricius, 1781)	N	S	Si	Sa	TUE
<b>Attageninae Laporte, 1840</b>					
<b>Attagenini Laporte, 1840</b>					
<b>Genus <i>Attagenus</i> Latreille, 1802</b>					
<b>Subgenus <i>Attagenus</i> Latreille, 1802</b>					
33. <i>bifasciatus</i> (A.G. Olivier, 1790)		S	Si	Sa	TUM
34. <i>brunneus</i> Faldermann, 1835	N	S	Si	Sa	OLA
35. <i>calabricus</i> Reitter, 1881		S	Si		APPE
36. <i>cypionoides</i> Reitter, 1881		S			SCO
37. <i>fallax</i> Gené, 1839		S	Si	Sa	MED
38. <i>fasciatus</i> (Thunberg, 1795)	N	S			COS
39. <i>lobatus</i> Rosenhauer, 1856		S	Si	Sa	CEM
40. <i>maritimus</i> (Gené, 1839)			Si	Sa	WME
41. <i>obtusus</i> (Gyllenhal, 1808)				Sa	TEM
42. <i>pantherinus</i> (Ahrens, 1814)	N			Sa	EUR
43. <i>pellio</i> (Linnaeus, 1758)	N	S	Si	Sa	COS
44. <i>pilosissimus</i> Roubal, 1932	N	S			ALAP
45. <i>punctatus</i> (Scopoli, 1772)	N	S	Si		EUR
46. <i>rossii</i> Ganglbauer, 1904		S	Si	Sa	MED
47. <i>schaefferi</i> (Herbst, 1792)					OLA
<i>s. schaefferi</i> (Herbst, 1792)	N	S	Si	Sa	
48. <i>silvaticus</i> Zhantiev, 1976		S	Si		SIE
49. <i>simplex</i> Reitter, 1881		S	Si	Sa	MED
50. <i>tigrinus</i> (Fabricius, 1792)					
<i>t. tigrinus</i> (Fabricius, 1792)	N	S	Si	Sa	MED
51. <i>trifasciatus</i> (Fabricius, 1787)	N		Si?		MED
52. <i>unicolor</i> (Brahm, 1790)					COS
<i>u. unicolor</i> (Brahm, 1790)	N	S	Si	Sa	
53. <i>uniformis</i> Fairmaire, 1860				Si	NAF
<b>Megatominae Leach, 1815</b>					
<b>Anthrenini Gistel, 1848</b>					
<b>Genus <i>Anthrenus</i> Geoffroy, 1762</b>					
<b>Subgenus <i>Anthrenops</i> Reitter, 1881</b>					
54. <i>coloratus</i> Reitter, 1881		S			OLA+AFR
<b>Subgenus <i>Anthrenus</i> Geoffroy, 1762</b>					
55. <i>angustefasciatus</i> Ganglbauer, 1904	N	S			EUR
56. <i>delicatus</i> Kiesenwetter, 1851	N	S	Si	Sa	MED
57. <i>festivus</i> Erichson, 1846		S?	Si	Sa	MED

Taxa	Distribution				Chorotype
	N	S	Si	Sa	COS
58. <i>flavipes</i> LeConte, 1854		S		Sa	
<i>f. flavipes</i> LeConte, 1854					
59. <i>goliath</i> Saulcy, 1868	N	S	Si	Sa	MED
60. <i>munroi</i> Hinton, 1943		S		Sa	MED
61. <i>mroczkowskii</i> Kalík, 1954		S			SEU
62. <i>pimpinellae</i> (Fabricius, 1775)					COS
<i>p. isabellinus</i> Küster, 1848	N	S	Si	Sa	
<i>p. pimpinellae</i> (Fabricius, 1775)	N	S	Si		
63. <i>scrophulariae</i> (Linnaeus, 1758)					COS
<i>s. scrophulariae</i> (Linnaeus, 1758)	N	S	Si	Sa	
<b>Subgenus Florilinus Mulsant &amp; Rey, 1868</b>					
64. <i>museorum</i> (Linnaeus, 1761)	N	S	Si	Sa	COS
65. <i>oberthueri</i> Reitter, 1881	N		Si		WME
<b>Subgenus Helocerus Mulsant &amp; Rey, 1868</b>					
66. <i>fuscus</i> A.G. Olivier, 1790	N	S	Si	Sa	SCO
67. <i>minutus</i> Erichson, 1846		S	Si	Sa	MED
<b>Subgenus Nathrenus Casey, 1900</b>					
68. <i>biskrensis</i> Reitter, 1887			Si		NAF
69. <i>molitor</i> Aubé, 1850	N			Sa	MED
70. <i>signatus</i> Erichson, 1846	N	S	Si		EUR
71. <i>verbasci</i> (Linnaeus, 1767)	N	S	Si	Sa	COS
72. <i>versicolor</i> Reitter, 1887			Si	Sa	MED
<b>Megatomini Leach, 1815</b>					
<b>Genus Anthrenocerus Arrow, 1915</b>					
73. <i>australis</i> (Hope, 1843)	N				AUST (i)
<b>Genus Ctesias Stephens, 1830</b>					
<b>Subgenus Ctesias Stephens, 1830</b>					
74. <i>serra</i> (Fabricius, 1792)	N	S	Si	Sa	EUR
<b>Genus Globicornis Latreille, 1829</b>					
<b>Subgenus Globicornis Latreille, 1829</b>					
75. <i>bifasciata</i> (Perris, 1866)			Si	Sa	WME
76. <i>fasciata</i> (Fairmaire & Brisout de Barnevile, 1859)	N	S	Si?	Sa	CEU
77. <i>luckowi</i> Herrmann, Háva & Kadej, 2011	N				ALPC
78. <i>nigripes</i> (Fabricius, 1792)	N	S			EUR
79. <i>picta</i> (Küster, 1851)		S	Si	Sa	EUR
80. <i>tristis</i> (Reitter, 1881)		S			EME
81. <i>variegata</i> (Küster, 1851)	N	S	Si	Sa	SEU
<b>Subgenus Hadrotoma Erichson, 1848</b>					
82. <i>corticalis</i> (Eichhoff, 1863)	N	S	Si		EUR
83. <i>emarginata</i> (Gyllenhal, 1808)	N	S	Si	Sa	EUR
84. <i>sulcata</i> (C.N.F. Brisout de Barnevile, 1866)		S	Si		WME
<b>Genus Megatoma Herbst, 1791</b>					
<b>Subgenus Megatoma Herbst, 1791</b>					
85. <i>ruficornis</i> Aubé, 1866		S			SEU
86. <i>undata</i> (Linnaeus, 1758)	N	S			EUR
<b>Genus Phradonoma Jacquelin du Val, 1859</b>					
87. <i>villosum</i> (C. Duftschmid, 1825)	N				CAE

Taxa	Distribution				Chorotype
<b>Genus <i>Reesa</i> Beal, 1967</b>					
88. <i>vespulae</i> (Milliron, 1939)	N				SCO
<b>Genus <i>Trogoderma</i> Dejean, 1821</b>					
89. <i>angustum</i> Solier, 1849	IT	IT			SCO
90. <i>glabrum</i> (Herbst, 1783)	N	S	Si?		SCO
91. <i>granarium</i> Everts, 1898	N	S	Si		COS
92. <i>inclusum</i> LeConte, 1854	N	S	Si	Sa	COS
93. <i>megatomoides</i> Reitter, 1881	N				OLA
94. <i>variabile</i> Ballion, 1878	N			Sa	COS
95. <i>versicolor</i> (Creutzer, 1799)	N	S	Si	Sa	COS
<b>Total species</b>	<b>61</b>	<b>67</b>	<b>57</b>	<b>59</b>	

## Discussion

The updated checklist of Italian Dermestidae includes 95 species, moreover *Anthrenus (A.) pimpinellae* occurs with two subspecies, but the taxonomic status of *A. (A.) p. isabellinus* needs to be clarified (cf. Háva and Nardi 2004). This total number is very probably still provisional since this family has been little studied in Italy, and the occurrence of further species recently described (e.g. Zhantiev 2001) or resurrected is possible. The previous checklist (Audisio et al. 1995), included only 81 species. Considering the four adopted geographic divisions, the relatively high species richness in the peninsular region (67) is unsurprising (cf. Minelli et al. 2002).

Some of the above records come from the following two protected areas of southern peninsular Italy: the Gargano National Park (Apulia) and the Pollino National Park (Basilicata and Calabria). 24 species of Dermestidae are now known from the former Park, while 23 species are now known from the Pollino National Park, but only 15 species are common to both (cf. Holdhaus 1911, Panganetti-Hummel 1918, Angelini 1987, 1996b, Háva and Nardi 2004). 24 and 23 species represent 37% and 34% of those currently recorded with certainty from peninsular Italy. These values confirm the importance of these Parks for insects (cf. Holdhaus 1911, Palm 1939, Grandi 1956, Angelini 1986, 1987, Rusek and Stumpp 1988, Agostini and Scali 1989, Rivosecchi 1989, Merz 1993, Angelini 1996b, Ruffo and Stoch 2006) and for biodiversity conservation.

## Acknowledgments

We are indebted to Claudio Angeloro (Corpo Forestale dello Stato - Ufficio Territoriale per la Biodiversità di Foresta Umbra, Italy) for the assistance in Apulia during the research of the CNBFVR; Daniele Birtele (CNBFVR), Paolo Cornacchia (Porto Mantovano, Italy), Carmine Esposito (Velletri, Italy), Gianluca Scaglioni (Porto Mantovano, Italy) and Ilaria Toni (CNBFVR) for the collection of specimens; to Luca Bartolozzi (Museo di Storia Naturale, Sezione di Zoologia, Università di Firenze, Florence, Italy) for the loan of specimens from the collections under his care; to

Franco Mason (Verona, Italy) for his constructive comments on a previous version of this paper; to Ottó Merkl (Hungarian Natural History Museum, Budapest, Hungary) for the information on the types of *Attagenus calabricus*; to Miloslav Rakovič (Prague, Czech Republic) and to Jonathan Cooter (Oxford, England) for the linguistic revision of the manuscript. We thank Riaan Stals (Pretoria, South Africa) and an anonymous referee for their reviews.

## References

- Agostini N, Scali V (1989) Segnalazione di nuove stazioni italiane di *Bacillus atticus* Brunner, 1882: Isole Tremiti, Gargano e Sicilia (Phasmatodea). Bollettino della Società entomologica italiana 121(1): 10–12.
- Alziari G, Lemaire J-M (2012) Les missions entomologiques du Muséum d'Histoire naturelle de Nice (France), 2004–2010. Biocosme Mésogén 29(3–4): 57–104.
- Angelini F (1986) Coleotterofauna del Massiccio del Pollino (Basilicata–Calabria) (Coleoptera). Entomologica 21: 37–125.
- Angelini F (1987) Coleotterofauna del Promontorio del Gargano (Coleoptera). Atti del Museo civico di Storia naturale di Grosseto 11–12: 5–84.
- Angelini F (1991) Coleotterofauna dell'Altipiano della Sila (Calabria, Italia) (Coleoptera). Memorie della Società entomologica italiana 70(1): 171–254.
- Angelini F (1996a) Coleotterofauna della riserva naturale WWF Lago di Pignola. Basilicata, Potenza. Alfagrafica Volonnino Lavello, 135 pp.
- Angelini F (1996b) I Coleotteri in Basilicata. Risorsa Natura in Basilicata. Le Aree protette regionali. Ufficio Stampa Consiglio Reg. Basilicata, Potenza 5–6: 165–170.
- Angelini F (1998) Coleotterofauna reperita mediante trappola luminosa in due stazioni umide della Basilicata (Italia meridionale). Quaderni della Stazione di Ecologia del civico Museo di Storia naturale di Ferrara 11: 7–37.
- Angelini F, Montemurro F (1986) Coleotterofauna del Bosco di Policoro (Matera). Biogeographia, Lavori della Società italiana di Biogeografia 10 (1984): 545–604.
- Audisio P, Liberti G, Nardi G, Poggi R (1995) Coleoptera Polyphaga VIII (Cantharoidea, Dermestoidea). In: Minelli A, Ruffo S, La Posta S (Eds) Checklist delle specie della fauna Italiana, 53. Edizioni Calderini, Bologna, 1–17.
- Bargagli P (1872) Materiali per la fauna entomologica dell'isola di Sardegna. Coleotteri. Bullettino della Società entomologica italiana 4: 97–104.
- Baudi F (1890) Catalogo dei Coleotteri del Piemonte. Annali della reale Accademia d'Agricoltura di Torino 32 (1889): 1–274.
- Beal RS (1998) Taxonomy and Biology of Nearctic Species of *Anthrenus* (Coleoptera: Dermestidae). Transactions of the American Entomological Society 124: 271–332.
- Bertolini S (1874) Catalogo sinonimico e topografico dei Coleotteri d'Italia. Supplemento del Bullettino della Società entomologica italiana 6(1): 93–108.
- Bertolini S (1890) Contribuzione alla fauna trentina dei Coleotteri. Bullettino della Società entomologica italiana 21 (1899): 157–205.

- Bertolini S (1904) Catalogo dei Coleotteri d'Italia. Ed. Rivista italiana di Scienze naturali, Siena (1899–1904), 144 pp.
- Biscaccianti AB, Audisio P, Monguzzi R (2008) Aggiornamenti sulla distribuzione di *Cucujus cinnaberinus* e di altri Cucujoidea (Coleoptera: Nitidulidae, Cucujidae, Laemophloeidae). Bollettino dell'Associazione Romana di Entomologia 63(1–4): 47–57.
- Bordoni A, Rocchi S, Cuoco S (2006) Ricerche sulla Coleottero fauna delle zone umide della Toscana. VI. Piana di Guasticce – Livorno (Coleoptera). Quaderni della Stazione di Ecologia del civico Museo di Storia naturale di Ferrara 16: 43–179.
- Cerretti P, Tagliapietra A, Tisato M, Vanin S, Mason F, Zapparoli M (Eds) (2003) Artropodi dell'orizzonte del faggio nell'Appennino settentrionale, Primo contributo. Conservazione Habitat Invertebrati 2. Gianluigi Arcari Editore, Mantova, 256 pp.
- Cerretti P, Mason F, Minelli A, Nardi G, Whitmore D (Eds) (2009) Research on the Terrestrial Arthropods of Sardinia. Zootaxa 2318: 1–602.
- Cerruti M (1954) Coleoptera. In: Zavattari E (Ed) Biogeografia dell'isola di Zannone (Arcipelago Pontino). Rendiconti dell'Accademia Nazionale dei XL 4(4–5) (76–77) (1953–1954): 108–116.
- Dal Monte G (1972) Generalità sui principi scientifici e tecnici per una razionale conservazione dei cereali nei magazzini. In: Domenichini G (Ed) Atti del 1° Simposio “La difesa antiparassitaria nelle industrie alimentari e la protezione degli alimenti”, Piacenza, 18–20 ottobre 1972. Tipografia Editoriale, Piacentina, Piacenza, 91–107.
- Dalla Torre KW (1911) Coleopterorum Catalogus. Pars 33: Nosodendridae, Byrrhidae, Dermestidae. In: Junk W, Schenkling S (Eds) Coleopterorum Catalogus. W. Junk, Berlin, 1–196.
- Denux O, Zagatti P (2010) Coleoptera families other than Cerambycidae, Curculionidae sensu lato, Chrysomelidae sensu lato and Coccinellidae. Chapter 8.5. In: Roques A, Kenis M, Lees D, Lopez-Vaamonde C, Rabitsch W, Rasplus J-Y, Roy DB (Eds) Alien terrestrial arthropods of Europe. BioRisk 4(1): 315–406.
- Depoli G (1928) I coleotteri della Liburnia. Parte III. Diversicornia. Fiume, Società di Studi Fiumani 6: 208–243.
- Domenichini G, Fogliazza D, Pagani M (1993) Studies on the transmission of *Listeria* by means of arthropods. Italian Journal of Food Science 4(4): 269–278.
- Faggioli D (1956) Campagna di ricerche dell'Istituto di Entomologia dell'Università di Bologna nella "Foresta Umbra" (Gargano). II. Elenco delle specie raccolte (primo lotto). Bollettino dell'Istituto di Entomologia della Università degli Studi di Bologna 21: 167–177.
- Failla-Tedaldi L (1887) Escursione entomologica all'Isola di Lampedusa. Il Naturalista siciliano 6: 157–162.
- Falci A, Livreri Console S, Giardina G, Lo Duca R, La Mantia T (2012) I Lepidotteri Rapaloceri (Insecta Lepidoptera) del Bosco della Ficuzza: stato delle conoscenze e possibili relazioni tra il loro status e i cambiamenti nell'uso del suolo. Il Naturalista siciliano, S. IV 36(3): 435–451.
- Ganglbauer L (1904) Die Käfer von Mitteleuropa. Die Käfer der österreichisch-ungarischen Monarchie, Deutschlands, der Schweiz, sowie des französischen und italienischen Alpengebietes. Vierter Band, erste Hälfte. Dermestidae, Byrrhidae, Nosodendridae, Georyssidae, Dryopidae, Heteroceridae, Hydrophilidae. Karl Gerold's Sohn, Wien, 286 pp.

- Ghiliani V (1842) Insetti di Sicilia determinati dal Sig. F. Ghiliani nel suo viaggio in quest'isola nell'anno 1839. Atti dell'Accademia Gioenia di Scienze naturali di Catania 19: 19–48.
- Ghiliani V (1847) Mémoire sur la station de quelques Coléoptères dans les différentes régions du Piémont. Annales de la Société entomologique de France, Deuxième Série 5: 83–142.
- Ghiliani V (†) (1887) Elenco delle specie di Coleotteri trovate in Piemonte. Opera postuma pubblicata per cura del membro ordinario Dott. Lorenzo Camerano. Annali della reale Accademia d'Agricoltura di Torino 29 (1886): 1–381.
- Giachino PM (1982) Collezione Coleotterologica di Massimiliano Spinola. Cataloghi – Museo Regionale di Scienze Naturali (Torino) 3: 1–616.
- Grandi G (1956) Campagna di ricerche dell'Istituto di Entomologia dell'Università di Bologna alla "Foresta Umbra" (Gargano). I: Ricapitolazione dei risultati ottenuti. Annali. Accademia italiana di Scienze forestali 4: 405–418.
- Gridelli E (1949) Raccolte faunistiche compiute nel Gargano da A. Ghigi e F. P. Pomini. IX. – Coleotteri. Acta – Pontificia Academia Scientiarum 13(18): 145–196.
- Gridelli E (1961) Coleoptera. In: Zavattari E (Ed) Biogeografia delle Isole Pelagie. Rendiconti dell'Accademia nazionale dei XL, ser. IV, 11 (1960): 369–407.
- Grosso-Silva JM, Serrano ARM (2000) Registros interessantes de coleópteros (Insecta, Coleoptera) para Portugal (2<sup>a</sup> nota). Boletín de la Sociedad Entomológica Aragonesa (S.E.A.) 27: 1–5.
- Gulli G (1961) Contributo alla conoscenza dei Coleotteri etnei. Bollettino dell'Associazione Romana di Entomologia 16(1): 1–6.
- Háva J (1999) Contribution to the knowledge of the Old World Dermestidae (Coleoptera) Part. 2: Genus *Dermestes*. Folia Heyrovskyana 7: 141–150.
- Háva J (2003) World Catalogue of the Dermestidae (Coleoptera). Studie a zprávy oblastního muzea Praha – východ v Brandýse nad Labem a Staré Boleslaví, Supplementum 1: 1–196.
- Háva J (2004) World keys to the genera and subgenera of Dermestidae (Coleoptera), with descriptions, nomenclature and distributional records. Acta Musei nationalis Pragae, Series B, natural History 60: 149–164.
- Háva J (2007) Dermestidae (Thorictinae excepted). In: Löbl I, Smetana A (Eds) Catalogue of Palaearctic Coleoptera. Volume 4. Elateroidea – Derodontooidea – Bostrichoidea – Lymexyloidea – Cleroidea – Cucujoidea. Apollo Books, Stenstrup, 57, 299–302, 306–320.
- Háva J (2008) Errata for volume 4 [Dermestidae]. In: Löbl I, Smetana A (Eds) Catalogue of Palaearctic Coleoptera. Volume 5. Tenebrionoidea. Apollo Books, Stenstrup, 25.
- Háva J (2011a) Errata for volume 4 [Dermestidae]. In: Löbl I, Smetana A (Eds) Catalogue of Palaearctic Coleoptera. Volume 6. Chrysomeloidea. Apollo Books, Stenstrup, 25.
- Háva J (2011b) Brouci čeledi kožojedovití (Dermestidae) České a Slovenské republiky. Beetles of the family Dermestidae of the Czech and Slovak Republics. Nakladatelství Academia, Praha, 104 pp. [in Czech and English]
- Háva J (2013) A new synonymy in the myrmecophilous beetle genus *Thorictus* Germar, 1834 (Coleoptera: Dermestidae: Thorictinae). Heteropterus, Revista de Entomología 13(1): 27–29.
- Háva J, Herrmann A (2009) New faunistic records of Dermestidae (Coleoptera) – Part 3. Latvijas Entomolgs 47: 3–5.

- Háva J, Herrmann A (2011) New faunistic records of Dermestidae (Coleoptera) – Part 6. Boletín de la Sociedad Entomológica Aragonesa (S.E.A.) 49: 227–228.
- Háva J, Herrmann A, Kadej M (2010) New faunistic records of Dermestidae (Coleoptera) – Part 5. Latvijas Entomologs 49: 28–31.
- Háva J, Herrmann A, Kadej M (2011) Faunistic contribution to the family Dermestidae (Coleoptera) from Ukraine and Russia. Amurian Zoological Journal III(3): 252–254.
- Háva J, Herrmann A, Kadej M (2013) New faunistic records of Dermestidae Coleoptera) – Part 7. Arquivos Entomológicos 8: 135–140.
- Háva J, Kalík V (2005) Contribution to the *Dermestes peruvianus* species group from the Neotropical region (Coleoptera: Dermestidae). Baltic Journal of Coleopterology 5: 87–98.
- Háva J, Legalov AA (2010) Contribution to the knowledge to the Dermestidae (Coleoptera) from Siberia and adjacent territories. Amurian Zoological Journal 2(4): 294–297.
- Háva J, Nardi G (2004) Preliminary data on the Nosodendridae and Dermestidae (Coleoptera). In: Cerretti P, Hardersen S, Mason F, Nardi G, Tisato M, Zapparoli M (Eds) Invertebrati di una foresta della Pianura Padana, Bosco della Fontana, Secondo contributo. Conservazione Habitat Invertebrati, 3. Cierre Grafica Editore, Verona, 117–123.
- Háva J, Nardi G (2007) Notes on some *Attagenus* from Italy (Insecta, Coleoptera: Dermestidae). Aldrovandia 3: 121–124.
- Háva J, Nardi G (2011) First contribution to the knowledge of the Dermestidae of Sardinia (Coleoptera). In: Nardi G, Whitmore D, Bardiani M, Birtele D, Mason F, Spada L, Cerretti P (Eds) Biodiversity of Marganai and Montimannu (Sardinia). Research in the framework of the ICP Forests network. Conservazione Habitat Invertebrati 5. Cierre Edizioni, Sommacampagna, Verona, 407–446.
- Háva J, Zahradník P (2011) *Anthrenus (Anthrenus) angustefasciatus* Ganglbauer, 1904 (Coleoptera: Dermestidae), in the Czech Republic. Elateridarium 5: 1–4.
- Henry J (1938) Autre contribution à l'étude des Coléoptères de la Vallée d'Aoste. Bulletin de la Société de la Flore Valdôtainne 23: 65–71.
- Herrmann A, Bahillo De La Puebla P (2003) Sobre la presencia de *Dermestes (Dermestinus) hankae* Háva, 1999 y *Dermestes (Dermestinus) aurichalceus* Küster, 1846 (Coleoptera, Dermestidae) en la Península Ibérica. Boletín de la Sociedad Entomológica Aragonesa (S.E.A.) 33: 91–93.
- Herrmann A, Háva J (2007) The first record of the Australian immigrant *Anthrenocerus australis* (Hope, 1843) (Coleoptera: Dermestidae) from Italy. Gredleriana 7: 412–413.
- Herrmann A, Háva J, Kadej M (2011) A new species of *Globicornis* Latreille in Cuvier (Coleoptera: Dermestidae: Megatominae) from Switzerland. Studies and Reports, Taxonomical Series 7: 139–144.
- Holdhaus K (1911) Über die Coleopteren- und Molluskenfauna des Monte Gargano (Unter besonderer berücksichtigung der Adriatisfrage). Denkschriften der Mathematisch-Naturwissenschaftlichen Klasse der Kaiserlichen Akademie der Wissenschaften 87: 431–465.
- Holdhaus K (1923) Elenco dei Coleotteri dell'Isola d'Elba con studii sul problema della Tirreneide. Memorie della Società entomologica italiana 2: 77–175.
- Horion AD (1955) Faunistik der Mitteleuropäischen Käfer. Band IV: Sternoxia (Buprestidae), Fossilipedes, Macrodactylia, Brachymera. Entomologische Arbater Museum Frey München. Sonderband, Tutzing, I–XXI + 249 pp.

- ICZN (International Commission on Zoological Nomenclature) (1999) International Code of Zoological Nomenclature. Fourth edition adopted by the International Union of Biological Sciences. The International Trust for Zoological Nomenclature, London, I–XXIX + 306 pp.
- Iehl F (1909) Contribution à la Faune des Coléoptères valdôtains. Bulletin. Société de la Flore Valdôtaine 5: 25–31.
- John H (1963) Revision der Gattung *Thorictus* Germar (Thorictidae, Clavicornia Col.). Eine Bestandaufnahme, Kritik und Neubeschreibungen. Entomologische Abhandlungen und Berichte aus dem Staatlichen Museum für Tierkunde in Dresden 28: 177–255.
- John H (1965) Revision der Gattung *Thorictus* Germar (Thorictidae, Clavicornia, Coleoptera). Eine Bestandaufnahme, Kritik und Neubeschreibungen. II. Teil. Entomologische Abhandlungen und Berichte aus dem Staatlichen Museum für Tierkunde in Dresden 31: 319–395.
- John H (1966) Neue Spezies von *Thorictus* Germ. (Col., Thorictidae). The Entomologist's Monthly Magazine 102: 19–21.
- Kadej M (2005) Morphological variability of *pimpinellae*-group (Coleoptera: Dermestidae: *Anthrenus*) with a key to identification. Part 1. Polskie Pismo Entomologiczne 74: 117–135.
- Kadej M, Háva J, Kalík V (2007) Review of the *Anthrenus pimpinellae* species group from Palaearctic region (Coleoptera: Dermestidae: Anthrenini). Genus 18(4): 721–750.
- Kenis M (2005) 4 Insects – Insecta. In: Wittenberg R (Ed) An inventory of alien species and their threat to biodiversity and economy in Switzerland. CABI Bioscience Switzerland Centre report to the Swiss Agency for Environment, Forests and Landscape, Delémont, 131–212.
- Kraatz G (1858) Beitrag zur Käferfauna Griechenlands. Dritter Stück: Staphylinidae (Schluss), Trichopterygia, Histeridae, Phalacridae, Nitidulariae, Trogositarii, Colydi, Cucujidae, Cryptophagidae, Thorictidae, Mycetophagidae, Dermestini, Byrrhini. Berliner entomologische Zeitschrift 2: 123–148.
- La Mantia T, Bellavista M, Giardina G, Sparacio I (2010) Longhorn beetles of the Ficuzza woods (W Sicily, Italy) and their relationship with plant diversity (Coleoptera, Cerambycidae). Biodiversity Journal 1(1–4): 15–44.
- Liebmann W (1962) Ein Beitrag zur Käferfauna von Pantelleria. Stuttgarter Beiträge zur Naturkunde aus dem Staatlichen Museum für Naturkunde in Stuttgart 87: 1–6.
- Lo Cascio P, Cecchi B, Abbazzi P, Arnone M (2006) A contribution to the knowledge of the Coleoptera of the Aeolian islands. Il Naturalista siciliano, S. IV 30(2): 315–341.
- Löbl I (2007) Dermestidae Thorictinae. In: Löbl I, Smetana A (Eds) Catalogue of Palaearctic Coleoptera. Volume 4. Elateroidea – Derodontoidea – Bostrichoidea – Lymexyloidea – Cleroidea – Cucuoidea. Apollo Books, Stenstrup, 302–306.
- Lohse GA (1979) 45. Familie: Dermestidae. In: Freude H, Harde KW, Lohse GA (Eds) Die Käfer Mitteleuropas. Bd. 6., Diversicornia. Goecke & Evers, Krefeld, 304–327.
- Luigioni P (1923a) Contributo allo studio della fauna entomologica Italiana. Coleotteri nuovi o poco noti per il Lazio. VI Elenco. Atti della Pontificia Accademia Romana dei Nuovi Lincei 76 (1922–1923): 132–145.
- Luigioni P (1923b) Contributo allo studio della fauna entomologica italiana. Coleotteri dell'isola di Capri. Annuario del Museo Zoologico della R. Università di Napoli (nuova serie) 5(6): 1–8.
- Luigioni P (1929) I Coleotteri d'Italia. Catalogo sinonimico-topografico-bibliografico. Memorie della Pontificia Accademia delle Scienze, I Nuovi Lincei (2) 13: [4+] 1–1159 [+ 1].

- Marcuzzi G (1985) Ulteriori dati per una conoscenza della Coleottero fauna pugliese. *Quaderni di Ecologia animale* 23: 5–21.
- Mason F, Cerretti P, Nardi G, Whitmore D, Birtele D, Hardersen S, Gatti E (2006) Aspects of biological diversity in the CONECOFOR plots. IV. The InvertebrateBiodiv pilot project. In: Ferretti M, Petriccione B, Bussotti F, Fabbio G (Eds) *Aspects of biodiversity in selected forest ecosystems in Italy: status and changes over the period 1996–2003*. Third report of the Task Force on Integrated and Combined (I&C) evaluation of the CONECOFOR programme. *Annali dell'Istituto Sperimentale per la Selvicoltura* 30, Supplemento 2: 51–70.
- Merz B (1993) Fruchtfliegen aus der Region des Mte. Gargano (Italien, Puglia), mit Beschreibung einer neuen *Tephritis*-Art (Diptera, Tephritidae). *Mitteilungen der Entomologischen Gesellschaft Basel* 43(3): 110–127.
- Minelli A, Chemini C, Argano R, Ruffo S (Eds) (2002) *La Fauna in Italia*. Touring Editore, Milano e Ministero dell'Ambiente e della Tutela del Territorio, Roma, 448 pp.
- Minelli A, Negrisolo E (1993) Ricerche zoologiche nell'alto Trevigiano II. Materiali faunistici (Annelidae, Arthropoda). *Lavori – Società Veneziana di Scienze naturali* 18: 59–132.
- Mroczkowski M (1958) Dermestidae (Coleoptera) collected in Bulgaria by members of the staff on the Zoological Institute of the Polish Academy of Sciences in the years 1950 and 1956. *Fragmenta Faunistica* 8(1): 1–9.
- Mroczkowski M (1968) Distribution of the Dermestidae (Coleoptera) of the world with a catalogue of all known species. *Annales Zoologici* 26: 15–191.
- Mulsant E, Rey C (1868) *Histoire Naturelle des Coléoptères de France. Tribu des Scuticolles*. F. Savy, Paris, 186 pp., 3 pls. [reprint from *Annales de la Société Linnéenne de Lyon* (N.S.), 15: 1–188, 3 pls.]
- Nardi G (1997) Coleoptera Dermestidae, Lyctidae, Bostrichidae, Ptinidae. In: Zapparoli M (Ed) *Gli Insetti di Roma. Comune di Roma, Dipartimento X Area Risorsa Suolo e Tutela Ambientale*, Quaderni dell'Ambiente 6, 177–179.
- Nardi G, Vomero V (Eds) (2007) *Artropodi del Parco Nazionale del Vesuvio: ricerche preliminari. Conservazione Habitat Invertebrati*, 4. Cierre Edizioni, Verona, 496 pp.
- Nardi G, Whitmore D, Bardiani M, Birtele D, Mason F, Spada L, Cerretti P (Eds) (2011) Biodiversity of Marganai and Montimannu (Sardinia). Research in the framework of the ICP Forests network. *Conservazione Habitat Invertebrati* 5. Cierre Edizioni, Sommacampagna, Verona, 896 pp., 1 map.
- Nardi R, Villani V, Infantino S, Meio S (1993) Risultati di difesa integrata in una industria alimentare. In: Domenichini G (Ed) *Atti del 5° Simposio “La difesa antiparassitaria nelle industrie alimentari e la protezione degli alimenti”*, Piacenza, 23–25 settembre 1992. Chirilli Editori, Pinerolo, 617–627.
- Nicoli Aldini R (1998) Osservazioni morfologiche e biologiche su *Trogoderma variabile* Ballion (Coleoptera Dermestidae), nuovo infestante di derrate alimentari in Italia. In: Cravedi P (Ed) *Atti del 6° Simposio “La difesa antiparassitaria nelle industrie alimentari e la protezione degli alimenti”*, Piacenza, 24–26 settembre 1997. Chirilli editori, Pinerolo, 115–126.
- Nicoli Aldini R (1999) Un Coleottero Dermestide inquilino dell'alveare, *Trogoderma inclusum* Leconte. *L'APE, Nostra Amica* 21: 6–13.

- Nicoli Aldini R (2003a) Catture con trappole luminose in un'industria di spezie e primo ritrovamento in Italia di *Attagenus fasciatus* (Thunberg) e *Reesa vespulae* (Milliron) (Coleoptera Dermestidae). Bollettino di Zoologia agraria e Bachicoltura 35: 153–162.
- Nicoli Aldini R (2003b) Insetti delle derrate e dell'ambiente antropico di recente introduzione in Italia. In: Cravedi P (Ed) Atti del 7° Simposio “La difesa antiparassitaria nelle industrie alimentari e la protezione degli alimenti”, Piacenza, 18–20 settembre 2002. Chiriotti Editori, Pinerolo, 118–143.
- Nicoli Aldini R (2004) Insetti delle derrate e dell'ambiente antropico di recente introduzione in Italia. Tecnica Molitoria 55(10): 977–999.
- Novak P (1952) Kornjaši jadranskog primorja (Coleoptera). Jugoslavenska Akademija Znanosti i Umjetnosti, Split, 524 pp.
- Osella G (1988) Gli Invertebrati. In: AA.VV., Collana Verde, 76. La riserva naturale valle dell'Orfento. Ministero dell'Agricoltura e delle Foreste, Corpo Forestale dello Stato, Roma, 62–68.
- Palm T (1939) En exkursionsdag på Monte Pollino. Entomologbladet 3: 58–69.
- Panganetti-Hummel [sic! = Paganetti-Hummler] G (1918) Beiträge zur Coleopterenfauna Italiens. Murgien. Neue Beiträge zur Systematischen Insektenkunde, Beilage zur Zeitschrift für wissenschaftliche Insektenbiologie 1: 85–88.
- Plata-Negrache P (1972) Las especie de la subfamilia Dermestinae Latr. (Col., Dermestidae), en el Archipelago Canario. Vieraea 2: 147–163.
- Pollini A (1998) Manuale di entomologia applicata. Edagricole, Bologna, I–X + 1426 pp.
- Porta A (1929) Fauna Coleopterorum Italica. III. Diversicornia. Stabilimento Tipografico Piacentino, Piacenza, 465 pp.
- Porta A (1934) Fauna Coleopterorum Italica. Supplementum. Stabilimento Tipografico Piacentino, Piacenza, I–VII + 208 pp.
- Prieto M, Háva J (2013) Dos nuevos *Dermestes* Linnaeus, 1758 para la fauna de Cataluña (NE Península Ibérica) (Coleoptera: Dermestidae: Dermestinae). Arquivos Entomológicos 8: 241–244.
- Ragusa E (1892) Catalogo ragionato dei Coleotteri di Sicilia. Il Naturalista siciliano 11(9–11): 185–209.
- Reiche ML (1860) Coléoptères de Sicilie récueillis par M. E. Bellier de la Chavignerie et déterminés par M. L. Reiche et description de dix espèces nouvelles. Annales de la Société Entomologique de France, ser. 3, 8: 717–722.
- Reitter E (1878) *Hadrotoma breviclavis* n. sp. In: Schneider O, Leder H (Ed) Beiträge zur Kenntniss der kaukasischen Käferfauna. Verhandlungen Naturforschenden Vereines Brünn 16: 179.
- Reitter E (1881) Bestimmungs-Tabellen der europäischen Coleopteren. III. Enthaltend die Familien: Scaphidiidae, Lathridiidae und Dermestidae. Verhandlungen der Kaiserlich-Königlichen Zoologisch-Botanischen Gesellschaft in Wien 30: 41–94.
- Reitter E (1883) Dermestidae. In: Heyden (von) L, Reitter E, Weise J (Eds) Catalogus Coleopterorum Europae et Caucasi. Editio Tertia. London, Berlin, Paris, 89–90.
- Reitter E (1887) Bestimmungs-Tabellen der europäischen Coleopteren. III Heft. Enthaltend die Familien: Scaphidiidae, Lathridiidae und Dermestidae. II. Auflage. Verlag von Edmund Reitter (1886), Mödling, 75 pp.
- Reitter E (1891) Dermestidae. In: Heyden (von) L, Reitter E, Weise J (Eds) Catalogus Coleopterorum Europeae, Caucasi et Armeniae Rossicae. Friedländer & Sohn, Berlin, 336–340.

- Reitter E (1906) Dermestidae. In: Heyden (von) L, Reitter E, Weise J (Eds) Catalogus Coleopterorum Europeae, Caucasi et Armeniae Rossicae. Editio secunda. Friedländer & Sohn, Berlin, 377–380.
- Rivosecchi L (1989) Note sugli Sciomyzidae IX. Due nuove specie del genere *Pherbellia* provenienti dall'Italia meridionale (Diptera, Cyclorrhapha, Acalyptera). Fragmenta entomologica 21(2): 153–161.
- Rossi P (1794) Mantissa Insectorum exhibens species nuper in Etruria, adiectis faunae Etruscae illustrationibus, ac emendationibus. Tom II. Typographia Prosperi, Pisis, 154 pp., 8 pls.
- Rottenberg A (1870) Beiträge zur Coleopteren-Fauna von Sicilien. (Zweites Stück.). Berliner entomologische Zeitschrift 14: 235–260, pl. II.
- Roubal J (1932) Addition à la liste des Coléoptères de Monte Gargano. Bolletino della Società entomologica italiana 64(4): 66.
- Ruffo S, Stoch F (Eds) (2006) Checklist and distribution of the Italian fauna. 10,000 terrestrial and inland water species. Memorie del Museo Civico di Storia Naturale di Verona, 2. serie, Sezione Scienze della Vita 17: 1–304, with data on CD-ROM.
- Rusek J, Stumpp J (1988) A new *Acerentulus* species from Southeast Italy (Protura: Acerentomidae). Acta entomologica bohemoslovaca 85: 153–158.
- Sabella G, Sparacio I (2004) Il ruolo dei Parchi siciliani nella conservazione di taxa di Insetti di particolare interesse naturalistico (Insecta Coleoptera et Lepidoptera Rhopalocera). Il Naturalista siciliano, S. IV 28(1): 447–508.
- Sainte-Claire Deville J (1937) (†) Catalogue raisonné des Coléoptères de France. (Complété et publié par A. Méquignon). L'Abeille 36(3): 265–372.
- Sapuppo G (2002) Contributo alla conoscenza dei Coleotteri della zona etnea. Collezione Gulli – Sapuppo. Atti e Memorie dell'Ente Fauna siciliana 8: 35–280.
- Schatzmayr A (1941) Coleotteri raccolti nella Grande Sila dal dott. E. Moltoni. Atti della Società italiana di Scienze naturali 80: 49–102.
- Sparacio I (1997) Coleotteri di Sicilia Parte II. Meditarraneo, Guide Naturalistiche, 4. L'Epos Ed., Palermo, 208 pp.
- Steck T (1887) Entomologie Ergebnisse einer Excursion nach Sizilien. Mittheilungen der Naturforschenden Gesellschaft in Bern 1143–1168 (1886): 178–188.
- Trematerra P (1987) Annotazioni biologiche e morfologiche su *Attagenus alfierii* Pic (Coleoptera Dermestidae) nuovo per la fauna italiana. Disinfestazione 4(6): 16–18.
- Vigna Taglianti A, Audisio PA, Belfiore C, Biondi M, Bologna MA, Carpaneto GM, De Biase A, De Felici S, Piattella E, Racheli T, Zapparoli M, Zoia S (1993) Riflessioni di gruppo sui corotipi fondamentali della fauna W-paleartica ed in particolare italiana. Biogeographia, Lavori della Società italiana di Biogeografia (n.s.) 16 (1992): 159–179.
- Vigna Taglianti A, Audisio PA, Biondi M, Bologna MA, Carpaneto GM, De Biase A, Fattorini S, Piattella E, Sindaco R, Venchi A, Zapparoli M (1999) A proposal for a chorotype classification of the Near East fauna, in the framework of the Western Palearctic region. Biogeographia, Lavori della Società italiana di Biogeografia (n.s.) 20: 31–59.
- Villa A, Villa GB (1844) Catalogo dei Coleotteri della Lombardia. In: Cattaneo C (Ed) Notizie naturali e civili sulla Lombardia, vol. 1. Coi tipi di Giuseppe Berdaroni di Giovanni, Milano, 77 pp. [reprint]. [In reality, this title is not present in the book; this reprint assem-

- ble two paragraphs of the chapter VIII of the book: "V. Insetti della Lombardia", 406–415 and "Catalogo degli Insetti Coleopteri della Lombardia", 416–477.]
- Vitale F (1904) I Coleotteri Messinesi (continuazione). Bollettino del Naturalista 24(8): 74–76.
- Winkler A (1926) Pars. 6. Dermestidae. In: Winkler A (Ed) Catalogus Coleopterorum Regionis Palaearcticae. A. Winkler, Wien, 1924–1932, 675–682.
- Zangheri P (1969) Repertorio sistematico e topografico della flora e fauna vivente e fossile della Romagna. Tomo III. Memorie fuori serie del Museo Civico di Storia Naturale di Verona 1: 855–1414.
- Zhantiev RD (2001) Zhuki-kozheedy roda *Orphilus* Er. (Coleoptera, Dermestidae) fauny Palearktiki. Entomologicheskoe Obozrenie 80: 611–620. [in Russian, English summary]
- Zhantiev RD (2009) New and little-Known dermestid beetles (Coleoptera, Dermestidae) from the Caucasus. Entomological Review 89(7): 874–877. doi: 10.1134/S0013873809070112
- Zhantiev RD (2011a) Fauna Europaea: Dermestidae. In: Audisio P (Ed) Fauna Europaea: Coleoptera 2. Fauna Europaea Version 2.4. <http://www.faunaeur.org> [accessed 29 March 2012 as version 2.4 of 27 January 2011]
- Zhantiev RD (2011b) New and Little Known Dermestid Beetles of the Genus *Thorictus* (Coleoptera, Dermestidae). Entomological Review 91(8): 1005–1010. doi: 10.1134/S0013873811080069
- Zilli A (2000) Lepidoptera species with a western and eastern distribution in the Mediterranean: reasons for a disjunction. Biogeographia, Lavori della Società italiana di Biogeografia (n.s.) 20 (1999): 223–228.
- Zinetti F, Terzani F (2009) Coleottero fauna di due Riserve Naturali della Provincia di Arezzo (Toscana) con particolare riguardo alle loro zone umide (Insecta, Coleoptera). Quaderni della Stazione di Ecologia del civico Museo di Storia naturale di Ferrara 19: 5–48.