

Two new species of Trichoceridae from the Middle Jurassic Jiulongshan Formation of Inner Mongolia, China

Fei Dong^{1,†}, Chungkun Shih^{2,‡}, Dong Ren^{1,§}

¹ College of Life Sciences, Capital Normal University, 105 Xisanhuanbeilu, Haidian District, Beijing 100048, China

† <http://zoobank.org/C2C462B8-8597-4F58-9F82-4FF511BD6BD6>

‡ <http://zoobank.org/A49AAC84-569A-4C94-92A1-822E14C97B62>

§ <http://zoobank.org/A35337D7-47AC-4F05-9E2B-2EFE05117BF0>

Corresponding author: Dong Ren (rendong@mail.cnu.edu.cn)

Academic editor: Jes Rust | Received 19 December 2013 | Accepted 7 May 2014 | Published 28 May 2014

<http://zoobank.org/E80CF090-FD5E-4F24-9E88-ED3B04ECD603>

Citation: Dong F, Shih C-K, Ren D (2014) Two new species of Trichoceridae from the Middle Jurassic Jiulongshan Formation of Inner Mongolia, China. ZooKeys 411: 145–160. doi: 10.3897/zookeys.411.6858

Abstract

Two new species, *Eotrichocera (Archaeotrichocera) longensis* sp. n. and *Eotrichocera (Archaeotrichocera) amabilis* sp. n. of Trichoceridae are described based on a combination of the following characters: Sc ending proximad of the forking of R₂, shape of d cell and A₂ rather short and bending sharply toward posterior margin. These fossil specimens were collected from the Middle Jurassic Jiulongshan Formation of Daohugou in Inner Mongolia, China.

Keywords

Diptera, *Eotrichocera*, *Archaeotrichocera*, Fossil, Daohugou

Introduction

Trichoceridae is a family of medium-sized dipterans, commonly called winter crane flies, Yang and Yang (1995) found that just a few adults can live in cold environment, even in winter, and indicated that the name of winter crane flies might not be proper. However, the adults (include the largest species of *Trichocera*) not only live in cold environment, but also mate and lay eggs under the snow cover in winter (Hågvar and Krzemińska 2007). Hence, the common name of winter crane flies is proper. The adults live in damp places close to lakes, rivers, or streams and most of them feed on plant fluids (Yang 2009); while the larvae live in moist or wet or terrestrial biotopes and feed on plant debris, decaying leaves in forests, mushrooms and animal droppings (Dahl and Alexander 1976) or take cankered plants or animal bodies as food (Yang 2009).

There are 77 species of fossil and amber trichocerids, which have been assigned into three subfamilies: Trichocerinae, Paracladurinae and Kovalevinae; and twelve genera: *Cladoneura* Scudder, 1894; *Eotrichocera* Kalugina, 1985; *Rasnitsynina* Krzemińska, Krzemiński & Dahl, 2009; *Mailotrichocera* Kalugina, 1985; *Paleotrichocera* Kalugina, 1986; *Karatina* Krzemińska, Krzemiński, Dahl & Lukashevich, 2009; *Trichocera* Meigen, 1803; *Tanyochoreta* Zhang, 2006; *Zherikhinina* Krzemińska, Krzemiński & Dahl, 2009; *Undaya* Krzemińska, Krzemiński & Dahl, 2009; *Kovaleva* Krzemińska, Krzemiński & Dahl, 2009; *Paracladura* Brunetti, 1911 (Krzemińska et al. 2009a). The oldest species of trichocerids, *Mailotrichocera variabilis*, *M. mikereichi* and *M. zessini* have been described from Lower Jurassic of Germany (Krzemińska et al. 2009a).

Among them, there are eight species in three genera described from the Daohugou locality of China: *Eotrichocera* (*Archaeotrichocera*) *ephemera* Zhang, 2006; *Tanyochoreta* *integera* Zhang, 2006; *Tanyochoreta* *chifengica* Zhang, 2006; *Tanyochoreta* (*Sinotrichocera*) *parva* Zhang, 2006; *Eotrichocera* (*Archaeotrichocera*) *conica* Krzemińska, Krzemiński & Ren, 2009; *Eotrichocera* (*Archaeotrichocera*) *rara* Krzemińska, Krzemiński & Ren, 2009; *Eotrichocera* (*Archaeotrichocera*) *spatiosa* Liu, Shih & Ren, 2012 and *Zherikhinina* *reni* Krzemińska, Krzemiński & Dahl, 2009.

Furthermore, *Sinotrichocera* Zhang, 2006 has been changed as a subgenus belonging to *Tanyochoreta*; *Oligotrichocera* Dahl, 1971 as a subgenus belonging to *Trichocera* Podenas, 2001; *Trichonomites* Kalugina, 1986 and *Paleotrichocera* Kalugina, 1986 are synonymized (Krzemińska et al. 2009a). All genera and species of Trichoceridae Kertész, 1902, after revisions and transfers, are summarized in Table 1, which is updated and expanded from the Tables 1 and 4 in Krzemińska et al. 2009a.

The specimens for this study were collected from the Jiulongshan Formation of the Daohugou Village in Inner Mongolia, China. The Daohugou fossil-bearing beds are considered as the late Middle Jurassic (Bathonian-Callovian boundary, 165 Mya) (Ren et al. 2002; Gao and Ren 2006; Ren et al. 2010a; Shi et al. 2011). Daohugou is one of the localities where the fossils of Yanliao biota were distributed. A huge number of fossil insects have been reported (Ren and Engel 2007; Engel and Ren 2008; Liu and Ren 2008; Ren et al. 2009; Wang and Ren 2009; Gu et al. 2010; Ren et al. 2010b; Wang et al. 2010; Wang et al. 2012; Yang et al. 2012).

Table I. Fossil species of Trichoceridae Kertész, 1902.

Genus	Species	Author(s)	Date	Age	Locality
<i>Cladoneura</i>	<i>C. willistoni</i>	Scudder	1894	Lower Oligocene	Florissant, USA
<i>Eotrichocera</i> (<i>Archaeotrichocera</i>)	<i>E. (A.) ephemera</i>	Zhang	2006	Middle Jurassic	Daohugou, China
	<i>E. (A.) conica</i>	Krzemińska, Krzemiński & Ren	2009a	Middle Jurassic	Daohugou, China
	<i>E. (A.) rara</i>	Krzemińska, Krzemiński & Ren	2009a	Lower Cretaceous	Kempendyay, Russia
	<i>E. (A.) spatiosa</i>	Liu, Shih & Ren	2012a	Middle Jurassic	Daohugou, China
<i>Eotrichocera</i> (<i>Eotrichocera</i>)	<i>E. (E.) christinae</i>	Kalugina	1985	Lower Jurassic or earlier Middle Jurassic	Novospasskoe, Russia
<i>Karatina</i>	<i>K. longipes</i>	Rohdendorf	1964	Upper Jurassic	Karatau, Kazakhstan
	<i>K. explorans</i>	Krzemińska, Krzemiński, Dahl & Lukashevich	2009a	Lower Cretaceous	Baissa, Russia
	<i>K. pellita</i>	Krzemińska, Krzemiński, Dahl & Lukashevich	2009a	Upper Jurassic	Karatau, Kazakhstan
<i>Kovaleva</i> (<i>Kovaleva</i>)	<i>K. (K.) fragmentosa</i>	Krzemińska, Krzemiński & Dahl	2009a	Jurassic/Cretaceous boundary	Unda and Daya, E. Transbaikalia
	<i>K. (K.) hirsuta</i>	Krzemińska, Krzemiński & Dahl	2009a	Jurassic/Cretaceous boundary	Daya, E. Transbaikalia
	<i>K. (K.) obscura</i>	Krzemińska, Krzemiński & Dahl	2009a	Jurassic/Cretaceous boundary	Daya, Unda and Shevia, E. Transbaikalia
	<i>K. (K.) sheviae</i>	Krzemińska, Krzemiński & Dahl	2009a	Jurassic/Cretaceous boundary	Shevia, E. Transbaikalia
	<i>K. (K.) volodii</i>	Krzemińska, Krzemiński & Dahl	2009a	Jurassic/Cretaceous boundary	Daya, E. Transbaikalia
<i>Kovaleva</i> (<i>Vladimirevna</i>)	<i>K. (V.) mirabilis</i>	Krzemińska, Krzemiński & Dahl	2009a	Jurassic/Cretaceous boundary	Daya, E. Transbaikalia
<i>Mailotrichocera</i>	<i>M. jurassica</i>	Kalugina	1985	Uppermost middle or earliest Upper Jurassic	Uda, E. Transbaikalia
	<i>M. gracilis</i>	Krzemińska, Krzemiński & Dahl	2009a	Jurassic/Cretaceous boundary	Daya, Unda and Shevia, E. Transbaikalia
	<i>M. mikereichi</i>	Krzemińska, Krzemiński & Ansorge	2009a	Lower Jurassic	Dobbertin, Germany; Grimmen, Germany
	<i>M. ovifera</i>	Krzemińska, Krzemiński & Dahl	2009a	Jurassic/ Cretaceous boundary	Unda, E. Transbaikalia
	<i>M. prisca</i>	Krzemińska, Krzemiński & Dahl	2009a	Jurassic/Cretaceous boundary	Unda and Shevia, E. Transbaikalia
	<i>M. sukachevae</i>	Krzemińska, Krzemiński & Dahl	2009a	Jurassic/Cretaceous boundary	Unda, E. Transbaikalia
	<i>M. variabilis</i>	Krzemińska, Krzemiński & Ansorge	2009a	Lower Jurassic	Dobbertin, Germany; Grimmen, Germany
	<i>M. zessini</i>	Krzemińska, Krzemiński & Ansorge	2009a	Lower Jurassic	Grimmen, Germany
<i>Paleotrichocera</i>	<i>P. mongolica</i>	Kalugina	1986	Lower Cretaceous	Gurvan Erenyi Nuru, Mongolia
<i>Panacladura</i>	<i>P. caucasiana</i>	Krzemińska, Krzemiński & Dahl	2009a	Middle Miocene	Stavropol, Caucasus

Genus	Species	Author(s)	Date	Age	Locality
<i>Rasnitsynina</i>	<i>R. collecta</i>	Krzemińska, Krzemiński & Dahl	2009a	Jurassic/Cretaceous boundary	Shevia, E. Transbaikalia
	<i>R. minae</i>	Krzemińska, Krzemiński & Dahl	2009a	Jurassic/Cretaceous boundary	Shevia and Daya, E Transbaikalia
<i>Tanyochoreta</i> (<i>Sinotrichocera</i>)	<i>T. (S.) parva</i>	Zhang	2006	Middle Jurassic	Daohugou, China
<i>Tanyochoreta</i> (<i>Tanyochoreta</i>)	<i>T. (T.) chifengica</i>	Zhang	2006	Middle Jurassic	Daohugou, China
	<i>T. (T.) integera</i>	Zhang	2006	Middle Jurassic	Daohugou, China
<i>Tanyochoreta</i> (<i>Trichokara</i>)	<i>T. (T.) composita</i>	Krzemińska, Krzemiński & Dahl	2009a	Upper Jurassic	Karatau, Kazakhstan
	<i>T. (T.) fracta</i>	Krzemińska, Krzemiński & Dahl	2009a	Upper Jurassic	Karatau, Kazakhstan
	<i>T. (T.) minuta</i>	Krzemińska, Krzemiński & Dahl	2009a	Upper Jurassic	Karatau, Kazakhstan
	<i>T. (T.) tenuis</i>	Krzemińska, Krzemiński & Dahl	2009a	Upper Jurassic	Karatau, Kazakhstan
	<i>T. (T.) zagadka</i>	Krzemińska, Krzemiński & Dahl	2009a	Jurassic/Cretaceous boundary	Unda, E Transbaikalia
	<i>T. (T.) zbulwami</i>	Krzemińska, Krzemiński & Dahl	2009a	Jurassic/Cretaceous boundary	Daya and Unda, E Transbaikalia
<i>Trichocera</i>	<i>T. scudderri</i>	Meunier	1915	Upper Oligocene	Rott, Germany
	<i>T. miocaenica</i>	Statz	1934	Upper Oligocene	Rott, Germany
	<i>T. antiqua</i>	Dahl	1971	Upper Eocene	Baltic
	<i>T. primaeva</i>	Dahl	1971	Upper Eocene	Baltic
	<i>T. fujiyamai</i>	Gentilini	1984	Upper Miocene	Monte Castellaro, Italy
	<i>T. anbar</i>	Podenas	2001	Upper Eocene	Baltic
	<i>T. bona</i>	Podenas	2001	Upper Eocene	Baltic
	<i>T. cerea</i>	Podenas	2001	Upper Eocene	Baltic
	<i>T. diluta</i>	Podenas	2001	Upper Eocene	Baltic
	<i>T. ebenos</i>	Podenas	2001	Upper Eocene	Baltic
	<i>T. christelae</i>	Krzemińska, Krzemiński & Dahl	2009a	Upper Eocene	Baltic
<i>Undaya</i>	<i>U. conami</i>	Krzemińska, Krzemiński & Dahl	2009a	Lower Cretaceous	Purbeck, UK
	<i>U. cretacea</i>	Krzemińska, Krzemiński & Dahl	2009a	Lower Cretaceous	Baissa, Russia
	<i>U. hanswernerri</i>	Krzemińska, Krzemiński & Dahl	2009a	Upper Eocene	Baltic
	<i>U. turgana</i>	Krzemińska, Krzemiński & Dahl	2009a	earlier Lower Cretaceous	Turga, E. Transbaikalia
	<i>U. alata</i>	Krzemińska, Krzemiński & Dahl	2009a	Jurassic/Cretaceous boundary	Unda and Shevia, E. Transbaikalia
	<i>U. comis</i>	Krzemińska, Krzemiński & Dahl	2009a	Jurassic/Cretaceous boundary	Unda, E. Transbaikalia
	<i>U. gargantuina</i>	Krzemińska, Krzemiński & Dahl	2009a	Jurassic/Cretaceous boundary	Daya and Unda, E. Transbaikalia
	<i>U. hilara</i>	Krzemińska, Krzemiński & Dahl	2009a	Jurassic/Cretaceous boundary	Unda and Shevia, E. Transbaikalia
	<i>U. kaluginae</i>	Krzemińska, Krzemiński & Dahl	2009a	Jurassic/Cretaceous boundary	Daya, E. Transbaikalia
	<i>U. lenae</i>	Krzemińska, Krzemiński & Dahl	2009a	Upper Jurassic	Shar-Teg, Mongolia

Genus	Species	Author(s)	Date	Age	Locality
<i>U.</i>	<i>U. lukashevichae</i>	Krzemińska, Krzemiński & Dahl	2009a	Upper Jurassic	Shar-Teg, Mongolia
	<i>U. maxima</i>	Krzemińska, Krzemiński & Dahl	2009a	Lower Cretaceous	Kempendyay, Russia
	<i>U. mitis</i>	Krzemińska, Krzemiński & Dahl	2009a	Jurassic/Cretaceous boundary	Daya and Unda, E. Transbaikalia
	<i>U. molesta</i>	Krzemińska, Krzemiński & Dahl	2009a	Jurassic/Cretaceous boundary	Unda and Daya, E. Transbaikalia
	<i>U. namdyriensis</i>	Krzemińska, Krzemiński & Dahl	2009a	Lower Cretaceous	Kempendyay, Russia
	<i>U. parvula</i>	Krzemińska, Krzemiński & Dahl	2009a	Jurassic/Cretaceous boundary	Daya and Unda, E. Transbaikalia
	<i>U. pura</i>	Krzemińska, Krzemiński & Dahl	2009a	Jurassic/Cretaceous boundary	Unda and Daya, E. Transbaikalia
	<i>U. salsa</i>	Krzemińska, Krzemiński & Dahl	2009a	Lower Cretaceous	Kempendyay, Russia
	<i>U. savina</i>	Krzemińska, Krzemiński & Dahl	2009a	Jurassic/Cretaceous boundary	Savina, E. Transbaikalia
	<i>U. saxeae</i>	Krzemińska, Krzemiński & Dahl	2009a	Lower Cretaceous	Kempendyay, Russia
	<i>U. triangula</i>	Krzemińska, Krzemiński & Dahl	2009a	Lower Cretaceous	Kempendyay, Russia
Zherikhinina	<i>Z. itatica</i>	Kalugina	1985	Middle Jurassic	Kubekovo, Russia
	<i>Z. alastos</i>	Krzemińska & Lukashevich	2009b	Upper Jurassic	Shar Teg, Mongolia
	<i>Z. baissana</i>	Krzemińska, Krzemiński & Dahl	2009a	Lower Cretaceous	Baissa, Russia
	<i>Z. bontsagana</i>	Krzemińska, Krzemiński & Dahl	2009a	Lower Cretaceous	Bon Tsagan, Mongolia
	<i>Z. karatavica</i>	Krzemińska, Krzemiński & Dahl	2009a	Upper Jurassic	Karatau, Kazakhstan
	<i>Z. novospasskaya</i>	Krzemińska, Krzemiński & Dahl	2009a	later Lower or early Middle Jurassic	Novospasskoe, Russia
	<i>Z. reni</i>	Krzemińska, Krzemiński & Dahl	2009a	Middle Jurassic	Daohugou, China
	<i>Z. tola</i>	Krzemińska, Krzemiński & Dahl	2009a	Lower Cretaceous	Onokhoy, Mongolia
	<i>Z. zherikhini</i>	Krzemińska, Krzemiński & Dahl	2009a	Upper Jurassic	Karatau, Kazakhstan

Materials and methods

The wing venation nomenclature used in this paper is based on the interpretations and system proposed by Lukashevich (2004) and Krzemińska et al. (2009a). The fossil specimens were examined under a Leica MZ7.5 dissecting microscope and illustrated with the aid of a drawing tube attachment. Line drawings were prepared with Adobe Photoshop CS3 Extended graphics software.

All specimens studied in the paper are housed in the Key Lab of Insect Evolution and Environmental Changes, College of Life Sciences, Capital Normal University, Beijing, China.

Systematic paleontology

Family Trichoceridae Kertész, 1902

Genus *Eotrichocera* Kalugina, 1985

Subgenus *Archaeotrichocera* Krzemińska, Krzemiński & Dahl, 2009

Type species. *Eotrichocera (Archaeotrichocera) ephemera* Zhang, 2006

Other included species. *Eotrichocera (Archaeotrichocera) conica* Krzemińska, Krzemiński & Ren, 2009; *Eotrichocera (Archaeotrichocera) rara* Krzemińska, Krzemiński & Ren, 2009; *Eotrichocera (Archaeotrichocera) spatiosa* Liu, Shih & Ren, 2012.

Key to the species of *Eotrichocera (Archaeotrichocera)*

- | | | |
|---|--|---|
| 1 | Sc ending at anterior margin distad of R ₂ | 2 |
| — | Sc ending at anterior margin proximad of R ₂ | 3 |
| 2 | Large size (wing length 12.0 mm)..... | |
| | <i>E. (A.) spatiosa</i> Liu, shih & Ren, 2012 (Daohugou, J ₂) | |
| — | Medium size (wing length 5.5 mm)..... | |
| | <i>E. (A.) rara</i> Krzemińska, Krzemiński & Ren, 2009 (Daohugou, J ₂) | |
| 3 | Crossvein sc-r distad of 1/2 (at 2/3) of length of Rs | 4 |
| — | Crossvein sc-r proximad of or at 1/3 of length of Rs | 5 |
| 4 | Rs forking distad of 2/3 (at 0.77) times of wing length | |
| | <i>E. (A.) ephemera</i> Zhang, 2006 (Daohugou, J ₂) | |
| — | Rs forking proximad of 2/3 (at about 0.53) times of wing length..... | |
| | <i>E. (A.) longensis</i> sp. n. (Daohugou, J ₂) | |
| 5 | A ₂ long (0.22 times as long as wing), d cell narrow and long (W/L=0.43 | |
| | <i>E. (A.) conica</i> Krzemińska, Krzemiński & Ren, 2009 (Daohugou, J ₂) | |
| — | A ₂ short (0.13 times as long as wing), d cell broad (W/L=0.58) | |
| | <i>E. (A.) amabilis</i> sp. n. (Daohugou, J ₂) | |

***Eotrichocera (Archaeotrichocera) longensis* sp. n.**

<http://zoobank.org/8A0D358E-7BCA-476B-A0A9-6EECA93FCBA8>

http://species-id.net/wiki/Eotrichocera_longensis

Etymology. “*longensis*” is a Latin word, referring to the long leg of this specimen.

Diagnosis. Sc rather short about 0.65 times as long as the wing and ending at anterior margin proximad of R₂; Rs forking proximad of 2/3 (at about 0.55) times of wing length; the d-cell narrow and long (about 2.5 times as long as wide); A₂ short and bending sharply toward anterior margin (angle about 128°).

Holotype. An almost complete female specimen with well-preserved body, wings and head. Specimen number CNU-DIP-NN2013133. Wing length 9.0 mm, width 3.8 mm (Figs 1A, 2, 3A).




Figure 1. *Eotrichocera (Archaeotrichocera) longensis* sp. n. Holotype, specimen CNU-DIP-NN2013131
A Photograph. Paratype, specimen CNU-DIP-NN2013131 **B** Photograph. Scale bars = 1 mm.



A**B**

Figure 2. *Eotrichocera (Archaeotrichocera) longensis* sp. n. Holotype, specimen CNU-DIP-NN2013133
A Line drawing **B** Tarsus of the mid leg. Scale bars = 1 mm; t_1 = the first segment of tarsus; t_2 = the second segment of tarsus; t_3 = the third segment of tarsus; t_4 = the fourth segment of tarsus.




Figure 3. *Eotrichocera (Archaeotrichocera) longensis* sp. n. Holotype, specimen CNU-DIP-NN201313. **A** Line drawing of left wing. Paratype, specimen CNU-DIP-NN201313. **B** Line drawing of left wing. Scale bars = 1 mm

Paratype. A female specimen with body and wings, specimen number CNU-DIP-NN2013131. Wing length 7.7 mm, width 3 mm (Figs 1B, 3B).

Locality and horizon. Jiulongshan Formation, Late Middle Jurassic, Daohugou Village, Ningcheng County, Inner Mongolia Autonomous Region, China.

Description. Based on Holotype, different characters of the paratype CNU-DIP-NN2013131 in brackets. Medium-sized winter crane flies, body length (including head) 13 mm with well preserved wings, body and head, [paratype body length (excluding head) 10.5 mm].

Head: Antenna very long, about 3.5 times as long as the head length, palpi about two times as long as the head length, compound eyes preserved.

Table 2. Comparison of key characters among the two new species and other species of *Eotrichocera* (*Archaeotrichocera*) Krzemińska, Krzemiński & Dahl, 2009. L/W = ratio of length/width; W/L = ratio of width/length.

Key characters	<i>E. (A.) ephemera</i> Zhang, 2006	<i>E. (A.) conica</i> Krzemińska, Krzemiński & Ren, 2009	<i>E. (A.) rara</i> Krzemińska, Krzemiński & Ren, 2009	<i>E. (A.) spatiosa</i> Liu, Shih & Ren, 2012	<i>E. (A.) longensis</i> sp. n.	<i>E. (A.) amabilis</i> sp. n.
Wing length, in mm and (L/W)	7.1 (L/W=2.8)	10.0 (L/W=3)	5.5	12.0 (L/ W=2.7)	9.0 (L/W=2.3)	5.2 (L/W=2.2)
Sc length and ending at anterior margin	0.77 times of wing length, ending proximad of R ₂	0.71 times of wing length, ending proximad of R ₂	0.77 times of wing length, ending distad of R ₂	0.84 times of wing length, and ending distad of R ₂	about 0.65 times of wing length, ending proximad of R ₂	about 0.71 times of wing length, ending proximad of R ₂
sc-r position	at 2/3 of length of Rs	at 1/3 of length of Rs	at 1/2 of length of Rs	at 2/3 of length of Rs	at 2/3 of length of Rs	at 1/3 of length of Rs
Position of Rs forking	0.77 times of wing length	0.64 times of wing length	0.57 times of wing length	0.58 times of wing length	0.53 times of wing length	0.55 times of wing length
d-cell W/L (length)	W/L=0.53 (1/6 of wing length)	W/L=0.43 (0.2 times of wing length)	W/L=0.39 (0.21 times of wing length)	W/L=0.47 (0.19 times of wing length)	W/L= 0.4 (almost 1/5 of wing length)	W/L=0.58 (almost 0.17 of wing length)
A ₂ length	long (about 1/4 of wing length), curved evenly to posterior margin	medium (0.22 times of wing length), not reaching posterior margin	rather short (1/5 of wing length) and not reaching posterior margin	short (about 0.21 times of wing length) and curving to posterior margin	short (0.14) times of wing length) and curved to posterior margin	short (0.13) times of wing length) and curved to posterior margin
r-m length	1/5 of length of d-cell	about 1/3 of length of d-cell	1/5 of length of the d-cell	0.24 or 0.15 of length of the d-cell

Thorax: Much higher, in lateral view, than that of the abdomen, subcircular in shape, with robust and well-developed mesonotum. The halters spoon-type and the length of halters as long as thorax.

Wings: Wing is shorter than abdomen, not covering the end of the abdomen. Length 9.0 mm [Paratype with wing length of 7.7 mm], narrow and long (about 2.5 times as long as wide); venation clear, Sc rather short about 0.68 times as long as the wing [Paratype Sc rather short, about 0.65 times as long as the wing] and ending at anterior margin proximad of R₂; crossvein sc-r locating at 2/3 of Rs; Rs arising about one-fifth from the base of the wing; R₂₊₃ about 0.8 times as long as R₂₊₃₊₄; R₂ about one-tenth of length of R₃; R₃ almost three times as long as the R₂₊₃; dM₁₊₂ 0.6 times as long as mM₁₊₂, while M₁ 2.5 times of the dM₁₊₂; a well developed m-m crossvein about three-fourth length of bM₃, closing the d-cell and nearly 0.3 length of d-cell; bM₁₊₂ nearly 1.0 times as long as the length of the r-m and the latter at one-fifth of the d-cell; d-cell narrow and long (about 2.5 times as long as wide) and almost one-fifth length of wing; both crossveins m-m and m-cu intersecting with M₄ at the same point; Cu long, curved (angle about 135°) and reaching the wing posterior margin at 0.6 from the base of the wing; the stem of A divided into A₁ and A₂; A₁ long, slightly curving and reaching the wing posterior margin; A₂ short, 0.15 (right wing) [Paratype 0.14] times as long as

wing and almost 0.3 times as long as length of A_1 , bending sharply (angle about 128°) and reaching the wing posterior margin.

Legs: Legs slender and long; the hind leg nearly 1.2 times as long as the abdomen and 1.3 times as long as the wing. Tarsus with five segments; the first segment of tarsus (t_1) is 1.2 times as long as t_2 in mid leg.

Abdomen: Abdomen relatively long and thin, with ten segments. Female genital discernible.

Remarks. *Eotrichocera (Archaeotrichocera) longensis* sp. n. is assigned to Thrichocerinae based on the following characters: d-cell medium, m-cu present; A_2 short, antennae long, flagellomeres thin, much longer than two times of the head length. It belongs to *Eotrichocera (Archaeotrichocera)* because of wing length from 7.7 to 9.0 mm and d-cell almost one-fifth of wing length. In addition, it differs from all other known Thrichocerinae by its A_2 rather short and bending sharply toward anterior margin (angle about 128°), R_2 relatively long, Sc forking proximad of 2/3 (at about 0.55) times of wing length, and d-cell narrow and long. To compare the key characters among the new species and other species of *Eotrichocera (Archaeotrichocera)*, we set up the Table 2.

Eotrichocera (Archaeotrichocera) amabilis sp. n.

<http://zoobank.org/D32A4E4B-EDF1-4684-802E-92F9617DEAB2>

http://species-id.net/wiki/Eotrichocera_amabilis

Etymology. The specific name of “*amabilis*” is a Latin word, meaning lovely.

Diagnosis. Body small and wing short; Sc 0.71 times as long as wing; the d-cell broad (about 1.7 times as long as wide); A_2 short and bending sharply toward posterior margin (angle about 128°).

Holotype. An almost complete female specimen with well-preserved body, wings and head. Specimen number CNU-DIP-NN2013134, Wing length 5.2 mm, width 2.2 mm (Figs 4A–D, 5A, 6).

Paratype. A specimen with body and wings with partial venation, specimen number CNU-DIP-NN2013132, (Figs 4E, 5B).

Locality and horizon. Jiulongshan Formation, Late Middle Jurassic, Daohugou Village, Ningcheng County, Inner Mongolia Autonomous Region, China.

Description. Based on Holotype, different characters of the paratype CNU-DIP-NN2013132 in brackets. Medium-sized winter crane flies, head length 0.47 mm, body length (including head) 5.8 mm with well preserved body and wings. [Paratype with partial body and wings with partial venation].

Head: antenna very long, about 5.7 times as long as the head length, palpi about two times as long as the head length, compound eyes preserved (Figs 4C, 6).

Thorax: Much higher, in lateral view, than that of the abdomen, subcircular in shape, with robust and well-developed mesonotum.

Wings: Wing is shorter than abdomen, not covering the end of the abdomen. Wing length of 5.2 mm [Paratype with wing length 5.0 mm], narrow and long ($L/W=2.2$); ve-




Figure 4. *Eotrichocera (Archaeotrichocera) amabilis* sp. n. Holotype, specimen CNU-DIP-NN2013134 **A** Photograph **B** Photograph, under alcohol **C** Line drawing **D** Tarsus of the mid leg. Paratype, specimen CNU-DIP-NN2013132 **E** Photograph. Scale bars = 1 mm; t_1 = the first segment of tarsus; t_2 = the second segment of tarsus.

nation clear, Sc rather short about 0.71 times as long as the wing and terminating clearly proximad of R_2 ; crossvein sc-r locating at 1/3 [Paratype 1/2] of Rs, and distad to the Sc ending; [Paratype Rs arising about one-fourth from the base of the wing]; Rs forking at 0.55 [Paratype 0.64] times of wing length; R_{2+3} about 1.9 times as long as R_{2+3+4} ; R_2 about 0.18 of length of R_3 ; R_3 almost 3.7 times as long as the R_{2+3} ; R_5 9.0 times as long as R_{2+3+4} ; M_1 1.6 times of the dM_{1+2} ; crossvein m-m well developed about 0.73 times as long as bM_3 , closing the d-cell and nearly 0.17 [Paratype 0.21] times as long as wing; bM_{1+2} nearly 2.1 times as long as the length of the r-m and the latter at one-fourth of the d-cell; d-cell broad ($W/L=0.58$ [Paratype 0.56]) and almost 0.17 times of length of wing; Cu long, curved (angle about 121°) and reaching the wing posterior margin at 0.67 from the base of the wing; the stem of A divided into A_1 and A_2 ; A_1 long, slightly curving and




Figure 5. *Eotrichocera (Archaeotrichocera) amabilis* sp. n. Holotype, specimen CNU-DIP-NN2013134
A Line drawing of left wing. Paratype, specimen CNU-DIP-NN2013132 **B** Line drawing of left wing.
Scale bars = 1 mm.

reaching the wing posterior margin; A₂ short, 0.13 times as long as wing and almost 0.25 times as long as length of A₁, bending sharply and reaching the wing posterior margin.

Abdomen: Abdomen relatively long and thin, with ten segments. Female genitalia discernible (Figs 4A–C) [Paratype genitalia indiscernible].

Legs: Legs slender and long; the hind leg nearly 1.5 times as long as the abdomen and 1.4 times as long as the wing. Tarsus with five segments; the first segment of tarsus (t₁) is 1.2 times as long as t₂ in mid leg.

Remarks. The new species is compared and differentiated from all other species in *Eotrichocera (Archaeotrichocera)* in Table 2.




Figure 6. *Eotrichocera (Archaeotrichocera) amabilis* sp. n. Holotype, specimen CNU-DIP-NN2013134, Photograph of head, under alcohol. Scale bar = 1 mm.

Due to limitation of fossil preservation, some of the morphological characters of previously described fossil are not objective or clear. We set up an aforementioned key based on the Sc length and ending location at anterior margin, wing length, crossvein sc-r position, Rs forking location and A₂ length, to differentiate the species of subgenus *Archaeotrichocera*. These characters may help future morphological and taxonomic studies in differentiating fossil species of Trichoceridae.

Acknowledgements

We sincerely appreciate valuable comments and constructive suggestion from Dr. Ewa Krzemińska. We thank Dr. Wang Y-J, Dr. Gao T-P, Shi C-F, Peng Y-Y and Li L-F, Shi G-F for their helpful comments, fruitful suggestion and improvement of our manuscript. This research was supported by grants from the National Basic Research Program of China (973 Program) (Grant 2012CB821906), the National Nature Science Foundation of China (No. 31230065, 41272006), Great Wall Scholar and KEY project of Beijing Municipal Commission of Education (Grant KZ201310028033), Program for Changjiang Scholars and Innovative Research Team in University (IRT13081).

References

- Dahl C, Alexander C-P (1976) A world catalogue of Trichoceridae Kertész, 1902 (Diptera). Entomol Scand 7:7–18. doi: 10.1163/187631276X00027
- Engel MS, Ren D (2008) New snakeflies from the Jiulongshan Formation of Inner Mongolia, China (Raphidioptera). Journal of the Kansas Entomological Society 81(3): 188–193. doi: 10.2317/JKES-802.19.1
- Gao K-Q, Ren D (2006) Radiometric Dating of Ignimbrite from Inner Mongolia Provides no Indication of a Post-Middle Jurassic Age for the Daozugou Beds. Acta Geologica Sinica (English Edition) 80: 42–45. doi: 10.1111/j.1755-6724.2006.tb00793.x
- Gu J-J, Qiao G-X, Ren D (2010) Revision and new taxa of fossil Prophalangopsidae (Orthoptera: Ensifera). Journal of Orthoptera Research 19(1): 41–56. doi: 10.1665/034.019.0110
- Hågvar S, Krzemińska E (2007) Contribution to the winter phenology of Trichoceridae (Diptera) in snow-covered southern Norway. Stud Dipt 14: 271–283.
- Kalugina NS, Kovalev VG (1985) Diptera Insects of Jurassic of Siberia. Nauka, Moscow, 198 pp. [In Russian]
- Krzemińska E, Krzemiński W, Dahl C (2009a) Monograph of Fossil Trichoceridae (Diptera) over 180 Million Years of Evolution. Institute of Systematics and Evolution of Animals of the Polish Academy of Sciences Press, Krakow, Poland, 35–40.
- Krzemińska E, Lukashevich E (2009b) The Late Jurassic fossil from Asia ancestry to a recent genus, Nothotrichocera Alexander 1926 (Diptera: Trichoceridae). Zoosymposia 3: 103–110.
- Liu L-X, Shih C-K, Ren D (2012) Two new species of Ptychopteridae and Trichoceridae from the Middle Jurassic of northeastern China (Insecta: Diptera: Nematocera). Zootaxa 3501: 55–62.

- Liu Y-S, Ren D (2008) Two new Jurassic stoneflies (Insecta: Plecoptera) from Daohugou, Inner Mongolia, China. *Progress in Natural Science* 18: 1039–1042. doi: 10.1016/j.pnsc.2008.03.014
- Lukashevich ED (2004) A revision of the Genus *Eoptychopterina* (Diptera: Eoptychopteridae). *Paleontological Journal* 38 (3): 294–306.
- Ren D, Gao K-Q, Guo Z-G, Ji S-A, Tan J-J, Song Z (2002) Stratigraphic division of the Jurassic in the Daohugou Area, Ningcheng, Inner Mongolia. *Geological Bulletin of China* 21: 584–591.
- Ren D, Engel MS (2007) A split-footed lacewing and two episomylines from the Jurassic of China (Neuroptera). *Annales Zoologici (Warszawa)* 57(2): 211–219.
- Ren D, Labandeira CC, Santiago-Blay JA, Rasnitsyn AP, Shih C-K, Bashkuev A, Logan MAV, Hotton CL, Dilcher D (2009) A probable pollination mode before angiosperms: Eurasian, long-proboscid scorpionflies. *Science* 326: 840–847. doi: 10.1126/science.1178338
- Ren D, Shih C-K, Gao T-P, Yao Y-Z, Zhao Y-Y (2010a) Silent Stories – Insect Fossil Treasures from Dinosaur Era of the Northeastern China. Science Press, Beijing, 324 pp.
- Ren D, Shih C-K, Labandeira CC (2010b) New Jurassic Pseudopolycentropodids from China (Insecta: Mecoptera). *Acta Geologica Sinica (English Edition)* 84(1): 22–30. doi: 10.1111/j.1755-6724.2010.00166.x
- Shi C-F, Yang Q, Ren D (2011) Two new fossil lacewing species from the Middle Jurassic of Inner Mongolia, China (Neuroptera: Grammolingiidae). *Acta Geologica Sinica (English Edition)* 85(2): 482–489. doi: 10.1111/j.1755-6724.2011.00416.x
- Wang Y-J, Liu Z-Q, Wang X, Shih C-K, Zhao Y-Y, Engel MS, Ren D (2010) Ancient pinnate leaf mimesis among lacewings. *Proceedings of the National Academy of Sciences USA* 107(37): 16212–16215. doi: 10.1073/pnas.1006460107
- Wang Y-J, Labandeira CC, Shih C-K, Ren D, Ding Q-L, Wang C, Zhao Y-Y (2012) Jurassic mimicry between a hanging fly and a ginkgo from China. *Proceedings of The National Academy of Sciences USA* 109(50): 20514–10519. doi: 10.1073/pnas.1205517109
- Wang Y, Ren D (2009) New fossil palaeointinids from the Middle Jurassic of Daohugou, Inner Mongolia, China (Insecta, Hemiptera). *Acta Geologica Sinica (English Edition)* 83(1): 33–38. doi: 10.1111/j.1755-6724.2009.00004.x
- Yang D (2009) Diptera. In: Yang D (Ed) Fauna of Hebei, China. Agricultural Science and Technology Press, Beijing, China, 14–25.
- Yang Q, Makarkin VN, Winterton SL, Khramov AV, Ren D (2012) A remarkable new family of Jurassic insects (Neuroptera) with primitive wing venation and its phylogenetic position in neuropterida. *PLoS ONE* 7(9): e44762. doi: 10.1371/journal.Pone.0044762
- Yang J-K, Yang D (1995) Insecta: Diptera: Trichoceridae. In: Yang J-K, Yang D (Eds) Insects and Large-Scale Fungi in Gutianshan of Zhejiang Province. Zhejiang Science and Technology Press, Zhejiang, China, 175–179.
- Zhang J-F (2006) New winter crane flies (Insecta: Diptera: Trichoceridae) from the Jurassic Daohugou Formation (Inner Mongolia, China) and their associated biota. *Canadian Journal of Earth Sciences* 43: 9–22. doi: 10.1139/e05-092