

Checklist of the family Sciaridae (Diptera) of Finland

Pekka Vilkamaa¹

¹ Finnish Museum of Natural History, Zoology Unit, P.O. Box 17, FI-00014 University of Helsinki, Finland

Corresponding author: Pekka Vilkamaa (pekka.vilkamaa@helsinki.fi)

Academic editor: J. Salmela | Received 26 February 2014 | Accepted 21 March 2014 | Published 19 September 2014

<http://zoobank.org/9F25385D-67C7-48ED-A7AB-6A65DCED753F>

Citation: Vilkamaa P (2014) Checklist of the family Sciaridae (Diptera) of Finland. In: Kahanpää J, Salmela J (Eds) Checklist of the Diptera of Finland. ZooKeys 441: 151–164. doi: 10.3897/zookeys.441.7381

Abstract

A checklist of the family Sciaridae (Diptera) recorded from Finland is provided. The genus *Sciarosoma* Chandler with a disputed family placement is also included in the list.

Keywords

Checklist, Finland, Diptera, Sciaridae, *Sciarosoma*

Introduction

Sciaridae, the black fungus gnats, is one of the large families in Sciaroidea, little studied and notoriously difficult in its taxonomy. Up-to-date keys are not available for all European species, but various publications must be consulted for identification. Our knowledge of Finnish fauna stems from two classic publications: Frey (1948) and Tuomikoski (1960) but Menzel and Mohrig (2000) made fundamental nomenclatorial changes in their study of the type materials of all Palearctic species.

As no records of Sciaridae in Finland were published between the publication of Tuomikoski's (1960) monograph on the family and Hackman's (1980) check-list

of Finnish nematoceran Diptera, the species lists in both works are essentially unchanged, despite some obvious lapses in the latter. Our knowledge of Finnish fauna has since grown mainly in the taxonomic treatment of various genera and species groups (see Vilkamaa et al. 2013). All species recorded in Finland in publications thus far, besides the most recent ones, appear in H. Silfverberg's compilations of changes to the Finnish insect fauna published every five years (for the latest one, see Silfverberg 2012).

With regard to nomenclature and generic classification, the present list follows the revisions of Menzel and Mohrig (2000) and Mohrig et al. (2013), although on the basis of phylogenetic analyses *Baeosciara* Tuomikoski, *Dolichosciara* Tuomikoski, *Peyerimhoffia* Kieffer, *Prosciara* Frey and *Mouffetina* Frey are regarded as genera (see Vilkamaa 2000, 2003, Vilkamaa and Hippa 2005, Shin et al. 2013). For names currently understood as junior synonyms, mostly only the names listed as valid in Tuomikoski (1960) are mentioned. For complete synonym lists, one can consult Menzel and Mohrig (2000) and Mohrig et al. (2013). Although some genera can be regarded as relatively well-known for the Finnish species they contain, for some others, such as the species-rich *Bradyzia* Winnertz, many new species including undescribed ones may be present in the native fauna. Even so, together with Germany, Sweden and the United Kingdom, Finland can be considered one of the countries best known for its sciarid fauna.

Number of species:

World: 2600 species

Europe: 700 species

Finland: 338 species

Faunistic knowledge level in Finland: average

Checklist

Nematocera Dumeril, 1805

infraorder Bibionomorpha Hennig, 1954

superfamily Sciaroidea Billberg, 1820

SCIARIDAE Billberg, 1820

BAEOSCIARA Tuomikoski, 1960

Baeosciara discolor (Lengersdorf, 1928)

= *pusillima* (Frey, 1948)

Baeosciara sinuata (Menzel & Mohrig, 1997)

Baeosciara scotica (Edwards, 1925)

BRADYSIA Winnertz, 1867

Bradyzia affinis (Zetterstedt, 1838)

= *pratincola* Tuomikoski, 1960

- Bradysia albanensis* (Lengersdorf, 1926)
Bradysia alpicola (Winnertz, 1867)
 = *mutabilis* (Lengersdorf, 1926)
 = *morio* auct. nec (Fabricius, 1794)
Bradysia angustata Tuomikoski, 1960
Bradysia angustostylata Menzel, 2005
Bradysia aprica (Winnertz, 1867)
Bradysia arcana Menzel & Mohrig, 1998
 = *fenestralis* auct. nec (Zetterstedt, 1838)
Bradysia arcula Vilkamaa, Salmela & Hippa, 2007
Bradysia ascenda Rudzinski, 1994
Bradysia bicolor (Meigen, 1818)
Bradysia bispinifera Mohrig & Krivosheina, 1983
Bradysia brevispina Tuomikoski, 1960
Bradysia browni (Shaw, 1935)
 = *diversiabdominalis* (Lengersdorf, 1941)
 = *laurencei* Menzel & Mohrig, 2000
Bradysia cinerascens (Grzegorzek, 1884)
 = *lanicauda* Tuomikoski, 1960
Bradysia confinis (Winnertz, 1867)
Bradysia excelsa Menzel & Mohrig, 1998
Bradysia flavipila Tuomikoski, 1960
Bradysia forcipulata (Lundbeck, 1898)
Bradysia forficulata (Bezzi, 1914)
 = *nocturna* Tuomikoski, 1960
Bradysia fungicola (Winnertz, 1867)
Bradysia giraudii (Egger, 1862)
Bradysia globulifera (Lengersdorf, 1934)
Bradysia hilariformis Tuomikoski, 1960
Bradysia hilaris (Winnertz, 1867)
Bradysia holsatica Heller, 2004
Bradysia hortensis Heller, 2000
Bradysia impatiens (Johannsen, 1912)
 = *diformis* (Frey, 1948)
 = *paupera* Tuomikoski, 1960
Bradysia inusitata Tuomikoski, 1960
Bradysia iridipennis (Zetterstedt, 1838)
Bradysia lapponica (Lengersdorf, 1926)
Bradysia latiterga Tuomikoski, 1960
Bradysia leptoptera Tuomikoski, 1960
Bradysia lilienthalae Mohrig & Menzel, 1990
Bradysia lobulifera Frey, 1948
Bradysia longicauda Mohrig & Menzel, 1990

- Bradysia longicubitalis* (Lengersdorf, 1924)
= *cinereovittata* Frey, 1948
- Bradysia minima* Mohrig & Mamaev, 1989
- Bradysia moesta* Frey, 1948
= *albosetosa* Frey, 1948
- Bradysia moestula* Tuomikoski, 1960
- Bradysia nervosa* (Meigen, 1818)
- Bradysia nitidicollis* (Meigen, 1818)
= *atroparva* Frey, 1948
- Bradysia normalis* Frey, 1948
- Bradysia ocellaris* (Comstock, 1882)
- Bradysia pallipes* (Fabricius, 1787)
= *brunnipes* (Meigen, 1804)
= *picipes* (Zetterstedt, 1838)
- Bradysia pauperata* (Winnertz, 1867)
- Bradysia peraffinis* Tuomikoski, 1960
- Bradysia pilistriata* Frey, 1948
- Bradysia placida* (Winnertz, 1867)
= *fimbriacauda* Tuomikoski, 1960
- Bradysia polonica* (Lengersdorf, 1929)
- Bradysia praecox* (Meigen, 1818)
- Bradysia rectinervis* Frey, 1948
- Bradysia reflexa* Tuomikoski, 1960
- Bradysia regularis* (Lengersdorf, 1934)
= *subnervosa* Frey, 1948
- Bradysia rufescens* (Zetterstedt, 1852)
- Bradysia scabriculum* Tuomikoski, 1960
= *subscabriculum* Mohrig & Menzel, 1990
- Bradysia siberica* Komarova, 2001
- Bradysia spinostyla* Mohrig & Menzel, 1990
- Bradysia strigata* (Staeger, 1840)
- Bradysia subalpina* Frey, 1948
- Bradysia subamoena* Mohrig & Krivosheina, 1989
- Bradysia submoesta* Mohrig & Krivosheina, 1989
- Bradysia tilicola* (Loew, 1850)
= *amoena* (Winnertz, 1867)
= *cellarum* Frey, 1948
- Bradysia trivittata* (Staeger, 1840)
- Bradysia vagans* (Winnertz, 1868)
= *callicera* Frey, 1948
- Bradysia vernalis* (Zetterstedt, 1851)
- Bradysia zonata* Rudzinski, 1993
- BRADYSIOPSIS** Tuomikoski, 1960

- Bradysiopsis vittata* (Meigen, 1830)
= *leucotricha* (Tuomikoski, 1960)
- Bradysiopsis vittigera* (Zetterstedt, 1851)
- CAMPTOCHAETA** Hippa & Vilkamaa, 1994
- Camptochaeta austriaca* Heller, 2012
- Camptochaeta bournei* (Shaw, 1941)
- Camptochaeta camptochaeta* (Tuomikoski, 1960)
- Camptochaeta consimilis* (Holmgren, 1869)
- Camptochaeta delicata* (Lengersdorf, 1935)
- Camptochaeta duplicata* Hippa & Vilkamaa, 1994
- Camptochaeta fallax* Hippa & Vilkamaa, 1994
- Camptochaeta hirtula* (Lengersdorf, 1934)
= *fulvicollis* (Tuomikoski, 1960)
- Camptochaeta propria* Hippa & Vilkamaa, 1994
- Camptochaeta scanica* Hippa & Vilkamaa, 1994
- Camptochaeta sicilicula* Hippa & Vilkamaa, 1994
- Camptochaeta simulator* Hippa & Vilkamaa, 1994
- Camptochaeta stammeri* (Lengersdorf, 1940)
- Camptochaeta tenuipalpalis* (Mohrig & Antonova, 1978)
- Camptochaeta uniformis* (Mohrig & Menzel, 1990)
- Camptochaeta vivax* (Frey, 1948)
- Camptochaeta xystica* Hippa & Vilkamaa, 1994
- CHAETOSCIARA** Frey, 1942
- Chaetosciara estlandica* (Lengersdorf, 1929)
- CLAUSTROPYGA** Hippa, Vilkamaa & Mohrig, 2003
- Claustropya acanthostyla* (Tuomikoski, 1960)
- Claustropya brevichaeta* (Mohrig & Antonova, 1978)
- Claustropya clausa* (Tuomikoski, 1960)
- Claustropya corticis* (Mohrig & Antonova, 1978)
- Claustropya ctenophora* Hippa, Vilkamaa & Mohrig, 2003
- Claustropya heteroclausa* (Rudzinski, 1991)
- Claustropya refrigerata* (Lengersdorf, 1930)
- Claustropya subcorticis* (Mohrig & Krivosheina, 1985)
- CORYNOPTERA** Winnertz, 1867
- Corynoptera barbata* Tuomikoski, 1960
- Corynoptera bicuspidata* (Lengersdorf, 1926)
= *gymnops* Tuomikoski, 1960
- Corynoptera bipartita* Mohrig & Krivosheina, 1985
- Corynoptera blanda* (Winnertz, 1867)
- Corynoptera boletiphaga* (Lengersdorf, 1940)
= *geogenia* Tuomikoski, 1960
- Corynoptera breviformis* Mohrig & Krivosheina, 1983
- Corynoptera cracentis* Vilkamaa, Hippa & Heller, 2013

- Corynoptera cuniculata* (Lengersdorf, 1942)
= *caldariorum* Tuomikoski, 1960
- Corynoptera defecta* (Frey, 1948)
- Corynoptera dentata* (Bukowski & Lengersdorf, 1936)
- Corynoptera deserta* Heller & Menzel, 2006
= *minutula* (Bukowski & Lengersdorf, 1936)
- Corynoptera dubitata* Tuomikoski, 1960
- Corynoptera fera* Mohrig & Heller, 1992
- Corynoptera flavicauda* (Zetterstedt, 1855)
- Corynoptera forcipata* (Winnertz, 1867)
- Corynoptera furcifera* Mohrig & Mamaev, 1987
- Corynoptera globiformis* (Frey, 1945)
- Corynoptera hypopygialis* (Lengersdorf, 1926)
= *piniphila* (Lengersdorf, 1940)
- Corynoptera inari* Vilkamaa, Hippa & Heller, 2013
- Corynoptera inexpectata* Tuomikoski, 1960
- Corynoptera irmgardis* (Lengersdorf, 1930)
- Corynoptera levigata* Tuomikoski, 1960
- Corynoptera luteofusca* (Bukowski & Lengersdorf, 1936)
- Corynoptera mariniae* Mohrig & Krivosheina, 1986
- Corynoptera melanochaeta* Mohrig & Menzel, 1992
- Corynoptera membranigera* (Kieffer, 1903)
= *trispina* Tuomikoski, 1960
- Corynoptera montana* (Winnertz, 1869)
- Corynoptera ninae* Hippa, Vilkamaa & Heller, 2010
- Corynoptera obscuripila* Tuomikoski, 1960
- Corynoptera parvula* (Winnertz, 1867)
= *uncinata* (Hippa & Vilkamaa, 1994)
- Corynoptera parvulaformis* Mohrig, 1985
- Corynoptera penna* (Pettey, 1918)
= *alneti* Hippa, Vilkamaa & Heller, 2010
- Corynoptera perochaeta* (Mohrig & Menzel, 1990)
- Corynoptera plusiochaeta* Hippa, Vilkamaa & Heller, 2010
- Corynoptera polana* Rudzinski, 2009
- Corynoptera postforcipata* Rudzinski, 1993
- Corynoptera postglobiformis* Mohrig, 1993
- Corynoptera praeforcipata* Mohrig & Mamaev, 1987
- Corynoptera quantula* (Hippa & Vilkamaa, 1994)
- Corynoptera saccata* Tuomikoski, 1960
- Corynoptera saetistyla* Mohrig & Krivosheina, 1985
- Corynoptera salmelai* Vilkamaa, Hippa & Heller, 2013
- Corynoptera sphenoaptera* Tuomikoski, 1960
- Corynoptera spiciforceps* Vilkamaa, Hippa & Heller, 2013

- Corynoptera spinifera* Tuomikoski, 1960
Corynoptera subblanda Tuomikoski, 1960
Corynoptera subdentata Mohrig, 1985
Corynoptera subparvula Tuomikoski, 1960
Corynoptera subsedula Mohrig & Mamaev, 1987
Corynoptera subtetrachaeta Komarova, 1995
Corynoptera subtilis (Lengersdorf, 1929)
 = *longicornis* (Bukowski & Lengersdorf, 1936)
Corynoptera subvariegata Rudzinski, 1992
Corynoptera tetrachaeta Tuomikoski, 1960
Corynoptera trepida (Winnertz, 1867)
 = *clinochaeta* Tuomikoski, 1960
Corynoptera triacantha Tuomikoski, 1960
Corynoptera tumidula Hippa, Vilkamaa & Heller, 2010
Corynoptera tuomikoskii Hippa, Vilkamaa & Heller, 2013
Corynoptera unidentata (Hippa & Vilkamaa, 1994)
Corynoptera vagula Tuomikoski, 1960
Corynoptera verrucifera (Lengersdorf, 1952)
Corynoptera voluptuosa Mohrig & Mamaev, 1987
Corynoptera waltraudis Mohrig & Mamaev, 1987
Corynoptera winnertzi Mohrig, 1993
COSMOSCIARA Frey, 1942
Cosmosciara perniciosa (Edwards, 1922)
CRATYNA Winnertz, 1867
 = *Plastosciara* Berg, 1899
 = *Decembrina* Frey, 1948
 = *Dendrosciara* Frey, 1948
sg. Cratyna Winnertz, 1867
Cratyna ambigua (Lengersdorf, 1934)
 = *latiforceps* (Bukowski & Lengersdorf, 1936)
Cratyna atra Winnertz, 1867
 = *pictiventris* (Kieffer, 1898)
Cratyna betulae (Mohrig & Mamaev, 1992)
Cratyna breviflagellata (Mohrig & Mamaev, 1985)
Cratyna monumenta Rudzinski, 2009
Cratyna pernitida (Edwards, 1915)
Cratyna schineri (Winnertz, 1867)
Cratyna sicata Vilkamaa, Hippa & Heller, 2013
Cratyna symplecta (Rudzinski, 1991)
Cratyna uliginosa (Lengersdorf, 1929)
Cratyna vaporariorum (Frey, 1948)
sg. Diversicratyna Menzel & Mohrig, 1998
Cratyna spiculososa (Rudzinski, 1993)

sg. *Spathobdella* Frey, 1948

Cratyna colei (Freeman, 1990)

= *brachialis* auct. nec (Winnertz, 1871)

Cratyna falcata (Tuomikoski, 1960)

Cratyna falcifera (Lengersdorf, 1933)

Cratyna longispina (Pettey, 1918)

= *tuberculata* (Tuomikoski, 1960)

Cratyna nobilis (Winnertz, 1867)

= *brachialis* (Winnertz, 1867)

Cratyna perplexa (Winnertz, 1867)

= *socialis* (Winnertz, 1871)

= *brevicornis* (Tuomikoski, 1957)

CTENOSCIARA Tuomikoski, 1960

Ctenosciara exigua Salmela & Vilkamaa, 2005

Ctenosciara hyalipennis (Meigen, 1804)

DICHOPYGINA Vilkamaa, Hippa & Komarova, 2004

Dichopygina aculeata Vilkamaa, Hippa & Komarova, 2004

Dichopygina intermedia (Mohrig & Krivosheina, 1982)

Dichopygina nigrohalteralis (Frey, 1948)

Dichopygina ramosa Vilkamaa, Hippa & Komarova, 2004

DOLICOSCIARA Tuomikoski, 1960

Dolichosciara flavipes (Meigen, 1804) 185

Dolichosciara hippai Komarova & Vilkamaa, 2006

Dolichosciara nigrovittata (Strobl, 1910)

Dolichosciara orcinia (Tuomikoski, 1960)

Dolichosciara ornata (Winnertz, 1867)

Dolichosciara saetosa (Lengersdorf, 1929)

Dolichosciara spissispina Vilkamaa, Hippa & Heller, 2013

EPIDAPUS Haliday, 1851

sg. *Epidapus* Haliday, 1851

Epidapus alnicola (Tuomikoski, 1957)

Epidapus atomarius (De Geer, 1778)

Epidapus gracilis (Walker, 1848)

Epidapus ignotus (Lengersdorf, 1942)

= *gracilior* (Tuomikoski, 1960)

Epidapus microthorax (Börner, 1903)

= *gracilicornis* (Lengersdorf, 1926)

Epidapus schillei (Börner, 1903)

= *titan* Frey, 1948

= *intermittens* Tuomikoski, 1959

sg. *Pseudoaptanogyna* Vimmer, 1926

Epidapus abieticola Frey, 1948

Epidapus bispinulosus Mohrig & Kauschke, 1994

- Epidapus echinatum* Mohrig & Kozánek, 1992
Epidapus ignavus (Lengersdorf, 1941)
KEILBACHIA Mohrig, 1987
Keilbachia ferrata (Hippa & Vilkamaa, 1994)
LEPTOSCIARELLA Tuomikoski, 1960
sg. **Hirtipennia** Mohrig & Menzel, 1997
Leptosciarella hirtipennis (Zetterstedt, 1838)
Leptosciarella holotricha Mohrig & Menzel, 1997
sg. **Leptosciarella** Tuomikoski, 1960
Leptosciarella brevior (Tuomikoski, 1960)
Leptosciarella brevipalpa (Mohrig & Menzel, 1992)
Leptosciarella claviforceps (Tuomikoski, 1960)
Leptosciarella dimera (Tuomikoski, 1960)
Leptosciarella fuscipalpa (Mohrig & Mamaev, 1979)
Leptosciarella helvetica (Rudzinski, 1992)
Leptosciarella hispida (Winnertz, 1871)
Leptosciarella ignis Heller, 2012
Leptosciarella krille Heller, 2012
Leptosciarella melanoma (Mohrig & Menzel, 1990)
Leptosciarella nudinervis (Tuomikoski, 1960)
Leptosciarella pilosa (Staeger, 1840)
Leptosciarella reducta Heller & Menzel, 2013
Leptosciarella rejecta (Winnertz, 1867)
Leptosciarella scutellata (Staeger, 1840)
 = *elegans* (Winnertz, 1867)
Leptosciarella subcoarctata Mohrig & Menzel, 1997
Leptosciarella subpilosa (Edwards, 1925)
Leptosciarella subspinulosa (Edwards, 1925)
Leptosciarella subviatrica Mohrig & Menzel, 1997
Leptosciarella trochanterata (Zetterstedt, 1851)
 = *coarctata* (Winnertz, 1867)
Leptosciarella truncata (Tuomikoski, 1960)
Leptosciarella viatica (Winnertz, 1867)
Leptosciarella viaticella Mohrig & Krivosheina, 1979
Leptosciarella yerburyi (Freeman, 1983)
sg. **Leptospina** Mohrig & Menzel, 1997
Leptosciarella atricha (Tuomikoski, 1960)
sg. **Trichosiopsis** Tuomikoski, 1960
Leptosciarella tuberculigera (Tuomikoski, 1960)
LYCORIELLA Frey, 1942
sg. **Coelostylina** Tuomikoski, 1960
Lycoriella eflagellata Tuomikoski, 1960
Lycoriella freyi Tuomikoski, 1960

sg. *Hemineurina* Tuomikoski, 1960

Lycoriella algida (Frey, 1948)

Lycoriella cochleata (Rübsamen, 1898)

Lycoriella conspicua (Winnertz, 1867)

Lycoriella inflata (Winnertz, 1867)

 = *venosa* auct. nec (Staeger, 1840)

Lycoriella modesta (Staeger, 1840)

Lycoriella piristylata Vilkamaa, Hippa & Heller, 2013

Lycoriella thuringiensis Menzel & Mohrig, 1991

Lycoriella vitticollis (Holmgren, 1883)

 = *permutata* (Lundbeck, 1900)

sg. *Lycoriella* Frey, 1942

Lycoriella aberrans Tuomikoski, 1960

Lycoriella acutostylia Mohrig & Menzel, 1990

Lycoriella agraria (Felt, 1897)

 = *cellaris* (Lengersdorf, 1934)

Lycoriella approximatonervis (Frey, 1948)

Lycoriella brevipila Tuomikoski, 1960

Lycoriella inconspicua Tuomikoski, 1960

Lycoriella ingenua (Dufour, 1839)

 = *solani* (Winnertz, 1871)

Lycoriella latilobata Menzel & Mohrig, 2000

Lycoriella lundstromi (Frey, 1948)

Lycoriella micria Mohrig & Menzel, 1990

Lycoriella minutula Mohrig & Krivosheina, 1987

Lycoriella pallidior Tuomikoski, 1960

Lycoriella parva (Holmgren, 1869)

 = *obscuratipes* (Frey, 1948)

 = *curvispina* Tuomikoski, 1960

Lycoriella sativae (Johannsen, 1912)

=*castanescens* (Lengersdorf, 1940)

 = *fucorum* (Frey, 1948)

Lycoriella subterranea (Märkel, 1844)

 = *vanderwielii* (Schmitz, 1920)

Lycoriella tenera Vilkamaa, Hippa & Heller, 2013

Lycoriella weberi Menzel & Heller, 2013

MOUFFETINA Frey, 1942

Mouffetina expolita (Coquillet, 1900)

Mouffetina pulchricornis (Edwards, 1925)

Mouffetina silvestris (Mohrig & Antonova, 1978)

PEYERIMHOFFIA Kieffer, 1903

Peyerimhoffia crassistylata (Frey, 1948)

Peyerimhoffia infera Vilkamaa & Hippa, 2005

- Peyerimhoffia menzeli* Vilkamaa & Hippa, 2005
Peyerimhoffia quadrifera Vilkamaa, Hippa & Heller, 2013
Peyerimhoffia sepei Hippa & Vilkamaa, 2005
Peyerimhoffia thula Vilkamaa & Hippa, 2005
Peyerimhoffia vagabunda (Winnertz, 1867)
 = *brachyptera* Kieffer, 1903
- PHYTOSCIARA** Frey, 1942
- Phytosciara halterata* (Lengersdorf, 1926)
Phytosciara macrotricha (Lengersdorf, 1926)
- PNYXIA** Johannsen, 1912
- Pnyxia scabiei* (Hopkins, 1895)
- PNYXIOPSIS** Tuomikoski, 1960
- Pnyxiopsis aliger* Tuomikoski, 1960
Pnyxiopsis degener (Tuomikoski, 1957)
- PROSCIARA** Frey, 1942
- Prosciara furtiva* Vilkamaa & Hippa, 1996
Prosciara plusiochaeta Hippa & Vilkamaa, 1991
Prosciara porrecta (Lengersdorf, 1929)
Prosciara producta (Tuomikoski, 1960)
Prosciara prosciaroides (Tuomikoski, 1960)
Prosciara ungulata (Winnertz, 1867)
- PSEUDOLYCORIELLA** Menzel & Mohrig, 1998
- Pseudolycoriella brunnea* (Bukowski & Lengersdorf, 1936)
Pseudolycoriella japonensis (Mohrig & Menzel, 1992)
Pseudolycoriella koreensis (Mohrig & Menzel, 1992)
Pseudolycoriella monticula (Mohrig & Menzel, 1992)
Pseudolycoriella nodulosa (Mohrig & Krivosheina, 1985)
Pseudolycoriella paludum (Frey, 1948)
Pseudolycoriella subbruckii (Mohrig & Hövemeyer, 1992)
- SCATOPSCIARA** Edwards, 1927
- sg. Scatopsciara** Edwards, 1927
- Scatopsciara atomaria* (Zetterstedt, 1851)
 = *vivida* (Winnertz, 1867)
- Scatopsciara bucera* Rudzinski, 1994
Scatopsciara calamophila Frey, 1948
Scatopsciara curvitorceps (Bukowski & Lengersdorf, 1936)
 = *myrmecophila* Frey, 1948
- Scatopsciara edwardsi* Freeman, 1983
Scatopsciara fluviatilis (Lengersdorf, 1940)
Scatopsciara geophila (Tuomikoski, 1960)
Scatopsciara multispina (Bukowski & Lengersdorf, 1936)
Scatopsciara neglecta Menzel & Mohrig, 1998
Scatopsciara pusilla (Meigen, 1818)

Scatopsciara subcalamophila Menzel & Mohrig, 1991

Scatopsciara subciliata Tuomikoski, 1960

Scatopsciara tricuspidata (Winnertz, 1867)

= *degenerans* (Frey, 1948)

Scatopsciara vitripennis (Meigen, 1818)

sg. *Xenopygina* Frey, 1948

Scatopsciara gabyae (Heller, 1998)

Scatopsciara obliqua Vilkamaa, Hippa & Heller, 2013

Scatopsciara paradoxa (Frey, 1948)

Scatopsciara simillima (Tuomikoski, 1960)

SCHWENCKFELDINA Frey, 1942

Schwenckfeldina carbonaria (Meigen, 1830)

Schwenckfeldina pectinea Menzel & Mohrig, 1991

SCIARA Meigen, 1803

Sciara flavimana Zetterstedt, 1851

Sciara hebes (Loew, 1869)

= *mendax* Tuomikoski., 1960

= *nursei* Freeman, 1983

= *ulrichi* Menzel & Mohrig, 1998

Sciara helvola Winnertz, 1867

Sciara hemeroboides (Scopoli, 1763)

= *thomae* (Linnaeus, 1767)

Sciara humeralis Zetterstedt, 1851

Sciara lackschewitzii (Lengersdorf, 1934)

SCYTHROPOCHROA Enderlein, 1911

Scythropochroa quercicola (Winnertz, 1869)

Scythropochroa radialis Lengersdorf, 1926

TRICHOSIA Winnertz, 1867

= *Leptosciara* Frey, 1942

= *Lestremioides* Frey, 1942

Trichosia Winnertz, 1867

Trichosia acrotricha Tuomikoski, 1960

Trichosia borealis (Frey, 1942)

Trichosia confusa Menzel & Mohrig, 1997

Trichosia flavicoxa Tuomikoski, 1960

Trichosia diota (Garrett, 1925)

Trichosia glabra (Meigen, 1830)

Trichosia habilis (Johannsen, 1912)

= *edwardsi* (Lengersdorf, 1930)

Trichosia morio (Fabricius, 1794)

= *caudata* (Walker, 1848)

Trichosia splendens Winnertz, 1867

Trichosia ussurica Mohrig & Antonova, 1978

XYLOSCIARA Tuomikoski, 1957

sg. *Protoxylosciara* Tuomikoski, 1960

Xylosciara longiforceps (Bukowski & Lengersdorf, 1936)

sg. *Xylosciara* Tuomikoski, 1957

Xylosciara heptacantha Tuomikoski, 1957

Xylosciara lignicola (Winnertz, 1867)

Xylosciara microdon (Frey, 1948)

Xylosciara misella (Frey, 1948)

Xylosciara phryganophila (Frey, 1948)

Xylosciara senta Vilkamaa, Hippa & Heller, 2013

Xylosciara spinata (Pettey, 1918)

= *betulae* Tuomikoski, 1960

Xylosciara steleocera Tuomikoski, 1960

Xylosciara trimera Tuomikoski, 1960

Xylosciara validinervis Tuomikoski, 1960

ZYGONEURA Meigen, 1830

sg. *Allozygoneura* Menzel & Mohrig, 1998

Zygoneura calthae Tuomikoski, 1960

sg. *Zygoneura* Meigen, 1830

Zygoneura sciarina Meigen, 1830

UNPLACED IN SCIAROIDEA

SCIAROSOMA Chandler, 2002

Sciarosoma nigriclava (Strobl, 1898)

= *borealis* Chandler, 2002

Excluded species

Lycoriella auripila (Winnertz, 1867), misidentified in Tuomikoski (1960).

Bradysia betuleti (Lengersdorf, 1940), misidentified in Tuomikoski (1960).

Bradysia subbetuleti Mohrig & Krivosheina, 1989, misidentified in Salmela and Vilkamaa (2005).

Corynoptera concinna (Winnertz, 1867), misidentified in Tuomikoski (1960).

Corynoptera bistrispina (Bukowski & Lengersdorf, 1936), misidentified in Tuomikoski (1960).

Trichosia trochanterata (Zetterstedt, 1851), misidentified in Tuomikoski (1960).

Acknowledgements

I thank my colleagues Kai Heller, Heikki Hippa, Frank Menzel, Werner Mohrig and Jukka Salmela for their long-term help in identifying Sciaridae.

References

- Frey R (1948) Entwurf einer neuen Klassifikation der Mückenfamilie Sciaridae. II. Die nord-europäischen Arten. Notulae Entomologique 27: 1–92.
- Hackman W (1980) A check list of the Finnish Diptera I. Nematocera and Brachycera (s.str.). Notulae Entomologique 60: 17–48.
- Menzel F, Mohrig W (2000) Revision der paläarktischen Trauermücken (Diptera: Sciaridae). Studia dipterologica Supplement 6: 1–761.
- Mohrig W, Heller K, Hippa H, Vilkamaa P, Menzel F (2013) Revision of the black fungus gnats (Diptera: Sciaridae) of North America. Studia dipterologica 19: 141–286.
- Salmela J, Vilkamaa, P (2005) Sciaridae (Diptera) from central Finland: faunistics and taxonomy. Entomologica Fennica 16: 287–300.
- Shin S, Jung S, Menzel F, Heller K, Lee H, Lee S (2013) Molecular phylogeny of black fungus gnats (Diptera: Sciaroidea: Sciaridae) and the evolution of larval habitats. Molecular Phylogenetics and Evolution 66: 833–846. doi: 10.1016/j.ympev.2012.11.008
- Silfverberg H (2012) Changes 2006–2010 in the list of Finnish insects. Entomologica Fennica 23: 18–41.
- Tuomikoski R (1960) Zur Kenntnis der Sciariden (Dipt.) Finnlands. Annales Zoologici Societatis Zoologicae Botanicae Fennicae 'Vanamo' 21: 1–164.
- Vilkamaa P (2000) Phylogeny of *Prosciara* Frey and related genera (Diptera: Sciaridae). Systematic Entomology 25: 47–72. doi: 10.1046/j.1365-3113.2000.00094.x
- Vilkamaa P (2003) Review of the genus *Baeosciara* Tuomikoski (Diptera: Sciaridae). Entomologica Fennica 14: 220–226.
- Vilkamaa P, Hippa H (2005) Phylogeny of *Peyerimhoffia* Kieffer, with the revision of the species. Insect Systematics and Evolution 35: 457–480. doi: 10.1163/187631204788912445
- Vilkamaa P, Heller K, Hippa H (2013) New species of Sciaridae (Diptera) to the Finnish fauna. Sahlbergia 19: 20–29.