

Checklist of the families Scathophagidae, Fanniidae and Muscidae of Finland (Insecta, Diptera)

Jere Kahanpää¹, Antti Haarto²

1 Finnish Museum of Natural History, Zoology Unit, P.O. Box 17, FI-00014 University of Helsinki, Finland

2 Zoological Museum, Section of Biodiversity and Environmental Science, Department of Biology, University of Turku, FI-20014 Turku, Finland

Corresponding author: Jere Kahanpää (jere.kahanpaa@helsinki.fi)

Academic editor: J. Salmela | Received 5 February 2014 | Accepted 19 May 2014 | Published 19 September 2014

<http://zoobank.org/776A4380-DB69-43C6-96AA-B6E336E68E1B>

Citation: Kahanpää J, Haarto A (2014) Checklist of the families Scathophagidae, Fanniidae and Muscidae of Finland (Insecta, Diptera). In: Kahanpää J, Salmela J (Eds) Checklist of the Diptera of Finland. ZooKeys 441: 347–367. doi: [10.3897/zookeys.441.7142](https://doi.org/10.3897/zookeys.441.7142)

Abstract

A revised checklist of the Scathophagidae, Fanniidae and Muscidae recorded from Finland is presented. *Phaonia amicula* Villeneuve, 1922 is noted from Finland for the first time.

Keywords

Species list, Finland, Diptera, biodiversity, faunistics

Introduction

Four families make up the traditional superfamily Muscoidea: Scathophagidae, Anthomyiidae, Fanniidae and Muscidae. The monophyly of the superfamily has been strongly questioned (Nirmala et al. 2001, Kutty et al. 2010) on the basis of DNA sequence analyses.

Three of the four families of Muscoidea are treated in this paper, the fourth (Anthomyiidae) is covered in a separate paper in this issue of ZooKeys. All three are important components of the Finnish fly fauna as can be seen from the large number of species involved and the fact that the species recorded from the country represent up to one fifth of the world fauna in the case of Scathophagidae.

Table 1. Number of species by family.

Family	Number of species in			Level of knowledge
	World (Pape et al. 2011)	Europe	Finland	
Scathophagidae	419	158	85–86	good
Fanniidae	359	83	61	average
Muscidae	5218	572	307–309	average

Unfortunately the muscoid flies have been poorly collected and studied in Finland in comparison with many other Diptera families. Lauri Tiensuu concentrated on these families during the 1930s (1935, 1938, 1939) and Walter Hackman (1956) reviewed the Finnish Scathophagidae. During the last ten years, both the fanniids (Kahanpää and Haarto 2013) and the muscids (Kahanpää 2013) have attracted some attention in the country. The number of species known from the country has increased markedly since the year 2000.

The Finnish muscoid species were last listed by Hackman (1980).

Checklist

suborder Brachycera Macquart, 1834
 clade Eremoneura Lameere, 1906
 clade Cyclorrhapha Brauer, 1863
 infraorder Schizophora Becher, 1882
 clade Muscaria Enderlein, 1936
 parvorder Calyptatae Robineau-Desvoidy, 1830
 superfamily Muscoidea Latreille, 1802

SCATHOPHAGIDAE Robineau-Desvoidy, 1830

DELININAE Séguy, 1952

DELINA Robineau-Desvoidy, 1830

Delina nigrita (Fallén, 1819)

LEPTOPA Zetterstedt, 1838

Leptopa filiformis Zetterstedt, 1838

MICROPSELAPHA Becker, 1894

Micropselapha filiformis (Zetterstedt, 1846)

PARALLELOMMA Strobl, 1894

= *Chylizosoma* Hendel, 1924

Parallelomma medium Becker, 1894

? *Parallelomma paridis* Hering, 1923 (see Notes)

Parallelomma sellatum (Hackman, 1956)

Parallelomma vittatum (Meigen, 1826)

PHROSIA Robineau-Desvoidy, 1830

Phrosia albilabris (Fabricius, 1805)

SCATHOPHAGINAE Robineau-Desvoidy, 1830

ACANTHOCNEMA Becker, 1894

sg. *Clinoceroides* Hendel, 1917

Acanthocnema glaucescens (Loew, 1864)

= *nigripes* Ringdahl, 1936

ACEROCNEMA Becker, 1894

Acerocnema macrocera (Meigen, 1826)

= *tiefi* Becker, 1894

= *pokornyi* Becker, 1894

ALLOMYELLA Malloch, 1923

Allomyella albipennis (Zetterstedt, 1838)

Allomyella frigida (Holmgren, 1883)

= *portenkoi* (Stackelberg, 1952) misid.

BOSTRICHOPYGA Becker, 1894

Bostrichopyga crassipes (Zetterstedt, 1838)

CHAETOSA Coquillett, 1898

Chaetosa punctipes (Meigen, 1826)

CLEIGASTRA Macquart, 1835

= *Cnemopogon* Rondani, 1856

Cleigastra apicalis (Meigen, 1826)

CONISTERNUM Strobl, 1894

= *Coniosternum* Becker, 1894

Conisternum lapponicum (Ringdahl, 1920)

Conisternum obscurum (Fallén, 1819)

Conisternum tinctinerve (Becker, 1894)

CORDILURA Fallén, 1810

= *Cordylura* Meigen, 1826 emend.

sg. *Cordilura* Fallén, 1810

Cordilura aberrans Becker, 1894

Cordilura aemula Collin, 1958

Cordilura atrata Zetterstedt, 1846

Cordilura ciliata Meigen, 1826

Cordilura picipes Meigen, 1826

Cordilura picticornis Loew, 1864

Cordilura proboscidea Zetterstedt, 1838

Cordilura pubera (Linnaeus, 1758)

Cordilura pudica Meigen, 1826

Cordilura rufimana Meigen, 1826

Cordilura socialis (Becker, 1894)

= *freyi* Hackman, 1956

sg. *Cordilurina* James, 1955

= *Parallelomma* Becker, 1894 preocc.

- Cordilura albipes* Fallén, 1819
Cordilura fuscipes Zetterstedt, 1838
sg. *Scoliaphleps* Becker, 1894
Cordilura ustulata Zetterstedt, 1838
 = *melanacra* Loew, 1873
COSMETOPUS Becker, 1894
Cosmetopus dentimana (Zetterstedt, 1838)
 = *fulvipes* (Zetterstedt, 1838)
Cosmetopus longa (Walker, 1849)
 = *bergrothi* Becker, 1900 in part
 = *fulvipes* auct. nec (Zetterstedt, 1838)
Cosmetopus ringdahli Andersson, 1974
 = *bergrothi* Becker, 1900 in part
ERNONEURA Becker, 1894
Ernoneura argus (Zetterstedt, 1838)
GONARCTICUS Becker, 1894
Gonarcticus abdominalis (Zetterstedt, 1846)
GONATHERUS Rondani, 1856
Gonatherus planiceps (Fallén, 1826)
GIMNOMERA Rondani, 1866
 = *Cochliarium* Becker, 1894
Gimnomera albipila (Zetterstedt, 1846)
Gimnomera cuneiventris (Zetterstedt, 1846)
Gimnomera dorsata (Zetterstedt, 1838)
Gimnomera hirta Hendel, 1930
Gimnomera tarsaea (Fallén, 1819)
HEXAMITOCERA Becker, 1894
Hexamitocera loxoceratum (Fallén, 1826)
HYDROMYZA Fallén, 1823
Hydromyza livens (Fabricius, 1794)
MEGAPHTHALMA Becker, 1894
Megaphtalma pallida (Fallén, 1819)
MEGAPHTHALMOIDES Ringdahl, 1936
Megaphtalmoides unilineatus (Zetterstedt, 1838)
MICROPROSOPA Strobl, 1894
Microprosopa haemorrhoidalis (Meigen, 1826)
Microprosopa lacteipennis Ringdahl, 1920
Microprosopa lineata (Zetterstedt, 1838)
Microprosopa pallidicauda (Zetterstedt, 1838)
NANNA Strobl, 1894
 = *Amaurosoma* Becker, 1894
Nanna armillata (Zetterstedt, 1846)
Nanna articulata (Becker, 1894)

- Nanna bispinosa* (Malloch, 1920)
Nanna brevifrons (Zetterstedt, 1838)
Nanna fasciata (Meigen, 1826)
Nanna flavipes (Fallén, 1819)
 = *minuta* (Becker, 1894)
 ?= *multisetosa* (Hackman, 1956)
Nanna inermis (Becker, 1894)
Nanna leucostoma (Zetterstedt, 1846)
Nanna tibiella (Zetterstedt, 1838)
 = *nigripes* (Zetterstedt, 1846)
NORELLIA Robineau-Desvoidy, 1830
Norellia tipularia (Fabricius, 1794) (see Notes)
NORELLISOMA Hendel, 1910
Norellisoma lituratum (Wiedemann, 1826)
Norellisoma spinimanum (Fallén, 1819)
OKENIELLA Hendel, 1907
Okeniella caudata (Zetterstedt, 1838)
Okeniella dasyprocta (Loew, 1864)
ORTHACHETA Becker, 1894
Orthacheta pilosa (Zetterstedt, 1838)
PLEUROCHAETELLA Vockeroth, 1965
Pleurochaetella simplices (Becker, 1900)
POGONOTA Zetterstedt, 1860
sg. Pogonota Zetterstedt, 1860
Pogonota barbata (Zetterstedt, 1838)
sg. Lasioscelus Becker, 1894
Pogonota immunda (Zetterstedt, 1838)
 = *clavatus* (Zetterstedt, 1846)
Pogonota sahlbergi Becker, 1900
SCATHOPHAGA Meigen, 1803
 = *Scatophaga* Fabricius, 1805 emend.
 = *Scopeuma* Meigen, 1800 suppr.
 = *Scatomyza* Fallén, 1810
Scathophaga apicalis Curtis in Ross, 1835
Scathophaga furcata (Say, 1823)
Scathophaga incola Becker, 1900
Scathophaga inquinata (Meigen, 1826)
 = *analis* Meigen, 1826
Scathophaga litorea (Fallén, 1819)
Scathophaga lutaria (Fabricius, 1794)
Scathophaga obscurinervis Becker, 1900
Scathophaga pictipennis Oldenberg, 1923
Scathophaga scybalaria (Linnaeus, 1758)

Scathophaga stercoraria (Linnaeus, 1758)

Scathophaga suilla (Fabricius, 1794)

SPAZIPHORA Rondani, 1856

Spaziphora hydromyzina (Fallén, 1819)

= *fascipes* (Becker, 1894)

STAEGERIA Rondani, 1856

Staegeria kunzei (Zetterstedt, 1821)

TRICHOPALPUS Rondani, 1856

Trichopalpus fraternus (Meigen, 1826)

Trichopalpus nigribasis Curran, 1927

= *pilirostris* (Ringdahl, 1936)

Trichopalpus obscurellus (Zetterstedt, 1846)

= *subarcticus* (Ringdahl, 1936)

FANNIIDAE Schnabl & Dziezicki, 1911

FANNIA Robineau-Desvoidy, 1830

= *Homalomyia* Bouché, 1834

= *Coelomyia* Haliday, 1840

Fannia aethiops Malloch, 1913

Fannia alpina Pont, 1970

Fannia armata (Meigen, 1826)

Fannia atra (Stein, 1895)

Fannia canicularis (Linnaeus, 1761)

Fannia carbonaria (Meigen, 1826)

Fannia carbonella (Stein, 1895)

Fannia coracina (Loew, 1873)

Fannia corvina (Verrall, 1892)

= *halterata* Ringdahl, 1918

Fannia cothurnata (Loew, 1873)

Fannia difficilis (Stein, 1895)

Fannia fuscitibia Stein, 1920

Fannia fuscula (Fallén, 1825)

Fannia genualis (Stein, 1895)

Fannia gotlandica Ringdahl, 1926

Fannia hirticeps (Stein, 1892)

Fannia hirundinis Ringdahl, 1948

Fannia immutica Collin, 1939

Fannia incisurata (Zetterstedt, 1838)

Fannia latifrontalis Hennig, 1955

Fannia latipalpis (Stein, 1892)

Fannia lepida (Wiedemann, 1817)

= *mutica* (Zetterstedt, 1845)

Fannia leucosticta (Meigen, 1838)

- Fannia limbata* (Tiensuu, 1938)
= *berolinensis* Hennig, 1955
- Fannia lucidula* (Zetterstedt, 1860)
= *glaucescens* auct. nec (Zetterstedt, 1845)
- Fannia lugubrina* (Zetterstedt, 1838)
- Fannia lustrator* (Harris, 1780)
= *hamata* (Macquart, 1835)
- Fannia manicata* (Meigen, 1826)
- Fannia melania* (Dufour, 1839)
= *ciliata* (Stein, 1895)
- Fannia metallipennis* (Zetterstedt, 1838)
= *kowarzi* (Verrall, 1892)
- Fannia minutipalpis* (Stein, 1895)
- Fannia mollissima* (Haliday, 1840)
= *spathulata* (Zetterstedt, 1845)
- Fannia monilis* (Haliday, 1838)
- Fannia pallitibia* (Rondani, 1866)
- Fannia parva* (Stein, 1895)
- Fannia pauli* Pont, 1997
= *nitida* (Stein, 1895) preocc.
- Fannia polychaeta* (Stein, 1895)
- Fannia postica* (Stein, 1895)
- Fannia posticata* (Meigen, 1826)
= *pretiosa* (Schiner, 1862)
- Fannia rabdionata* Karl, 1940
- Fannia ringdahlana* Collin, 1939
= *umbrosa* auct. nec (Stein, 1895)
- Fannia rondanii* (Strobl, 1893)
= *carbonaria* (Rondani, 1871) preocc.
= *aerea* misid.
- Fannia scalaris* (Fabricius, 1794)
= *subscalaris* Zimin, 1946
- Fannia serena* (Fallén, 1825)
- Fannia similis* (Stein, 1895)
- Fannia slovaca* Gregor & Rozkošný, 2005
- Fannia sociella* (Zetterstedt, 1845)
- Fannia spathiophora* Malloch 1918
= *nodulosa* Ringdahl, 1926
- Fannia speciosa* (Villeneuve, 1898)
- Fannia stigi* Rognes, 1982
- Fannia subatripes* d'Assis-Fonseca, 1967
- Fannia subpellucens* (Zetterstedt, 1845)
- Fannia subpubescens* Collin, 1958

Fannia tuberculata (Zetterstedt, 1849)

Fannia umbratica Collin, 1939

Fannia umbrosa (Stein, 1895)

= *subumbrosa* Ringdahl, 1934

Fannia verrallii (Stein, 1895)

Fannia vesparia (Meade, 1891)

Fannia vespertilionis Ringdahl, 1934

PIEZURA Rondani, 1844

= *Platycoenosia* Strobl, 1894

Piezura graminicola (Zetterstedt, 1846)

= *boletorum* (Rondani, 1866)

= *mikii* (Strobl, 1894)

Piezura pardalina (Rondani, 1866)

= *graminicola* auct. nec (Zetterstedt, 1846)

MUSCIDAE Latreille, 1802

ACHANTHIPTERINAE Hennig, 1962

ACHANTHIPTERA Rondani, 1856

Achanthiptera rohrelliformis (Robineau-Desvoidy, 1830)

= *inanis* (Fallén, 1825) preocc.

COENOSIINAE Verrall, 1888

tribe Coenosiini Verrall, 1888

COENOSIA Meigen, 1826

= *Caricea* Robineau-Desvoidy, 1830

= *Oplogaster* Rondani, 1856

= *Dexiopsis* Pokorný, 1893

Coenosia acuminata Strobl, 1898

= *annulipes* Ringdahl, 1932

Coenosia agromyzina (Fallén, 1825)

Coenosia ambulans Meigen, 1826

Coenosia bilineella (Zetterstedt, 1838)

Coenosia campestris (Meigen, 1830)

= *sexnotata* auct. nec Meigen, 1826

? *Coenosia comita* (Huckett, 1936) (see Notes)

= *ovulifera* Tiensuu, 1938

Coenosia dealbata (Zetterstedt, 1838)

= *fulvicornis* (Zetterstedt, 1845)

Coenosia flavimana (Zetterstedt, 1845)

= *albatella* (Zetterstedt, 1849)

Coenosia humilis Meigen, 1826

Coenosia intermedia (Fallén, 1825)

Coenosia lacteipennis (Zetterstedt, 1845)

Coenosia lineatipes (Zetterstedt, 1845)

- = *albicornis* misid.
Coenosia means Meigen, 1826
Coenosia mollicula (Fallén, 1825)
Coenosia octopunctata (Zetterstedt, 1838)
Coenosia paludis Tiensuu, 1939
Coenosia pedella (Fallén, 1825)
= *decipiens* Meigen, 1826
Coenosia perpusilla Meigen, 1826
Coenosia pudorosa Collin, 1953
Coenosia pulicaria (Zetterstedt, 1845)
Coenosia pumila (Fallén, 1825)
Coenosia pygmaea (Zetterstedt, 1845)
Coenosia ruficornis Macquart, 1835
= *litoralis* (Zetterstedt, 1846)
Coenosia rufipalpis Meigen, 1826
= *flavicauda* Ringdahl, 1937
Coenosia sallae Tiensuu, 1938
Coenosia testacea (Robineau-Desvoidy, 1830)
= *tricolor* (Zetterstedt, 1845)
= *alleni* d'Assis-Fonseca, 1966
Coenosia tigrina (Fabricius, 1775)
Coenosia trilineella (Zetterstedt, 1838)
= *trilineata* emend.
Coenosia verralli Collin, 1953
= *steini* Verrall, 1912 preocc.
LIMNOSPILA Schnabl, 1902
Limnospila albifrons (Zetterstedt, 1849)
LISPOCEPHALA Pokorný, 1893
Lispocephala alma (Meigen, 1826)
Lispocephala erythrocerata (Robineau-Desvoidy, 1830)
Lispocephala falculata Collin, 1963
Lispocephala fuscitibia Ringdahl, 1944
Lispocephala pallipalpis (Zetterstedt, 1845)
Lispocephala spuria (Zetterstedt, 1838)
= *vitrifennis* Ringdahl, 1951
Lispocephala verna (Fabricius, 1794)
MACRORCHIS Rondani, 1877
Macrorchis meditata (Fallén, 1825)
PSEUDOCOENOSIA Stein, 1916
Pseudocoenosia abnormis Stein, 1916
Pseudocoenosia solitaria (Zetterstedt, 1838)
= *longicauda* (Zetterstedt, 1860)
SCHOENOMYZA Haliday, 1833

- Schoenomyza litarella* (Fallén, 1823)
tribe Limnophoriini Villeneuve, 1902
LIMNOPHORA Robineau-Desvoidy, 1830
Limnophora nigripes (Robineau-Desvoidy, 1830)
Limnophora pandellei Séguy 1923 (see Notes)
Limnophora riparia (Fallén, 1824)
Limnophora rotundata Collin, 1930
Limnophora tigrina (Am Stein, 1860)
 = *notata* (Fallén, 1823) preocc.
Limnophora triangula (Fallén, 1825)
Limnophora uniseta Stein, 1916
LISPE Latreille, 1796
Lispe consanguinea Loew, 1858
Lispe hydromyzina Fallén, 1825
Lispe litorea Fallén, 1825
Lispe melaleuca Loew, 1847
Lispe pygmaea Fallén, 1825
Lispe tentaculata (De Geer, 1776)
Lispe uliginosa Fallén, 1825
SPILOGONA Schnabl, 1911
Spilogona aerea (Fallén, 1825)
Spilogona albisquama (Ringdahl, 1932)
Spilogona alpica (Zetterstedt, 1845)
Spilogona arenosa (Ringdahl, 1918)
Spilogona atrisquamula Hennig, 1959
Spilogona baltica (Ringdahl, 1918)
Spilogona brunneifrons Ringdahl, 1931
Spilogona brunneisquama (Zetterstedt, 1845)
Spilogona carbonella (Zetterstedt, 1845)
Spilogona contractifrons (Zetterstedt, 1838)
Spilogona denigrata (Meigen, 1826)
Spilogona depressiuscula (Zetterstedt, 1838)
Spilogona depressula (Zetterstedt, 1845)
Spilogona dispar (Fallén, 1823)
 = *funeralis* Rondani, 1866
Spilogona falleni Pont, 1984
 = *litorea* auct. nec (Fallén, 1823)
Spilogona krogerusi (Ringdahl, 1941)
 = *micans* misid.
Spilogona leucogaster (Zetterstedt, 1838)
Spilogona malaisei (Ringdahl, 1920)
Spilogona marginifera Hennig, 1959
 = *marginalis* (Fallén, 1824) preocc.

- Spilogona meadei* (Schnabl, 1915)
Spilogona megastoma (Bohemian, 1866)
Spilogona nigriventris (Zetterstedt, 1845)
Spilogona nitidicauda (Schnabl, 1911)
Spilogona novemmaculata (Zetterstedt, 1860)
Spilogona opaca (Schnabl, 1915)
Spilogona pacifica (Meigen, 1826)
 = *vana* (Zetterstedt, 1845)
Spilogona palmeni (Ringdahl, 1935)
Spilogona pseudodispar (Frey, 1915)
 = *spinitibia* (Ringdahl, 1918)
Spilogona pusilla (Huckett, 1932)
Spilogona quinquelineata (Zetterstedt, 1838)
Spilogona semiglobosa (Ringdahl, 1916)
Spilogona sororcula (Zetterstedt, 1845)
 = *zetterstedti* (Ringdahl, 1918) preocc.
Spilogona spectabilis (Tiensuu, 1938)
Spilogona surda (Zetterstedt, 1845)
Spilogona tenuis Hennig, 1959
Spilogona tornensis (Ringdahl, 1926)
Spilogona triangulifera (Zetterstedt, 1838)
Spilogona trianguligera (Zetterstedt, 1838)
 = *insularis* (Collin, 1921)
Spilogona trigonata (Zetterstedt, 1838)
Spilogona tundrae (Schnabl, 1915)
 = *macropyga* (Frey, 1915)
Spilogona tundrica (Schnabl & Dziedzicki, 1911)
Spilogona varsaviensis (Schnabl & Dziedzicki, 1911)
 = *glaucha* (Stein, 1916)
Spilogona veterrima (Zetterstedt, 1845)
VILLENEUVIA Schnabl & Dziedzicki, 1911
Villeneuvia aestuum (Villeneuve, 1902)
AZELIINAE Robineau-Desvoidy, 1830
tribe Azeliini Robineau-Desvoidy, 1830
AZELIA Robineau-Desvoidy, 1830
Azelia aterrima (Meigen, 1826)
Azelia cilipes (Haliday, 1838)
Azelia gibbera (Meigen, 1826)
Azelia monodactyla Loew, 1874
Azelia nebulosa Robineau-Desvoidy, 1830
 = *macquarti* (Staeger in Schiødte, 1843)
Azelia trigonica Hennig, 1956
 = *nuda* Hennig, 1956

Azelia triquetra (Wiedemann, 1817)

Azelia zetterstedtii Rondani, 1866

DRYMEIA Meigen, 1826

= *Pogonomyia* Rondani, 1871

= *Trichopticoides* Ringdahl, 1931

Drymeia hamata (Fallén, 1823)

Drymeia tetra (Meigen, 1826)

Drymeia vicana (Harris, 1780)

= *decolor* (Fallén, 1824)

HYDROTAEA Robineau-Desvoidy, 1830

= *Ophyra* Robineau-Desvoidy, 1830

= *Lasiops* Meigen, 1838

Hydrotaea aenescens (Wiedemann, 1830)

Hydrotaea albipuncta (Zetterstedt, 1845)

Hydrotaea anxia (Zetterstedt, 1838)

= *bispinosa* (Zetterstedt, 1845)

Hydrotaea armipes (Fallén, 1825)

= *occulta* (Meigen, 1826)

Hydrotaea basdeni Collin, 1939

Hydrotaea borussica Stein, 1899

Hydrotaea cyrtoneurina (Zetterstedt, 1845)

Hydrotaea dentipes (Fabricius, 1805)

Hydrotaea diabolus (Harrpis, 1780)

= *ciliata* (Fabricius, 1794) preocc.

= *bimaculata* (Meigen, 1826)

Hydrotaea floccosa Macquart, 1835

= *armipes* auct. nec (Fallén, 1825)

Hydrotaea ignava (Harrpis, 1780)

= *leucostoma* (Wiedemann, 1817)

Hydrotaea irritans (Fallén, 1823)

Hydrotaea meridionalis Portschinsky, 1882

Hydrotaea meteorica (Linnaeus, 1758)

Hydrotaea militaris (Meigen, 1826)

Hydrotaea nidicola Malloch, 1925

Hydrotaea palaestrica (Meigen, 1826)

Hydrotaea pandellei Stein, 1899

Hydrotaea parva Meade, 1889

Hydrotaea pellucens Portschinsky, 1879

Hydrotaea pilipes Stein, 1903

Hydrotaea pilitibia Stein, 1916

Hydrotaea ringdahli Stein, 1916

Hydrotaea scambus (Zetterstedt, 1838)

Hydrotaea similis Meade, 1887

Hydrotaea tuberculata Rondani, 1866

Hydrotaea velutina Robineau-Desvoidy, 1830

POTAMIA Robineau-Desvoidy, 1830

= *Dendrophaonia* Malloch, 1923

Potamia littoralis Robineau-Desvoidy, 1830

= *querceti* (Bouché, 1834)

THRICOPS Rondani, 1856

= *Alleostylus* Schnabl, 1888

Thricops aculeipes (Zetterstedt, 1838)

Thricops albibasalis (Zetterstedt, 1849)

= *sudeticus* auct. nec (Schnabl, 1888)

Thricops coquilletti (Malloch, 1920)

Thricops cunctans (Meigen, 1826)

= *hirsutulus* auct. nec (Zetterstedt, 1838)

Thricops diaphanus (Wiedemann, 1817)

Thricops foveolatus (Zetterstedt, 1845)

Thricops genarum (Zetterstedt, 1838)

= *sundewalli* (Zetterstedt, 1845)

Thricops hirtulus (Zetterstedt, 1838)

= *subrostratus* (Zetterstedt, 1845)

Thricops innocuus (Zetterstedt, 1838)

Thricops lividiventris (Zetterstedt, 1845)

Thricops longipes (Zetterstedt, 1845)

Thricops nigrifrons (Robineau-Desvoidy, 1830)

Thricops nigrithellus (Zetterstedt, 1838)

Thricops rostratus (Meade, 1882)

Thricops rufisquamus (Schnabl, 1915)

= *penicillatus* (Ringdahl, 1926)

Thricops semicinereus (Wiedemann, 1817)

Thricops separ (Zetterstedt, 1845)

Thricops simplex (Wiedemann, 1817)

tribe Reinwardtiini Brauer & Bergenstamm, 1889

MUSCINA Robineau-Desvoidy, 1830

Muscina angustifrons (Loew, 1858)

Muscina levida (Harris, 1780)

= *assimilis* (Fallén, 1823)

Muscina pascuorum (Meigen, 1826)

Muscina prolapsa (Harris, 1780)

= *pabulorum* (Fallén, 1817)

Muscina stabulans (Fallén, 1817)

MUSCINAE Latreille, 1802

tribe Muscini Latreille, 1802

EUDASYPHORA Townsend, 1911

Eudasypnora cyanicolor (Zetterstedt, 1845)

Eudasypnora zimini (Hennig, 1963)

MESEMBRINA Meigen, 1826

= *Hypodermodes* Townsend, 1912

Mesembrina intermedia Zetterstedt, 1848?

Mesembrina meridiana (Linnaeus, 1758)

Mesembrina mystacea (Linnaeus, 1758)

Mesembrina resplendens Wahlberg, 1844

MORELLIA Robineau-Desvoidy, 1830

sg. Morellia Robineau-Desvoidy, 1830

Morellia aenescens Robineau-Desvoidy, 1830

Morellia hortorum (Fallén, 1817)

Morellia podagrifica (Loew, 1857)

sg. Ziminiella Nihei & Carvalho, 2007

Morellia simplex (Loew, 1857)

MUSCA Linnaeus, 1758

Musca autumnalis De Geer, 1776

= *corvina* Fabricius, 1781

Musca domestica Linnaeus, 1758

Musca tempestiva Fallén, 1817

NEOMYIA Walker, 1859

= *Orthellia* Robineau-Desvoidy, 1863

Neomyia cornicina (Fabricius, 1781)

= *caesarion* (Meigen, 1826)

= *fennica* (Frey, 1909)

Neomyia viridescens (Robineau-Desvoidy, 1830)

= *cornicina* auct. nec (Fabricius, 1781)

POLIETES Rondani, 1866

Polietes domitor (Harris, 1780)

= *albolineatus* (Fallén, 1823)

Polietes lardarius (Fabricius, 1781)

Polietes nigrolimbatus (von Bonsdorff, 1866)

Polietes steinii (Ringdahl, 1913)

PYRELLIA Robineau-Desvoidy, 1830

Pyrellia vivida Robineau-Desvoidy, 1830

= *cadaverina* auct. nec (Linnaeus, 1758)

tribe Stomoxyini Meigen, 1824

HAEMATOBIA Le Peletier & Serville, 1828

= *Lyperosia* Rondani, 1856

Haematobia irritans (Linnaeus, 1758)

HAEMATOBOSCA Bezzi, 1907

Haematobosca alcis Snow, 1891

= *crassipalpis* (Ringdahl, 1920)

Haematobosca stimulans (Meigen, 1824)

STOMOXYS Geoffroy, 1762

Stomoxys calcitrans (Linnaeus, 1758)

MYDAEINAE Verrall, 1888

GRAPHOMYA Robineau-Desvoidy, 1830

Graphomya maculata (Scopoli, 1763)

Graphomya minor Robineau-Desvoidy, 1830

GYMNODIA Robineau-Desvoidy, 1863

= *Brontaea* Kowarz, 1873

Gymnodia humilis (Zetterstedt, 1860)

HEBECNEMA Schnabl, 1889

Hebecnema fumosa (Meigen, 1826)

Hebecnema nigra (Robineau-Desvoidy, 1830)

= *vespertina* auct. nec (Fallén, 1823)

Hebecnema nigricolor (Fallén, 1825)

Hebecnema umbratica (Meigen, 1826)

Hebecnema vespertina (Fallén, 1823)

= *affinis* Malloch, 1921

MYDAEA Robineau-Desvoidy, 1830

Mydaea affinis Meade, 1891

= *discimana* Malloch, 1920

Mydaea ancilla (Meigen, 1826)

Mydaea anicula (Zetterstedt, 1860)

Mydaea corni Scopoli, 1763

= *pagana* (Fabricius, 1794) preocc.

= *scutellaris* Robineau-Desvoidy, 1830

Mydaea deserta (Zetterstedt, 1845)

Mydaea detrita (Zetterstedt, 1845)

= *electa* (Zetterstedt, 1860)

Mydaea humeralis Robineau-Desvoidy, 1830

= *tincta* (Zetterstedt, 1845)

Mydaea nebulosa (Stein, 1893)

Mydaea obscurella Malloch, 1921

= *bengtsoni* Ringdahl, 1924

Mydaea orthonevra (Macquart, 1835)

= *detrita* auct. nec (Zetterstedt, 1845)

Mydaea palpalis Stein, 1916

Mydaea setifemur Ringdahl, 1924

Mydaea sootryeni Ringdahl, 1928

Mydaea urbana (Meigen, 1826)

MYOSPILA Rondani, 1856

Myospila bimaculata (Macquart, 1834)

Myospila meditabunda (Fabricius, 1781)

OPSOLASIA Coquillett, 1910*Opsolasia orichalcea* (Zetterstedt, 1849)*PHAONIINAE* Malloch, 1917tribe *Phaoniini* Malloch, 1917**HELINA** Robineau-Desvoidy, 1830*Helina abdominalis* (Zetterstedt, 1846)*Helina allotalla* (Meigen, 1830)*Helina annosa* (Zetterstedt, 1838)*Helina atricolor* (Fallén, 1825)= *denudata* (Zetterstedt, 1845)= *denutata* emend.*Helina bohemani* (Ringdahl, 1916)*Helina celsa* (Harris, 1780)= *quadrimaculata* (Fallén, 1823) preocc.= *quadrimaculella* Hennig, 1957? *Helina ciliata* Karl, 1929 (see Notes)*Helina ciliatocosta* (Zetterstedt, 1845)= *ciliatocostata* emend.*Helina cilipes* (Schnabl, 1902)*Helina cinerella* (van der Wulp, 1867)= *vanderwulpi* (Schnabl, 1888)= *calceata* misid.*Helina confinis* (Fallén, 1825)= *anceps* (Zetterstedt, 1838)*Helina consimilis* (Fallén, 1825)*Helina cothurnata* (Rondani, 1866)= *obscuripes* auct. nec (Zetterstedt, 1845)*Helina daicles* (Walker, 1849)= *binotata* (Zetterstedt, 1845)*Helina depuncta* (Fallén, 1825)*Helina erecta* (Harris, 1780)= *lucorum* (Fallén, 1823) preocc.= *laetifica* (Robineau-Desvoidy, 1830)*Helina flavisquama* (Zetterstedt, 1849)*Helina fratercula* (Zetterstedt, 1845)*Helina fulvisquama* (Zetterstedt, 1845)*Helina impuncta* (Fallén, 1825)*Helina latitarsis* Ringdahl, 1924*Helina laxifrons* (Zetterstedt, 1860)*Helina longicornis* (Zetterstedt, 1838)*Helina luteisquama* (Zetterstedt, 1845)*Helina maculipennis* (Zetterstedt, 1845)= *obscuripes* (Zetterstedt, 1845)

- Helina obscurata* (Meigen, 1826)
Helina pertusa (Meigen, 1826)
Helina protuberans (Zetterstedt, 1845)
Helina pubiseta (Zetterstedt, 1845)
Helina quadrinotata (Meigen, 1826)
Helina quadrum (Fabricius, 1805)
Helina reversio (Harris, 1780)
 = *duplicata* (Meigen, 1826)
 = *duplaris* auct. nec (Zetterstedt, 1845)
 = *communis* (Robineau-Desvoidy, 1830)
Helina setiventris Ringdahl, 1924
Helina sexmaculata (Preyssler, 1791)
 = *uliginosa* (Fallén, 1825) preocc.
 = *punctata* (Robineau-Desvoidy, 1830)
Helina spinicosta (Zetterstedt, 1845)
Helina squalens (Zetterstedt, 1838)
 = *borealis* (Zetterstedt, 1838)
Helina subvittata (Séguy, 1923)
 = *rothi* Ringdahl, 1939
 = *marmorata* auct. nec (Zetterstedt, 1860)
Helina tetrastigma (Meigen, 1826)
 = *flagripes* (Rondani, 1866)
Helina trivittata (Zetterstedt, 1860)
 = *atripes* (Meade, 1889)
Helina veterana (Zetterstedt, 1838)
Helina vicina (Czerny, 1900)
 = *suecica* Ringdahl, 1924
- LOPHOSCELES** Ringdahl, 1922
- Lophosceles cinereiventris* (Zetterstedt, 1845)
 = *cristata* (Zetterstedt, 1845)
Lophosceles frenatus (Holmgren, 1872)
Lophosceles hians (Zetterstedt, 1838)
Lophosceles mutatus (Fallén, 1825)
- PHAONIA** Robineau-Desvoidy, 1830
- Phaonia aeneiventris* (Zetterstedt, 1845)
 = *cinctinervis* (Zetterstedt, 1860)
Phaonia alpicola (Zetterstedt, 1845)
Phaonia amabilis (Meigen, 1826)
Phaonia amicula Villeneuve, 1922 (see Notes)
Phaonia angelicae (Scopoli, 1763)
 = *basalis* (Zetterstedt, 1838)

- Phaonia angulicornis* (Zetterstedt, 1838)
= *erinacea* (Fallén, 1824)
- Phaonia apicalis* Stein, 1914
- Phaonia atriceps* (Loew, 1858)
- Phaonia atrocyanea* Ringdahl, 1916
- Phaonia canescens* Stein, 1916
- Phaonia consobrina* (Zetterstedt, 1838)
- Phaonia czernyi* Hennig, 1963
= *steinii* Czerny, 1900 preocc.
- Phaonia errans* (Meigen, 1826)
= *erratica* (Fallén, 1825) preocc.
- Phaonia erronea* (Schnabl, 1887)
- Phaonia falleni* (Michelsen, 1977)
= *vagans* (Fallén, 1825) preocc.
- Phaonia fugax* Tiensuu, 1946
- Phaonia fuscata* (Fallén, 1825)
- Phaonia gobertii* (Mik, 1881)
- Phaonia gracilis* Stein, 1916
- Phaonia grandaeva* (Zetterstedt, 1845)
- Phaonia halterata* (Stein, 1893)
- Phaonia hybrida* (Schnabl, 1888)
- Phaonia incana* (Wiedemann, 1817)
- Phaonia jaroschewskii* (Schnabl, 1888)
= *crinipes* Stein, 1913
- Phaonia kowarzii* (Schnabl, 1886)
= *fulvicornis* Tiensuu, 1936
- Phaonia laeta* (Fallén, 1823)
- Phaonia latipalpis* Schnabl, 1911
= *umbraticola* d'Assis-Fonseca, 1957
- Phaonia longicornis* Stein, 1916
- Phaonia lugubris* (Meigen, 1826)
= *morio* (Zetterstedt, 1845)
- Phaonia magnicornis* (Zetterstedt, 1845)
- Phaonia meigeni* Pont, 1986
= *lugubris* auct. nec (Meigen, 1826)
- Phaonia mystica* (Meigen, 1826)
= *vittifera* (Zetterstedt, 1845)
- Phaonia nymphaearum* (Robineau-Desvoidy, 1830)
= *nitida* (Macquart, 1835)
- Phaonia pallida* (Fabricius, 1787)
- Phaonia pallidisquama* (Zetterstedt, 1849)
- Phaonia palpata* (Stein, 1897)
- Phaonia perdita* (Meigen, 1830)
- Phaonia pratensis* (Robineau-Desvoidy, 1830)

- = *laeta* auct. nec (Fallén, 1923)
Phaonia rufipalpis (Macquart, 1835)
Phaonia rufiventris (Scopoli, 1763)
= *populi* (Meigen, 1826)
Phaonia serva (Meigen, 1826)
Phaonia steinii (Strobl, 1898)
Phaonia subfuscinervis (Zetterstedt, 1838)
Phaonia subventa (Harris, 1780)
= *variegata* (Meigen, 1826)
Phaonia taigensis Zinovjev, 1987
Phaonia tiefii Schnabl, 1888
Phaonia trimaculata (Bouché)
= *servaeformis* Ringdahl, 1916
Phaonia tuguriorum (Scopoli, 1763)
= *signata* (Meigen, 1826)
Phaonia valida (Harris, 1780)
= *viarum* (Robineau-Desvoidy, 1830)
= *erratica* auct. nec (Fallén, 1825)
Phaonia villana Robineau-Desvoidy, 1830
= *mystica* auct. nec (Meigen, 1826)
Phaonia vivida (Rondani, 1870)
= *austriaca* (Czerny, 1900)
Phaonia wahlbergi Ringdahl, 1930
Phaonia zugmayeriae (Schnabl, 1888)
?= *humeralis* (Zetterstedt, 1845)
= *humarella* (Stein, 1900)

Excluded species

- Coenosia femoralis* (Robineau-Desvoidy, 1830) misidentified
Fannia barbata (Stein, 1892) not found within present borders
Fannia lineata (Stein, 1895) not found within present borders
Fannia nigra Malloch, 1910 not found within present borders
Hydrotaea glabricula (Fallén, 1825)
Limnophora pollinifrons Stein, 1916 not found within present borders
Phaonia bitincta (Rondani, 1866) misidentified
Scathophaga calida Haliday in Curtis, 1832 not found within present borders
Spilogona acrostichalis (Stein, 1916)
Spilogona norvegica (Ringdahl, 1932)
Spilogona obscuripennis (Stein, 1916) not found within present borders
Spilogona septemnotata (Zetterstedt, 1845) misidentified
Spilogona setigera (Stein, 1907)
Spilogona spininervis (Villeneuve, 1922)

Notes

***Coenosia comita* (Huckett, 1936).** Collected only from Salla. The type locality of *C. ovulivera* Tiensuu, Vuorijärvi, is now Russian territory (Tiensuu 1938). A second specimen, also taken in 1936 by R. Krogerus (in MZH), bears only the label 'Salla' and probably also originates from the vicinity of Vuorijärvi, but it is possible that it was collected from the Finnish part of Salla, a municipality split between Russia and Finland in 1944.

***Helina ciliata* Karl, 1929.** Ringdahl (1952), quoting "Tiensuu's papers", lists this species from Finland. We could not locate the original record in Tiensuu's published works or among the available collection specimens. It seems possible Ringdahl made a mistake in his tables. The species is also listed as Finnish by Hackman (1980), again without any details.

***Limnophora pandellei* Séguy, 1923.** Only females have been found in Finland. The identification remains slightly doubtful (Kahanpää 2013).

***Norellia tipularia* (Fabricius, 1794).** This species was first recorded from Finland by Winqvist (2011). Šifner (2003, 2008) considers *N. tipularia* a nomen dubium and probably synonymous with *N. spinipes* Meigen, but Ozerov and Krivosheina (2011) disagree and provide a detailed diagnosis.

***Parallelomma paridis* Hering, 1923.** This name was synonymized with *P. vittatum* (Meigen) by Šifner (1978). Ozerov (2010, 2014) examined the types of *P. paridis* and upheld the synonymy. Not everyone agrees with this synonymy (Nelson 1990). Finnish CO1 DNA fingerprints (Kahanpää, unpublished) of larvae on *Paris* (Liliaceae) and adults caught among *Dactylorhiza* (Orchidaceae) suggest that two species are involved.

***Phaonia amicula* Villeneuve, 1922.** New to Finland. Recorded as *Phaonia* sp. nr. *halterata* by Winqvist (2011): Ab: Turku, Ruissalo (67093:32336), 3.vii.2008, leg. K. Winqvist, 1 male.

Acknowledgements

I would like to express my appreciation to Dr. Verner Michelsen for his constructive feedback and for identifying the first Finnish specimen of *Phaonia amicula*. Mr. Kaj Winqvist kindly allowed us to include his record of *P. amicula* in this paper.

References

- Hackman W (1956) The Scathophagidae (Dipt.) of Eastern Fennoscandia. Fauna fennica 2: 1–66.
Hackman W (1980) A Check List of the Finnish Diptera. Notulae entomologicae 60: 17–48, 117–162.
Kahanpää J (2013) Viisikymmentä Suomelle uutta käpälslajia (Diptera: Brachycera). Sahlbergia 19(1–2): 63–71.

- Kahanpää J, Haarto A (2013) Notes on Fanniidae (Diptera) of Finland, with a description of the female of *Fannia stigi* Rognes, 1982. *Entomologica fennica* 24: 179–185.
- Kutty SN, Pape T, Wiegmann BM, Meier R (2010) Molecular phylogeny of the Calyptratae (Diptera: Cyclorrhapha) with an emphasis on the superfamily Oestroidea and the position of Mystacinobiidae and McAlpine's fly. *Systematic Entomology* 35(4): 614–635. doi: 10.1111/j.1365-3113.2010.00536.x
- Nelson JM (1990) Observations on the biology and status of British dung-flies of the genus *Parallelolomma* Becker (Dipt., Scathophagidae). *Entomologist's Monthly Magazine* 126: 187–189.
- Nirmala X, Hypša V, Žurovec M (2001) Molecular phylogeny of Calyptratae (Diptera: Brachycera): the evolution of 18S and 16S ribosomal rDNAs in higher dipterans and their use in phylogenetic inference. *Insect Molecular Biology* 10(5): 475–485.
- Ozerov AL (2010) Flies of the genus *Parallelolomma* Becker in Strobl, 1894 (Diptera, Scathophagidae) of Russia. *Russian Entomological Journal* 18[2009](4): 309–317.
- Ozerov A (2014) The Primary Types of Scathophagidae (Diptera) in the Museum für Naturkunde Berlin, and Senckenberg Deutsches Entomologisches Institut, Germany. *Zoosystematics and Evolution* 90(1): 33–43. doi: 10.3897/zse.90.7121
- Ozerov AL, Krivosheina MG (2011) The first record of *Norellia tipularia* (Fabricius, 1794) (Diptera: Scathophagidae) for the fauna of Russia. *Russian Entomological Journal* 20(1): 101–103.
- Pape T, Blagoderov V, Mostovski MB (2011) Order Diptera Linnaeus, 1758. In: Zhang Z-Q (Ed) Animal biodiversity: An outline of higher-level classification and survey of taxonomic richness. *Zootaxa* 3148: 222–229. <http://www.mapress.com/zootaxa/2011/f/zt03148p229.pdf>
- Ringdahl O (1952) Catalogus Insectorum Sueciae. XI. Diptera Cyclorrhapha: Muscaria Schizometopa. *Opuscula entomologica* 17: 129–186.
- Šifner F (1978) La révision synonymique des espèces du genre *Americina* Malloch, 1923 (Diptera, Scatophagidae). In: Országh I (Ed) Proceedings of the IVth Meeting of Czechoslovak Dipterists. *Dipterologica bohemoslovaca* 1. Veda, Bratislava, 283–302.
- Šifner F (2003) The family Scathophagidae (Diptera) of the Czech and Slovak republics (with notes on selected Palaearctic taxa). *Acta Musei Nationalis Pragae, Series B, Historia Naturalis* 59(1–2): 1–90.
- Šifner F (2008) A catalogue of the Scathophagidae (Diptera) of the Palaearctic region, with notes on their taxonomy and faunistics. *Acta entomologica Musei Nationalis Pragae* 48(1): 111–196. http://www.aemnp.eu/pdf/48_1/48_1_111.pdf
- Tiensuu L (1935) Die bisher aus Finnland bekannten Musciden. *Acta Societatis pro Fauna et Flora Fennica* 58(4): 1–56.
- Tiensuu L (1938) Beiträge zur Kenntnis der Musciden (Dipt.) Finnländs. *Annales entomologici fennici* 4(1): 21–33.
- Tiensuu L (1939) Beiträge zur Kenntnis der Musciden (Dipt.) Finnländs. 2. *Annales entomologici fennici* 5(4): 241–255.
- Winqvist K (2011) Ruissalon kärpästä (Diptera, Brachycera). w-album 11: 3–23. http://org.utu.fi/harrastus/TEKS/w-album/w_album_11.pdf