

Two new species of the leafhopper genus *Mitjaevia* Dworakowska from China (Hemiptera, Cicadellidae, Typhlocybinae)

Xiaoxiao Chen¹, Yuehua Song¹, Mick D. Webb²

1 School of Karst Science, Guizhou Normal University / State Key Laboratory Cultivation Base for Guizhou Karst Mountain Ecology Environment of China, Guiyang, Guizhou 550001, China **2** Insect Division, Department of Life Sciences, the Natural History Museum, London, SW7 5BD, UK

Corresponding author: Yuehua Song (songyuehua@163.com)

Academic editor: Mick Webb | Received 21 November 2019 | Accepted 30 July 2020 | Published 27 August 2020

<http://zoobank.org/63EFF2A4-4B7D-46B4-B31F-C3D2C543A924>

Citation: Chen X, Song Y, Webb MD (2020) Two new species of the leafhopper genus *Mitjaevia* Dworakowska from China (Hemiptera, Cicadellidae, Typhlocybinae). ZooKeys 964: 31–40. <https://doi.org/10.3897/zookeys.964.48655>

Abstract

In the present paper, two new species of the leafhopper genus *Mitjaevia* Dworakowska, 1970 from Guizhou Province China are described and illustrated, i.e., *Mitjaevia shibingensis* sp. nov. and *Mitjaevia dworakowskiae* sp. nov. A checklist to species of the genus and a key to distinguish the Chinese species of the genus are given and the female valvulae are described and figured for the first time.

Keywords

Checklist, distribution, Homoptera, identification key, morphology, taxonomy

Introduction

The leafhopper genus *Mitjaevia* Dworakowska, 1970 belongs to the tribe Erythroneurini of Typhlocybinae, with *Erythroneura amseli* Dlabola, 1961 as its type species; seventeen species are known, seven from China (see Checklist). Two new species from Guizhou Province, China are described and illustrated in this paper together with a checklist and key to species from China.

Materials and methods

Specimens for this study were collected by sweep-net. Morphological terminology used follows Dietrich (2005) and Song and Li (2013) and observations and drawings were made using Olympus SZX16 and BX53 microscopes. Habitus photos were taken using a KEYENCE VHX-5000 digital microscope. Body measurements are from the apex of the vertex to the tip of the forewing. All specimens examined are deposited in the collection of the School of Karst Science, Guizhou Normal University, China (GZNU).

Taxonomy

Mitjaevia Dworakowska, 1970

Mitjaevia Dworakowska, 1970: 763.

Type species. *Erythroneura amseli* Dlabola, 1961, by original designation.

Diagnosis. Species with distinctive dark brown markings; head distinctly narrower than pronotum; male pygofer with simple dorsal appendage and sometimes ventral appendage; subgenital plate with a series of lateral peg-like setae basally or subbasally and a few macrosetae centrally at midlength; style elongate with subapical extension and variably developed lateral lobe; aedeagus with shaft cylindrical or laterally compressed, with or without processes, with ventral gonopore, basal apodeme variably developed and preatrium distinct.

Distribution. Palaearctic and Oriental Regions.

Remarks. Dworakowska (1970: 763–765) gave a detailed description of this genus based on the three included species known at that time; based on subsequently included species a modified description was given by Song et al. (2011: 26–27) and Dmitriev (2020). In addition, the female valvulae are described and figured here for the first time. Although a diagnosis is given above, clearly further studies are needed to elucidate fully the diagnostic characters of the genus and to test if the genus is monophyletic in the light of the observed variation in male genitalia between species.

Checklist to species of the genus *Mitjaevia*

- 1 *Mitjaevia amseli* (Dlabola, 1961: 297, figs 137–141, *Erythroneura*. Uzbekistan); Dlabola 1964: 248, Afghanistan; Dworakowska 1970: 765, figs 33–44, transferred to *Mitjaevia*. Kazakhstan; Korolevskaya, 1976: 42–43, figs 7, 8.
- 2 *Mitjaevia atropictila* (Ahmed, 1970a: 35; fig. 5: A–F, *Erythroneura*. Pakistan); Sharma 1984: 33, figs 19–29, transferred to *Mitjaevia*. India.

- 3 *Mitjaevia aurantiaca* (Mitjaev, 1969: 1045; figs 1, 2, *Erythroneura*. Kazakhstan); Dworakowska 1970: 765, transferred to *Mitjaevia*; Korolevskaya 1976: 42, figs 9, 10.
- 4 *Mitjaevia aurea* Dworakowska, 1994: 118; figs 407–414. India.
- 5 *Mitjaevia bibichanae* (Dlabola, 1961: 296, figs 131–135, *Erythroneura*. Uzbekistan); Korolevskaya 1976: 43–44, figs 11–13, transferred to *Mitjaevia*. Tadzhikistan.
- 6 *Mitjaevia callosa* Dworakowska, 1980: 179; figs 263–272. India.
- 7 *Mitjaevia diana* (Distant, 1918: 100, *Typhlocyba*. India); Dworakowska 1970: 765; 1980: 179, figs 252–262, transferred to *Mitjaevia*. India, Kazakhstan.
- 8 *Mitjaevia elegantula* Dworakowska, 1994: 119; figs 415–425. India.
- 9 *Mitjaevia korolevskayae* Dworakowska, 1979: 44; figs 349–358. Vietnam.
- 10 *Mitjaevia maculata* (Ahmed, 1970b: 175; fig. 6: A–H, *Helionidia*. Pakistan); Dworakowska and Viraktamath 1975: 529, transferred to *Mitjaevia*. India.
- 11 *Mitjaevia nanaoensis* Chiang & Knight, 1990: 223; fig. 18: 1–7. China.
- 12 *Mitjaevia narzikulovi* Korolevskaya, 1976: 43; figs 1–6. Tadzhikistan.
- 13 *Mitjaevia notata* (Ahmed & Khokhar, 1971: 70; fig. 4a–f, *Helionidia*. Pakistan); Dworakowska 1980: 179, transferred to *Mitjaevia*. India.
- 14 *Mitjaevia protuberanta* Song, Li & Xiong, 2011: 27; figs 1–10. China.
- 15 *Mitjaevia shibingensis* sp. nov. China.
- 16 *Mitjaevia sikkimensis* Dworakowska, 1994: 119; figs 426–434. India.
- 17 *Mitjaevia dworakowskiae* sp. nov. China.
- 18 *Mitjaevia tappana* Chiang & Knight, 1990: 224; fig. 19: 1–7. China.
- 19 *Mitjaevia wangwushana* Song, Li & Xiong, 2011: 29; figs 11–19. China.

Key to species of *Mitjaevia* from China (males)

- | | | |
|---|---|---|
| 1 | Aedeagus with process..... | 2 |
| — | Aedeagus without process | 3 |
| 2 | Processes arising from aedeagal shaft subbasally (Figs 42, 43) ... | <i>M. protuberanta</i> |
| — | Processes arising from aedeagal shaft subapically (Figs 44, 45) | <i>M. wangwushana</i> |
| 3 | Aedeagus with shaft cylindrical, evenly tapered from base to apex (Figs 20, 34) | 4 |
| — | Aedeagus with shaft laterally compressed, abruptly tapered subapically to apex (Figs 45, 49) | 5 |
| 4 | Style lateral lobe small (Fig. 18); aedeagal shaft tapered to narrowly rounded apex in lateral view (Fig. 20) | <i>M. shibingensis</i> sp. nov. |
| — | Style lateral lobe large (Fig. 31); aedeagal shaft tapered to acute apex in lateral view (Fig. 34)..... | <i>M. dworakowskiae</i> sp. nov. |
| 5 | Subgenital plate with few long macrosetae; aedeagus as in Figs 48, 49 | <i>M. nanaoensis</i> |
| — | Subgenital plate with several long macrosetae; aedeagus as in Figs 46, 47..... | <i>M. tappana</i> |

***Mitjaevia shibingensis* sp. nov.**

<http://zoobank.org/A8734F83-DBCD-4741-92C5-F2CF7AA083AC>

Figs 1–7, 15–27

Description. Vertex pale yellow, with pair of small black apical spots and two irregular markings at sides of coronal suture (Figs 1, 3). Face pale brownish yellow, anteclypeus with apical half dark brown; frontoclypeus with brownish black patches at sides basally (Fig. 4). Pronotum mostly dark brown, with pair of symmetrical brownish yellow oval impressed patches medially, showing brownish yellow near anterior margin (Figs 1, 3). Scutellum orange yellow, with brown irregular elliptical spot at base medially (Figs 1, 3). Forewing with orangey and gray patches (Fig. 6).

Abdominal apodemes small, not extended to hind margin of 3rd sternite (Fig. 15). Male genitalia with subgenital plate relatively short, broadened basally, provided with two long macrosetae at midlength on lateral surface and numerous peg-like setae along dorsal margin basally to near midlength; several microsetae scattered on apical portion (Fig. 17). Style elongate, with subapical extension laterally, lateral lobe moderately large (Fig. 18). Aedeagal shaft narrow tapered to narrowly rounded apex in lateral view, gonopore arising near midlength on ventral surface; basal apodeme reduced; preatrium well developed (Figs 19, 20). Connective moderately broadly Y-shaped, central lobe well developed (Fig. 21). Female 7th sternite as in Fig. 24. Valvula I elongate, curved dorsad and evenly tapered from base to apex, finely striate along dorsal margin of apical 1/5 (Fig. 25). Valvulae II elongate, slightly expanded blade-like to near apex, thereafter tapered to down-turned apex, with few dorsal roundish teeth distally on right branch (Fig. 26). Valvula III tapered distally to narrowly rounded apex (Fig. 27).

Measurement. Body length, males 2.6–2.8 mm, females 2.7–2.8 mm.

Specimen examined. *Holotype* ♂: CHINA, Guizhou Prov., Shibing, 27 V 2019, coll. Zhouwei Yuan, Chao Tan and Xiaowei Yuan. *Paratypes*: 14♂♂, 55♀♀, same data as holotype.

Remarks. This species has a similar shaped aedeagus to *M. korolevskaya* but the style has a preapical extension (“heel”) and a smaller lateral lobe.


Etymology. The new species is named after its type locality: “Shibing”, Guizhou Province.

***Mitjaevia dworakowskae* sp. nov.**


<http://zoobank.org/AE8B70FC-D3C5-4F7E-ACA6-1B23044AAB49>

Figs 8–14, 28–41


Description. Vertex light yellow, with two pairs of irregular black preapical spots distributed symmetrically (Figs 8, 10). Face milky yellow, anteclypeus with central area brownish; frontoclypeus with brownish black patches at sides basally (Fig. 11). Pronotum mostly black, with pair of symmetrical pale-yellow oval impressed patches medially, also showing pale yellow near anterior margin (Figs 8, 10). Scutellum milky



Figures 1–14. Species of *Mitjaevia* **1–7** *Mitjaevia shibingensis* sp. nov. **1** habitus, dorsal view **2** habitus, lateral view **3** head and thorax, dorsal view **4** face **5** style and connective, ventral view, aedeagus lateral view **6** forewing **7** hindwing **8–14** *Mitjaevia dworakowskiae* sp. nov. **8** habitus, dorsal view **9** habitus, lateral view **10** head and thorax, dorsal view **11** face **12** style and connective, ventral view, aedeagus lateral view **13** forewing **14** hindwing.



Figures 15–27. *Mitjaevia shibingensis* sp. nov. **15** abdominal apodemes **16** male pygofer, lateral view **17** subgenital plate, lateral view **18** style **19** aedeagus, ventral view **20** aedeagus, lateral view **21** connective **22** male pygofer dorsal appendage **23** male pygofer dorsal appendage **24** female 7th sternite **25** valvula I **26** valvulae II **27** valvula III.




Figures 28–41. *Mitjaevia dworakowskai* sp. nov. **28** abdominal apodemes **29** male pygofer lobe, lateral view **30** subgenital plate **31** style **32** style **33** aedeagus, ventral view **34** aedeagus, lateral view **35** connective **36** pygofer dorsal appendage **37** pygofer dorsal appendage **38** female 7th sternite **39** valvula I **40** valvulae II **41** valvula III.

yellow, with longitudinal black stripe between scutellar suture and apex (Figs 8, 10). Forewing with brown and brownish yellow patches (Fig. 13).

Abdominal apodemes small, not extended beyond hind margin of 3rd sternite (Fig. 28). Male genitalia with subgenital plate laterally with 3 macrosetae at midlength and three more distal shorter macrosetae, dorsal peg-like setae restricted to central part (Fig. 30). Style elongate with preapical extension on inner surface, lateral lobe large (Figs 31, 32). Aedeagal shaft narrow slightly sinuate and tapered to acute apex in lateral view with gonopore arising near midlength of ventral surface; basal apodeme reduced (Figs 33, 34). Connective broadly Y-shaped, central lobe slender (Fig. 35). Female 7th as in Fig. 38. Valvulae as in previous species (Figs 39–41).

Measurement. Body length, males 2.3–2.4 mm, females 2.4–2.5 mm.



Figures 42–49. Species of Chinese *Mitjaevia* **42, 43** *M. protuberanta* Song, Li & Xiong **42** aedeagus, ventral view **43** aedeagus, lateral view **44, 45** *M. wangwushana* Song, Li & Xiong **44** aedeagus, ventral view **45** aedeagus, lateral view **46, 47** *M. tappana* Chiang & Knight **46** aedeagus, ventral view **47** aedeagus, lateral view **48, 49** *M. nanaoensis* Chiang & Knight **48** aedeagus, ventral view **49** aedeagus, lateral view (Figs 42–45, from original; Figs 46–49, redrawn from Chiang and Knight 1990).

Specimen examined. *Holotype* ♂: CHINA, Guizhou Prov., Shibing, 27.V.2019, coll. Zhouwei Yuan, Chao Tan and Xiaowei Yuan. **Paratypes:** 14♂♂, 19♀♀, same data as holotype.

Remarks. This species can be distinguished by the narrow and slightly sinuate aedeagal shaft in lateral view and the style with a subapical extension on the inner surface with a greatly enlarged lateral lobe.

Etymology. This species is named for Dr Irina Dworakowska in recognition of her immense contribution to taxonomy of World Typhlocybinae.

Acknowledgements

We thank Chandra Viraktamath for critically reading an early draft of the ms. The study was partly supported by the Guizhou Provincial Science and Technology Foundation ([2018]1411), the Guizhou Science and Technology Support Project ([2019]2855), the Science and Technology Project of Guiyang City ([2020]7–18), the Training Program for High-level Innovative Talents of Guizhou Province ([2016]4020), the Program for Academician Workstation in Guiyang University ([2019]5605), The World Top Discipline Program of Guizhou Province: Karst Ecoenvironment Sciences (No.125 2019 Qianjiao Keyan Fa) and the Project for Regional Top Discipline Construction of Guizhou Province: Ecology in Guiyang University (Qian Jiao Keyan Fa [2017]85).

References

- Ahmed M (1970a) Studies of the genus *Erythroneura* Fitch (Erythroneurini: Cicadellidae) in West Pakistan. *Pakistan Journal of Zoology* 2(1): 29–42.
- Ahmed M (1970b) Further studies on the leafhoppers of tribe Erythroneurini (Cicadellidae: Typhlocybinae) from West Pakistan. *Pakistan Journal of Zoology* 2(2): 167–184.
- Ahmed M, Khokhar KF (1971) Genus *Helionidia* Zachvatkin (Erythroneurini-Cicadellidae) with four new species from West Pakistan. *Sind University Research Journal* 5(1): 63–72.
- Chiang CC, Knight WJ (1990) Studies on Taiwanese Typhlocybinae (Homoptera: Cicadellidae) (IV) tribe Erythroneurini. *Bulletin of the National Museum of Natural Science Number 2:* 223–225.
- Dietrich CH (2005) Keys to the families of Cicadomorpha and subfamilies and tribes of Cicadellidae (Hemiptera: Auchenorrhyncha). *Florida Entomologist* 88: 502–517. [https://doi.org/10.1653/0015-4040\(2005\)88\[502:KTTFOC\]2.0.CO;2](https://doi.org/10.1653/0015-4040(2005)88[502:KTTFOC]2.0.CO;2)
- Distant WL (1918) Rhynchota. VII. Homoptera: appendix. Heteroptera: addenda. The fauna of British India, including Ceylon and Burma. London, Taylor & Francis [VIII+] 210 pp.
- Dlabola J (1961) Zikaden von Zentralasien, Dagestan und Transkaukasien (Homopt. Auchenorrhyncha). *Acta Entomologica Musei Nationalis Pragae* 34: 241–358.

- DLabola J (1964) Die Zikaden Afghanistans (Homoptera, Auchenorrhyncha II. Teil). Ergebnisse der Sammelreisen von Dr. H.G. Amsel, G. Ebert, Dr. Erichson, J. Klapperich und Dr. K. Lindberg. Mitteilungen der Münchener Entomologischen Gesellschaft 54: 237–255.
- Dmitriev D (2020) An online interactive key and searchable database of Auchenorrhyncha (Hemiptera). <http://dmitriev.speciesfile.org/>
- Dworakowska I (1970) On the genera *Asianidia* Zachv. and *Singapora* Mahm. with the description of two new genera (Auchenorrhyncha, Cicadellidae, Typhlocybinae). Bulletin de l'Academie Polonaise des Sciences, Serie des Sciences Biologiques 18(12): 759–765.
- Dworakowska I (1979) On some Erythroneurini from Vietnam (Typhlocybinae, Cicadellidae). Annotationes Zoologicae et Botanicae 131: 44–45.
- Dworakowska I (1980) On some Typhlocybinae from India (Homoptera, Auchenorrhyncha, Cicadellidae). Entomologische Abhandlungen Staatliches Museum für Tierkunde in Dresden 43(8): 151–201.
- Dworakowska I (1994) Typhlocybinae (Auchenorrhyncha: Cicadellidae) of Sikkim, a preliminary survey. Folia Entomologica Hungarica Rovartani K zlem ények 55: 93–215.
- Dworakowska I, Viraktamath CA (1975) On some Typhlocybinae from India (Auchenorrhyncha, Cicadellidae). Bulletin de l'Academie Polonaise des Sciences. (Serie des Sciences Biologiques) 23(8): 521–530.
- Korolevskaya LL (1976) Leafhoppers of the genus *Mitjaevia* Dworakowska, I 1970 (Homoptera, Typhlocybinae) from Tadzhikistan. Izvestiya Akademii nauk Tadzhikskoi SSR (Otdelenie biologicheskikh nauk) 3(64): 41–44.
- Mitjaev ID (1969) New species of leaf-hoppers (Homoptera, Cicadinea) from Tien Shan and Karatau. Zoologicheskii zhurnal 48(7): 1041–1047.
- Sharma B (1984) On some Typhlocybinae from Kishtwar, Jammu & Kashmir (India), with description of five new species (Homoptera, Auchenorrhyncha, Cicadellidae). Reichenbachia 22(3): 31–40.
- Song YH, Li ZZ (2013) Two new species of *Empoascanara* Distant (Hemiptera: Cicadellidae: Typhlocybinae) from Yunnan Province, China. Zootaxa 3637(1): 089–093. <https://doi.org/10.11646/zootaxa.3637.1.11>
- Song YH, Li ZZ, Xiong KN (2011) Two new species of the genus *Mitjaevia* Dworakowska from China (Hemiptera: Cicadellidae: Typhlocybinae). Zootaxa 2805: 26–30. <https://doi.org/10.11646/zootaxa.2805.1.2>