

A new species and additional records of *Rugilus* Leach from Qinling, China (Coleoptera, Staphylinidae, Paederinae)

Jia-Yao Hu¹, Chen-Zu Song¹, Li-Zhen Li¹

¹ Department of Biology, College of Life and Environmental Sciences, Shanghai Normal University, 100 Guilin Road, Shanghai, 200234 P. R. China

Corresponding author: Li-Zhen Li (lizhenli@shnu.edu.cn)

Academic editor: V. Assing | Received 24 March 2015 | Accepted 20 May 2015 | Published 25 May 2015

<http://zoobank.org/2081783A-6949-4DE5-967C-11B5D301C00C>

Citation: Hu J-Y, Song C-Z, Li L-Z (2015) A new species and additional records of *Rugilus* Leach from Qinling, China (Coleoptera, Staphylinidae, Paederinae). ZooKeys 505: 147–152. doi: 10.3897/zookeys.505.9681

Abstract

A new species of *Rugilus* Leach, *R. (Rugilus) huanghaoi* sp. n. from Qinling, Shaanxi Province, China, is described and illustrated. Additional records of seven species from Qinling are reported.

Keywords

Coleoptera, Staphylinidae, Paederinae, *Rugilus*, Qinling, China, new species

Introduction

According to a series of revisions of the genus *Rugilus* Leach from Palaearctic and Oriental regions (Assing 2012a, 2012b, 2013, 2014, 2015), 32 species of the genus have been recorded from China, 25 of them are placed in the nominotypical subgenus, and seven in the subgenus *Eurystilicus* Fagel. Seven species have been reported from the Qinling Shan, one of the most diverse areas in China: *R. (Eurystilicus) rufescens* (Fauvel, 1874), *R. (E.) simlaensis* (Cameron, 1931), *R. (E.) velutinus* (Fauvel, 1895), *R. (Rugilus) dabaicus* Assing, 2012, *R. (R.) fodens* Assing, 2012, *R. (R.) gansuensis* Rougemont, 1998, and *R. (R.) reticulatus* Assing, 2012. During several recent field trips to this region conducted by the authors and their colleagues, all known and an additional new species were collected.

Material and methods

The type material listed in the present study is deposited in the Insect Collection of Shanghai Normal University, Shanghai, P. R. China (SNUC).

The dissected body parts were mounted in Euparal on plastic slides. The habitus photos were taken using a Canon 7D camera. The photos of the sternites and aedeagi were taken using a Canon G9 camera mounted on an Olympus CX31 microscope.

Measurements:

Body length: measured from the anterior margin of the labrum to the apex of the abdomen.

Length of forebody: measured from anterior margin of the labrum to the posterior margin of the elytra.

Eye length: longitudinal length of eye in dorsal view.

Postocular length: measured from posterior margin of eye to posterior constriction of head.

Head width: width of head across (and including) eyes.

Head length: measured from the clypeal anterior margin to head base.

Pronotum width: maximal width of pronotum.

Pronotum length: measured in midline from front margin to posterior margin.

Width of elytra: combined width of elytra at posterior margin.

Length of elytra: measured from apex of scutellum to posterior margin.

Description of new species

Rugilus (Rugilus) huangbaoi sp. n.

<http://zoobank.org/8E11E667-B5DC-4D85-A334-153047FE5BDC>


Figs 1–7

Type material. Holotype: male: “China: Shaanxi Prov., Zhouzhi County, Houzhenzi, Qinling, Qinlingliang, N33.48.963, E107.44.483, alt. 2018 m, 7.V.2008, HUANG Hao & XU Wang leg.” (SNUC). Paratypes: 1 male, 1 female: “China: Shaanxi Prov., Mei County, Taibai Shan, Kaitianguan, N34.00.692, E107.51.415, alt. 1853 m, 22–23.V.2008, HUANG Hao & XU Wang leg.”.

Description. Body length 5.4–6.4 mm; forebody length: 4.1–4.3 mm.

Body (Fig. 1) dark brown; lateral margins of elytra widely yellowish brown; legs and antennae reddish brown.

Head (Fig. 2) 0.96–0.97 times as long as wide; punctation umbilicate, very dense, rather coarse and partly confluent, interstices reduced to very narrow ridges; without microsculpture. Eyes large and convex; approximately 0.8–0.9 times as long as pos-



Figures 1–7. *Rugilus (Rugilus) huanghaoi* sp. n. **1** habitus **2** forebody **3** male sternite VII **4** male sternite VIII **5** aedeagus in ventral view **6** aedeagus in lateral view **7** median portion of tergite III. Scale bars: 1 mm (1, 2), 0.25 mm (3–7).

ocular portion. Anterior margin of labrum with two pronounced teeth on either side of the median incision.

Pronotum (Fig. 2) 1.16–1.17 times as long as wide, 0.75–0.77 times as broad and 0.92–0.94 times as long as head; punctuation similar to that of head; midline with narrow and short impunctate elevation in posterior half; interstices without microsculpture. Elytra (Fig. 2) 0.90–0.98 times as long as wide, 1.13–1.26 times as long and 1.47–1.52 times as broad as pronotum; punctuation dense, distinctly finer than that of head and pronotum; interstices without microsculpture.

Abdomen narrower than elytra; tergites III–VI with transverse impressions anteriorly. Punctuation of these impressions coarse and dense; punctuation of remaining surfaces fine and dense; interstices with distinct microsculpture (Fig. 7); posterior margin of tergite VII with distinct palisade fringe.

Male. Sternite VII (Fig. 3) with broad and trapezoidal excision in the middle of posterior margin; on either side of this excision with a tuft of long black setae. Sternite VIII (Fig. 4) with triangular excision posteriorly. Aedeagus (Figs 5, 6) long and narrow; ventral process widest near middle and gradually narrowed apically in ventral view; abruptly narrowed and slightly curved ventrally in apical third in lateral view.

Comparative notes. Based on the similar external characters, especially the bi-coloured elytra and the male sexual characters, the new species is most similar to *R. morvani* (Rougemont, 1987) from Nepal, from which it is distinguished by the deeper excision of the male sternite VII, with a more prominently produced centre, longer

and denser setae on either side of the excision, and by the longer and narrower apical portion of the aedeagal ventral process.

Distribution and habitat data. The species was found in two localities in the Qinling Shan. The specimens were collected by sifting decaying leaf litter in mixed forests at altitudes from ca. 1850 to 2020 m. The paratypes were collected together with *R. reticulatus*.

Etymology. The species is named in honor of Hao Huang, one of the collectors of the type material.

New records

Rugilus (Eurystilicus) rufescens (Fauvel, 1874)

Material examined. CHINA: Shaanxi: 2 males, Foping, 850–950 m, 20.VII.2004, Hu, Tang & Zhu leg.

Comment. The species is widespread in the East Palaearctic and Oriental regions (Assing 2012a).

Rugilus (Eurystilicus) simlaensis (Cameron, 1931)

Material examined. CHINA: Shaanxi: 1 female, Ankang City, Ningshaan County, Huoditang Forestry Centre, 33°26'N, 108°27'E, 1500–1700 m, 12.VII.2012, Li-Zhen Li leg.

Comment. The distribution of this species ranges from the Himalaya to Mainland China and Taiwan (Assing 2012a). The above material was collected together with *R. velutinus*.

Rugilus (Eurystilicus) velutinus (Fauvel, 1895)

Material examined. CHINA: Shaanxi: 3 males, Ankang City, Ningshaan County, Huoditang Forestry Centre, 33°26'N, 108°27'E, 1500–1700 m, 12.VII.2012, Li-Zhen Li leg.; 1 female, Hanzhong City, Nanzheng County, Yuanba Town, Liping National Forest Park, 32°50'N, 106°36'E, 1400–1600 m, 16.VII.2012, Yu-Hong Pan leg.

Comment. The species is widespread in the East Palaearctic and Oriental regions (Assing 2012a). Some of the above material was collected together with *R. simlaensis*.

Rugilus (Rugilus) dabaicus Assing, 2012

Material examined. CHINA: Shaanxi: 5 males, 16 females, Ningshaan County, Qinling, Huoditang Linchang, N33.26.060, E108.26.291, 1724 m, 24–25.V.2008, Hao

Huang & Wang Xu leg.; 19 females, same locality, 1500–1700 m, 12.VII.2012, Yan Chen, Li-Zhen Li, Wen-Rong Li, Wen-Li Ma, Yu-Hong Pan & Jie-Qiong Zhao leg.; 15 females, Foping, 1250–1400 m, 18.VII.2004, Hu, Tang & Zhu leg.

Comment. The species was originally described from Daba Shan in Hubei (Assing 2012a) and recently recorded from Qinling Shan in Shaanxi (Assing 2015).

***Rugilus (Rugilus) fodens* Assing, 2012**

Material examined. CHINA: Shaanxi: 2 males, 5 females, Hanzhong City, Nanzheng County, Yuanba Town, Liping National Forest Park, 32°50'N, 106°36'E, 1400–1600 m, 15.VII.2012, Chen, Li, Ma & Zhao leg.; 2 males, 2 females, same locality, 16.VII.2012.

Comment. This species was previously known from Daba Shan in Hubei and Mi-cang Shan in Sichuan and Shaanxi (Assing 2012a). The above material was collected together with *R. gansuensis*.

***Rugilus (Rugilus) gansuensis* Rougemont, 1998**

Material examined. CHINA: Shaanxi: 1 male, 2 females, Zhouzhi County, Houzhenzi, Qinling, West Sangongli Gou, N33.50.613, E107.48.524, 1336 m, 17–19.V.2008, Hao Huang & Wang Xu leg.; 1 male, Zhouzhi County, Houzhenzi, Qinling, N33.51.203, E107.50.183, 1260 m, 5–10.V.2008, Hao Huang & Wang Xu leg.; 1 male, 2 females, Hanzhong City, Nanzheng County, Yuanba Town, Liping National Forest Park, 32°50'N, 106°36'E, 1400–1600 m, 15.VII.2012, Chen, Li, Ma, Pan & Zhao leg.

Comment. This species is widespread in Qinling from Gansu to Shaanxi (Assing 2012a, 2013). Some of the above material was collected together with *R. fodens*.

***Rugilus (Rugilus) reticulatus* Assing, 2012**

Material examined. CHINA: Shaanxi: 2 males, 10 females, Zhouzhi County, Qinling, Daoban, N38.43.645, E107.58.147, 1900 m, 4.V.2008, Hao Huang & Wang Xu leg.; 1 male, Mei County Taibai Shan, Kaitianguan, N34.00.692, E107.51.415, 1853 m, 22–23.V.2008, Hao Huang & Wang Xu leg.; 11 females, Mt. Taibai, 1450–1750 m, 15.VII.2004, Hu & Tang leg.; 3 females, Foping, 2065 m, 21.VII.2004, Hu, Tang & Zhu leg.

Comment. The species was known from Qinling in Shaanxi and Funiu Shan in Henan (Assing 2012a). Some of the above material was collected together with *R. huanghabei*.

Acknowledgments

We thank our colleagues for their field work of collecting specimens. We are also most grateful to Volker Assing (Hannover, Germany) and two anonymous reviewers for their helpful comments on an earlier version of the manuscript. The present study is supported by the National Natural Science Foundation of China (No. 31201734), Shanghai Municipal Education Commission (13YZ062).

References

- Assing V (2012a) The *Rugilus* species of the Palaearctic and Oriental regions (Coleoptera: Staphylinidae: Paederinae). Stuttgarter Beiträge zur Naturkunde A, Neue Serie 5: 115–190.
- Assing V (2012b) A revision of Palaearctic and Oriental *Rugilus* Leach, 1819. II. Three new species from China and additional records (Coleoptera: Staphylinidae: Paederinae). Koleopterologische Rundschau 82: 137–149.
- Assing V (2013) A revision of Palaearctic and Oriental *Rugilus*. III. Five new species from the Palaearctic region and additional records (Coleoptera: Staphylinidae: Paederinae). Linzer Biologische Beiträge 45(1): 171–190.
- Assing V (2014) A revision of Palaearctic and Oriental *Rugilus*. IV. Three new species from Nepal and additional records (Coleoptera: Staphylinidae: Paederinae). Linzer Biologische Beiträge 45(1): 449–459.
- Assing V (2015) A revision of Palaearctic and Oriental *Rugilus*. V. Two new species from China and additional records (Coleoptera: Staphylinidae: Paederinae). Linzer Biologische Beiträge 47(1). (in press)