

Two new species of *Paramesosciophilodes* (Diptera, Nematocera, Mesosciophilidae) from the Middle Jurassic of China

Jiaqi Gao¹, Guifeng Shi¹, Chungkun Shih¹, Dong Ren¹

¹ College of Life Sciences, 105 Xisanhuanbeilu, Haidian District, Capital Normal University, Beijing 100048, China

Corresponding author: Dong Ren (rendong@mail.cnu.edu.cn)

Academic editor: V. Blagoderov | Received 12 August 2014 | Accepted 10 June 2015 | Published 2 July 2015

<http://zoobank.org/AAAE4E77-00F0-40AA-8362-800D3B4148CF>

Citation: Gao J, Shi G, Shih C, Ren D (2015) Two new species of *Paramesosciophilodes* (Diptera, Nematocera, Mesosciophilidae) from the Middle Jurassic of China. ZooKeys 511: 117–129. doi: 10.3897/zookeys.511.8425

Abstract

Two new species, *Paramesosciophilodes bellus* sp. n. and *Paramesosciophilodes rarissima* sp. n., from the Jiulongshan Formation at Daohugou Village, Inner Mongolia, China, are described in the extinct family Mesosciophilidae. Altogether seven genera with 21 species of mesosciophilids have been described from the Jurassic of Siberia and Kazakhstan, the Lower Cretaceous of Transbaikalia, and the Middle Jurassic of Inner Mongolia. An emended generic diagnosis of *Paramesosciophilodes* and a list of known taxa of mesosciophilids are provided.

Keywords

Paramesosciophilodes, Daohugou, Inner Mongolia, China

Introduction

Mesosciophilidae is one of the extinct dipteran families of the suborder Nematocera. Rohdendorf (1946) described a species, *Mesosciophila venosa*, which was assigned to a new subfamily, Mesosciophilinae, within the family Allactoneuridae, along with Fungivoritinae. Later he implicitly synonymized Mesosciophilinae with Fungivoritidae and excluded *Allactoneura* DeMejere, 1907 from the family (Rohdendorf 1957,

1962). Kovalev (1985) elevated Mesosciophilinae to family level; and synonymized Fungivoritidae under the First Reviser Rule. Blagoderov (1993) erected the genus *Mesosciophilopsis* with three species within the family Mesosciophilidae, and also revised the diagnosis of the Mesosciophilidae. Two important generic characters of *Mesosciophilina* Kovalev, 1985, reported from the Middle Jurassic, are cell r distinctly large, longer than 1/6 of wing length, and r-m significantly shorter than bRs, which are regarded as “obvious ancestral characters” (Kovalev 1985). On the other hand, the generic features of *Mesosciophilopsis* Blagoderov, 1993, described from the Early Cretaceous, are cell r distinctly small, shorter than 1/6 of wing length, and r-m significantly longer than bRs, which are regarded as “derived characters” (Blagoderov 1993, Zhang 2002). Zhang (2007) established a monotypic genus *Paramesosciophilodes* for his new species, *P. ningchengensis*, and described another species within the genus *Mesosciophila*. The generic diagnosis of *Paramesosciophilodes* includes cell r 0.16–0.18 times as long as wing length, bRs markedly shorter than r-m and R_{4+5} is strongly arched near its midlength. Later, Zhang (2008) assigned three new species to three genera, including *Paramesosciophilodes eximia* Zhang, 2008, and reviewed all the records of mesosciophilids. Li and Ren (2009) described two species of *Jurasciophila* from the late Middle Jurassic Jiulongshan Formation of Daohugou in southeastern Inner Mongolia, China. Species of *Jurasciophila* Li & Ren, 2009 have cell r small, shorter than 1/6 of wing length, and r-m significantly shorter than bRs, which are regarded as “transitional characters” (Li and Ren 2009). Wang et al., in 2012, assigned two species respectively to *Mesosciophila* and *Paramesosciophilodes* of Mesosciophilidae (Wang et al. 2012). Shi et al. recently described a new genus with two new species, *Similsciophila singularis* and *S. sinuate*, from the late Middle Jurassic of Jiulongshan Formation (Shi et al. 2014). To date, 7 genera and 19 species of mesosciophilids have been described from the Jurassic of Siberia and Kazakhstan, the Lower Cretaceous of Transbaikalia, and the Middle Jurassic of Inner Mongolia, which are summarized in Table 1. In addition, an emended generic diagnosis of *Paramesosciophilodes*, based on the new findings, is provided.

There have been many transfers and corrections regarding species belonging to the Mesosciophilidae. *Eoboletina gracilis* Rohdendorf, 1946 from the Upper Jurassic of Kazakhstan might belong to the family Mesosciophilidae (Blagoderov 1993). The Mongolian genus *Mesosciophilites* Kovalev, 1985 of the Lower Cretaceous should be transferred to the Mycetophilidae (Blagoderov 1993). The Australian species *Pseudalysiinia fragmenta* Jell & Duncan, 1986 should be transferred to an unnamed genus of Mesosciophilidae rather than to the extant genus of *Pseudalysiinia* Tonnoir, 1929 of Mycetophilidae (Blagoderov 1993), and we agree with this change. The genus *Sciophilites* Kovalev, 1990 from the Lower Cretaceous of Transbaikalia might belong to either the Mesosciophilidae or to the Mycetophilidae (Blagoderov 1993). *Sinosciophila meileyingziensis* Hong, 1992 from the Lower Cretaceous of Kezuo has been transferred to the Sciophilidae (Zhang 2008), but it might be a representative of Mesosciophilidae. The other three species *Liaoxifungivora simplicis* Hong, 1992, *Atalosciophila yanensis* Ren, Lu, Guo & Ji, 1995 and *Huaxiasciophilites jingxiensis* Zhang, Hong & Li, 2001 from the Lower Cretaceous of China might belong to the family Mycetophilidae,

Table 1. A list of the described fossil Mesosciophilidae.

| Genus | Species | Locality | Age |
|-----------------------------|--|--|---------------------------------|
| <i>Mesosciophila</i> | <i>Mesosciophila venosa</i> Rohdendorf, 1946 | Karatau, Chimkent Oblast, Kazakhstan | Karabastau Fm., J ₃ |
| | <i>Mesosciophila eucalla</i> Zhang, 2007 | Daohugou, Ningcheng, Inner Mongolia, China | Jiulongshan Fm., J ₂ |
| | <i>Mesosciophila abstracta</i> Zhang, 2008 | Daohugou, Ningcheng, Inner Mongolia, China | Jiulongshan Fm., J ₂ |
| | <i>Mesosciophila sigmoidea</i> Wang, Zhao & Ren, 2012 | Daohugou, Ningcheng, Inner Mongolia, China | Jiulongshan Fm., J ₂ |
| <i>Mesosciophilodes</i> | <i>Mesosciophilodes augustipennis</i> Rohdendorf, 1946 | Karatau, Chimkent Oblast, Kazakhstan | Karabastau Fm., J ₃ |
| | <i>Mesosciophilodes similis</i> Rohdendorf, 1964 | Karatau, Chimkent Oblast, Kazakhstan | Karabastau Fm., J ₃ |
| | <i>Mesosciophilodes synchrona</i> Zhang, 2008 | Daohugou, Ningcheng, Inner Mongolia, China | Jiulongshan Fm., J ₂ |
| <i>Mesosciophilina</i> | <i>Mesosciophilina bolsakovi</i> Kovalev, 1985 | Siberia, Russia | Itat Fm., J ₂ |
| | <i>Mesosciophilina irinae</i> Kovalev, 1985 | Siberia, Russia | Itat Fm., J ₂ |
| <i>Mesosciophilopsis</i> | <i>Mesosciophilopsis curtus</i> Blagoderov, 1993 | Baissa, Buryat, Yevravnenskiy, Transbaikalia | Zaza Fm., K ₁ |
| | <i>Mesosciophilopsis expletus</i> Blagoderov, 1993 | Baissa, Buryat, Yevravnenskiy, Transbaikalia | Zaza Fm., K ₁ |
| | <i>Mesosciophilopsis minor</i> Blagoderov, 1993 | Baissa, Buryat, Yevravnenskiy, Transbaikalia | Zaza Fm., K ₁ |
| <i>Paramesosciophilodes</i> | <i>Paramesosciophilodes ningchengensis</i> Zhang, 2007 | Daohugou, Ningcheng, Inner Mongolia, China | Jiulongshan Fm., J ₂ |
| | <i>Paramesosciophilodes eximia</i> Zhang, 2008 | Daohugou, Ningcheng, Inner Mongolia, China | Jiulongshan Fm., J ₂ |
| | <i>Paramesosciophilodes aequus</i> Wang, Zhao & Ren, 2012 | Daohugou, Ningcheng, Inner Mongolia, China | Jiulongshan Fm., J ₂ |
| | <i>Paramesosciophilodes bellus</i> Gao, Shi, Shih & Ren, sp. n. | Daohugou, Ningcheng, Inner Mongolia, China | Jiulongshan Fm., J ₂ |
| | <i>Paramesosciophilodes rarissima</i> Gao, Shi, Shih & Ren, sp. n. | Daohugou, Ningcheng, Inner Mongolia, China | Jiulongshan Fm., J ₂ |
| <i>Jurasciophila</i> | <i>Jurasciophila curvula</i> Li & Ren, 2009 | Daohugou, Ningcheng, Inner Mongolia, China | Jiulongshan Fm., J ₂ |
| | <i>Jurasciophila lepida</i> Li & Ren, 2009 | Daohugou, Ningcheng, Inner Mongolia, China | Jiulongshan Fm., J ₂ |
| <i>Similsciophila</i> | <i>Similsciophila singularis</i> Shi, Shih & Ren, 2014 | Daohugou, Ningcheng, Inner Mongolia, China | Jiulongshan Fm., J ₂ |
| | <i>Similsciophila sinuate</i> Shi, Shih & Ren, 2014 | Daohugou, Ningcheng, Inner Mongolia, China | Jiulongshan Fm., J ₂ |

(Notes: J₂-Middle Jurassic, J₃-Late Jurassic, K₁-Early Cretaceous)

rather than to its previous assignment to the family of Pleciofungivoridae or the family Mesosciophilidae (Zhang 2007).

Here, based on a combination of unique wing venational characters of two recently collected specimens, we describe *Paramesosciophilodes bellus* sp. n. and *Parame-*

sosciophilodes rarissima sp. n. These specimens with bodies and complete wings were collected from the late Middle Jurassic Jiulongshan Formation of Daohugou Village in the Ningcheng County, Chifeng City, southeastern Inner Mongolia, China. Many well-preserved fossil insects have been described from this locality recently (Ren et al. 2010, 2012), such as dipterans, neuropterans, orthopterans, heteropterans, etc. (Zhang et al. 2008, 2011; Wang et al. 2010; Gu et al. 2012; Yao et al. 2012).

Materials and methods

This study is based on two specimens housed in the Key Lab of Insect Evolution & Environmental Changes, Capital Normal University, Beijing, China (Curator: Dong Ren). The specimens were examined under a LEICA MZ12.5 dissecting microscope. The photos of fossils were taken with a Nikon SMZ1000 stereo microscope. Line drawings were prepared with the aid of CorelDraw 12 graphic software. The method of calculating the ratio of cell r length vs. wing length is as follows: the length of cell r is the length along R_1 , while the length of wing is the length from wing base to wing apex. Wing venation nomenclature follows that of Wootton and Ennos (1989) and Shcherbakov et al. (1995): bRs or dRs = section of R_{4+5} basal or distal to r-m, respectively; bM_{1+2} or dM_{1+2} = section of M_{1+2} basal or distal to r-m, respectively.

Systematic paleontology

Order Diptera Linnaeus, 1758

Suborder Nematocera Latreille, 1825

Family Mesosciophilidae Rohdendorf, 1946

Genus *Paramesosciophilodes* Zhang, 2007

Type species. *Paramesosciophilodes ningchengensis* Zhang, 2007.

Included species. Type species; *Paramesosciophilodes eximia* Zhang, 2008; *Paramesosciophilodes aequus* Wang, Zhao & Ren, 2012; *Paramesosciophilodes bellus* Gao, Shi, Shih & Ren sp. n., *Paramesosciophilodes rarissima* Gao, Shi, Shih & Ren, sp. n.

Emended diagnosis. Medium (sized mesosciophilid gnats. Body (including legs) covered with long, dense pubescence. Mesonotum convex. Scutellum sharp, clearly projecting. Wing, Sc_1 elongate, slightly shorter than one-half of wing length (0.43–0.47 times as long as wing length); Sc_2 situated distinctly basad to Rs origin, arising near midway between h to Sc_1 ending; bRs shorter than r-m; R_1 slightly curved; both R_1 and R_{4+5} divergent terminally; Rs furcated distad or at level of fork of M_{1+2} ; R_{2+3} oblique and curved; R_{4+5} arched near its midlength; cell r 0.16–0.19 times as long as wing length; stem of M not developed; M_{1+2} furcated slightly distad, or basad, to level of Sc_1 ending. Tibiae and tarsi with sparse, short setae.

***Paramesosciophilodes bellus* Gao, Shi, Shih & Ren, sp. n.**

<http://zoobank.org/8FEE85B5-4556-40CA-8B60-B8309F5B1504>

Figs 1, 2

Etymology. The specific name is from the Latin of *bellus*, meaning beautiful and delicate, for the well-preserved and beautiful specimen.

Material. Holotype No. CNU-DIP-NN2013631 p/c, part and counterpart. A well-preserved insect with complete body and two wings but poorly preserved halter, without head, in dorsoventral aspect.

Locality and horizon. Daohugou Village, Shantou Township, Ningcheng County, Inner Mongolia, China, Jiulongshan Formation, late Middle Jurassic.

Diagnosis. The Sc_1 ending proximad of the midlength of cell r; bRs 0.7 times of the length of r-m; R_{4+5} strongly curved; M_{1+2} forking basad of forking of Rs, and distad of the level of Sc_1 ending; CuA strongly arched, reaching the posterior margin of the wing markedly basad of Rs forking to R_{2+3} and R_{4+5} .

Description of holotype. Medium-sized mesosciophilid with dark body, adult male, in dorsal aspects (Figs 1 and 2A). Wings out-spread, length 5.4 mm, width 2.0 mm. Body length 7.2 mm. Head and antennae not preserved. Thorax convex, length 2.0 mm, width 1.3 mm. Scutellum clearly projecting. Abdomen thin, subcylindrical, length 5.2 mm, width 1.7 mm, approx. 2.6 times as long as head and thorax combined, with eight abdominal segments, first four segments gradually widened distally, last four segments gradually narrowed terminally. Partially preserved male genitalia relatively small, distinctly narrower than eighth abdominal segment. Halteres poorly preserved. Legs relatively thin and long, femora clearly thicker in the middle; femora, tibiae and tarsi with two rows of sparse and short setae. Hind leg length 6.3 mm (femur 1.7 mm, tibia 2.4 mm, tarsus 2.2 mm).

Wings membranous, oblong, darker in color in costal area, moderately wide (length 2.7 times of width), and not reaching the apex of abdomen at rest (Fig. 2). C strong, ending beyond wing apex, at which R_{4+5} ending. Sc_1 relatively long, approx. 0.4 times





Figure 1. *Paramesosciophilodes bellus* sp. n., holotype, Photographs of habitus (dorsoventral aspect): **A** part No. CNU-DIP-NN2013631 p **B** counterpart CNU-DIP-NN2013631 c.

A



B




Figure 2. *Paramesosciophilodes bellus* sp. n., Line drawings of holotype: **A** part **B** wing venation.

the length of wing, ending far distad of the intersection of bRs and r-m. Humeral vein distinct and oblique. Sc₂ well developed, starting in front of bRs. Cell r distinctly small (0.89 mm), approx. 0.165 times the wing length (5.4 mm). Section of R from Sc₂ to bRs origin approx. 2.4 times as long as bRs. R forking into three branches: R₁, R₂₊₃ and R₄₊₅. R₁ and R₄₊₅ somewhat divergent terminally; R₂₊₃ and R₄₊₅ arched. Forking of Rs

distad of the level of M forking. Rs strong, arising from beyond the basal one-third of length of wing, bRs+dRs nearly 0.4 times the R_{4+5} . Rs forking to R_{2+3} and R_{4+5} distad of forking of M_{1+2} . Section bRs 0.7 times the r-m. R_1 slightly curved, relatively long (nearly 0.5 times the length of wing), slightly deflected after junction with R_{2+3} ; R_{2+3} slightly curved, shifted toward wing base, beyond the level of M_{1+2} forking. Vein R_{4+5} strongly arched near its midway, almost parallel with R_1 , but slightly oblique at apex. Stem of M completely reduced basad of crossvein m-cu, with only a short segment distad of m-cu. Stem of M forking into M_{1+2} and M_{3+4} . M_{1+2} forking into M_1 and M_2 near R_{2+3} level. M_1 arched anteriorly, M_2 nearly straight. Crossvein r-m short, curved, slightly oblique, shorter than bRs, nearly perpendicular to M_{1+2} , almost parallel to R_{2+3} , intersected at M_{1+2} , forking to bM_{1+2} and dM_{1+2} . bM_{1+2} approx. 6.6 times as long as m-cu. dM_{1+2} approx. as long as bM_{1+2} , and longer than r-m. CuA running parallel close to M_{3+4} basally. CuA reaching the posterior margin of wing at approx. the same level of M_{1+2} forking to M_1 and M_2 . CuP short, slightly curved at its midway, not reaching the posterior margin of wing.

Remarks. *Paramesosciophilodes bellus* sp. n. resembles most closely *P. ningchengensis*, but can be distinguished from the latter in having Sc_1 ending at C proximad of the midlength of cell r (vs. at the midlength of cell r for *P. ningchengensis*) and CuA reaching the posterior margin of the wing markedly basad of Rs forking to R_{2+3} and R_{4+5} (vs. slightly basad of Rs forking to R_{2+3} and R_{4+5}).

This new species is differentiated from *P. ningchengensis*, *P. eximia*, *P. aequus*, and *P. rarissima* sp. n. based on a combination of characters listed in Table 2.

***Paramesosciophilodes rarissima* Gao, Shi, Shih & Ren, sp. n.**

<http://zoobank.org/2DC54917-79F7-4919-8B03-3BDA9BAF3B00>

Figs 3, 4

Etymology. The specific name is from the Latin word of *rarissimus*, meaning rare.

Material. Holotype No. CNU-DIP-NN2013145 p/c, part and counterpart. A well-preserved insect with complete body with two wings, without head and halteres, in dorsoventral aspect.

Locality and horizon. Daohugou Village, Shantou Township, Ningcheng County, Inner Mongolia, China, Jiulongshan Formation, late Middle Jurassic.

Diagnosis. Sc_1 ending near the midlength of cell r; bRs 0.8 times the r-m; R_{4+5} strongly curved; M_{1+2} forking basad of R_{2+3} level and distad of level of Sc_1 ending at C; CuA strongly arched, reaching the posterior margin of the wing at the level of intersection of Rs forking to R_{2+3} and R_{4+5} .

Description of holotype. Medium-sized mesosciophilid gnats, in dorsal aspect (Figs 3 and 4A). Body length (without head and part of thorax) 7.2 mm as preserved. Legs covered with long, dense pubescence. Head, antennae, and halteres not preserved. Thorax length 1.8 mm, width 1.5 mm. Mesonotum convex. Scutellum sharp, clearly projecting. Wings membranous, oblong, length 5.0 mm, width 2.2

Table 2. Comparison of seven key characters of five species of *Paramesocitophylodes*.

| | <i>P. ningchengensis</i> | <i>P. eximia</i> | <i>P. aequus</i> | <i>P. bellus</i> sp. n. | <i>P. rarissima</i> sp. n. |
|--|--|---|--|--|---|
| Length ratio of cell r and the wing | 0.167 (left wing) 0.180 (right wing) | 0.183 (left wing) 0.172 (right wing) | 0.22 as described. But, the missing wing base was not included in wing length measurement. | 0.165 | 0.184 |
| Length of Sc ₁ | 46–47% of the wing length | 46% of the wing length | 24% of the wing length as described. But, the missing wing base was not included in wing length measurement. | 46–47% of the wing length | 43% of the wing length |
| Sc ₁ ending at C | at the midlength of cell r | distad of midlength of cell r | proximad of midlength of cell r | proximad of the midlength of cell r | near the midlength of cell r |
| bRs vs r-m | 0.6–0.7 times of length of r-m | 0.5 times of the length of r-m | 0.9 times of the length r-m | 0.7 times of the length of r-m | 0.8 times of the length of r-m |
| R ₄₊₅ | slightly curved | slightly curved | strongly curved | strongly curved | strongly curved |
| The position of base of M ₁₊₂ forking vs the forking of Rs | M ₁₊₂ forking distinctly basad of forking of Rs | M ₁₊₂ forking almost at level of forking of Rs | M ₁₊₂ forking basad of the forking of Rs | M ₁₊₂ forking basad of forking of Rs, | M ₁₊₂ forking basad of forking of Rs |
| The position of base of M ₁₊₂ forking vs the level of Sc ₁ ending | M ₁₊₂ forking basad or distad of the level of Sc ₁ ending | M ₁₊₂ forking at the level of Sc ₁ ending | M ₁₊₂ forking distad of the level of the Sc ₁ ending | M ₁₊₂ forking slightly distad of the level of Sc ₁ ending | M ₁₊₂ forking slightly distad of the level of Sc ₁ ending |
| CuA shape | CuA strongly arched | CuA smoothly arched | CuA smoothly arched | CuA strongly arched | CuA smoothly arched |
| CuA ending at the posterior margin vs. Rs forking to R ₂₊₃ and R ₄₊₅ | CuA ending slightly basad of Rs forking to R ₂₊₃ and R ₄₊₅ | CuA ending slightly distad of Rs forking to R ₂₊₃ and R ₄₊₅ | CuA ending slightly distad of Rs forking to R ₂₊₃ and R ₄₊₅ | CuA ending markedly basad of Rs forking to R ₂₊₃ and R ₄₊₅ | CuA ending slightly distad of Rs forking to R ₂₊₃ and R ₄₊₅ |




Figure 3. *Paramesosciophilodes rarissima* sp. n., holotype, Photographs of habitus (dorsoventral aspect): **A** No. CNU-DIP-NN2013145 p **B** No. CNU-DIP-NN2013145 c.

mm, darker in color in costal area, length 2.3 times width, and not reaching the apex of abdomen at rest. Abdomen thin, subcylindrical, length 5.4 mm, width 1.5 mm, with first five segments gradually widened distally, other segments gradually narrowed terminally. Legs poorly preserved, femora thicker in the middle, covered with numerous setae.

C strong, ending beyond wing apex, at which R_{4+5} ending (Fig. 4). Sc converging with C before the level of R_{4+5} . Sc_1 elongate, slightly shorter than one-half of wing length (0.43–0.47 times the wing length), and ending far distad of the intersection of bRs and r-m. Vein h distinct and oblique. Sc_2 developed well, starting in front of Rs, situated distinctly basal to Rs origin, arising beyond midway between h to Sc_1 ending. Cell r relatively large (0.92 mm), approx. 0.18 times the wing length (5.0 mm). The section of R from Sc_2 to Rs origin approx. 0.7 times the section bRs. R forking to R_1 and Rs, then Rs to R_{2+3} and R_{4+5} . Both R_1 and R_{4+5} somewhat divergent terminally; R_{2+3} and R_{4+5} arched. Rs usually strong, arising from basal one-half of length of wing, forking to R_{2+3} and R_{4+5} beyond the forking of M_{1+2} . Section bRs 0.8 times the r-m. R_1 slightly curved, relatively long, nearly 0.5 times the wing. Both R_1 and R_{4+5} divergent terminally. R_{2+3} curved, beyond the level of M_1 and M_2 forking. R_{4+5} strongly arched near its midlength. Stem of M, basad to crossvein m-cu completely reduced, with only a short segment distal to m-cu. Stem of M forking into M_{1+2} and M_{3+4} . M_{1+2} forking into M_1 and M_2 basad of R_{2+3} level and distad to level of Sc_1 ending at C. M_1 arched cephalad. M_2 nearly straight. Crossvein r-m short, curved, slightly oblique, shorter than bRs, r-m intersecting M_{1+2} and dividing M_{1+2} into bM_{1+2} and dM_{1+2} . Section bM_{1+2} approx. 4.3 times the crossvein m-cu. Section dM_{1+2} approx. 1.2 times the section bM_{1+2} , and longer than r-m. CuA running parallel and close to M_{3+4} basally. CuP short, slightly curved midway, reaching the posterior margin of wing at the same level as Sc_1 ending at C.

Remarks. *Paramesosciophilodes rarissima* sp. n. is distinguished from all other species of *Paramesosciophilodes* based on a combination of characters listed in Table 2.




Figure 4. *Paramesosciophilodes rarissima* sp. n., Line drawings of holotype: **A** part **B** wing venation.

Discussion

As shown in Table 1, a total of 7 genera and 21 species of mesosciophilids have been reported from various localities in the Jurassic of Siberia and Kazakhstan, Lower Cretaceous of Transbaikalia, and Middle Jurassic of Inner Mongolia. One genus with 2

species was described in the Middle Jurassic Itat Formation, Siberia; 6 genera with 13 species were reported from the Middle Jurassic Jiulongshan Formation of Daohugou, Inner Mongolia, China; 2 genera with 3 species were described from the Late Jurassic Karabastau Formation in Kazakhstan; and one genus with 3 species was documented from the Early Cretaceous Zaza Formation at Baissa, Transbaikalia.

The data show that the known earliest mesosciophilids have been reported from the Middle Jurassic, while the latest ones are described from the Early Cretaceous. It seems that mesosciophilids became less diverse in the Early Cretaceous, and were possibly replaced by Mycetophilidae (Blagoderov 1993), which is supported by Zhang's data, who listed all the mesosciophilids and mycetophilids from Daohugou, and compared them with other faunas (Zhang 2002).

Acknowledgements

We sincerely appreciate comments and suggestion from Dr. Blagoderov and two anonymous reviewers in improving this manuscript. We are grateful to Dr. Taiping Gao in the Key Lab of Capital Normal University for his valuable comments on the manuscript. This research is supported by National Basic Research Program of China (973 Program) (2012CB821906), the National Natural Science Foundation of China (No. 31230065, 41272006), Project of Great Wall Scholar of Beijing Municipal Commission of Education, Program for Changjiang Scholars and Innovative Research Team in University (IRT13081).

References

- Blagoderov VA (1993) Dipterans (Mesosciophilidae) from the Lower Cretaceous of Transbaikalia. *Paleontological Journal* 27(1A): 123–130.
- Gu JJ, Montealegre ZF, Robert D, Engel MS, Qiao GX, Ren D (2012) Wing stridulation in a Jurassic katydid (Insecta, Orthoptera) produced low-pitched musical calls to attract females. *Proceedings of the National Academy of Sciences* 109(10): 3868–3873. doi: 10.1073/pnas.1118372109
- Hong YC (1992) The study of Early Cretaceous Coleoptera, Raphidioptera, Diptera (Insecta) of Kezuo, West Liaoning Province. *Acta Geologica Gansu* 1(1): 1–13. [In Chinese with English Summary]
- Jell PA, Duncan PM (1986) Invertebrates, mainly insects, from the freshwater, Lower Cretaceous, Koonwarra Fossil Bed (Korumburra Group), South Gippsland, Victoria. *Memoirs of the Association for Australasian Palaeontologists* 3: 111–205.
- Kovalev VG (1985) Infraorder Bibionomorpha. *Dipterous Insects of Jurassic Siberia*. Nauka, Moscow, 113–189. [In Russian]
- Kovalev VG (1990) Flies. Muscida. In: Rasnitsyn AP (Ed.) *Late Mesozoic Insects of Eastern Transbaikalia*. *Trudy Paleont Inst. Akad. Nauk SSSR, Russia*, 123–177. [In Russian]

- Latreille PA (1825) Familles naturelles du règne animal, exposées succinctement et dans un ordre analytique, avec l'indication de leurs genres. J.-B. Baillière, Paris.
- Li TT, Ren D (2009) A new fossil genus of Mesosciophilidae (Diptera, Nematocera) with two new species from the Middle Jurassic of Inner Mongolia, China. *Progress in Natural Science* 19(12): 1837–1841. doi: 10.1016/j.pnsc.2009.06.011
- Linnaeus C (1758) *Systema naturae per regna tria naturae, secundum classes, ordines, genera, species, cum caracteribus, differentiis, synonymis, locis*. 10th ed. Salvii, Holmiae [= Stockholm]
- Ren D, Lu LW, Guo ZG, Ji S (1995) *Faunae and Stratigraphy of Jurassic–Cretaceous in Beijing and the adjacent areas*. Seismic Publishing House, Beijing, 222 pp. [In Chinese and English]
- Ren D, Shih CK, Gao TP, Yao YZ, Zhao YY (2010) *Silent Story: Insect Fossil Treasures from Dinosaur Era of the Northeastern China*. Science Press, Beijing, 332 pp.
- Ren D, Shih CK, Gao TP, Yao YZ, Zhao YY (2012) *Insect Fossil Treasures from the Mesozoic of the Northeastern China*. Science Press, Beijing, 409 pp. [In Chinese]
- Rohdendorf BB (1946) The evolution of the wing and phylogeny of Oligoneura (Diptera, Nematocera). *Trudy Paleontologicheskogo Instituta, Akademii Nauk SSSR* 13(2): 1–108. [In Russian]
- Rohdendorf BB (1957) Paleontological investigations in the USSR. *Trudy Paleontologicheskogo Instituta* 66: 1–102. [In Russian]
- Rohdendorf BB (1962) *Fundamentals of palaeontology. Tracheate and Chelicerate Arthropods*. Shcherbakov DE, Lukashevich ED, Blagoderov VA (1995) Triassic Diptera and initial radiation of the order. *An International Journal of Dipterological Research* 6(2): 75–115.
- Shi GF, Shih CK, Ren D (2014) A new genus with two new species of mesosciophilids from the Middle Jurassic of China (Diptera: Nematocera: Mesosciophilidae). *Journal of Natural History* 49(19–20): 1147–1158. doi: 10.1080/00222933.2014.951085
- Tonnoir AL (1929) Australian Mycetophilidae. Synopsis of the genera. *Proceedings Linnean Society New South Wales* 54(5): 584–614.
- Wang Q, Zhao YY, Ren D (2012) Two new species of Mesosciophilidae (Insecta: Diptera: Nematocera) from the Yanliao biota of Inner Mongolia, China. *Alcheringa: An Australasian Journal of Palaeontology* 36(4): 509–514. doi: 10.1080/03115518.2012.685434
- Wang YJ, Liu ZQ, Wang X, Shih CK, Zhao YY, Engel MS, Ren D (2010) Ancient pinnate leaf mimicry among lacewings. *Proceedings of the National Academy of Sciences* 107(37): 16212–16215. doi: 10.1073/pnas.1006460107
- Wootton RJ, Ennos AR (1989) The implications of function on the origin and homologies of the dipterous wing. *Systematic Entomology* 14(4): 507–520. doi: 10.1111/j.1365-3113.1989.tb00300.x
- Yao YZ, Ren D, Rider DA, Cai WZ (2012) Phylogeny of the Infraorder Pentatomomorpha Based on Fossil and Extant Morphology, with Description of a New Fossil Family from China. *PLoS ONE* 7(5): e37289. doi: 10.1371/journal.pone.0037289
- Zhang JF (2002) Discovery of Daohugou Biota (pre-Jehol Biota) with a discussion on its geological age. *Journal of Stratigraphy* 26(3): 173–177. [In Chinese with English abstract]

- Zhang JF (2007) New mesosciophilid gnats (Insecta: Diptera: Mesosciophilidae) in the Daohugou biota of Inner Mongolia, China. *Cretaceous Research* 28(2): 297–301. doi: 10.1016/j.cretres.2006.05.007
- Zhang JF (2008) Three new species of Mesosciophilid Gnats from the Middle-Late Jurassic of China (Insecta: Diptera: Nematocera: Mesosciophilidae). *Pakistan Journal of Biological Sciences* 11(22): 2567–2572. doi: 10.3923/pjbs.2008.2567.2572
- Zhang KY, Yang D, Ren D, Ge FC (2008) New Middle Jurassic tangle-veined flies from Inner Mongolia, China. *Acta Palaeontologica Polonica* 53(1): 161–164. doi: 10.4202/app.2008.0112
- Zhang KY, Yang D, Ren D (2011) A new brachyceran family Origoasilidae fam. n. from the Late Mesozoic of China (Insecta: Diptera). *Acta Geologica Sinica (English Edition)* 85(5): 994–997. doi: 10.1111/j.1755-6724.2011.00533.x
- Zhang ZJ, Hong YC, Li ZY (2001) Description of a new fossil genus and species *Huaxiasciophilites jingxiensis* (Diptera: Mesosciophilidae) from Early Cretaceous Jingxi Basin of Beijing, China. *Entomologia Sinica* 8(3): 193–198.