

The bees of the family Halictidae (Hymenoptera) described by Ferdinand Morawitz from the collection of Aleksey Fedtschenko

Yulia V. Astafurova¹, Maxim Yu. Proshchalykin²

1 Zoological Institute, Russian Academy of Sciences, Universitetskaya Nab., 1, Saint Petersburg 199034, Russia

2 Federal Scientific Centre for East Asian Terrestrial Biodiversity, Far Eastern Branch of Russian Academy of Sciences, Vladivostok 690022, Russia

Corresponding author: Maxim Yu. Proshchalykin (proshchalykin@biosoil.ru)

Academic editor: M.S. Engel | Received 7 September 2020 | Accepted 2 November 2020 | Published 17 November 2020

<http://zoobank.org/532128F5-8058-479E-82FE-6C4B7C7A73FC>

Citation: Astafurova YV, Proshchalykin MYu (2020) The bees of the family Halictidae (Hymenoptera) described by Ferdinand Morawitz from the collection of Aleksey Fedtschenko. ZooKeys 994: 35–104. <https://doi.org/10.3897/zookeys.994.58441>

Abstract

The type specimens of the family Halictidae, described by Ferdinand Morawitz from the collection of Aleksey Fedtschenko deposited in the Zoological Museum of the Moscow State University and in the Zoological Institute, Russian Academy of Sciences, St. Petersburg (Russia), are critically reviewed. Precise information with illustrations of types for 43 taxa is provided. Lectotypes are here designated for the following seven nominal taxa: *Halictus aprilinus* Morawitz, 1876, *H. cingulatus* Morawitz, 1876, *H. laevinodis* Morawitz, 1876, *H. limbellus* Morawitz, 1876, *H. nasica* Morawitz, 1876, *H. rhynchites* Morawitz, 1876 and *H. vulgaris* Morawitz, 1876.

Keywords

Anthophila, Apiformes, lectotypes, Palaearctic Region, taxonomy

Introduction

More than 140 years ago (1876), the second part of Ferdinand Morawitz's critical study on the bees collected by the Aleksey Fedtschenko 1869–1871 Expeditions in "Turkestan" was published. In the prior volume, "Apidae genuinae" (1875), Morawitz treated

a total of 255 species of numerous genera, of which many species were described as new. In this second part, “Andrenidae” (1876), the remaining bees were dealt with, including the species of the difficult genera *Andrena*, *Halictus* and *Hylaeus*, totalling 183 species (Pesenko and Astafurova 2003). The species treatments are of a high professional standard, the localities are precisely documented (A. Fedtschenko 1871, O. Fedtschenko 1874, Baker 2004, Kuhlmann 2005, Dathe and Proshchalykin 2017) and the type series have been carefully conserved over a long period, generally in the collections of the Zoological Museum of the Moscow State University, Moscow, Russia (ZMMU) and in the Zoological Institute of the Russian Academy of Sciences, St. Petersburg, Russia (ZISP). To this day, these remain some of the most important manuscripts on bees of this region.

The entomological literature uses various, often obscure, terms and names for Central Asian regions and countries. The term “Turkestan” has a particularly special use in entomology, widely adopted by Morawitz (1875, 1876), Dalla Torre (1896), Meade-Waldo (1923) and in other fundamental papers, regardless of its imprecise assignment to countries (Proshchalykin and Dathe 2018).

The territory of Central Asia, described as the “Western Regions” (Xi Yui) in Chinese sources, was referred to in the Russian and European historiography of the 18th and early 19th Centuries as Lesser Bukharia, as opposed to Greater Bukharia, where the Bukhara Khanate was situated. In Europe, these lands came to be referred to in the 18th and 19th Centuries as Turkestan, i.e. “the Land of Turks,” which was the original Iranian name for the territory east of Fergana and Bukhara where nomadic Turkic tribes roamed. Subsequently, when the Turkic tribes occupied the enormous territory from the Caspian Sea to Lop Nor, the name Turkestan acquired a new meaning, so broad that it was deemed necessary to distinguish such areas as Western—Bukhara, or Russian Turkestan – and Eastern or Chinese Turkestan (Murzayev 1957). According to current views, Fedtschenko’s “Turkestan” comprises the countries Uzbekistan, Tajikistan, Kyrgyzstan and southern parts of Kazakhstan.

The family Halictidae is represented in Morawitz’s 1876 publication by four genera (*Nomioides* Schenck, 1866, *Halictus* Latreille, 1804, *Sphecodes* Latreille, 1804 and *Nomia* Latreille, 1804), comprising 73 species (Nos. 327–399). Only 30 species were previously known, while the remaining 43 were newly described (*Nomioides* – 2 species; *Halictus* – 36, *Sphecodes* – 2, *Nomia* – 2) (Table 1). Few of these taxa were mentioned in subsequent publications, remaining enigmatic for decades. Since the 1930s, V. Popov [ZISP] based his studies on the taxonomy and ecology of the Central Asian bee fauna using these collections. Popov involved other mellitologists in his research, so that specimens also reached specialists in other European museums. For example, P. Blüthgen took part in such work on *Halictus* and published several taxonomic papers, based on the Fedtschenko collection (Blüthgen 1931a, 1934a, 1955, Popov 1935).

The second attempt to recognise the true type material of Fedtschenko’s Halictidae was made by K. Warncke, a teacher in Dachau (Germany), who visited ZMMU from 26.03.1975 to 01.04.1975 (Dathe and Proshchalykin 2017). He worked his way

Table 1. The nominal taxa of Fedtschenko's Halictidae, described by F. Morawitz, and their current status.

N	Species name	Sex	Current status	Depositories of types
1.	<i>Halictus albifarsis</i>	♀	Homonym	LT (ZMMU); PLT (ZISP/ZMMU)
2.	<i>Halictus annulipes</i>	♀	Valid	LT (ZMMU); PLT (ZISP/ZMMU)
3.	<i>Halictus aprilinus</i>	♀	Valid	LT, PLT (ZMMU)
4.	<i>Halictus atomarius</i>	♀	Valid	LT, PLT (ZISP)
5.	<i>Halictus cariniventris</i>	♂	Valid	LT, PLT (ZMMU)
6.	<i>Halictus cingulatus</i>	♀	Valid	LT (ZMMU); PLT (ZISP/ZMMU)
7.	<i>Halictus croceipes</i>	♀, ♂	Valid	LT (ZMMU); PLT (ZISP/ZMMU)
8.	<i>Halictus desertorum</i>	♀	Valid	LT (ZMMU)
9.	<i>Halictus determinatus</i>	♀	Homonym	LT (ZMMU); PLT (ZISP/ZMMU)
10.	<i>Halictus equestris</i>	♀	Valid	LT (ZMMU)
11.	<i>Halictus ferganicus</i>	♂	Synonym	LT (ZMMU); PLT (ZISP)
12.	<i>Halictus fuscous</i>	♂	Synonym	HT (ZMMU)
13.	<i>Halictus fulvitarsis</i>	♂	Valid	HT (ZMMU)
14.	<i>Halictus funerarius</i>	♀	Valid	LT (ZISP); PLT (ZISP/ZMMU)
15.	<i>Halictus fuscicollis</i>	♀	Valid	LT (ZMMU); PLT (ZISP)
16.	<i>Halictus hyalinipennis</i>	♀, ♂	Valid	LT (ZISP); PLT (ZISP/ZMMU)
17.	<i>Halictus laevinodis</i>	♀	Valid	LT (ZISP); PLT (ZMMU)
18.	<i>Halictus limbellus</i>	♀	Valid	LT (ZMMU); PLT (ZISP/ZMMU)
19.	<i>Halictus longirostris</i>	♀, ♂	Valid	LT (ZMMU); PLT (ZISP/ZMMU)
20.	<i>Halictus maculipes</i>	♀	Valid	LT (ZMMU)
21.	<i>Halictus melanarius</i>	♂	Valid	HT (ZMMU)
22.	<i>Halictus minor</i>	♀	Valid	LT (ZISP); PLT (ZISP/ZMMU)
23.	<i>Halictus modernus</i>	♀	Valid	HT (ZMMU)
24.	<i>Halictus nasica</i>	♀, ♂	Valid	LT (ZMMU); PLT (ZISP/ZMMU)
25.	<i>Halictus nigrilabris</i>	♂	Valid	LT (ZMMU); PLT (ZISP)
26.	<i>Halictus nigripes</i>	♂	Homonym	LT (ZMMU); PLT (ZISP/ZMMU)
27.	<i>Halictus obscuratus</i>	♀	Valid	LT (ZMMU); PLT (ZISP/ZMMU)
28.	<i>Halictus palustris</i>	♀	Valid	LT (ZMMU); PLT (ZISP)
29.	<i>Halictus pectoralis</i>	♀	Homonym	HT (ZMMU)
30.	<i>Halictus picipes</i>	♀	Valid	LT (ZMMU); PLT (ZISP/ZMMU)
31.	<i>Halictus rhynchites</i>	♀, ♂	Valid	LT (ZMMU); PLT (ZISP/ZMMU)
32.	<i>Halictus scutellaris</i>	♀	Valid	LT (ZMMU); PLT (ZISP/ZMMU)
33.	<i>Halictus sogdianus</i>	♀	Synonym	LT, PLT (ZMMU)
34.	<i>Halictus trifasciatus</i>	♀	Homonym	LT (ZMMU)
35.	<i>Halictus varipes</i>	♀, ♂	Synonym	LT, PLT (ZMMU)
36.	<i>Halictus vulgaris</i>	♀	Synonym	LT (ZMMU); PLT (ZISP/ZMMU)
37.	<i>Sphecodes nigripennis</i>	♀	Synonym	LT, PLT (ZMMU)
38.	<i>Sphecodes pectoralis</i>	♀	Valid	LT, PLT (ZMMU)
39.	<i>Sphecodes rufithorax</i>	♀, ♂	Synonym	LT (ZMMU); PLT (ZISP/ZMMU)
40.	<i>Nomia edentata</i>	♀, ♂	Valid	LT (ZMMU); PLT (ZISP/ZMMU)
41.	<i>Nomia rufescens</i>	♀	Valid	LT (ZMMU)
42.	<i>Nomiooides parviceps</i>	♂	Valid	NT (ZISP)
43.	<i>Nomiooides turanica</i>	♀, ♂	Valid	LT, PLT (ZISP/ZMMU)

Comment. LT – lectotype, PLT – paralectotype/s, NT – neotype.

through the drawers containing the Fedtschenko material and labelled specimens of all nominal taxa found there as "Lectotypen." Thereby, certain specimens from some groups of bees (*Hylaeus*, *Andrena*, *Anthidium* etc.) received a red label with the inscription "Lectotypus Warncke 1975." However, of all Fedtschenko's Halictidae described

by Morawitz, only five specimens received such labels: *Sphecodes nigripennis* Morawitz, 1876, *S. pectoralis* Morawitz, 1876, *S. rufithorax* Morawitz, 1876, *Nomia edentata* Morawitz, 1876 and *N. rufescens* Morawitz, 1876 (Figs 32A, 33B, 34E, 35F, 36A). All other type specimens were left without nomenclatural status labels.

In his publication on the genus *Halictus*, which appeared seven years later (Warncke 1982), only the sex and locality for each lectotype is cited, for example: “*Halictus limbellus* Mor. / ♀ Lectotypus / Samarkand.” The selected specimen is not further individually identified and data on syntypes and paralectotypes are completely missing from the text. Thus, of the twenty seven designated lectotypes, fifteen are either invalid or unnecessary (when the type series includes only the holotype) and require corrections by subsequent authors (Pesenko 1986a, Ebmer 1997, current publication).

In the 1980's, Yu. Pesenko continued the study of Halictidae in the Fedtschenko's collection of the ZISP and ZMMU, designating lectotypes for seven of Morawitz's nominal taxa in the genera *Halictus*, *Nomia* and *Nomiooides* (Pesenko 1983, 1984, 1986a).

As a part of a detailed types inventory of the ZISP collection, all primary types of Halictidae, including seven species described by F. Morawitz from the collection of A. Fedtschenko, are being progressively photographed and catalogued (Astafurova and Proshchalykin 2018, 2019, 2020). The present paper is the first complete, illustrated summary of all species of the family Halictidae, described by F. Morawitz from the collection of A. Fedtschenko, an invaluable reference for researchers across this region who otherwise could not easily assign names to these difficult bees.

Materials and methods

All of the material listed below was examined for this study. In the following list, the taxa are treated in alphabetical order of the names used in the original descriptions. Each entry includes the name of the taxon in its original combination, the complete reference to the original description of the species (including the original combination and spelling of the name and the author, year and page of the description) and a list of type specimens present in the collections of the ZMMU and ZISP. The data from each label are separated by two slashes (//). Square brackets are used for English translations and when information is added to specimen label information (e.g. geographical coordinates) or published data (e.g. current name of an old place name; affiliation to a present-day country). Photographs were made using a combination of a stereomicroscope Olympus SZX10 and a digital camera (Olympus OM-D and Canon EOS70D).

Illustrations were obtained by montaging from an image series that covers different focal planes into a single in-focus image with the Helicon Focus 6. The final illustrations were post-processed for contrast and brightness using Adobe® Photoshop® software.

The classification and current species status for *Halictus* and *Lasioglossum* follow Michener (2007) and Ascher and Pickering (2020), for *Sphecodes* follow Astafurova et al. (2018a, b, 2019), for Nomiinae follow Astafurova (2014) and for Nomiooidinae follow Pesenko (1983).

Taxonomy

List of species

Subfamily Halictinae

Genus *Halictus* Latreille, 1804

1. *Halictus albitarsis* Morawitz, 1876

Figure 1

Halictus albitarsis Morawitz, 1876: 217 (key to females), 246, ♀.

Type locality. Samarkand (Uzbekistan).

Published (original) locality. Uzbekistan: Samarkand, Tashkent.

Lectotype. ♀, designated by Warncke 1982: 117, <golden circle> // 21.[III.1869] // Самаркандъ [Uzbekistan, Samarkand, 39°39'N, 66°57'E] // *Halictus albitarsis* Mor., [N]372 [handwritten by F. Morawitz] // *Lasioglossum Erylaeus albitarsis* Mor., ♀ = lectotype, det A.W. Ebmer 1993 // Syntypes Lectotypus <red label> [ZMMU].

Paralectotypes (12 ♀). 6 ♀, 11.[III.1871], 24., 25.[V.1871] // Ташкентъ [Tashkent] // [N]372; 1 ♀, 23.[III.1871] // Ташкентъ [Tashkent] // [N]372 // *Halictus albitarsis* Mor., F. Morawitz det. [handwritten by F. Morawitz] [ZMMU]; 1 ♀, <golden circle>, 11.[III.1871] // Ташкентъ [Tashkent] // *albitarsis* Mor., Typ. [handwritten by F. Morawitz]; 3 ♀, 11.[III.1871] // Ташкентъ [Tashkent] // к. [о.лекция] Ф. Моравица [Collection of F. Morawitz] // *Halictus albitarsis* Mor. [handwritten by F. Morawitz]; 1 ♀, 11.[III.1871] // Ташкентъ [Tashkent] // Paralectotypus *Halictus albitarsis* Mor., design. Warncke, [19]82 <identical red label on each paralectotype specimens, labelled by Yu. Astafurova> [ZISP].

Current status. *Lasioglossum (Sphecodogastra) leucopymatum* (Dalla Torre, 1896), replacement name for *Halictus albitarsis* Morawitz, 1876 (nec *Hylaeus albitarsis* Schenck, 1853, nec *Halictus albitarsis* Cresson, 1872).

Remarks. The secondary designation of the lectotype by Ebmer 1995: 585 is unnecessary.

Description of male. Ebmer 1995: 585.

Distribution. Kazakhstan, Turkmenistan, Uzbekistan, Afghanistan (Ebmer 1995).

2. *Halictus annulipes* Morawitz, 1876

Figure 2

Halictus annulipes Morawitz, 1876: 217 (key), 221, ♀.

Type locality. Karatyube Mt., 15 km S Samarkand (Uzbekistan).

Published (original) locality. Uzbekistan: Samarkand, Karatyube Mt. (= 15 km S Samarkand).




Figure 1. *Halictus albitalris* Morawitz, 1876, lectotype, female **A** habitus, lateral view and labels **B** head, frontal view **C** mesosoma, dorsal view **D** metasoma, dorsal view. Scale bars: 1.0 mm (**A, D**), 0.5 mm (**B, C**).



Figure 2. *Halictus annulipes* Morawitz, 1876, lectotype, female **A** habitus, lateral view **B** head, frontal view **C** mesosoma, dorsal view **D** metasoma, dorsal view **E** labels. Scale bars: 1.0 mm (**A**), 0.5 mm (**B-E**).

Lectotype. designated by Warncke 1982: 81, ♀, 20[V.1869] // Зеравшан.[сказа]
дол.[иши] [Uzbekistan, Zeravshan River valley, Karatyube Mt., 39°30'N, 66°52'E] //
Halictus annulipes Mor., [N]332 [handwritten by F. Morawitz] // F. Morawitz det.
18.7.5. // Lectotypus *Halictus annulipes* Mor. 1876, design. Warncke, 1982 <red label,
labelled by Yu. Astafurova> [ZMMU].

Paralectotypes (2 ♀). 1 ♀, 17. [V.1869] // Самарканда́й [Uzbekistan, Samarkand] // [N]332 // *H. annulipes* F. Morawitz det. 1875 [ZMMU]; 1 ♀, 20[V.1869]
// Зеравшан.[сказа] дол.[иши] [Zeravshan River valley] // *annulipes* Mor. Typ. [hand-
written by F. Morawitz] // Paralectotypus *Halictus annulipes* Mor., 1876, design.
Warncke, 1982 <identical red label on each paralectotype specimen, labelled by Yu.
Astafurova> [ZISP].

Current status. *Lasioglossum (Dialictus) annulipes* (Morawitz, 1876).

Remarks. Description of male. Kohl 1905: 238, as *Halictus metopias* Vachal (syn-
onymised by Warncke 1975: 89).

Distribution. Bulgaria, Armenia, Turkey, Russia (North Caucasus), Afghanistan,
Iran, Tajikistan, Uzbekistan, Kazakhstan, Pakistan (Astafurova and Proshchalykin
2017, Ascher and Pickering 2020).

3. *Halictus aprilinus* Morawitz, 1876

Figure 3

Halictus aprilinus Morawitz, 1876: 216 (key), 228, ♀.

Type locality. Kattakurgan (Uzbekistan).

Published (original) locality. Uzbekistan: Katty-Kurgan.

Lectotype (designated here). ♀, <golden circle>, 28. [IV.1869] [Uzbekistan, Kat-
takurgan, 39°53'N, 66°15'E] // ♀ // *H. aprilinus* n. sp., ♀, F. Morawitz det., typus
[handwritten by F. Morawitz] // Lectotypus *Halictus aprilinus* Mor., design. Astafurova
et Proshchalykin, 2020 <red label> [ZMMU].

Paralectotypes. 3 ♀, 28. [IV.1869] // Каттыкургань [Kattakurgan] // [N]343 //
Paralectotypus *Halictus aprilinus* Mor., design. Astafurova et Proshchalykin, 2020 <red
label> [ZMMU].

Current status. *Lasioglossum (Sphecodogastra) aprilinum* (Morawitz, 1876).

Remarks. Warncke (1982: 76) did not designate the lectotype, but only wrote
“Typus Mus. Moskau”.

Description of male. Blüthgen 1925a: 119, as *Halictus inexpectatus* (synonymised
by Blüthgen 1931a: 212).

Distribution. Southern Kazakhstan, Turkmenistan, Uzbekistan, Mongolia, China
(Xinjiang) (Pesenko 2007, Murao et al. 2017).

4. *Halictus atomarius* Morawitz, 1876

Figure (see Astafurova and Proshchalykin 2018: 11, figs 7a–e).

Halictus atomarius Morawitz, 1876: 218 (key), 254, ♀.

Type locality. Tashkent (Uzbekistan).

Published (original) locality. Uzbekistan: Tashkent.

Lectotype. ♀, designated by Ebmer 1985: 290, 8.[VIII.1870] // Ташкентъ [Uzbekistan, Tashkent, 41°18'N, 69°16'E] // *Halictus atomarius* Mor. [handwritten by F. Morawitz] // к.[оллекция] Ф. Моравица [Collection of F. Morawitz] // Syntypus <red label> // *Lasioglossum Evylaeus atomarium* (Mor.), ♀, Lectotypus, det. A.W. Ebmer 1985 // Lectotypus <red label> [ZISP].

Paralectotype. 1 ♀, 8.[VIII.1870] // Ташкентъ [Tashkent] // *Halictus atomarius* Mor. [handwritten by F. Morawitz] // к.[оллекция] Ф. Моравица [Collection of F. Morawitz] [ZISP].

Current status. *Lasioglossum (Evylaeus s.l.) politum atomarium* (Morawitz, 1876) (subspecies status according to Ebmer 1988b: 667).

Remarks. Description of male. Bytinski-Salz and Ebmer 1974: 211, as *Lasioglossum politum* ssp. *aramaeum* Ebmer 1974 (synonymised by Ebmer 1985: 290).

Distribution. Egypt, Palestine, Syria, Jordan, Israel, Turkey, Iran, Central Asia (Bytinski-Salz and Ebmer 1974, Warncke 1982, Pesenko 2007).

5. *Halictus cariniventris* Morawitz, 1876

Figure 4

Halictus cariniventris Morawitz, 1876: 220 (key), 226, ♂.

Type locality. Osh (Kyrgyzstan).

Published (original) locality. Uzbekistan: Samarkand, Dzhyuzak [=Jizzakh], Sokh; Kyrgyzstan: Osh.

Lectotype. ♂, designated by Blüthgen 1955: 19, 1.[VIII.1871] // Ошъ [Kyrgyzstan, Osh, 40°32'N, 72°47'E] // *Halictus cariniventris* Mor., [N]341 [handwritten by F. Morawitz] // Lectotypus *Halictus cariniventris* Mor., design. Blüthgen <red label, labelled by Yu. Astafurova> [ZMMU].

Paralectotypes (4 ♂). 2 ♂, 4., 7.[VII.1869] // Самарканда [Samarkand] // [N]341; 2 ♂, 18., 22.[VII.1870] // Джюзакъ [Dzhyuzak] // [N]341 // Paralectotypus *Halictus cariniventris*, design. Blüthgen <identical red labels on each paralectotype specimen, labelled by Yu. Astafurova> [ZMMU].

Current status. *Halictus (Mucoreochalictus) pollinosa cariniventris* Morawitz, 1876 (subspecies status according to Ebmer 1988b: 578).

Remarks. The lectotype designation by Warncke (1982: 138) is unnecessary.




Figure 3. *Halictus aprilinus* Morawitz, 1876, lectotype, female **A** habitus, lateral view and labels **B** head, frontal view **C** mesosoma, dorsal view **D** metasoma, dorsal view. Scale bars: 1.0 mm (**A**), 0.5 mm (**B–D**).



Figure 4. *Halictus cariniventris* Morawitz, 1876, lectotype, male **A** habitus, lateral view **B** metasoma, dorsal view **C** head, frontal view **D** mesosoma, dorsal view **E** labels. Scale bars: 1.0 mm.

Description of female: Strand 1921: 314, as *Halictus cariniventris* var. *creticola* (synonymised by Ebmer 1988b: 578).

Distribution. Europe (except north), Russia (European part, except north), Turkey, Israel, Iran, Afghanistan, Pakistan, Central Asia, Mongolia, North China (Pesenko 2006a, Astafurova and Proshchalykin 2017).

6. *Halictus cingulatus* Morawitz, 1876

Figure 5

Halictus cingulatus Morawitz, 1876: 218 (key), 245, ♀.

Type locality. Samarkand (Uzbekistan).

Published (original) locality. Uzbekistan: Samarkand, Dzham [near Samarkand], Aksay [near Samarkand].

Lectotype (designated here). ♀, 18.[III.1869] // Самарканда́й [Uzbekistan, Samarkand, 39°39'N, 66°57'E] // *Halictus cingulatus* Mor., [N]371 [handwritten by F. Morawitz] // Lectotypus *Halictus cingulatus* Mor., design. Astafurova et Proshchalykin, 2020 <red label> [ZMMU].

Paralectotypes (22 ♀). 11 ♀, 27., 28. [II. 1869], 18., 21. [V. 1869] // Самарканда́й [Samarkand] // [N]371; 1 ♀, 12.[V.1969] // Заравшан.[ская] дол.[иная] [Zeravshan River valley, Dzham] // [N]370; 1 ♀, 16.[V.1969] // Заравшан.[ская] дол. [иная] [Zeravshan River valley, Aksay] // [N]370 [ZMMU]; 1 ♀, the same labels, but 21.[V.1869] [ZMMU]; 1 ♀, <golden circle> // 18.[V.1869] // Самарканда́й [Samarkand] // *cingulatus* Mor. Typ. [handwritten by F. Morawitz]; 1 ♀, 18.[III.1869] // Самарканда́й [Samarkand] // к.[оллекция] Ф. Моравица [Collection of F. Morawitz]; 1 ♀, 20.[III.1869] // Самарканда́й [Samarkand] // к.[оллекция] Ф. Моравица [Collection of F. Morawitz] // *Halictus cingulatus* Mor., ♀, CoType, F. Morawitz det. [handwritten by F. Morawitz]; 1 ♀, 27.[III.1869] // Самарканда́й [Samarkand] // к.[оллекция] Ф. Моравица [Collection of F. Morawitz]; 1 ♀, 16.[III.1869] // Самарканда́й [Samarkand] // к.[оллекция] Ф. Моравица [Collection of F. Morawitz] // *Halictus cingulatus* Mor., ♀, F. Morawitz det. [handwritten by F. Morawitz]; 1 ♀, 30.[III.1869] // Заравшан.[ская] дол.[иная] [Zeravshan River valley, Samarkand] // *Halictus cingulatus* Mor. [handwritten by F. Morawitz]; 2 ♀, 12.[III.1869] // Заравшан.[ская] дол.[иная] [Zeravshan River valley, Samarkand] // к.[оллекция] Ф. Моравица [Collection of F. Morawitz] // Paralectotypus *Halictus cingulatus* Mor., design. Astafurova et Proshchalykin, 2020 <identical red labels on each paralectotype specimen> [ZISP].

Current status. *Lasioglossum (Sphecodogastra) cingulatum* (Morawitz, 1876).

Remarks. Male unknown.

The lectotype designation by Warncke (1982: 116) is invalid because he labelled none of the eleven females from Samarkand deposited in ZMMU.

Distribution. South Kazakhstan, Central Asia, Pakistan (Ebmer 1995, Murao et al. 2017).

7. *Halictus croceipes* Morawitz, 1876

Figure 6

Halictus croceipes Morawitz, 1876: 217 (key to ♀), 220 (key to ♂), 224, ♀, ♂.

Type locality. Yeri, Sughd Province (Tajikistan).

Published (original) locality. Kyrgyzstan: Taka; Uzbekistan: Karatyube [Mt., 15 km S Samarkand], Dzham Gorge, Urgut; Tajikistan: Iori.

Lectotype. ♀, designated by Ebmer 1997: 956, <golden circle> // 1.[VI.1869] // Заравшан.[анская] дол.[и на] [Tajikistan, Zeravshan River valley, Iori (= Yeri), 39°30'N, 67°52'E] // *croceipes* Mor. Typ [handwritten by F. Morawitz] // *croceipes* Mor. Blüthgen det. 1935 // Lectotype <red label> // *Lasioglossum (Erylaeus) croceipes* (Mor.), ♀, Lectotypus, det. A.W. Ebmer 1974 [ZMMU].

Paralectotypes (6 ♀). 1 ♀, 20.[V.1869] // Заравшан.[ская] дол.[ина] [Zeravshan River valley, Karatyube] // *croceipes* Mor., [N]338 [handwritten by F. Morawitz] // *croceipes* Mor. Blüthgen det. 1935; 3 ♀, 8.[VIII.1871] // Така [Taka] // [N]338; 1 ♀, 13.[V.1869] // Джамское ущ. [Dzham Gorge] // [N]338 [ZMMU]; 1 ♀, 1.[VI.1869] // Заравшан.[ская] дол.[ина] [Zeravshan River valley, Iori] // к.[оллекция] Ф. Моравица [Collection of F. Morawitz] // *croceipes* Mor., Blüthgen det. 1935 // Paralectotypus *Halictus croceipes* Mor., design. Ebmer 1997 <identical red labels on each paralectotype specimen, labelled by Yu. Astafurova> [ZISP].

Current status. *Lasioglossum (Hemihalictus) croceipes* (Morawitz, 1876).

Remarks. The lectotype designation by Warncke (1982: 106) is invalid because there are two females in ZMMU from the Zeravshan River valley, neither with Warncke's lectotype label.

Distribution. Turkey, Iran, Afghanistan, Central Asia, Kazakhstan (Ebmer 1997, Murao et al. 2017).

8. *Halictus desertorum* Morawitz, 1876

Figure 7

Halictus desertorum Morawitz, 1876: 217 (key), 228, ♀.

Type locality. Kattakurgan (Uzbekistan).

Published (original) locality. Uzbekistan: Katty-Kurgan.

Lectotype. ♀, designated by Warncke 1982: 107, 28.[IV.1869] // Катты-Курганъ [Uzbekistan, Katty-Kurgan (=Kattakurgan), 39°53'N, 66°15'E] // *Halictus desertorum* Mor., [N]344 [handwritten by F. Morawitz] // Lectotypus *Halictus desertorum* Mor., design. Warncke <red label, labelled by Yu. Astafurova> [ZMMU].

Current status. *Halictus (Placidothalictus) desertorum* Morawitz, 1876.

Remarks. Description of male. Blüthgen 1929: 84, figs 9a, b.

Distribution. Turkey, southern Kazakhstan, Turkmenistan, Uzbekistan, Pakistan (Blüthgen 1929, Ebmer 1988a).




Figure 5. *Halictus cingulatus* Morawitz, 1876, lectotype, female **A** habitus, lateral view and labels **B** head, frontal view **C** mesosoma, dorsal view **D** metasoma, dorsal view. Scale bars: 1.0 mm.



Figure 6. *Halictus croceipes* Morawitz, 1876, lectotype, female **A** habitus, lateral view and labels **B** head, frontal view **C** mesosoma, dorsal view **D** metasoma, dorsal view. Scale bars: 1.0 mm (**A, D**), 0.5 mm (**B, C**).




Figure 7. *Halictus desertorum* Morawitz, 1876, lectotype, female **A** habitus, lateral view and labels **B** head, frontal view **C** mesosoma, dorsal view **D** metasoma, dorsal view. Scale bars: 1.0 mm (**A**), 0.5 mm (**B-D**).

9. *Halictus determinatus* Morawitz, 1876

Figure 8

Halictus determinatus Morawitz, 1876: 217 (key), 233, ♀.

Type locality. 30 km SSE Samarkand, Sangu-dzhuman Pass (Uzbekistan).

Published (original) locality. Uzbekistan: “On the road to Sangy-Dzhuman [pass] and the Kulbasy Mountain”.

Lectotype. ♀, designated by Pesenko 1984: 20, 25.[V.1869] // Санғы-Джуманъ [Uzbekistan, Sangu-dzuman Pass, Zeravshan Ridge, 30 km SSE Samarkand, 39°27'N, 67°14'E] // *Halictus determinatus* Mor., [N]351 [handwritten by F. Morawitz] // Lectotypus *H. determinatus* Mor., ♀, design. Pesenko [1]981 <red label> [ZMMU].

Paralectotypes (6 ♀). 3 ♀, the same label as in the lectotype; 1 ♀, 25.[V.1869] // Заравшан.[сказ] дол.[иина] [Zeravshan River valley] // [N]351 [ZMMU]; 2 ♀, Санғы-Джуманъ [Sangu-Dzuman] // к.[оллекция] Ф. Моравица [Collection of F. Morawitz] // *determinatus* F. Mor., ♀ [handwritten by F. Morawitz] // Paralectotypus *H. determinatus* Mor., ♀, design. Pesenko [1]981 <identical red labels on each paralec-totype specimen, labelled by Yu. Pesenko> [ZISP].

Current status. *Halictus (Platyhalictus) determinandus* Dalla Torre, 1896, replacement name for *H. determinatus* Morawitz, 1876 (nec *H. determinatus* Walker, 1871).

Remarks. Description of male. Ebmer 1980: 471, Fig. 1.

Distribution. A rare montane Central Asian species. Northern Afghanistan, east-ern Uzbekistan, Tajikistan and Kyrgyzstan (Pesenko 2005a).

10. *Halictus equestris* Morawitz, 1876

Figure 9

Halictus equestris Morawitz, 1876: 217 (key), 242, ♀.

Type locality. Khozyay-Dun (Uzbekistan).

Published (original) locality. Uzbekistan: Khodzhaduk [= Khozyay-Dun].

Lectotype. ♀, designated by Warncke 1982: 105, 21.[V.1869] // Заравшан.[сказ] дол.[иина] [Uzbekistan, Zeravshan River valley, Khozyay-Dun, 39°24'N, 67°01'E] // *Halictus equestris* Mor., [N]366 [handwritten by F. Morawitz] // Lectotypus *Halictus equestris* Mor., design. Warncke [19]82 <red label> [ZMMU].

Current status. *Lasioglossum (Lasioglossum) equestre* (Morawitz, 1876).

Remarks. Description of male. Morawitz 1876: 219 (key), 243, as *Halictus ferghanicus* (synonymised by Blüthgen 1926: 391).

According to the Fedchenko list (Baker 2004: 243), the locality given in Warncke's lectotype designation is in Uzbekistan, not Tajikistan.

Distribution. Turkey, Uzbekistan, Tajikistan, Kyrgyzstan, south-eastern Kazakh-stan (Pesenko 1986a, Ascher and Pickering 2020).




Figure 8. *Halictus determinatus* Morawitz, 1876, lectotype, female **A** habitus, lateral view **B** metasoma, dorsal view **C** mesosoma, dorsal view **D** head, frontal view **E** labels. Scale bars: 1.0 mm.




Figure 9. *Halictus equestris* Morawitz, 1876, lectotype, female **A** habitus, lateral view and labels **B** head, frontal view **C** mesosoma, dorsal view **D** metasoma, dorsal view. Scale bars: 1.0 mm.

11. *Halictus ferghanicus* Morawitz, 1876

Figure 10

Halictus ferghanicus Morawitz, 1876: 219 (key), 243, ♂.

Type locality. Shakhimardan (Uzbekistan).

Published (original) locality. Uzbekistan: near Shakhimardan.

Lectotype. ♂, designated by Warncke 1982: 105, <golden circle> // 6.[VII.1871] // Шагимарданъ [Shakhimardan in the Uzbek enclave in the territory of Kyrgyzstan, Alai Ridge; 39°58'N, 71°47'E] // *Halictus ferghanicus* Mor., [N]367 [handwritten by F. Morawitz] // Lectotypus *Halictus ferghanicus* Mor., ♂, design. Warncke [19]82 <red label, labelled by Yu. Pesenko > [ZMMU].

Paralectotypes (2 ♂). 1 ♂, <golden circle> // 2.[VII.1871] // Шагимарданъ [Shagimardan] // *ferghanicus* Mor., Typ. [handwritten by F. Morawitz]; 1 ♂, 2.[VII.1871] // Шагимарданъ [Shagimardan] // к.[оллекция] Ф. Моравица [Collection of F. Morawitz] // *Halictus ferghanicus* M. [handwritten by F. Morawitz] // *Lasioglossum equestre* Mor., Pesenko det., 1985 // Paralectotypus, *Halictus ferghanicus* Mor., design. Warncke, [19]82 <identical red labels on each paralecotype specimen, labelled by Yu. Pesenko> [ZISP].

Current status. *Lasioglossum (Lasioglossum) equestre* (Morawitz, 1876) (synonymised by Blüthgen 1926: 391).

Distribution. See *Halictus equestris* Morawitz, 1876.

12. *Halictus fucusus* Morawitz, 1876

Figure 11

Halictus fucusus Morawitz, 1876: 219 (key), 230, ♂.

Type locality. 30 km SE Kozhatogai, Turkistan Province (Kazakhstan).

Published (original) locality. Uzbekistan: steppe between Tashkent and Syrdarya River.

Holotype. ♂, 18.[V.1871] // Степь м.[еждау] С.[ыр] А.[арьей] и Т.[ашкентом] [Kazakhstan, Turkistan Province, steppe between Syrdarya River and Tashkent, 30 km SE Kozhatogai, 41°47'N, 68°23'E] // *Halictus fucusus* Mor., [N]345 [handwritten by F. Morawitz] // *Halictus senilis* Evers. v. *fucusus* Mor., ♂, P. Blüthgen det. // Holotypus <red label> [ZMMU].

Current status. *Halictus (Argalictus) senilis* (Eversmann, 1852) (synonymised by Blüthgen 1922: 47).

Remarks. The lectotype designation of Warncke (1982: 148) is unnecessary as the species was described from a single male that was directly written about by Morawitz (1876: 231).

Distribution. North Africa, South and East Europe, Russia (south to Urals Mountains on the east), Caucasus, Turkey, Near East, Iraq, Iran, Afghanistan, Central Asia, Kazakhstan, Mongolia, China (Astafurova and Proshchalykin 2017).




Figure 10. *Halictus ferghanicus* Morawitz, 1876, lectotype, male **A** habitus, lateral view and labels **B** head, frontal view **C** mesosoma, dorsal view **D** metasoma, dorsal view. Scale bars: 1.0 mm.




Figure 11. *Halictus fucusus* Morawitz, 1876, holotype, female **A** habitus, lateral view and labels **B** metasoma, dorsal view **C** head, frontal view **D** mesosoma, dorsal view. Scale bars: 1.0 mm.

13. *Halictus fulvitarsis* Morawitz, 1876

Figure 12

Halictus fulvitarsis Morawitz, 1876: 219 (key), 239, ♂.

Type locality. Khodzha-Chiburgan River (Tajikistan/Kyrgyzstan, not Uzbekistan as was indicated in Warncke 1982: 91).

Published (original) locality. Tajikistan/Kyrgyzstan: Khodzha-Chiburgan River.

Holotype. ♂, 26.[VI.1871] // Чибурганъ [Tajikistan/Kyrgyzstan: Khodzha-Chiburgan River (near Vorukh, ≈ 39°48'N, 70°41'E)] // *Halictus fulvitarsis* Mor., [N]361 [handwritten by F. Morawitz] // Holotypus <red label> [ZMMU].

Current status. *Lasioglossum (Lasioglossum) fulvitarse* (Morawitz, 1876).

Remarks. Description of male. Blüthgen 1934b: 147.

The lectotype designation by Warncke (1982: 91) is unnecessary as the species was described from a single male that was directly written about by Morawitz (1876: 240).

Distribution. Tajikistan, Kyrgyzstan, Kazakhstan (Morawitz 1876, Pesenko 1986a, Murao et al. 2017).

14. *Halictus funerarius* Morawitz, 1876

Figure (see Astafurova and Proshchalykin 2020: 414, figs 11a–e).

Halictus funerarius Morawitz, 1876: 217 (key), 235, ♀.

Type locality. 30 km SSE Samarkand, Sangu-dzhuman Pass (Uzbekistan).

Published (original) locality. Uzbekistan: Sangu Dzuman.

Lectotype. ♀, designated by Pesenko 1984: 21, <golden circle> // Санғы Джуманъ [Uzbekistan, Sangydzhuman Pass, 30 km SSE Samarkand, 39°22'N, 67°00'E] // 25.[V.1869] // *funerarius* Mor. Typ. [handwritten by F. Morawitz] // Lectotypus *H. funerarius* Mor., ♀, design. Pesenko [1]981 <red label> // Zoological Institute St. Petersburg INS_HYM_0000159 [ZISP].

Paralectotypes (8 ♀). 8 ♀, the same label as in lectotype // Paralectotypus *H. funerarius* Mor. design. Pesenko [1]981 <red labels> [3 ♀ – ZISP; 5 ♀ – ZMMU].

Current status. *Halictus (Protohalictus) funerarius* Morawitz, 1876.

Distribution. A rare montane Central Asian species: Kazakhstan, Uzbekistan, Tajikistan, Western Kyrgyzstan, Iran, north-eastern Afghanistan and north-western China (Pesenko 2005a).

15. *Halictus fuscicollis* Morawitz, 1876

Figure 13

Halictus fuscicollis Morawitz, 1876: 217 (key), 229, ♀.



Figure 12. *Halictus fulvitarsis* Morawitz, 1876, holotype, female **A** habitus, lateral view and labels **B** metasoma, dorsal view **C** head, frontal view **D** mesosoma, dorsal view. Scale bars: 1.0 mm.

Type locality. 50 km NW Chardara, Kyzylkum Desert (Turkistan Province, Kazakhstan).

Published (original) locality. Kazakhstan: “Kyzyl-Kum Steppe, near Baybek”.

Lectotype. ♀, designated by Warncke 1982: 137, 30.[IV.1871] // Кизилькумъ [Kazakhstan, Baybek Well, Kyzylkum Desert, ca. 50 km NW Chardara, $\approx 41^{\circ}44'N$, $67^{\circ}54'E$] // *Halictus fuscicollis* Mor., [N]345 [handwritten by F. Morawitz] // *Halictus Vestitohalictus fuscicollis* Mor., ♀, Lectotypus, design. A.W. Ebmer 1994 //




Figure 13. *Halictus fuscicollis* Morawitz, 1876, lectotype, female **A** habitus, lateral view and labels **B** head, frontal view **C** mesosoma, dorsal view **D** metasoma, dorsal view. Scale bars: 0.5 mm.

Lectotypus *Halictus fuscicollis* Mor., design. Warncke <red label, labelled by Yu. Pesenko> [ZMMU].

Paralectotype. 1 ♀, <golden circle> Kisilkum [handwritten by F. Morawitz] // *fuscicollis* Mor. Typ. [handwritten by F. Morawitz] // Paralectotypus <red label>, labelled by Yu. Astafurova [ZISP].

Current status. *Halictus (Placidochalictus) fuscicollis* Morawitz, 1876.

Remarks. Description of male. Morawitz 1894: 68, as *Halictus flavocallosus* (synonymised by Blüthgen 1931a: 214).

Distribution. Southern Kazakhstan, Turkmenistan, Iran, China (Xinjiang) (Morawitz 1876, 1894, Murao et al. 2017, Ascher and Pickering 2020).

16. *Halictus hyalinipennis* Morawitz, 1876

Figure (see Astafurova and Proshchalykin 2018: 23, figs 21a–e).

Halictus hyalinipennis Morawitz, 1876: 218 (key to females), 220 (key to males), 253–254, ♀, ♂.

Type locality. Tashkent (Uzbekistan).

Published (original) locality. Kazakhstan: Chardara, steppe between Tashkent and Syr-Darya; Uzbekistan: Dzhamsk Gorge, Dzhizmansk Gorge, Ulus, Dzham, Urgut, Keles, Samarkand, Soch, Shakhimardan, Uch-Kurgan; Kyrgyzstan: Alay, Osh, Gulsha, Taka.

Lectotype. ♀, designated by Astafurova and Proshchalykin 2018: 22, Ташкентъ [Uzbekistan, Tashkent, 41°18'N 69°16'E] // *hyalinipennis* F. Mor., ♀ [handwritten by F. Morawitz] // к.[оллекция] Ф. Моравица [Collection of F. Morawitz] // Lectotypus *Halictus hyalinipennis* Morawitz, 1876, ♀, design. Astafurova & Proshchalykin 2018 <red label> [ZISP].

Paralectotypes (29 ♀, 10 ♂). 1 ♀, Шагимарданъ [Shagimardan] // 3.[VII.1871] // к.[оллекция] Ф. Моравица [Collection of F. Morawitz]; 1 ♂, Така [Taka] // к.[оллекция] Ф. Моравица [Collection of F. Morawitz]; 1 ♀, 1 ♂, Сохъ [Sokh] // 29. [29.VI.1871] // к.[оллекция] Ф. Моравица [Collection of F. Morawitz]; 1 ♂, Уч-Курганъ [Uch- Kurgan] // к.[оллекция] Ф. Моравица [Collection of F. Morawitz]; 2 ♀, Ташкентъ [Tashkent] // к.[оллекция] Ф. Моравица [Collection of F. Morawitz] [ZISP]; 1 ♂, Ошъ [Osh] // 1.[VIII.1871]; 1 ♂, Самарканъ [Samarkand] // 4.[VII.1869] // [N]384; 3 ♂, Самарканъ [Samarkand] // 7.[VII.1869]; 2 ♀, Самарканъ [Samarkand] // 3., 21.[III.1869]; 16 ♀, Ташкентъ [Tashkent] // 26., 27.[III.1871], 21., 23.[V.1871] and 1.,3.,5.,10.[VI.1871]; 1 ♀, Чардара [Chardara] // 27.[IV.1871]; 1 ♀, Учъ-Курганъ [Uch-Kurgan] // 15.[VII.1871]; 2 ♂, Шагимарданъ [Shagimardan] // 2.[VII.1871] 5 ♀, Заравшан.[ская] дол.[ина] [Zeravshan River valley], 3., 11., 18., 23.[III.1871] // Paralectotypus *Halictus hyalinipennis* Morawitz, 1876, design. Astafurova & Proshchalykin 2018 <identical red labels on each paralectotype specimen> [ZMMU].

Current status. *Lasioglossum (Sphecodogastra) hyalinipenne* (Morawitz, 1876).

Distribution. Iran, Afghanistan, Kazakhstan, Uzbekistan, Kyrgyzstan, Tajikistan (Morawitz 1876, Ebmer 1974, Warncke 1982, Murao et al. 2017).

17. *Halictus laevinodis* Morawitz, 1876

Figure (see Astafurova and Proshchalykin 2018: 24, figs 23a–e).

Halictus laevinodis Morawitz, 1876: 218 (key to females), 248, ♀.

Type locality. 30 km SSE Samarkand, Sangu-dzhuman Pass (Uzbekistan).

Published (original) locality. Uzbekistan: Sangy Dzhuman.

Lectotype (designated here). ♀, <golden circle> // Санғы Джуманъ [Uzbekistan, Sangy-dzhuman Pass, 30 km SSE Samarkand, Zeravshan Ridge, 39°27'N, 67°14'E] // 25.[III.1869] // *laevinodis* Mor., Typ. [handwritten by F. Morawitz] // Lectotype *Halictus laevinodis*, design. Astafurova et Proshchalykin, 2020 <red label> [ZISP].

Paralectotypes (2 ♀). 1 ♀, 25.[III.1869] // Санғы Джуманъ [Sangy Dzhuman] // *Halictus laevinodis* Mor. [handwritten by F. Morawitz]; 1 ♀ 25.[III.1869] // Санғы Джуманъ [Sangy Dzhuman] // [N]375 // Paralectotype *Halictus laevinodis*, design. Astafurova et Proshchalykin, 2020 <identical red labels on each paralectotype specimen> [ZMMU].

Current status. *Lasioglossum (Hemihalictus) laevinode* (Morawitz, 1876).

Remarks. Astafurova and Proshchalykin (2018: 23) indicated the lectotype specimen as «Holotype», but Morawitz (1876) did not directly indicate a single specimen and two specimens from the type series are deposited in ZMMU.

Description of male. Blüthgen, 1934b: 154, Fig. 3.

Distribution. Kazakhstan, Uzbekistan, Tajikistan, Kyrgyzstan, Afghanistan (Morawitz 1876, Blüthgen in Popov 1935, Ebmer 1980, Murao et al. 2017).

18. *Halictus limbellus* Morawitz, 1876

Figure 14

Halictus limbellus Morawitz, 1876: 218 (key), 249, ♀.

Type locality. Samarkand (Uzbekistan).

Published (original) locality. Uzbekistan: Samarkand; Tajikistan: Peti.

Lectotype (designated here). ♀, 5.[IV.1869] // Самарканда [Uzbekistan, Samarkand, 39°39'N, 66°57'E] // *Halictus limbellus* Mor., [N]377 [handwritten by F. Morawitz] // Lectotypus *Halictus limbellus* Mor., Astafurova et Proshchalykin, 2020 <red label> [ZMMU].

Paralectotypes (3 ♀). 1 ♀, 5.[IV.1869] // Самарканда [Samarkand] // [N]377 // Typus <red label> [ZMMU]; 1 ♀, <golden circle> // 5.[IV.1869] // Самарканда [Samarkand] // [N]377 // Typus <red label> [ZMMU].




Figure 14. *Halictus limbellus* Morawitz, 1876, lectotype, female **A** habitus, lateral view and labels **B** head, frontal view **C** mesosoma, dorsal view **D** metasoma, dorsal view. Scale bars: 1.0 mm (**A**), 0.5 mm (**B–D**).

kand] // *Halictus limbellus* F. Mor., Typ. [handwritten by F. Morawitz]; 1 ♀, Самаркандъ [Samarkand] // к.[оллекция] Ф. Моравица [Collection of F. Morawitz] // *limbellus* Mor., Typ. [handwritten by F. Morawitz] // Paralectotype *Halictus limbellus* Mor., Astafurova et Proshchalykin, 2020 <identical red labels on each paralectotype specimen> [ZISP].

Current status. *Lasioglossum (Hemihalictus) limbellum* (Morawitz, 1876).

Remarks. Description of male. Blüthgen 1930: 763.

The lectotype designation by Warncke (1982: 69) is invalid because he labelled no female of the two females from Samarkand deposited in ZMMU.

Distribution. Central and eastern Europe, Turkey, Caucasus, Russia (North Caucasus), Israel, Iran, Afghanistan, Tajikistan, Uzbekistan, Kazakhstan, China (Gansu) (Pesenko 2007, Astafurova and Proshchalykin 2017, Murao et al. 2017).

19. *Halictus longirostris* Morawitz, 1876

Figure 15

Halictus longirostris Morawitz, 1876: 216 (key to females), 219 (key to males), 236, ♀, ♂.

Type locality. Shakhimardan (Uzbekistan).

Published (original) locality. Uzbekistan: on the road to Sangy-dzhuman Pass, Shakhimardan; Tajikistan: Peti.

Lectotype. ♂, designated by Warncke 1982: 80, 3.[VII.1871] // Шагимарданъ [Shakhimardan in the Uzbek enclave in the territory of Kyrgyzstan, Alai Ridge; 39°58'N, 71°47'E] // *Halictus longirostris* Mor., [N]356 [handwritten by F. Morawitz] // Lectotype *Halictus longirostris* Mor., design. Warncke [19]82 <red label, labelled by Yu. Astafurova> [ZMMU].

Paralectotypes (2 ♀). 1 ♀, 25.[V.1869] // Санғы Джуманъ [Sangy Dzhuman] // [N]356; 1 ♀, 12.[VII.1870] // Фанъ [Fan] // [N]356 [ZMMU]; 1 ♀, 2.[VII.1871] // Шагимарданъ [Shaghimardan] // к.[оллекция] Ф. Моравица [Collection of F. Morawitz] // *Halictus longirostris* Mor. [handwritten by F. Morawitz] // Paralectotype *Halictus longirostris* Mor., design. Warncke < identical red labels on each paralectotype specimen, labelled by Yu. Astafurova> [ZISP].

Current status. *Lasioglossum (Hemihalictus) longirostre* (Morawitz, 1876).

Distribution. Greece, Israel, Lebanon, Turkey, Caucasus, Iran, Afghanistan, Central Asia, Kazakhstan, China (Xinjiang) (Murao et al. 2017, Niu et al. 2020).

20. *Halictus maculipes* Morawitz, 1876

Figure 16

Halictus maculipes Morawitz, 1876: 218 (key to females), 247, ♀.



Figure 15. *Halictus longirostris* Morawitz, 1876, lectotype, male **A** head, frontal view **B** habitus, lateral view **C** mesosoma, dorsal view **D** metasoma, dorsal view **E** labels. Scale bars: 1.0 mm (**A, B**), 0.5 mm (**C, D**).

Type locality. Sokh District [Uzbekistan].

Published (original) locality. Kekh [Sokh District, Uzbekistan].

Lectotype. ♀ designated by Warncke 1982: 68, <golden circle> // 27.[VI.1871] // Сохъ [Uzbekistan, Sokh District, $\approx 39^{\circ}57'N$, $71^{\circ}07'E$] // *Halictus maculipes* Mor., [N]373 [handwritten by F. Morawitz] // Lectotypus *Halictus maculipes* Mor., design. Warncke <red label, labelled by Yu. Astafurova> [ZMMU].

Current status. *Lasioglossum (Hemihalictus) maculipes* (Morawitz, 1876).

Remarks. Male unknown.

Distribution. Turkey, Turkmenistan, Tajikistan, Uzbekistan, Iran, Afghanistan (Morawitz 1876, Ebmer 1984, 1986, Warncke 1982, Ascher and Pickering 2020).

21. *Halictus melanarius* Morawitz, 1876

Figure 17

Halictus melanarius Morawitz, 1876: 241, ♂.

Type locality. Shakhimardan (Uzbekistan).

Published (original) locality. Near Shakhimardan.

Holotype. ♂, 9.[VII.1871] // Шахимарданъ [Shakhimardan in the Uzbek enclave in the territory of Kyrgyzstan, Alai Ridge; $39^{\circ}58'N$, $71^{\circ}47'E$] // *Halictus melanarius* Mor., [N]364 [handwritten by F. Morawitz] // *Lasioglossum fallax* (Mor.) syn: *melanarium* (Mor.) det. A.W. Ebmer 1979 // Holotypus <red label> [ZMMU].

Current status. *Lasioglossum (Lasioglossum) fallax* ssp. *melanarium* (Morawitz, 1876) (subspecies status according to Ebmer 1998: 382).

Remarks. Description of female: Ebmer 1980: 493, Figs 12 and 13, as *Lasioglossum melanum* Ebmer, 1980 (synonymised by Ebmer 1998: 382).

The lectotype designations by Ebmer (1980: 495) and by Warncke (1982: 91) were unnecessary as the species was described from a single male that was directly written about by Morawitz (1876: 241).

Distribution. Kazakhstan, Uzbekistan, Tajikistan, Kyrgyzstan, Afghanistan, Mongolia (Hovd) (Pesenko 2006b).

22. *Halictus minor* Morawitz, 1876

Figure (see Astafurova and Proshchalykin 2020: 418, figs 15a–e).

Halictus minor Morawitz, 1876: 217 (key), 233, ♀.

Type locality. 30 km SSE Samarkand, Sangu-dzhuman Pass (Uzbekistan).

Published (original) locality. Uzbekistan: Gus [near Urgut], Sangy-Dzhuman; Tajikistan: Pyandzhikent.




Figure 16. *Halictus maculipes* Morawitz, 1876, lectotype, female **A** habitus, lateral view and labels **B** head, frontal view **C** mesosoma, dorsal view **D** metasoma, dorsal view. Scale bars: 1.0 mm (**A**), 0.5 mm (**B–D**).



Figure 17. *Halictus melanarius* Morawitz, 1876, holotype, male **A** view habitus, lateral view and labels **B** head, frontal **C** genitalia, dorsal view **D** mesosoma, dorsal view **E** metasoma, dorsal view. Scale bars: 1.0 mm (**A, B, D, E**), 0.5 mm (**C**).

Lectotype. ♀, designated by Pesenko 1984: 23, <golden circle> // Сангы Джуманъ [Uzbekistan, Sangy-dzhuman Pass, 30 km SSE Samarkand, 39°20'N, 67°19'E] // 25.[V.1869] // *minor* Mor. [handwritten by F. Morawitz] // Lectotypus *H. minor* Mor., design. Pesenko, [1]1981, ♀ <red label> // Zoological Institute St. Petersburg INS_HYM_0000164 [ZISP].

Paralectotypes (3 ♀). 1 ♀, Сангы Джуманъ [Sangy Dzhuman] // к. [коллекция] Ф. Моравица [Collection of F. Morawitz] // *minor* Mor. [handwritten by F. Morawitz] // Syn.: *jarkandensis* Strand, ♀ // Paralectotypus *H. minor* Mor., design. Pesenko, [1]1981, ♀ <red label> [ZISP]; 1 ♀, 25.[V.1869] // Сангы Джуманъ [Sangy Dzhuman] // *Halictus minor* Mor., [N]352 [handwritten by F. Morawitz]; 1 ♀, 24.[V.1869] // Заравшан. [сказа] дол. [иши] [Zeravshan River valley, Gus] // [N]352 // Paralectotypus *H. minor* Mor., design. Pesenko, [1]1981, ♀ <identical red labels on each paralectotype specimen> [ZMMU].

Current status. *Halictus (Platyhalictus) minor* Morawitz, 1876.

Remarks. Description of male. Blüthgen 1936: 295, fig. 11.

Distribution. Azerbaijan, Afghanistan, Iran, Kazakhstan, Central Asia, Altai, Pakistan, north-western and northern China, northern India (Pesenko 2005a, b).

23. *Halictus modernus* Morawitz, 1876

Figure 18

Halictus modernus Morawitz, 1876: 217 (key), 235, ♀.

Type locality. Samarkand (Uzbekistan).

Published (original) locality. Uzbekistan: near Samarkand.

Holotype. ♀, 5.[VII.1870] // Самарканда [Uzbekistan, Samarkand, 39°39'N, 66°57'E] // *Halictus modernus* Mor., [N]354 [handwritten by F. Morawitz] // Holotype <red label> [ZMMU].

Current status. *Halictus (Lampralictus) modernus* Morawitz, 1876.

Remarks. The lectotype designation of Warncke (1982: 147) is unnecessary as the species was described from a single female that was directly written about by Morawitz (1876: 235).

Description of male. Ebmer 1984: 315, figs 3–5.

Distribution. Turkmenistan, Uzbekistan, Kyrgyzstan, Afghanistan, Pakistan (Pesenko 2005a).

24. *Halictus nasica* Morawitz, 1876

Figure 19

Halictus nasica Morawitz, 1876: 216 (key to females), 219 (key to males), 229, ♀, ♂.

Type locality. Samarkand (Uzbekistan).




Figure 18. *Halictus modernus* Morawitz, 1876, holotype, female **A** habitus, lateral view **B** head, frontal view **C** mesosoma, dorsal view **D** labels **E** metasoma, dorsal view. Scale bars: 1.0 mm.

Published (original) locality. Southern Kazakhstan: Kysyl-Kum [desert] near draw-well Chakany; Uzbekistan: steppe between Katty-Kurgan and Ulus, Samarkand, Murzarabat, Chinaz, Sokh.

Lectotype (designated here). ♀, 9.[VI.1869] // Заравшан.[сказа] дол.[иина] [Uzbekistan, Zeravshan River valley, near Samarkand, 39°39'N, 66°57'E] // *Halictus nasica* Mor., [N]346 [handwritten by F. Morawitz] // Lectotypus *Halictus nasica* Mor., design. Astafurova et Proshchalykin, 2020 <red label> [ZMMU].

Paralectotypes (28 ♀, 44 ♂). 14 ♀, the same label as in the lectotype; 11 ♀, 2 ♂, 9., 13.[VI.1869], 4., 7.[VII.1869] // Самарканда [Samarkand] // [N]346; 12 ♂, 24.[VII.1870], 29.[VIII.1870] // Мурзарабад [Murzababad] // [N]346; 1 ♀, 9 ♂, 28.[IV.1871] // Кизилькумъ [Kizilkum] // [N]346; 17 ♂, 25.[VII.1870] // Чиназъ [Chinaz] // [N]346 [ZMMU]; 1 ♀, 28.[IV.1871] // Кизилькумъ [Kizilkum] // к.[оллекция] Ф. Моравица [Collection of F. Morawitz] // *Halictus nasica* Mor. [handwritten by F. Morawitz]; 1 ♀, <golden circle> // 9.[VI.1869] // Заравшан.[сказа] дол. [иина] [Zeravshan River valley, near Samarkand] // *nasica* Mor. Typ. [handwritten by F. Morawitz]; 3 ♂, 28.[VI.1871] // Сохъ [Sokh] // к.[оллекция] Ф. Моравица [Collection of F. Morawitz]; 1 ♂, 9.[VI.1869] // Самарканда [Samarkand] // к.[оллекция] Ф. Моравица [Collection of F. Morawitz] // Paralectotypus *Halictus nasica* Mor., design. Astafurova et Proshchalykin, 2020 <identical red labels on each paralectotype specimen> [ZISP].

Current status. *Halictus (Vestitohalictus) nasica* Morawitz, 1876.

Remarks. The lectotype designation by Warncke (1982: 138) is invalid because he labelled none of the 15 females from “valle Serafshan” deposited in ZMMU.

Distribution. Morocco, Kazakhstan, Uzbekistan, Turkmenistan, Iran, Afghanistan, Pakistan (Murao et al. 2017, Ascher and Pickering 2020).

25. *Halictus nigrilabris* Morawitz, 1876

Figure 20

Halictus nigrilabris Morawitz, 1876: 249, ♂.

Type locality. Sarafschan River valley, Yeri (Tajikistan).

Published (original) locality. Tajikistan: valley Sarafschan between Iori and Dashty-Kazy.

Lectotype. ♂, designated by Warncke 1982: 91, <golden circle> // 31.[V.1869] // Заравшан.[сказа] дол.[иина] [Tajikistan, Zeravshan River valley, near Iori (= Yeri), 39°29'N, 67°53'E] // *Halictus nigrilabris* Mor., [N]378 [handwritten by F. Morawitz] // Lectotypus *Halictus nigrilabris* Mor. design. Warncke [1]1982 <red label, labelled by Yu. Pesenko> [ZMMU].

Paralectotype. 1 ♂, Заравшан.[сказа] дол.[иина] [Zeravshan River valley] // *Halictus nigrilabris* F. Morawitz [handwritten by F. Morawitz] // к.[оллекция] Ф. Моравица [Collection of F. Morawitz] // Paralectotypus *Halictus nigrilabris* Mor., design. Warncke




Figure 19. *Halictus nasica* Morawitz, 1876, lectotype, female **A** habitus, lateral view and labels **B** head, frontal view **C** mesosoma, dorsal view **D** metasoma, dorsal view. Scale bars: 1.0 mm (**A,D**), 0.5 mm (**B,C**).




Figure 20. *Halictus nigrilabris* Morawitz, 1876, lectotype, male **A** habitus, lateral view and labels **B** head, frontal view **C** mesosoma, dorsal view **D** antennae, dorsal view **E** metasoma, dorsal view. Scale bars: 1.0 mm.

[1]982 <red label, labelled by Yu. Pesenko> // Zoological Institute St. Petersburg, INS_HYM 0000076 [ZISP].

Current status. *Lasioglossum (Lasioglossum) nigrilabre* (Morawitz, 1876).

Remarks. Description of female. Blüthgen 1931b: 336, as *Halictus subprasinus* (synonymised by Ebmer 1978: 39).

Distribution. Iran, Afghanistan, Turkmenistan, Uzbekistan, Tajikistan, Kyrgyzstan (Blüthgen 1931b, Pesenko 1986a).

26. *Halictus nigripes* Morawitz, 1876

Figure 21

Halictus nigripes Morawitz, 1876: 251, ♂.

Type locality. Karazuk, vicinity of Shakhimardan (Uzbekistan).

Published (original) locality. Tajikistan: Iori George, Iskander River; Uzbek enclave in Kyrgyzstan: Karakazuk; Kyrgyzstan: Alay.

Lectotype. ♂, designated by Blüthgen 1934a: 302, 11.[VII.1871] // Каразукъ [Uzbekistan, Karazuk, vicinity of Shakhimardan 39°60'N, 71°50'E] // *Halictus nigripes* Mor., [N]380 [handwritten by F. Morawitz] // *nigripes* Mor., ♂, lecto-holotype, Blüthgen det. 1933 // Lectotypus *Halictus nigripes* Mor., design. Blüthgen [19]34 <red label> labelled by Yu. Astafurova [ZMMU].

Paralectotypes (3 ♂). 1 ♂, 23.[VII.1871] // Алай [Alay] // [N]380 // *nigripes* Mor. ♂, Lecto-Paratype, Blüthgen det., 1933 [ZMMU]; 1 ♂, <golden circle> // 23.[VII.1871] // Алай [Alay] // *nigripes* Mor. Typ., [N]380 [handwritten by F. Morawitz]; 1 ♂, 22.[VII.1871] // Алай [Alay] // к. [оллекция] Ф. Моравицца [Collection of F. Morawitz] // *Halictus nigripes* Mor. [handwritten by F. Morawitz] // Paralectotypus *Halictus nigripes* Mor., design. Blüthgen <identical red labels on each paralectotype specimen, labelled by Yu. Astafurova> [ZISP].

Current status. *Lasioglossum (Hemihalictus) melanopus* (Dalla Torre, 1896), replacement name for *Halictus nigripes* Morawitz, 1876 (nec *H. nigripes* Lepeletier, 1841).

Remarks. The specimens from Iskander and Iori George in the Morawitz type series are the holotype and paratypes of *Halictus pseudonigripes* Blüthgen, 1934.

Description of female: Vachal 1902: 229, as *Halictus attritus* (synonymised by Blüthgen 1934a: 301).

Distribution. Uzbekistan, Kyrgyzstan, Kazakhstan, Tajikistan, Afghanistan, China (Xinjiang) (Morawitz 1876, Murao et al. 2017, Ascher and Pickering 2020).

27. *Halictus obscuratus* Morawitz, 1876

Figure 22

Halictus obscuratus Morawitz, 1876: 218 (key), 245, ♀.




Figure 21. *Halictus nigripes* Morawitz, 1876, lectotype, male **A** habitus, lateral view **B** metasoma, dorsal view **C** head, frontal view **D** mesosoma, dorsal view **E** labels. Scale bars: 1.0 mm (**A**), 0.5 mm (**B-D**).




Figure 22. *Halictus obscuratus* Morawitz, 1876, lectotype, female **A** habitus, lateral view and labels **B** metasoma, dorsal view **C** mesosoma, dorsal view **D** head, frontal view **E**. Scale bars: 1.0 mm.

Type locality. Sangy-dzhuman Pass, 30 km SSE Samarkand [Uzbekistan].

Published (original) locality. Uzbekistan: Samarkand, Aksay; Tajikistan: Iori Gorge, Varzaminor [=Aynil], Sangy-Dzhuman Pass.

Lectotype. ♀, designated by Warncke 1982: 116, 25.[V.1869] // Санги Джуманъ [Uzbekistan, Sangy-dzhuman Pass, 30 km SSE Samarkand, 39°20'N, 67°19'E] // *Halictus obscuratus* Mor., [N]370 [handwritten by F. Morawitz] // Lectotypus *Halictus obscuratus* Mor., design. Warncke [19]82 <red label, labelled by Yu. Astafurova> [ZMMU].

Paralectotypes (7 ♀). 1 ♀, 3.[IV.1869] // Самаркандъ [Samarkand] // [N]370; 1 ♀, 27.[II.1869] // Самаркандъ [Samarkand] // [N]370 // *obscuratus* [handwritten by F. Morawitz]; 1 ♀, [7.VI.1870] // Варзаминаръ [Varzaminor] // [N]370; 1 ♀, 16.[V.1869] // Заравшан.[сказа] дол.[иная] [Zeravshan River valley, Aksay] // N[370]; 1 ♀, 2.[VI.1869] // Заравшан.[сказа] дол.[иная] [Zeravshan River valley, Iori Gorge] // N[370]; [ZMMU]; 1 ♀, <golden circle> // 25.[V.1869] // Санги Джуманъ [Sangy Dzhuman] // *obscuratus* Mor., Typ. [handwritten by F. Morawitz]; 1 ♀, Санги Джуманъ [Sangy Dzhuman] // к.[оллекция] Ф. Моравица [Collection of F. Morawitz] // *obscuratus* Mor. [handwritten by F. Morawitz]; 1 ♀, Заравшан.[сказа] дол.[иная] [Zeravshan River valley] // к.[оллекция] Ф. Моравица [Collection of F. Morawitz] // *Halictus obscuratus* F. Morawitz, ♀ [handwritten by F. Morawitz] // Paralectotypus *Halictus obscuratus* Mor., design. Warncke <identical red labels on each paralectotype specimen, labelled by Yu. Astafurova> [ZISP].

Current status. *Lasioglossum (Sphecodogastra) obscuratum* ssp. *obscuratum* (Morawitz, 1876).

Remarks. Description of male. Blüthgen 1923a: 277.

Distribution. Europe (except North), Cyprus, Azerbaijan, Russia (North Caucasus), Turkey, Syria, Jordan, Israel, Iran, Afghanistan, Central Asia, Kazakhstan (Astafurova and Proshchalykin 2017).

28. *Halictus palustris* Morawitz, 1876

Figure 23

Halictus palustris Morawitz, 1876: 217 (key), 234, ♀.

Type locality. Iskanderkul Lake (Tajikistan).

Published (original) locality. Tajikistan: “near Iskander-Kul Lake”.

Lectotype. ♀, designated by Warncke 1982: 147, <golden circle> // 15.[VI.1870] // Искандеръ [Tajikistan, Iskanderkul Lake, Hissar Ridge, 39°04'N, 68°22'E] // *Halictus palustris* Mor., [N]353 [handwritten by F. Morawitz] // Lectotypus *Halictus palustris* Mor., ♀, design. Warncke [1]982 <red label>, labelled by Yu. Pesenko [ZMMU].

Paralectotype. 1 ♀, <golden circle> // 15.[VI.1870] // Искандеръ [Iskander] // *palustris* Mor., Typ. [handwritten by F. Morawitz] // Paralectotypus *H. palustris* Mor., ♀, design. Pesenko [1]981 <red label> [ZISP].

Current status. *Halictus (Tytthhalictus) palustris* Morawitz, 1876.

Remarks. Description of male. Blüthgen 1936: 291.

Distribution. Kazakhstan, Uzbekistan, Tajikistan, Kyrgyzstan, China (Xinjiang) (Pesenko 1986b, Murao et al. 2017).




Figure 23. *Halictus palustris* Morawitz, 1876, lectotype, female **A** habitus, lateral view and labels **B** head, frontal view **C** mesosoma, dorsal view **D** metasoma, dorsal view. Scale bars: 1.0 mm.

29. *Halictus pectoralis* Morawitz, 1876

Figure 24

Halictus pectoralis Morawitz, 1876: 218 (key), 251, ♀.

Type locality. Gulcha (Kyrgyzstan).

Published (original) locality. Kyrgyzstan: Gulsha [Gulcha].

Holotype. ♀, 10.[VIII.1871] // Гульша [Kyrgyzstan, Gulcha, 40°19'N, 73°26'E] // *Halictus pectoralis* Mor., [N]381 [handwritten by F. Morawitz] // Holotype *H. pectoralis* Mor., 1876 <red label, labelled by Yu. Pesenko> [ZMMU].

Current status. *Lasioglossum (Hemihalictus) subaenescens* ssp. *asiaticus* (Dalla Torre, 1896), replacement name for *Halictus pectoralis* Morawitz, 1876 (nec *H. pectoralis* Smith, 1853) (subspecies status according to Ebmer 1997: 932).

Remarks. Description of male. Blüthgen 1923a: 271, as *Halictus proximus* (synonymised by Warncke 1975: 96).

The lectotype designation by Warncke (1982: 106) is unnecessary as the species was described from a single female that was directly written about by Morawitz (1876: 251).

Distribution. Egypt, Turkey, Near East, Iran, Azerbaijan, Central Asia, Mongolia (Hovd), China (Xinjiang) (Ebmer 1997, Pesenko 2007).

30. *Halictus picipes* Morawitz, 1876

Figure 25

Halictus picipes Morawitz, 1876: 218 (key), 244, ♀.

Type locality. Zeravshan River valley (Tajikistan).

Published (original) locality. between Panjakent and Iori (Tajikistan).

Lectotype. ♀, designated by Pesenko 1986a: 138, 30.[V.1869] // Заравшан.[анская] дол.[и на] [Tajikistan, Zeravshan River valley, between Panjakent and Iori (= Yeri)] // *Halictus picipes* Mor., [N]369 [handwritten by F. Morawitz] // Lectotype *Halictus picipes* Mor., ♀, design. Pesenko [1]985 <red label> [ZMMU].

Paralectotypes (3 ♀). 1 ♀, the same label as in the lectotype [ZMMU]; 1 ♀, <golden circle>, the same label as in the lectotype; 1 ♀, Заравшан.[скай] дол.[и на] [Zeravshan River valley] // к.[оллекция] Ф. Моравица [Collection of F. Morawitz] // *Halictus picipes* Mor., ♀ [handwritten by F. Morawitz] // Paralectotypus *Hal. picipes* Mor., design. Pesenko [1]985 <identical red labels on each paralectotype specimen> [ZISP].

Current status. *Lasioglossum (Leuchalictus) picipes* (Morawitz, 1876).

Remarks. The lectotype designation by Warncke (1982: 111) is invalid because he labelled neither of the two females from “valle Serafshan” deposited in ZMMU.

Description of male. Blüthgen in Popov 1935: 362.

Distribution. Israel, Turkey, Iraq, Iran, Afghanistan, Turkmenistan, Uzbekistan, Tajikistan (Pesenko 1986a, Ascher and Pickering 2020).



Figure 24. *Halictus pectoralis* Morawitz, 1876, holotype, female **A** habitus, lateral view and labels **B** head, frontal view **C** mesosoma, dorsal view **D** metasoma, dorsal view. Scale bars: 1.0 mm (**A**), 0.5 mm (**B-D**).




Figure 25. *Halictus picipes* Morawitz, 1876, lectotype, female **A** habitus, lateral view and labels **B** head, frontal view **C** mesosoma, dorsal view **D** metasoma, dorsal view. Scale bars: 1.0 mm (**A, C, D**), 0.5 mm (**B**).

31. *Halictus rhynchites* Morawitz, 1876

Figure 26

Halictus rhynchites Morawitz, 1876: 217 (key to females), 220 (key to males), 222, ♀, ♂.

Type locality. Shakhimardan (Uzbekistan).

Published (original) locality. Usbekistan: Khodzha-Chiburgan gorge, near Shakhimardan Usbekistan]; Kyrgyzstan: Alay, Kichi-alay.

Lectotype (designated here). ♂, 7.[VII.1871] // Шагимарданъ [Shakhimardan in the Uzbek enclave in the territory of Kyrgyzstan, Alai Ridge, 39°58'N, 71°47'E] // [N]334 // Lectotypus *Halictus rhynchites* Mor., design. Astafurova et Proshchalykin, 2020 <red label> [ZMMU].

Paralectotypes (8 ♀, 5 ♂). 1 ♂, the same label as in a lectotype // *Halictus rhynchites* Mor., [N]334 [handwritten by F. Morawitz]; 1 ♀, 2 ♂, 26.[VI.1871] // Чибурганъ [Chiburgan] // [N]334; 1 ♀, 21.[VI.1871] // Чибурганъ [Chiburgan] // [N]334; 2 ♀, 28.[VII.1871] // Кичи-Алай [Kichi-Alay] // [N]334 [ZMMU]; 2 ♀, 1 ♂, Чибурганъ [Chiburgan] // к. [оллекция] Ф. Моравица [Collection of F. Morawitz] // *rhynchites* F. Mor. [handwritten by F. Morawitz]; 1 ♂, Кчи-Алай [Kchi-Alay] // к. [оллекция] Ф. Моравица [Collection of F. Morawitz] // *rhynchites* F. Mor. [handwritten by F. Morawitz]; 1 ♀, < golden circle > // 22.[VII.1871] // Алай [Alay] // *rhynchites* Mor., Typ. [handwritten by F. Morawitz]; 1 ♀, 22.[VII.1871] // Алай [Alay] // к. [оллекция] Ф. Моравица [Collection of F. Morawitz] // *Halictus rhynchites* Mor. [handwritten by F. Morawitz] // Paralectotypus *Halictus rhynchites* Mor., design. Astafurova et Proshchalykin, 2020 <identical red labels on each paralectotype specimen> [ZISP].

Current status. *Lasioglossum (Sphecodogastra) rhynchites* (Morawitz, 1876).

Remarks. The lectotype designation by Warncke (1982: 81) is invalid because he labelled neither of the two females from "Shakhimardan" deposited in ZMMU.

Distribution. Turkey, Afghanistan, southern Kazakhstan, Turkmenistan, Uzbekistan, Kyrgyzstan (Ebmer 1995, Murao et al. 2017).

32. *Halictus scutellaris* Morawitz, 1876

Figure 27

Halictus scutellaris Morawitz, 1876: 218 (key), 238, ♀.

Type locality. Bairkum (Chimkent Province, Kazakhstan).

Published (original) locality. Kazakhstan: Bayrakum [= Bairkum]; Tajikistan: Pendzhikent, Iori.

Lectotype. ♀, designated by Pesenko 1986a: 140, 4.[V.1871] // Байракумъ [Kazakhstan, Chimkent Province, Bairkum, Syr-Darya River, 42°05'N, 68°10'E] //




Figure 26. *Halictus rhynchites* Morawitz, 1876, lectotype, male **A** habitus, lateral view and labels **B** metasoma, dorsal view **C** mesosoma, dorsal view **D** antenna. Scale bars: 1.0 mm (**A**), 0.5 mm (**B–D**).




Figure 27. *Halictus scutellaris* Morawitz, 1876, lectotype, female **A** habitus, lateral view and labels **B** metasoma, dorsal **C** view head, frontal view **D** mesosoma, dorsal view. Scale bars: 1.0 mm.

Halictus scutellaris Mor., [N]359 [handwritten by F. Morawitz] // Lectotypus *Halictus scutellaris* Mor., ♀, design. Pesenko [1]985 <red label> [ZMMU].

Paralectotypes (5 ♀). 1 ♀, the same label as in the lectotype; 1 ♀, 30.[V.1869] // Заравшан.[ская] дол.[иная] [Zeravshan River valley, Iori] // N[359] [ZMMU]; 1 ♀, Байракумъ [Bayrakum] // к.[оллекция] Ф. Моравица [Collection of F. Morawitz] // *scutellaris* F. Mor., ♀ [handwritten by F. Morawitz]; 2 ♀, 4.[V.1871] // Байракумъ [Bayrakum] // к.[оллекция] Ф. Моравица [Collection of F. Morawitz] // Paralectotypus *Halictus scutellaris* Mor., design. Pesenko [1]985 <identical red labels on each paralectotype specimen> [ZISP].

Current status. *Lasioglossum (Leuchalictus) scutellare* (Morawitz, 1876).

Remarks. Description of male. Blüthgen 1929: 53, as *Halictus scutellaris*.

The lectotype designation by Warncke (1982: 111) is invalid because he labelled neither of the two females from “Bayrakum” deposited in ZMMU.

Distribution. Southern Kazakhstan, Turkmenistan, Tajikistan, Kyrgyzstan, China (Xinnjiang) (Pesenko 1986a, Murao et al. 2017).

33. *Halictus sogdianus* Morawitz, 1876

Figures 28

Halictus sogdianus Morawitz, 1876: 216 (key), 227, ♀.

Type locality. Samarkand (Uzbekistan).

Published (original) locality. Uzbekistan: Samarkand, Dshyuzak [=Jizzakh], Iskander [River]; Kyrgyzstan: Osh.

Lectotype. ♀, designated by Blüthgen 1934a: 303, 7.[VII.1870] // Самаркандъ [Uzbekistan, Samarkand, 39°39'N, 66°57'E] // *Halictus sogdianus* Mor., [N]342 [handwritten by F. Morawitz] // *sogdianus* Mor., ♀, Lecto-Holotype, Blüthgen det. 1931 // Lecto-Type <red label> // Lectotypus *Halictus sogdianus* Mor., design. Blüthgen 1934 <red label> [ZMMU].

Paralectotypes (5 ♀). 1 ♀, 21.[VI.1870] // Искандеръ [Iskander] // [N]342; 1 ♀, 2.[VIII. 1871] // Ошъ [Osh] // [N]342; 2 ♀, 4.[VII.1869] // Самаркандъ [Samarkand] // [N]342; 1 ♀, 18.[VII.1870] // Джюзакъ [Dzhyuzak] // [N]342 // Paralectotype *Halictus sogdianus* Mor. design. Blüthgen <identical red labels on each paralectotype specimen, labelled by Yu. Astafurova> [ZMMU].

Current status. *Halictus (Vestitohalictus) pulvereus* Morawitz, 1874 (synonymised by Ebmer 1988b: 576).

Distribution. Russia (European part, North Caucasus, Crimea), Cyprus, Turkey, Iran, Afghanistan, Central Asia, Mongolia, north-western China (Astafurova and Proshchalykin 2017).




Figure 28. *Halictus sogdianus* Morawitz, 1876, lectotype, female **A** habitus, lateral view and labels **B** head, frontal view **C** metasoma, dorsal view **D** mesosoma, dorsal view. Scale bars: 1.0 mm (**A**), 0.5 mm (**B-D**).

34. *Halictus trifasciatus* Morawitz, 1876

Figure 29

Halictus trifasciatus Morawitz, 1876: 218 (key), 240, ♀.

Type locality. Bairkum (Chimkent Province, Kazakhstan).

Published (original) locality. Kazakhstan: Bayrakum.

Lectotype. ♀, designated by Warncke 1982: 90, <golden circle> // 4.[V.1871] // Байракумъ [Kazakhstan, Chimkent Province, Bairkum, Syr-Darya River, 42°05'N, 68°10'E] // *Halictus trifasciatus* Mor., [N]362 [handwritten by F. Morawitz] // Lectotypus *Halictus trifasciatus* Mor., design. Warncke [19]82 <red label, labelled by Yu. Pesenko> [ZMMU].

Current status. *Lasioglossum (Lasioglossum) lebedevi* Ebmer, 1972, replacement name for *Halictus trifasciatus* Morawitz, 1876 (nec *Hylaeus trifasciatus* Schenck, 1853).

Remarks. Male unknown.

Distribution. Southern Kazakhstan (Morawitz 1876, Mutao et al. 2017). The record from Azerbaijan (Aliyev et al. 2007) is doubtful and needs checking.

35. *Halictus varipes* Morawitz, 1876

Figure 30

Halictus varipes Morawitz, 1876: 217 (key to females), 220 (key to males), 223, ♀, ♂.

Type locality. Jizzakh (Uzbekistan).

Published (original) locality. Uzbekistan: Katty-Kurgan [= Kattakurgan], Dzhyuzak [= Jizzakh], Karatyube [= Karatepa near Samarkand], Urgut, Sangy-Dzhuman; Kyrgyzstan: near Osh.

Lectotype. ♀, designated by Blüthgen 1955: 17, 19.[VII.1870] // Джюзакъ [Uzbekistan, Dzhyuzak (= Jizzakh), 40°07'N, 67°51'E] // [N]337 // *Halictus varipes* Mor., ♀, Lecto-Holotype, P. Blüthgen det. // Typus <red label> // Lectotypus *Halictus varipes* Mor., 1876, design. Blüthgen, 1955 <red label> [ZMMU].

Paralectotypes (1 ♀, 2 ♂). 1 ♂, the same labels as in the lectotype // *Halictus varipes* Mor. ♂, lecto-Paratype, Blüthgen det.; 1 ♂, the same label, but 14.[VII.1870] // *Halictus varipes* Mor. ♂, lecto-Holotype, Blüthgen det.; 1 ♀, 20. [VI.1869] // Катты-Курганъ [Katty-Kurgan] // *Halictus varipes* Mor. ♀, lecto-Paratype, Blüthgen det. // Paralectotype *Halictus varipes* Mor., design. Blüthgen 1955 <identical red labels on each paralectotype specimen, labelled by Yu. Astafurova> [ZMMU].

Current status. *Halictus (Seladonia) lucidipennis* Smith, 1853 (synonymised by Sakagami and Ebmer 1987: 326).

Distribution. Southern Palaearctic and Oriental Regions. North Africa, Israel, Arabian Peninsula, Turkey, Central Asia, Iraq, Iran, Afghanistan, Pakistan, Mongolia, north-western China, India, Nepal, Myanma, Thailand, Sri Lanka (Pesenko 2006a).




Figure 29. *Halictus trifasciatus* Morawitz, 1876, lectotype, female **A** habitus, lateral view **B** labels **C** metasoma, dorsal view **D** head, frontal view **E** mesosoma, dorsal view. Scale bars: 2.0 mm (**A**), 1.0 mm (**C-E**).



Figure 30. *Halictus varipes* Morawitz, 1876, lectotype, female **A** head, frontal view **B** habitus, lateral view and labels **C** metasoma, dorsal view **D** mesosoma, dorsal view. Scale bars: 1.0 mm (**B**), 0.5 mm (**A, C, D**).

36. *Halictus vulgaris* Morawitz, 1876

Figure 31

Halictus vulgaris Morawitz, 1876: 218 (key), 250, ♀.**Type locality.** Samarkand (Uzbekistan).**Published (original) locality.** Uzbekistan: Tashkent, Samarkand, Katty-Kurgan [Kattakurgan].**Lectotype (designated here).** ♀, 3.[III.1869] [Uzbekistan, Samarkand, 39°39'N, 66°57'E// *Hylaeus* [sic!] *vulgaris* Mor., [N]379 [handwritten by F. Morawitz] // Lectotypus *Halictus vulgaris* Mor., design. Astafurova et Proshchalykin, 2020 <red label> [ZMMU].**Paralectotypes** (257 ♀). 26 ♀, 4., 20., 23., 30.[III.1869], 3., 11., 19.[IV.1869] // Самаркандъ [Samarkand]; 5 ♀, 28.[IV.1869] // Каттыкурганъ [Kattykurgan]; 222 ♀, 10., 11., 24., 26., 27., 28., [II.1871], 23., 24. [III.1871], 1., 2., 3., 5., 8., 10., 11.[IV.1871] // Ташкентъ [Tashkent] [ZMMU]; 1 ♀, 3.[IV.1871] // Ташкентъ [Tashkent] // к. Ф. Моравица // *Halictus vulgaris* Mor. [handwritten by F. Morawitz]; 1 ♀, 5.[IV.1871] // Ташкентъ [Tashkent] // к. [оллекция] Ф. Моравица [Collection of F. Morawitz]; 2 ♀, 8.[IV.1871] // Ташкентъ [Tashkent] // к. [оллекция] Ф. Моравица [Collection of F. Morawitz]; 1 ♀, 1.[IV.1871] // Ташкентъ [Tashkent] // к. [оллекция] Ф. Моравица [Collection of F. Morawitz] // Paralectotypus, *Halictus vulgaris* Mor., design. Astafurova et Proshchalykin <identical red labels on each paralectotype specimen> [ZISP].**Current status.** *Lasioglossum (Eviulaeus) marginatum* (Brullé, 1832) (synonymised by Blüthgen 1926: 391).**Distribution.** Europe (except North), North Africa, Caucasus, Russia (East of European part, North Caucasus), Turkey, Syria, Jordan, Israel, Iraq, Iran, Afghanistan, Pakistan, Central Asia, Kazakhstan, north India, Nepal (Astafurova and Proshchalykin 2017).**Remarks.** The lectotype designation by Warncke (1982: 116) is invalid because he labelled none of the 222 females from Tashkent deposited in ZMMU.**Genus *Sphecodes* Latreille, 1804****37. *Sphecodes nigripennis* Morawitz, 1876**

Figure 32

Sphecodes nigripennis Morawitz, 1876: 257, ♀.**Type locality.** Shardara District of Turkistan Province (Kazakhstan).**Published (original) locality.** Kazakhstan: coasts of Kosaral Lake; Uzbekistan: Khodzhaduk [= Khozyay-Dun].**Lectotype.** ♀, designation by Warncke 1992: 30, 24.IV.1871 [the original blue data label was damaged] // Косараалъ [Kazakhstan, "Kosaral Lake", Shardara (= Chardara) District of Turkistan (= South-Kazakhstan) Province, \approx 41°10'N, 68°06'E] // *Sphecodes nigripennis* Mor. [handwritten by F. Morawitz] // Lectotypus, Warncke 1975 <red label> [ZMMU].




Figure 31. *Halictus vulgaris* Morawitz, 1876, lectotype, female **A** head, frontal view **B** habitus, lateral view and labels **C** metasoma, dorsal view **D** mesosoma, dorsal view. Scale bars: 1.0 mm (**B, C**), 0.5 mm (**A, D**).




Figure 32. *Sphecodes nigripennis* Morawitz, 1876, lectotype, female **A** habitus, lateral view and labels **B** head, frontal view **C** mesosoma, dorsal view **D** metasoma, dorsal view. Scale bars: 2.0 mm (**A, D**), 1.0 mm (**B, C**).

Paralectotype. 1 ♀, 21.[V.1869] // Заравшан.[сказа] дол.[иная] [Zeravshan River valley] // Paralectotype *Sphecodes nigripennis* Mor., design. Warncke <red label, labelled by Yu. Astafurova> [ZMMU].

Current status. *Sphecodes gibbus* (Linnaeus 1758) (synonymised by Blüthgen 1923b: 510).

Distribution. North Africa, Europe (north to 63°), Israel, Jordan, Russia (east to Yakutia), Turkey, Iran, Pakistan, Central Asia, Kazakhstan, Mongolia, NW China, India (Warncke 1992, Bogusch and Straka 2012, Astafurova et al. 2019).

38. *Sphecodes pectoralis* Morawitz, 1876

Figure 33

Sphecodes pectoralis Morawitz, 1876: 256, ♀.

Type locality. Shardara District of Turkistan Province (Kazakhstan).

Published (original) locality. Kazakhstan: coasts of Kosaral Lake; Kyzylkum [desert] near Chakany Well.

Lectotype. ♀, designation by Warncke 1992: 24, 24.[IV.1871] // Косаралъ [Kazakhstan, “Kosaral Lake”, Shardara (= Chardara) District of Turkistan (= South-Kazakhstan) Province], ≈ 41°10'N, 68°06'E// *Sphecodes pectoralis* Mor. [handwritten by F. Morawitz] // Lectotypus, Warncke 1975 <red label> [ZMMU].

Paralectotype. 1 ♀, 28.[VI.1871] // Кызылъкумъ [Kyzylkum] // Paralectotype *Sphecodes pectoralis* Mor., design. Warncke <red label, labelled by Yu. Astafurova> [ZMMU].

Current status. *Sphecodes pectoralis* Morawitz, 1876.

Remarks. Description of male. Meyer 1919: 126, as *Sphecodes cristatus* sensu Meyer (non Hagens 1882) (see Blüthgen 1924: 475).

Distribution. South Kazakhstan, Central Asia, China (Gansu, Xinjiang) (Astafurova et al. 2018a, b, 2020).

39. *Sphecodes rufithorax* Morawitz, 1876

Figure 34

Sphecodes rufithorax Morawitz, 1876: 255, ♀, ♂.

Type locality. Bairkum (Chimkent Province, Kazakhstan).

Published (original) locality. Kazakhstan: Bayrakum [Bairkum]; steppe between Syr-Darya River and Tashkent.

Lectotype. ♀, designated by Warncke 1992: 24, 17. [V.1871] // Байракумъ [Kazakhstan, Chimkent Province, Bairkum, Syr-Darya River, 42°05'N, 68°10'E] // *Sphe-*




Figure 33. *Sphecodes pectoralis* Morawitz, 1876, lectotype, female **A** head, frontal view **B** habitus, lateral view and labels **C** mesosoma, dorsal view **D** metasoma, dorsal view. Scale bars: 2.0 mm (**B, D**), 1.0 mm (**A, C**).



Figure 34. *Sphecodes rufithorax* Morawitz, 1876, lectotype, female **A** habitus, lateral view **B** head, frontal view **C** metasoma, dorsal view **D** vertex, dorso-lateral view **E** labels **F** mesosoma, dorsal view. Scale bars: 1.0 mm (**A–C, F**), 0.5 mm (**D**).

codes rufithorax F. Moraw., ♀ [handwritten by F. Morawitz] // Lectotypus Warncke, 1975 <red label> [ZMMU].

Paralectotypes (2 ♀, 1 ♂). 1 ♀, Байракумъ [Bairkum] // кол.[лекция] Ф. Моравица [Collection of F. Morawitz] // *Sphecodes rufithorax* F. Moraw., ♀ [handwritten by F. Morawitz] // F. Morawitz det., Typ.; 1 ♀, 20.[V.1871] // Степь м.[ежду] С.[ыр] А.[арьей] и Т.[ашкентом] [Steppe between Syrdarya River and Tashkent] // *Sphecodes rufithorax* Mor., ♂ [handwritten by F. Morawitz] // Paralectotypus *Sphecodes rufithorax* Mor., design. Warncke, [19]92 <red label, labelled by Yu. Astafurova> [ZMMU]; 1 ♀, the same labels [ZISP].

Current status. *Sphecodes olivieri* Lepeletier, 1825 (synonymised by Warncke 1992: 24).

Distribution. North Africa, the Arabian Peninsula, Israel, Jordan, South Europe, Russia (south of European part), Turkey, Caucasus, Iran, Pakistan, Central Asia, Kazakhstan, NW China (Astafurova et al. 2019).

Subfamily Nomiinae

Genus *Nomia* Latreille, 1804

40. *Nomia edentata* Morawitz, 1876

Figure 35

Nomia edentata Morawitz, 1876: 259, ♀, ♂.

Type locality. Jizzakh (Uzbekistan).

Published (original) locality. Uzbekistan: Samarkand, Dzhyuzak.

Lectotype. ♀, designated by Warncke 1976: 104, 20.[VII.1871] // Джозакъ [Uzbekistan, Dzhyuzak (= Jizzakh), 40°07'N, 67°51'E] // *Nomia edentata* Mor. [handwritten by F. Morawitz] // Lectotypus, Warncke 1975 <red label> [ZMMU].

Paralectotypes (2 ♂). 1 ♂, the same labels as in the lectotype // Paralectotypus, *Nomia edentata* Mor., design. Warncke <red label> [ZISP]; 1 ♂, 8.[VII.1869] // Самаркандъ [Samarkand] // Paralectotypus *Nomia edentata* Mor., design. Warncke <red label> [ZMMU].

Current status. *Pseudapis edentata* (Morawitz, 1876).

Distribution. North Africa, Saudi Arabia, Turkey, Azerbaijan, Kazakhstan, Central Asia, Iraq, Iran, Afghanistan, Pakistan, India (Astafurova 2014).

41. *Nomia rufescens* Morawitz, 1876

Figure 36

Nomia rufescens Morawitz, 1876: 261, ♀.

Type locality. Zeravshan River valley (Uzbekistan).

Published (original) locality. Uzbekistan: “Aykul Lake” in Zeravshan River valley.



Figure 35. *Nomia edentata* Morawitz, 1876, lectotype, female **A** habitus, lateral view **B** head, frontal view **C** metapostnotum, dorsal view **D** metasoma, dorsal view **E** mesosoma, dorsal view **F** labels. Scale bars: 1.0 mm (**A, D**), 0.5 mm (**B, C, E**).

Lectotype. ♀, designated by Warncke 1976: 106, 5. [VIII.1869] // Заравшанская].
дол.[иная] [Uzbekistan, Zeravshan River valley, Aykul Lake near Chelek, 39°56'N,
66°49'E] // *Nomia rufescens* Mor. // Lectotypus, Warncke 1975 <red label> [ZMMU].

Current status. *Pseudapis rufescens* (Morawitz, 1876).

Remarks. Description of male. Morawitz 1893: 79.

Distribution. Turkey, Central Asia, Kazakhstan (Astafurova 2014).




Figure 36. *Nomia rufescens* Morawitz, 1876, lectotype, female **A** habitus, lateral view and labels **B** head, frontal view **C** mesosoma, dorsal view **D** metasoma, dorsal view. Scale bars: 1.0 mm.

Subfamily Nomioinae

Genus *Nomioides* Schenck, 1866

42. *Nomioides parviceps* Morawitz, 1876

Figure (see Astafurova and Proshchalykin 2019: 56, figs 51a–e)

Nomioides parviceps Morawitz, 1876: 215, ♂.

Type locality. Bairamali (Turkmenistan).

Published (original) locality. Uzbekistan: Samarkand.

Holotype. ♂, окр.[естности] Самарканда [Uzbekistan, Samarkand, 39°39'N, 66°57'E], 13.VI[1869], lost (also see Pesenko 1983: 168–170).

Neotype. ♂, designated by Pesenko 1983: 168, Байрам-Али Закасп.[ийская] обл. [асты] [Turkmenistan, Maryi Province, Bairamali, 37°37'N, 62°09'E], 14.VII.[1]928, B. Гуссаковский [V. Gussakovskij leg.] // *Nomioides conjungens* m., ♂, Blüthgen det. 1931 // Neotypus *Nomioides parviceps* Mor., ♂, design. Pesenko [1]980 [handwritten by Yu. Pesenko] <red label> // Zoological Institute St. Petersburg, INS_HYM_0000142 [ZISP].

Current status. *Nomioides (Nomioides) parviceps* Morawitz, 1876.

Remarks. Description of female: Blüthgen 1925b: 45, as *Nomioides conjungens* (synonymised by Blüthgen 1934c: 253).

Distribution. Asia Minor, Afghanistan, Armenia, Central Asia (Astafurova and Proshchalykin 2019).

43. *Nomioides turanica* Morawitz, 1876

Figure (see Astafurova and Proshchalykin 2019: 68, figs 62a–e).

Nomioides turanica Morawitz, 1876: 214, ♀, ♂.

Type locality. Samarkand (Uzbekistan).

Published (original) locality. Tajikistan: Murzarabat; Uzbekistan: Sokh, Samarkand.

Lectotype. ♂, designated by Pesenko 1983: 174, 5.[VII.1870] // Самарканда [Uzbekistan, Samarkand, 39°39'N, 66°57'E] // *turanica* Mor., Typ. [handwritten by F. Morawitz] // Lectotypus *Nom. turanica* Mor., ♂, design. Pesenko, 1976 // Zoological Institute St. Petersburg, INS_HYM_0000131 [ZISP].

Paralectotypes (5 ♂). 1 ♂, 28.[VI.1871] // Сохъ [Sokh] // *Nomioides turanica* n. sp. F. Morawitz det.; 1 ♂, <gold circle> 8.[VII.1870] // Самарканда [Samarkand] // *Nomioides turanica* Mor. [handwritten by F. Morawitz] [ZMMU]; 3 ♂, Сохъ [Sokh] // к. [оллекция] Ф. Моравица [Collection of F. Morawitz] // *Nomioides turanica* Mor. // Paralectotypus ♂ *Nom. turanica* Mor., design. Pesenko, 1976 <identical red labels on each paralectotype specimen> [ZISP].

Current status. *Nomioides (Nomioides) turanicus* Morawitz, 1876.

Distribution. North Africa, Central Asia, Iran, Pakistan (Astafurova and Proshchalykin 2019).

Acknowledgements

We are grateful to A.V. Antropov (ZMMU) for assisting during work in the ZMMU and Michael Orr (Beijing, China) for checking the English grammar. We also thank P. Bogusch (Hradec Králové, Czech Republic) and R. Murao (Fukuoka, Japan) for careful review of the manuscript and several comments improved the text.

This investigation was supported by the Russian Funds for Basic Research (grant numbers 19-04-00027 and 20-54-44014), the Presidium RAS programme no.41 “Biodiversity of natural systems and biological sources of Russia” and the state research project AAAA-A17-117030310210-3.

References

- Aliyev KhA, Guseinzade GA, Magerramov MM (2007) Towards the knowledge of the bee fauna of the family Halictidae (Hymenoptera: Apoidea) of Nakhichevan Autonomous Republic of Azerbaijan. Caucasian Entomological Bulletin 3(2): 251–256. <https://doi.org/10.23885/1814-3326-2007-3-2-251-256>
- Ascher JS, Pickering J (2020) Discover Life bee species guide and world checklist (Hymenoptera: Apoidea: Anthophila). http://www.discoverlife.org/mp/20q?guide=Apoidea_species [accessed 20 August 2020]
- Astafurova YuV (2014) Bees of the subfamilies Rophitinae and Nomiinae (Hymenoptera, Halictidae) of the Russia and adjacent territories. KMK Scientific Press Ltd., Moscow, 383 pp. [In Russian]
- Astafurova YuV, Proshchalykin MYu (2017) Family Halictidae. In: Lelej AS, Proshchalykin MYu, Loktionov VM (Eds) Annotated Catalogue of the Hymenoptera of Russia. Volume I. Symphyta and Apocrita: Aculeata. Proceedings of the Zoological Institute RAS, Supplement 6: 277–292.
- Astafurova YuV, Proshchalykin MYu (2018) The type specimens of bees (Hymenoptera, Apoidea) deposited in the Zoological Institute of the Russian Academy of Sciences, St. Petersburg. Contribution I. Family Halictidae, genus *Lasioglossum* Curtis, 1833. Zootaxa 4408(1): 1–66. <https://doi.org/10.11646/zootaxa.4408.1.1>
- Astafurova YuV, Proshchalykin MYu (2019) The type specimens of bees (Hymenoptera, Apoidea) deposited in the Zoological Institute of the Russian Academy of Sciences, St. Petersburg. Contribution II. Family Halictidae, subfamilies Rophitinae, Nomiinae, and Nomioiidae. Zootaxa 4650(1): 1–71. <https://doi.org/10.11646/zootaxa.4650.1.1>
- Astafurova YuV, Proshchalykin MYu (2020) The type specimens of bees (Hymenoptera, Apoidea) deposited in the Zoological Institute of the Russian Academy of Sciences, St. Petersburg. Contribution III. Family Halictidae, genera *Halictus* Latreille, 1804, and *Sphecodes* Latreille, 1804. Zootaxa 4790(3): 401–446. <https://doi.org/10.11646/zootaxa.4790.3.1>
- Astafurova YuV, Proshchalykin MYu, Engel MS (2018a) The cuckoo bee genus *Sphecodes* Latreille, 1804 in Kazakhstan (Hymenoptera: Halictidae). Far Eastern Entomologist 369: 1–47. <https://doi.org/10.3897/jhr.66.29269>

- Astafurova YuV, Proshchalykin MYu, Niu Z-Q, Orr MC, Zhu C-D (2020) New and little-known bees of the genus *Sphecodes* Latreille, 1804 (Hymenoptera, Apoidea, Halictidae) from Southern and South-Western China. *Journal of Hymenoptera Research* 79: 145–162. <https://doi.org/10.3897/jhr.79.57276>
- Astafurova YuV, Proshchalykin MYu, Niu Z-Q, Zhu C-D (2018b) New records of bees of the genus *Sphecodes* Latreille (Hymenoptera, Halictidae) in the Palaearctic part of China. *ZooKeys* 792: 15–44. <https://doi.org/10.3897/zookeys.792.28042>
- Astafurova YuV, Proshchalykin MYu, Schwarz M (2019) The distribution of the genus *Sphecodes* Latreille (Hymenoptera, Halictidae) of the Arabian Peninsula and surrounding countries with description of hitherto unknown female of *S. atlanticus* Warncke, 1992 and male of *S. dathei* Schwarz, 2010. *ZooKeys* 872: 13–40. <https://doi.org/10.3897/zookeys.872.35361>
- Baker DB (2004) Type material of Hymenoptera described by O.I. Radoszkowsky in the Natural History Museum, London, and the localities of A.P. Fedtschenko's Reise in Turkestan. *Deutsche Entomologische Zeitschrift* 51(2): 231–252. <https://doi.org/10.1002/mmnd.4810510207>
- Blüthgen P (1922) Beiträge zur Synonymie der Bienengattung *Halictus* Latr. *Deutsche Entomologische Zeitschrift* 1922(1): 46–66. <https://doi.org/10.1002/mmnd.48019220105>
- Blüthgen P (1923a) Beiträge zur Kenntnis der Bienengattung *Halictus* Latr. *Archiv für Naturgeschichte. Abteilung A* 89(5): 232–332.
- Blüthgen P (1923b) Beiträge zur Systematik der Bienengattung *Sphecodes* Latr. *Deutsche Entomologische Zeitschrift* 1923: 441–513.
- Blüthgen P (1924) Beiträge zur Systematik der Bienengattung *Sphecodes* Latr. II. *Deutsche Entomologische Zeitschrift* 1924: 457–516. <https://doi.org/10.1002/mmnd.48019240601>
- Blüthgen P (1925a) Beiträge zur Kenntnis der Bienengattung *Halictus* LATR. II. *Archiv für Naturgeschichte. Abteilung A* (1924) 90(10): 86–136.
- Blüthgen P (1925b) Die Bienengattung *Nomioides* Schenck. *Stettiner entomologische Zeitung* 85(1): 1–100.
- Blüthgen P (1926) Beiträge zur Synonymie der Bienengattung *Halictus* Latr. IV. *Deutsche Entomologische Zeitschrift* 1925(5): 385–419. <https://doi.org/10.1002/mmnd.192619250506>
- Blüthgen P (1929) Neue turkestanische *Halictus*-Arten (Hym., Apidae). *Konowia* 8(1): 51–86.
- Blüthgen P (1930) *Halictus* Latr. In: Schmiedeknecht O (Ed.) *Die Hymenopteren Nord- und itteleuropas*. G. Fischer, Jena, 729–767.
- Blüthgen P (1931a) Beiträge zur Synonymie der Bienengattung *Halictus* Latr. VII. (Hym., Apid.). *Deutsche Entomologische Zeitschrift* 1930(4): 209–215.
- Blüthgen P (1931b) Beiträge zur Kenntnis der Bienengattung *Halictus* Latr. III. Mitteilungen aus dem Zoologischen Museum in Berlin 17(3): 319–398.
- Blüthgen P (1934a) Beiträge zur Synonymie der Bienengattung *Halictus* Latr. IX. *Deutsche Entomologische Zeitschrift* 1933(2–3): 299–304. <https://doi.org/10.1002/mmnd.48019330206>
- Blüthgen P (1934b) Neue turkestanische *Halictus*-Arten. II. (Hym. Apidae). *Konowia* 13(3): 145–159.
- Blüthgen P (1934c) 1. Nachtrag zur Monographie der Bienengattung *Nomioides* Schck. (Hym., Apidae, Halictinae). *Stettiner entomologische Zeitung* 95(2): 238–283.
- Blüthgen P (1936) Neue paläarktische Binden-*Halictus* (Hym. Apidae). Mitteilungen aus dem Zoologischen Museum in Berlin 21(2): 270–313.

- Blüthgen P (1955) The Halictinae (Hymen., Apoidea) of Israel. I. Genus *Halictus* (subgenera *Halictus* s. str. and *Thrincohalictus*). Bulletin of the Research Council of Israel (B), 5(1): 5–23.
- Bogusch P, Straka J (2012) Review and identification of the cuckoo bees of central Europe (Hymenoptera: Halictidae: *Sphecodes*). Zootaxa 3311: 1–41. <https://doi.org/10.11646/zootaxa.3311.1.1>
- Bytinski-Salz H, Ebmer AW (1974) The Halictidae of Israel (Hymenoptera, Apoidea). II. Genus *Lasioglossum*. Israel Journal of Entomology 9: 175–217.
- Dalla Torre CG de (1896) Catalogus Hymenopterorum, hucusque descriptorum systematicus et synonymicus. Vol. X. Apidae (Anthophila). Engelmann, Lipsiae, 643 pp.
- Dathe HH, Proshchalykin MYu (2017) Type revision of Asiatic bees of the genus *Hylaeus* F. described by Ferdinand Morawitz (Hymenoptera: Apoidea, Colletidae). Zootaxa 4227(1): 1–48. <https://doi.org/10.11646/zootaxa.4227.1.1>
- Ebmer AW (1974) Beiträge zur Kenntnis der Fauna Afghanistans. *Halictus* Latr. et *Lasioglossum* Curt., Halictidae, Apoidea, Hymenoptera. Časopis Moravského Muzea 59: 183–210.
- Ebmer AW (1978) *Halictus*, *Lasioglossum*, *Rophites* und *Systropha* aus dem Iran (Halictidae, Apoidea) sowie neue Arten aus der Paläarktis. Linzer Biologische Beiträge 10(1): 1–109.
- Ebmer AW (1980) Asiatische Halictidae (Apoidea, Hymenoptera). Linzer Biologische Beiträge 12(2): 469–506.
- Ebmer AW (1984) Asiatische Halictidae, 2 (Apoidea, Hymenoptera). Annales Historico Naturales Musei Nationalis Hungarici 75: 313–325.
- Ebmer AW (1985) *Halictus* und *Lasioglossum* aus Marokko (Hymenoptera, Apoidea, Halictidae). Erster Nachtrag. Linzer Biologische Beiträge 17(2): 271–293.
- Ebmer AW (1986) Die Artgruppe des *Lasioglossum strictifrons* (Vachal 1895) mit einer Bestimmungstabelle der Weibchen (Hymenoptera, Apoidea, Halictidae). Linzer Biologische Beiträge 18(2): 417–443.
- Ebmer AW (1988a) Die europäischen Arten der Gattungen *Halictus* Latreille 1804 und *Lasioglossum* Curtis 1833 mit illustrierten Bestimmungstabellen (Insecta: Hymenoptera: Apoidea: Halictidae: Halictinae). 2 – Die Untergattung *Seladonia* Robertson 1918. Senckenbergiana Biologica 68(4/6): 323–375.
- Ebmer AW (1988b) Kritische Liste der nicht-parasitischen Halictidae Österreichs mit Berücksichtigung aller mitteleuropäischen Arten (Insecta: Hymenoptera: Apoidea: Halictidae). Linzer Biologische Beiträge 20(2): 527–711.
- Ebmer AW (1995) Asiatische Halictidae – 3. Die Artengruppe der *Lasioglossum carinate-Evylaeus* (Insecta: Hymenoptera: Apoidea: Halictidae: Halictinae). Linzer Biologische Beiträge 27(2): 525–652.
- Ebmer AW (1997) Asiatische Halictidae – 6. *Lasioglossum carinaless-Evylaeus*: Ergänzungen zu den Artengruppen von *L. nitidiusculum* und *L. punctatissimum* s. l., sowie die Artengruppe des *L. marginellum* (Insecta: Hymenoptera: Apoidea: Halictidae: Halictinae). Linzer Biologische Beiträge 29(2): 921–982.
- Ebmer AW (1998) Asiatische Halictidae – 7. Neue *Lasioglossum*-Arten mit einer Übersicht der *Lasioglossum* s. str.-Arten der nepalischen und yunnanischen Subregion, sowie des nördlichen Zentral-China (Insecta: Hymenoptera: Apoidea: Halictidae: Halictinae). Linzer Biologische Beiträge 30(1): 365–430.

- Fedtschenko A (1871*) List of localities visited by the Turkestan Scientific Expedition of the Imperial Society of Friends of Natural History, 1869–1871. Karkov & Co., Moscow, 9 pp.
[*privately printed and not dated; In Russian]
- Fedtschenko OA (1874) A. Fedtschenko's Reisen in Turkestan, 1868–71. Mitteilungen aus Justus Perthes' Geographischer Anstalt über wichtige neue Erforschungen auf dem Gesammtgebiete der Geographie 20: 201–206, 11 pls.
- Kohl FF (1905) Ergebnisse einer naturwissenschaftlichen Reise zum Erdschias-Dagh (Kleinasiens) Ausgeführt von Dr. Arnold Penther und Dr. Emerich Zederbauer auf Kosten der "Gesellschaft zur Förderung der naturhistorischen Erforschung des Orients in Wien" (nunmehr "Naturwissenschaftlicher Orientverein in Wien") im Jahre 1902: Hymenopteren. Annalen des k.k. Naturhistorischen Hofmuseums (Wien) 20: 220–246.
- Kuhlmann M (2005) Faunistik und Zoogeographie der Bienengattung *Colletes* Latreille 1802 (Hymenoptera: Apidae: Colletinae) in Mittelasien. Linzer Biologische Beiträge 37(2): 1353–1396.
- Meade-Waldo G (1923) Hymenoptera, Fam. Apidae, Subfam. Prosopidinae. In: Wytsman P (Ed.) Genera Insectorum 181: 1–45.
- Meyer R (1919) Apidae – Sphecodinae. Archiv für Naturgeschichte, Abteilung A, 85(1): 79–160.
- Michener CD (2007) The Bees of the World (2nd edn). Johns Hopkins University Press, Baltimore, xvi + [i] + 953 pp. [+ 20 pls]
- Morawitz F (1875) A Travel to Turkestan by the Member-Founder of the Society A. P. Fedtschenko, accomplished from the Imperial Society of Naturalists, Anthropologists, and Ethnographers on a Commission from the General-Governor of Turkestan K. P. von Kaufmann (Issue 9). Vol. II. Zoogeographical Investigations. Pt. V. (Division 7). Bees (Mellifera). Pt. I [Apidae genuinae]. Izvestiya Imperatorskogo Obshchestva Lyubiteley Estestvoznanija, Anthropologii i Ethnografii 21(3): 1–160. [In Russian]
- Morawitz F (1876) A Travel to Turkestan by the Member-Founder of the Society A.P. Fedtschenko, accomplished from the Imperial Society of Naturalists, Anthropologists, and Ethnographers on a Commission from the General-Governor of Turkestan K.P. von Kaufmann (Issue 13). Vol. II. Zoogeographical Investigations. Pt. V. (Division 7). Bees (Mellifera). Pt. II [Andrenidae]. Izvestiya Imperatorskogo Obshchestva Lyubiteley Estestvoznanija, Anthropologii i Ethnografii 21(3): 161–303. [In Russian]
- Morawitz F (1893) Supplement zur Bienenfauna Turkestans. Horae Societatis Entomologicae Rossicae 28(1): 7–87.
- Morawitz F (1894) Beitrag zur Bienenfauna Turkmeniens. Horae Societatis Entomologicae Rossicae 29(1/2): 1–76.
- Murao R, Tadauchi O, Miyanaga R (2017) The bee family Halictidae (Hymenoptera, Apoidea) from Central Asia collected by the Kyushu and Shimane Universities Expeditions. Biodiversity Data Journal 5: e15050. <https://doi.org/10.3897/BDJ.5.e15050>
- Murzayev EM (1957) Middle Asia. An Outline of Physical Geography. Geografgis, Moscow, 272 pp. [In Russian]
- Niu Z-Q, Zhang D, Zhu C-D (2020) Extraordinary bees of the genus *Lasioglossum* Curtis, 1833 (Hymenoptera: Apoidea: Halictidae) from China. Zoological Systematics 45(1): 50–58.

- Pesenko YuA (1983) Family Halictidae. Subfamily Halictinae. Tribe Nomoidini. (In Amount of the Palaearctic Region). Fauna of the USSR (New Series, no. 129). Hymenopterous Insects. Vol. XVII, No. 1, Nauka, Leningrad, 199 pp. [In Russian]
- Pesenko YuA (1984) A synominal annotated catalogue of species-group names of bees of the genus *Halictus* Latreille sensu stricto (Hymenoptera, Halictidae) in the World fauna. Proceedings of the Zoological Institute, USSR Academy of Sciences 128: 16–32. [In Russian]
- Pesenko YuA (1986a) An annotated key to the Palaearctic species of bees of the genus *Lasioglossum* sensu stricto (Hymenoptera, Halictidae) for females, with descriptions of new subgenera and species. Proceedings of the Zoological Institute, USSR Academy of Sciences 159: 113–151. [In Russian]
- Pesenko YuA (1986b) Systematics of the bee genus *Halictus* Latreille (Hymenoptera, Halictidae) with description of 7th and 8th metasomal sterna of males: subgenus *Tytthalcicus* Pesenko. Entomologicheskoe Obozrenie 65(3): 618–632. [In Russian]
- Pesenko YuA (2005a) New data on the taxonomy and distribution of the Palaearctic halictids: genus *Halictus* Latreille (Hymenoptera: Halictidae). Entomofauna 26(18): 313–348.
- Pesenko YuA (2005b) Contributions to the halictid fauna of the Eastern Palaearctic region: genus *Halictus* Latreille (Hymenoptera: Halictidae, Halictinae). Far Eastern Entomologist 150: 1–24.
- Pesenko YuA (2006a) Contributions to the halictid fauna of the Eastern Palaearctic Region: genus *Seladonia* Robertson (Hymenoptera: Halictidae, Halictinae). Esakia 46: 53–82.
- Pesenko YuA (2006b) Contributions to the halictid fauna of the Eastern Palaearctic Region: genus *Lasioglossum* Curtis (Hymenoptera: Halictidae, Halictinae). Zoosystematica Rossica 15(1): 133–166.
- Pesenko YuA (2007) Subfamily Halictinae. In: Lelej AS (Ed.) Key to the insects of Russian Far East. Vol. 4. Pt 5. Neuropteroidea, Mecoptera, Hymenoptera. Dalnauka, Vladivostok, 823–878. [In Russian]
- Pesenko YuA, Astafurova YuV (2003) Annotated bibliography of Russian and Soviet publications on the bees (Hymenoptera: Apoidea; excluding *Apis mellifera*): 1771–2002. Denisia 11: 1–616.
- Popov VB (1935) Beiträge zur Bienenfauna von Tadzhikistan. Proceedings of the Tajik Station of the USSR Academy of Sciences 5: 360–367. [In Russian]
- Proshchalykin MYu, Dathe HH (2018) In the footsteps of history: the bees of the genus *Hyphalaeus* Fabricius (Hymenoptera, Apoidea: Colletidae) collected by V.I. Roborovsky and P.K. Kozlov in Northwest China (1895–1926). Zootaxa 4434(3): 573–588. <https://doi.org/10.11646/zootaxa.4434.3.11>
- Sakagami SF, AW Ebmer (1987) Taxonomic notes on oriental halictine bees of the genus *Halictus* (subgen. *Seladonia*) (Hymenoptera Apoidea). Linzer Biologische Beiträge 19(2): 301–357.
- Strand E (1921) Apidologisches, insbesondere über paläarktische *Halictus*-Arten, auf Grund von Material des Deutschen entomologischen Museums. Archiv für Naturgeschichte. Abteilung A 87(3): 266–322.
- Vachal J (1902) *Halictus* nouveaux ou litigieux de la collection Radoszkowski (Hymenoptera, Apidae). Revue russe d'entomologie 2: 225–231.

- Warncke K (1975) Beitrag zur Systematik und Verbreitung der Furchenbienen in der Türkei (Hymenoptera, Apoidea, *Halictus*). Polskie Pismo Entomologiczne 45(1): 81–128.
- Warncke K (1976) Zur Systematik und Verbreitung der Bienengattung *Nomia* Latr. in der Westpaläarktis und dem turkestanischen Becken (Hymenoptera, Apoidea). Reichenbachia 16(7): 93–120.
- Warncke K (1982) Beitrag zur Bienenfauna des Iran. 14. Die Gattung *Halictus* Latr., mit Bemerkungen über unbekannte und neue *Halictus*-Arten in der Westpaläarktis und Zentralasien. Bollettino del Museo Civico di Storia Naturale di Venezia 32: 67–166.
- Warncke K (1992) Die westpaläarktischen Arten der Bienengattung *Sphecodes* Latr. Bericht der Naturforschende Gesellschaft Augsburg 52: 9–64.