

First checklist of the chrysidid wasps (Hymenoptera, Chrysidae) of Mongolia, with description of new species

Paolo Rosa¹, Maxim Yu. Proshchalykin², Marek Halada³, Ulykpan Aibek⁴

1 Via Belvedere 8/d, I–20881 Bernareggio (MB), Italy **2** Federal Scientific Centre for East Asian Terrestrial Biodiversity, Far Eastern Branch of Russian Academy of Sciences, Vladivostok 690022, Russia **3** Milady Horákové 74 37012 České Budějovice, Czech Republic **4** National University of Mongolia, Ulaanbaatar 210646, Mongolia

Corresponding author: Maxim Yu. Proshchalykin (proshchalykin@biosoil.ru)

Academic editor: M. Engel | Received 10 September 2020 | Accepted 5 November 2020 | Published 30 November 2020

<http://zoobank.org/34E6CD7A-EAD1-46D4-926A-61683DFFC740>

Citation: Rosa P, Proshchalykin MYu, Halada M, Aibek U (2020) First checklist of the chrysidid wasps (Hymenoptera, Chrysidae) of Mongolia, with description of new species. ZooKeys 999: 49–107. <https://doi.org/10.3897/zookeys.999.58536>

Abstract

An annotated checklist of the Chrysidae from Mongolia is provided. A revision of the bibliographical data is provided, since most of the collecting localities published for “Mongolia” refer to places currently located in China. The known Mongolian cuckoo wasp fauna counts 90 species in 18 genera and two sub-families. Four genera and 57 species are recorded for the first time, including two species here described as new for science: *Cleptes mongolicus* Rosa, Halada & Agnoli, **sp. nov.** (Dornod) and *Spinolia spinosa* Rosa & Halada, **sp. nov.** (Bayankhongor).

Keywords

Catalogue, Central Asia, new records, Palaearctic region

Introduction

Mongolia is a large landlocked country in eastern Central Asia, covering 1,564,100 km². Politically, Mongolia is divided into 21 provinces named “aimags” with the capital Ulaanbaatar (Fig. 1). It is bordered by Russia to the north and China




Figure 1. Administrative map of Mongolia (from Dathe and Proshchalykin (2016) and Proshchalykin (2017) modified). Aimags: **1** Bayan-Ulgii **2** Uvs **3** Khovd **4** Zavkhan **5** Govi-Altai **6** Khuvsgul **7** Arkhangai **8** Bayankhongor **9** Bulgan **10** Orkhon **11** Uvurkhangai **12** Umnugovi **13** Selenge **14** Darkhan-Uul **15** Tuv **16** Ulaanbaatar **17** Dundgovi **18** Khentii **19** Govi-Sümber **20** Dornogovi **21** Dornod **22** Sukhaahtar.

to the south, east, and west. Geographically and climatologically, it is an area of contrasts and extremes, between cold mountainous regions up to 4,000 m a.s.l. to the north and west and one of the largest deserts of the world in the south, the Gobi Desert. Most of the country is located on high plateaus, covered by steppes and extensive forested areas. It has an extreme continental climate with long, cold winters and short hot summers, during which most of its annual precipitation falls (Lavrenko 1979; Dathe and Proshchalykin 2016).

Mongolian cuckoo wasps are scarcely known and a few occasional records are found in the literature (Rosa 2017a). Only one article (Móczár 1967) deals with Mongolian material collected by Dr Z. Kaszab during his entomological excursions in this country (1963–1968). Other scattered findings have been published (du Buysson 1901; Semenov-Tian-Shanskij 1912, 1932, 1967; Semenov-Tian-Shanskij and Nikol'skaya 1954; Linsenmaier 1997a; Rosa et al. 2017a, b), while most of the remaining bibliographical data recorded for “Mongolia” actually refer to localities currently included in China (Inner Mongolia, Xinjiang, Gansu) (du Buysson 1893; Radoszkowski 1877, 1891; Mocsáry 1890; Dalla Torre 1892; Bischoff 1913; Hammer 1936; Tsuneki 1947, 1953a; Linsenmaier 1959, 1968; Semenov-Tian-Shanskij 1967; Kimsey and Bohart 1991; Rosa et al. 2014, 2015). Approximately 30 species were properly recorded from Mongolia so far (Rosa 2017a) and we here add 57 new records for this country, mostly based on the materials collected by Czech entomologists (M. Halada, J. Halada, J. Straka, and M. Kadlecová) in 2003–2007 and mainly housed in the private collections of MH (České Budějovice, Czech Republic) and PR (Bernareggio, Italy). Other new records were found during the examination of the Chrysidae collection housed at the Zoological Institute in St. Petersburg (Russia,

ZIN) and based on the material collected during the expeditions of V. Roborovskij and P. Kozlov in 1895 and P. Kozlov in 1926. Finally, a few specimens were examined from the material collected in Mongolia by Soviet-Mongolian expeditions in 1967–1982. Soviet-Mongolian expedition were conducted from 1967 to 1983 and led to the collection of extensive entomological material, which became the basis for the publication of numerous articles and books (including *Insects of Mongolia* in eleven volumes), devoted to the study of various insects families (Proshchalykin and Kuhlmann 2015), although the Chrysididae were never examined by anyone. Large part of the cuckoo wasps collected during these entomological expeditions is still unprepared and unidentified.

Unpublished distributional records from Mongolia were recently published in the volume on Russian Chrysididae (Rosa et al. 2019), for a better understanding of the distribution of the Asian species, but exact localities were omitted because they were not of interest for that publication. We here report the precise data of species recorded for the first time in Rosa et al. 2019, which are mostly based on material housed in the Linsenmaier collection (Luzern, Switzerland).

In the present paper, based on a comprehensive study of specimens (including primary types) deposited in various collections, we report additional records of 72 species, with two species described as new and 55 species recorded from Mongolia for the first time, resulting in a total number of 90 cuckoo wasps species known from this country (Table 1).

Materials and methods

Terminology follows Lanes et al. (2020), Hymenoptera Anatomy Ontology (HAO 2020), and partly Kimsey and Bohart (1991). Abbreviations used in the descriptions are as follows:

F1, F2, F3, etc.	flagellomeres 1, 2, 3, etc., respectively;
l/w	length/width;
MOD	anterior ocellus diameter;
MS	malar space, the shortest distance between base of mandible and lower margin of compound eye;
OOL	the shortest distance between posterior ocellus and compound eye;
P	pedicel;
PD	puncture diameter;
POL	the shortest distance between posterior ocelli;
T1–T5	metasomal terga numbered consecutively, starting with 1 at the second abdominal segment.

Pictures of the types were taken with Nikon D700 connected to the microscope Togal SCZ and stacked with the software Combine ZP.

Table 1. Records of Mongolian cuckoo wasp species by aimags.

No.	Species	Aimags
1.	<i>Chrysis aestiva</i> Dahlbom, 1854	7
2.	<i>Chrysis angustula</i> Schenck, 1856	7, 15
3.	<i>Chrysis asahinai</i> Tsuneki, 1950	8, 9, 12, 15, 20, 22
4.	<i>Chrysis belokobylskiji</i> Rosa, 2019	4, 12, 15
5.	<i>Chrysis brevitarsis</i> Thomson, 1870	9
6.	<i>Chrysis castigata</i> Linsenmaier, 1959	13, 15
7.	<i>Chrysis chinensis</i> Mocsáry, 1912	7, 13, 15
8.	<i>Chrysis consanguinea</i> Mocsáry, 1889	4, 7, 9, 13, 15, 16, 18, 21, 22
9.	<i>Chrysis dauriana</i> Linsenmaier, 1959	4, 7–9, 13, 18
10.	<i>Chrysis equestris</i> Dahlbom, 1854	7, 13
11.	<i>Chrysis fulgida</i> Linnaeus, 1761	7, 13, 15
12.	<i>Chrysis ignita</i> (Linnaeus, 1758)	9
13.	<i>Chrysis illecebrosa</i> Semenov, 1967	12
14.	<i>Chrysis illigeri</i> Wesmael, 1839	13, 15
15.	<i>Chrysis ismaeli</i> Semenov, 1967	12, 20, 21
16.	<i>Chrysis jactartis</i> Semenov, 1910	12, 13, 15, 18, 21
17.	<i>Chrysis leptomandibularis</i> Niehuis, 2000	15
18.	<i>Chrysis mane</i> Semenov, 1912	15
19.	<i>Chrysis matutina</i> Semenov, 1967	7
20.	<i>Chrysis mediata</i> Linsenmaier, 1951	15
21.	<i>Chrysis mocsaryi</i> Radoszkowski, 1889	3
22.	<i>Chrysis mysticalis</i> Linsenmaier, 1959	4, 7, 9, 15, 20
23.	<i>Chrysis nox</i> Semenov, 1954	5, 15
24.	<i>Chrysis pavesii</i> Rosa, 2017	5, 15
25.	<i>Chrysis priapus</i> Rosa, 2018	5
26.	<i>Chrysis pseudobrevitarsis</i> Linsenmaier, 1951	7, 15
27.	<i>Chrysis pupilla</i> Semenov, 1967	12
28.	<i>Chrysis rutilans</i> Olivier, 1791	15
29.	<i>Chrysis schencki</i> Linsenmaier, 1968	7, 9
30.	<i>Chrysis sibirica</i> Rosa, 2017	7
31.	<i>Chrysis solida</i> Haupt, 1957	21
32.	<i>Chrysis splendidula unica</i> Radoszkowski, 1891	7
33.	<i>Chrysis subcoriacea</i> Linsenmaier, 1959	7
34.	<i>Chrysis viridula</i> Linnaeus, 1761	15
35.	<i>Chrysura dichroa</i> (Dahlbom, 1854)	4
36.	<i>Chrysura ignifrons</i> (Brullé, 1833)	4
37.	<i>Cleptes dauriensis</i> Móczár, 1997	3, 8, 11
38.	<i>Cleptes mongolicus</i> Rosa, Halada, & Agnoli, sp. nov.	21
39.	<i>Colpopyga nesterovi</i> Rosa, 2017	21
40.	<i>Elampus albipennis</i> (Mocsáry, 1889)	7, 20
41.	<i>Elampus coloratus</i> Rosa, 2017	22
42.	<i>Elampus montanus</i> (Mocsáry, 1890)	20
43.	<i>Elampus panzeri</i> (Fabricius, 1804)	4, 7
44.	<i>Elampus sanzii</i> Gogorza, 1887	15
45.	<i>Elampus spinifemoris</i> (Móczár, 1967)	11
46.	<i>Euchreous mongolicus</i> Tsuneki, 1947	5, 11, 12
47.	<i>Euchreous orientis</i> Semenov, 1910	22
48.	<i>Hedychridium ardens</i> (Coquebert, 1801)	4, 7, 8, 11, 13, 16, 18, 21, 22
49.	<i>Hedychridium asianum</i> Linsenmaier, 1997	7–9, 16
50.	<i>Hedychridium belokobylskiji</i> Rosa, 2017	15
51.	<i>Hedychridium cypreum</i> (Dahlbom, 1845)	4, 5, 8, 11, 12, 15, 20
52.	<i>Hedychridium gabriellae</i> Rosa, 2017	8, 15, 20
53.	<i>Hedychridium longigena</i> Rosa, 2017	8, 9, 13, 15, 18, 20, 21
54.	<i>Hedychridium propodeale</i> Rosa, 2017	5
55.	<i>Hedychridium roseum</i> (Rossi, 1790)	7, 20–22
56.	<i>Hedychrumb chalybaeum</i> Dahlbom, 1854	5, 8, 13, 15, 16, 21, 22
57.	<i>Hedychrumb gerstaeckeri</i> Chevrier, 1869	13, 15, 18
58.	<i>Hedychrumb lama</i> du Buysson, 1891	3

No.	Species	Aimags
59.	<i>Hedybrum longicolle</i> Abeille de Perrin, 1877	9, 12, 15, 21, 22
60.	<i>Hedybrum nobile</i> (Scopoli, 1763)	4, 7, 13, 15
61.	<i>Hedybrum rutilans ermak</i> Semenov, 1967	7, 13, 15, 21, 22
62.	<i>Holopyga generosa asiatica</i> Trautmann, 1926	13
63.	<i>Holopyga kaszabi</i> Móczár, 1967	11, 12, 20
64.	<i>Holopyga minuma</i> Linsenmaier, 1959	21, 22
65.	<i>Ornalus aeneus</i> (Fabricius, 1787)	15, 16
66.	<i>Ornalus berezovskii</i> (Semenov, 1932)	16
67.	<i>Ornalus margianus</i> (Semenov, 1932)	7–9, 15, 22
68.	<i>Ornalus miramae</i> (Semenov, 1932)	8, 20, 22
69.	<i>Ornalus stella</i> (Semenov, 1932)	7, 11, 15
70.	<i>Parnopes glasunowi</i> Semenov, 1901	3
71.	<i>Parnopes popovii</i> Eversmann, 1858	7, 9, 12, 15, 20–22
72.	<i>Pentachrysis amoena</i> (Eversmann, 1858)	without locality
73.	<i>Philotetes bogdanovii</i> (Radoszkowski, 1877)	7
74.	<i>Philotetes cynthiae</i> Rosa, 2017	8, 11, 16, 22
75.	<i>Philotetes diakonovi</i> (Semenov, 1932)	20
76.	<i>Philotetes lyubae</i> Rosa, 2017	20
77.	<i>Philotetes mongolicus</i> (du Buysson, 1901)	7, 8, 11, 15, 16, 18, 22
78.	<i>Philotetes shokalskii</i> (Semenov, 1932)	8, 11, 12, 15, 16, 18–22
79.	<i>Pseudochrysis gengiskhan</i> Rosa, 2017	8, 9, 13, 15, 21, 22
80.	<i>Pseudochrysis neglecta</i> (Shuckard, 1837)	15
81.	<i>Pseudomalus auratus nigridorsus</i> (Tsuneki, 1953)	4, 9, 15, 18
82.	<i>Pseudomalus corensis</i> (Uchida, 1927)	9, 13, 15, 16, 18, 21
83.	<i>Pseudomalus punctatus</i> (Uchida, 1927)	9, 15, 18, 21
84.	<i>Pseudomalus pusillus</i> (Fabricius, 1804)	8, 9, 11–13, 15, 18, 21
85.	<i>Spinolia spinosa</i> Rosa & Halada, sp. nov.	8
86.	<i>Spinolia unicolor</i> (Dahlbom, 1831)	5
87.	<i>Stilbum calens</i> (Fabricius, 1781)	7, 9, 11, 15, 20
88.	<i>Trichrysis cyanea</i> (Linnaeus, 1758)	8, 13, 15
89.	<i>Trichrysis pellucida</i> (du Buysson, 1887)	without locality
90.	<i>Trichrysis secernenda</i> (Mocsáry, 1912)	13

Comment. Aimag designation as in Fig. 1.

The checklist follows the genera subdivision proposed by Kimsey and Bohart (1991), with few exceptions for some genera (e.g., *Euchroeus* Latreille, 1809, *Pseudochrysis* Semenov, 1891 and *Colpopyga* Semenov, 1954). The species are listed alphabetically. We have used the following abbreviations for collectors: **JH** – J. Halada; **JS** – J. Straka; **MH** – M. Halada; **MK** – M. Kadlecová. An asterisk (*) marks the new records.

Types and other specimens are deposited in the following Institutions and private collections:

- EIHU** Entomology Institute, Hokkaido University (Japan);
HNHM Hungarian Natural History Museum, Zoological Department, Budapest (Hungary);
ISEA-PAS Institute of Systematics and Evolution of Animals, Polish Academy of Sciences, Kraków (Poland);
LSL Linnean Society, London (England);
MCNM Museo National de Ciencias Naturales, Madrid (Spain);
MFN Museum für Naturkunde, Berlin (Germany);

MHNG	Museum d’Histoire Naturelle, Geneva (Switzerland);
MNHN	National Museum of Natural History, Paris (France);
MSNG	Museo di Storia Naturale, Genova (Italy);
NHMUK	British Museum of Natural History, London (UK);
NHMW	Museum of Natural History, Vienna (Austria);
NIAS	National Institute of Agro-Environmetal Science, Tsukuba (Japan);
NMLS	Natur Museum, Luzern (Switzerland);
OMNH	Osaka Museum of Natural History, Osaka (Japan);
ZIN	Zoological Institute, Russian Academy of Sciences, St. Petersburg (Russia);
ZMMU	Zoological Museum of Moscow Lomonosov State University (Russia);
ZMUL	Zoologiska Museet, Lund Zoological Museum, University of Lund (Sweden);
GLAC	G.L. Agnoli collection (Bologna, Italy);
MHC	M. Halada collection (České Budějovice, Czech Republic);
PRC	P. Rosa collection (Bernareggio, Italy);
UKC	U. Koschwitz collection (Eppenbraun, Germany).

Results

Taxa from Mongolia

Subfamily Cleptinae

Genus *Cleptes* Latreille, 1802

Cleptes Latreille, 1802: 316. Type species: *Sphex semiaurata* Linnaeus, 1761 [= *Cleptes semiauratus* (Linnaeus, 1761)], by monotypy.

Cleptes dauriensis Móczár, 1997

Cleptes (*Cleptes*) *dauriensis* Móczár, 1997: 36. Holotype ♀: Russia: Dauria, leg. F. Sahlb., “*Cleptes* n. sp. *nitidulo* Fbr. aff.”, Holotype *Cleptes dauriensis* ♀ Móczár n. sp. det. Móczár 1995” (Hym. Typ. No. 3845 Mus. Budapest) (HMNH).

Cleptes dauriensis: Rosa 2017a: 288. Rosa et al. 2019: 310 (Mongolia, Figs 4, 5).

Material examined. MONGOLIA: Khovd, 1 ♂, Bodongin-Gol River, 12 km SW Altai, 22.VII.1970, leg. M. Kozlov (ZIN); Uvurkhangai, 1 ♀, 12 km E of Arvaykheer, 46°22'N, 102°49'E, 1800 m, 3.VII.2004, leg. JH (GLAC); Bayankhongor, 1 ♂, 16 km SW of Bayankhongor, 46°13'N, 100°30'E, 2165 m, 10.VII.2004, leg. JH (GLAC).

Distribution. Mongolia (Bayankhongor, Khovd, Uvurkhangai); Russia (Zabaykalskii Terr.) (Rosa 2017a).

***Cleptes mongolicus* Rosa, Halada & Agnoli, sp. nov.**

<http://zoobank.org/73389B93-F683-41CC-84AC-3E16ED9B3000>

Figures 2, 3

Type material. Holotype: ♀, MONGOLIA: *Dornod*, 100 km W of Choibalsan, 820 m, 23.VII.2007, leg. M. Halada (ZIN). **Paratypes:** 1 ♂, same collecting locality and date (GLAC); 1 ♂, 20 km W of Choibalsan, 48°01'N, 114°14'E, 800 m, 24.VII.2007, leg. M. Halada (PRC).

Diagnosis. *Cleptes mongolicus* sp. nov. belongs to the *C. nitidulus* species group, based on the pronotum without posterior pit row and without longitudinal median sulcus or posterior median keel. It is closely related only to *C. margaritae* Móczár, 2000 from Tajikistan, for its general habitus and colouration. The latter belongs to the *C. satoi* group (Móczár 2000), for the modified pronotal structure, without posterior transversal groove, but with a posteromedian longitudinal keel. Besides the unmodified pronotum, the female of *C. mongolicus* sp. nov. can be easily separated from the female of *C. margaritae* by: a) pubescence whitish, shorter on metasoma (max 2.5 MOD) (vs. blackish, longer on metasoma, up to 3 MOD); b) punctuation on metasoma with polished T1, shallow and sparse tiny punctures on T2, double punctures on T3 (vs. scattered punctate on T1, densely and evenly punctate on T2 and T3); c) colouration: head entirely black; propodeum entirely blue; T3 and T4 laterally blue; pedicel and F1 yellow; femora apically, tibiae and tarsi yellow (vs. head blue; propodeum black with median blue spot; T3 and T4 fully black; pedicel and flagellum dark brown). The male of *Cl. margaritae* is currently unknown.

Description. Female. Holotype (Fig. 2A–F). Body length 4.6 mm. Forewing length 2.7 mm. POL = 2.2 MOD; OOL = 2.7 MOD. MS = 2.0 MOD. P:F1:F2:F3 = 1.0:1.0:0.7:0.7. F1 1.5 × as long as wide, F2 1.1 × as long as wide. **Head.** Head in frontal view 1.2 × as broad as long between lower edge of clypeus and vertex. Face and vertex with small, even, and sparse punctures (1–4 PD) (Fig. 2B). Clypeal lower margin simple, unmodified, 2 MOD width, without acute teeth at corners; lateral edges subparallel. Frontal sulcus broad and deep in the first part, from anterior ocellus to mid of face, faint in the second half, from mid-face to the clypeal margin (Fig. 2B). Mandibles tridentate. Ocellar triangle isosceles, without post-ocellar sulcus. Postero-lateral pits close to posterior ocelli deep and elongate. Pedicel as long as F1. Malar spaces elongate (2.0 MOD). **Mesosoma.** Pronotum unmodified; pronotal neck finely striated transversally; posterior margin of pronotum simple, without transverse row of pits or median keel. Pronotum with small punctures similar to those on vertex. Mesoscutum and mesoscutellum scarcely punctate, with tiny and scattered punctures (Fig. 2C), largely impunctate; notauli and parapsidal lines deep and complete. Mesopleuron with small, deep punctures; transversely aligned medially; with short, deep scrobal sulcus on posterior half (Fig. 2D). Metascutellum noticeably reduced by large metanotal trough and by deep and large anteromedian suture. Metapleuron transversely striate. Metapostnotum (dorsal surface of metapectal-propodeal complex) short, irregularly




Figure 2. *Cleptes mongolicus* sp. nov., female, holotype **A** habitus, dorsal view **B** head, frontal view **C** head and mesosoma, dorsal view **D** head and mesosoma, lateral view **E** metasoma, postero-lateral view **F** metasoma, dorso-lateral view. Scale bars: 1.0 mm.

reticulate, with large foveae along posterior margin, before the propodeal declivity. Propodeal posterior projections short, stout, and divergent. Wing veins and cells unmodified. **Metasoma.** All metasomal terga with impunctate, brownish stripe along posterior margin (Fig. 2F); T1 mostly impunctate, with a few, sparse, tiny punctures; T2 with even, sparse, small punctures (3–5 PD), posteromedially polished; T3 with dense, irregular and double punctuation; scattered to polished toward the apical margin; T4 with large, scattered punctures. **Colouration.** Head black, with violet reflections medially on clypeus; scapus dorsally violet, ventrally brownish without metallic reflections; P light brown and F1 yellow; other flagellomeres dark brown to blackish. Mandible dark brown, medially yellowish. Pronotal neck medially black; pronotum, mesonotum, mesopleuron, metanotum (excluding black anterior suture and axillary trough), metapleuron metallic red with purple reflections dorsally; propodeum dorsally blue, propodeal declivity black; body ventrally black. Metasoma entirely black; apical margin of each tergum with brownish stripe; laterally on T3 with feeble green reflections; laterally on T4 with extended blue reflections (Fig. 2E). Tegulae brown. Legs with tibiae and tarsi yellowish; coxae red to golden; profemur anteriorly metallic red excluding distal joint; metafemur posteriorly metallic; other parts brown.

Male. Paratypes. Body length 4.0–4.2 mm. POL = 1.6 MOD; OOL = 1.0 MOD. MS = 1.9 MOD. P:F1:F2:F3 = 1.0:1.4:0.9:0.9. F1 3.5 × as long as wide (width taken at




Figure 3. *Cleptes mongolicus* sp. nov., male, paratype **A** habitus, dorsal view **B** head, frontal view **C** head and mesosoma, dorsal view **D** mesosoma, lateral view **E** metasoma, postero-lateral view **F** metasoma, dorsal view. Scale bars: 1.0 mm.

distal apex), F2 1.5×. **Head.** Head in frontal view 1.3 × as broad as long between lower edge of clypeus and vertex. Face and vertex with small, even, and denser punctures (1–2 PD) compared to female (Fig. 3B). Frontal sulcus narrow and visible in the first part, from anterior ocellus to brow, faint in the second half, from mid-face to the clypeal margin (Fig. 3B). Lower face medially with punctures more spaced 4–5 PD. Ocellar triangle, post-ocellar sulcus, and posterolateral pits similar to female. F1 1.5 × as long as P. **Mesosoma.** Punctuation overall similar to that of female; metascutellum larger, with narrow anteromedian mesoscutellar-metascutal suture; metapleuron polished. Other characters as in female. **Metasoma.** T1 with denser (2–5 PD), tiny punctures; T2 with even, denser (1–3 PD), small punctures (3–5 PD), posteromedially sparser to polished; T3 with dense, irregular and double punctuation; scattered to polished toward the apical margin; T4 with similar punctures; T5 almost polished, with scattered punctures. **Colouration.** Species sexually dimorphic with head and mesosoma bright green, including ventral side; propodeum blue. Mandible metallic green from base to half length. Scapus green, pedicel and flagellum black. Metasoma entirely black, with terga apically brownish and laterally with feeble blue reflections on T3 and T4 (Fig. 3E). Tegulae brown. Coxae and femora medially green; trochanters brown, femora distally and tarsi yellowish.

Etymology. The specific epithet is named after the country of origin.

Distribution. Mongolia (Dornod).

Subfamily Chrysidinae**Tribe Chrysidini****Genus *Chrysis* Linnaeus, 1761**

Chrysis Linnaeus, 1761: 414. Type species: *Sphex ignita* Linnaeus, 1758 [= *Chrysis ignita* (Linnaeus, 1758)], by subsequent designation of Latreille 1810: 437.

Tetrachrysis Lichtenstein, 1876: 27. Type species: *Chrysis aeruginosa* Dahlbom, 1854, by subsequent designation of Ashmead 1902: 226. Synonymized by Lisenmaier 1959: 91.

***Chrysis aestiva* Dahlbom, 1854**

Chrysis aestiva Dahlbom, 1854: 286. Holotype ♀; Greece: Rhodes (Berlin?) (*aestiva* group).

Material examined. MONGOLIA: Arkhangai, 1 ♂, 90 km NE of Tsetserleg, 48°03'N, 102°25'E, 27.VII.2005, leg. JH (MHC).

Distribution. *Mongolia (Arkhangai); Asiatic-European, from Caucasus, Turkey, Greece, Iran, Palestine, European part of Russia to Mongolia (Rosa et al. 2019, present record).

Remarks. This is the most eastern record for *Chrysis aestiva*.

***Chrysis angustula* Schenck, 1856**

Chrysis angustula Schenck, 1856: 28. Lectotype ♀ (designated by Morgan 1984: 9); Germany: former Duchy of Nassau (Frankfurt) (*ignita* group).

Material examined. MONGOLIA: Arkhangai, 5 ♀♀, 1 ♂, Chuluut Gol River, 47°48'N, 100°19'E, 23.VII.2005, leg. JH (MHC); 4 ♀♀, 1 ♂, 70 km NE of Tsetserleg, 25.VII.2005, leg. JH (MHC); Tuv, 1 ♀, 2 ♂♂, 50 km E of Ulaanbaatar, Tuul River, 22.VI.2003, leg. JH (MHC).

Distribution. *Mongolia (Arkhangai, Tuv); Asiatic-European, from western Europe to China and Russia (Rosa et al. 2019).

***Chrysis asahinai* Tsuneki, 1950**

Chrysis (Tetrachrysis) asahinai Tsuneki, 1950: 80. Holotype ♀; China, Manchuria, 22.VIII.1938, leg. S. Asahina (OMNH) (*pulchella* group).

Chrysis asahinai: Móczár 1967: 189 (cat., Mongolia: 1 ♀, Estgobi aimag: Cagan Elis, 800 m, 30 km ESE von Zuun-Bajan, Exp. Dr. Z. Kasab, 1963, nr. 22, 23.VI.1963).

Material examined. MONGOLIA: *Bayankhongor*, 12 ♂♂, 130 km S of Bayankhongor, 45°03'N, 100°59'E, 1240 m, 6.VII.2004, leg. JH, MK (MHC, PRC); 1 ♀, ibid, Orog Nuur, 6–7.VII.2004, on *Saxaul*, leg. JS (PRC); *Bulgan*, 13 ♀♀, 4 ♂♂, Mongol Els Nat. Res., dunes, 47°24'N, 103°39'E, 31.VII.2005, leg. JH (MHC); *Sukhbaatar*, 1 ♂, 100 km SSW of Baruun-Urt, 1100 m, 30.VII.2007, leg. MH (MHC); *Tuv*, 2 ♀♀, 2 ♂♂, 75 km W of Ulaanbaatar, dunes, 2.VIII.2005, leg. JH (MHC); 39 ♀♀, 37 ♂♂, 70 km W of Ulaanbaatar, 1070 m, dunes, 16.VIII.2007, leg. JH, MH (MHC); *Umnugovi*, 1 ♂, Gobi, 100 km SW of Dalanzadgad, Bayanzag, on *Saxaul*, 1–2.VII.2003, leg. JH (MHC).

Distribution. Mongolia (**Bayankhongor*, **Bulgan*, Dornogovi, **Sukhbaatar*, **Tuv*, **Umnugovi*); China (Liaoning) (Rosa et al. 2014).

Chrysis belokobylskiji Rosa, 2019

Chrysis belokobylskiji Rosa, 2019: 2. Holotype ♀; Kyrgyzstan: Naryn River near Kara-kolka (ZIN) (examined); paratypes: 2 ♀♀, 1 ♂ [Mongolia: Nogon-kub, N. Gobi; 50 km E of Ulaanbaatar, Tuul River; 40 km SW of Uliastay] (*pulchella* group).

Material examined. MONGOLIA: *Umnugovi*, 1 ♀, Nogon-kub, N. Gobi, 1.VIII.1926, P. Kozlov (ZIN); *Tuv*, 1 ♂, 50 km E of Ulaanbaatar, Tuul River, 22.VI.2003, leg. JH (MHC); *Zavkhan*, 1 ♀, 40 km SW of Uliastay, dunes, 18.VII.2005, leg. JH (MHC).

Distribution. Mongolia (*Tuv*, *Umnugovi*, *Zavkhan*); China (Qinghai), Kyrgyzstan, Tajikistan (Rosa 2019).

Chrysis brevitarsis Thomson, 1870

Chrysis brevitarsis Thomson, 1870: 107. Holotype ♀; Sweden: Nerike [= Närke] (Lund) (examined) (*ignita* group).

Material examined. MONGOLIA: *Bulgan*, 1 ♀, 137 km NE of Aravaykheer, 47°20'N, 103°40.5'E, 1250 m, 26.VII.2004, leg. JH (MHC).

Distribution. *Mongolia (*Bulgan*); Asiatic-European, from western Europe to Russia (Rosa et al. 2019).

Chrysis castigata Linsenmaier, 1959

Chrysis (Chrysis) exsulans var. *asiatica* Linsenmaier, 1951: 82. Holotype ♀; Uzbekistan: Ferghana (Budapest) (examined) (*ignita* group), nom. praecocc., nec Radoszkowski 1889.

Chrysis (Chrysis) exsulans var. *castigata* Linsenmaier, 1959: 155. Replacement name for *C. asiatica* Linsenmaier, 1951.

Material examined. MONGOLIA: *Selenge*, 2 ♂♂, 90 km N of Ulaanbaatar, Segnez River, 1450 m, 6–8.VII.2003, leg. JH (MHC); *Tuv*, 1 ♂, 50 km N of Ulaanbaatar, E of Mandal, 1180 m, 8–13.VIII.2007, leg. MH (MHC).

Distribution. *Mongolia (Selenge, Tuv); Kazakhstan, Kyrgyzstan, Turkmenistan, Uzbekistan, Russia (Eastern Siberia) (Rosa et al. 2019).

Chrysis chinensis Mocsáry, 1912

Chrysis (Tetrachrysis) ignita var. *chinensis* Mocsáry, 1912: 589. Holotype ♀; China: Shanghai (HNHM) (examined) (*ignita* group).

Chrysis chinensis: Rosa et al. 2019: 109 (cat., Mongolia, without locality, see Material examined).

Material examined. Mongolia: *Arkhangai*, 24 ♂♂, Chuluut Gol River, 47°48'N, 100°19'E, 23.VII.2005, leg. JH (MHC); 2 ♀♀, 4 ♂♂, 90 km NE of Tsetserleg, 48°03'N, 102°25'E, 24.VII.2004, leg. JH (MHC); 2 ♀♀, ibid, 27.VII.2005, leg. JH (MHC); *Tuv*, 1 ♀, 1 ♂, Ulaanbaatar Bog Duul, 11.VII.1983, leg. Karl Bleyl (NMLS); 7 ♀♀, 28 ♂♂, 50 km E of Ulaanbaatar, Tuul River, 22.VI.2003, leg. JH (MHC); 6 ♂♂, Khan-gayn Mts, 5 km N of Khunt, 20.VII.2005, leg. JH (MHC); *Selenge*, 11 ♀♀, 19 ♂♂, 90 km N of Ulaanbaatar, Segnez River, 1450 m, 6–8.VII.2003, leg. JH (MHC).

Distribution. Mongolia (*Arkhangai, *Selenge, *Tuv); Asiatic-European, from western Europe (Switzerland) to China (Helsonjiang, Shanghai) (Linsenmaier 1959).

Remarks. This species was previously reported from Mongolia (Rosa et al. 2019) without exact locality.

Chrysis consanguinea Mocsáry, 1889

Chrysis (Gonochrysis) consanguinea Mocsáry, 1889: 299. Syntypes ♀♀; Italy: Sicily; Algeria (MHNG) (examined) (*viridula* group).

Material examined. MONGOLIA: *Arkhangai*, 1 ♂, Chuluut Gol River, 47°48'N, 100°19'E, 23.VII.2005, leg. JH (MHC); 3 ♀♀, 90 km NE of Tsetserleg, 48°03'N, 102°25'E, 27.VII.2005, leg. JH (MHC); *Bulgan*, 1 ♀, 137 km NE of Aravaykheer, 47°20'N, 103°40.5'E, 1250 m, 2.VII.2004, leg. JH (MHC); *Dornod*, 1 ♀, 100 km W of Choibalsan, 820 m, 23.VII.2007, leg. JH (MHC); *Khentii*, 4 ♂♂, 100 km NE of Ondorkhaan, Kerulen River, 970 m, 22.VII.2007, leg. MH (MHC); *Selenge*, 1 ♀, 2 ♂♂, 90 km N of Ulaanbaatar, Segnez River, 1450 m, 6–8.VII.2003, leg. JH (MHC); *Sukhbaatar*, 2 ♂♂, 100 km SSW of Baruun-Urt, 1100 m, 30.VII.2007, leg. MH (MHC); *Tuv*, 7 ♂♂, 50 km E of Ulaanbaatar, Tuul River, 22.VI.2003, leg. JH (MHC);

1 ♂, Khangaun Mts, 5 km N of Khunt, 20.VII.2005, leg. JH (MHC); 1 ♀, 50 km N of Ulaanbaatar, E of Mandal, 1180 m, 8–13.VIII.2007, leg. JH (MHC); *Ulaanbaatar*, 1 ♀, 7 km E of Ulaanbaatar, Gachuurt, 47°55'N, 107°06'E, 31.VII.2002, leg. JS (MHC); *Zavkhan*, 1 ♂, 40 km SW of Uliastay, dunes, 18.VII.2005, leg. JH (MHC).

Distribution. *Mongolia (Arkhangai, Bulgan, Dornod, Khentii, Selenge, Sukhbaatar, Tuv, Ulaanbaatar, Zavkhan); Palaearctic, from southern Europe and northern Africa to Eastern Siberia and Mongolia (Rosa et al. 2019, present records).

Chrysis dauriana Linsenmaier, 1959

Chrysis (Chrysis) cavaleriei ssp. *dauriana* Linsenmaier, 1959: 112. Holotype ♀; Russia: Dauria (NMLS) (examined) (*succincta* group). Elevated to species rank by Rosa et al. 2017a: 40.

Chrysis (Tetrachrysis) mongolica Semenov-Tian-Shanskij, 1967: 178, nec Mocsáry, 1914. Holotype ♀; Russia [not Mongolia]; Transbaikalia: Ingoda River (St. Petersburg) (examined). Rosa et al. 2017a: 39 (cat., type series), 155 (Plate 91). Synonymised by Rosa et al. 2017a: 40.

Chrysis mongoliana Bohart in Kimsey and Bohart 1991: 440. Replacement name for *Chrysis mongolica* Semenov-Tian-Shanskij, 1967: 178, nec Mocsáry 1914.

Material examined. MONGOLIA: *Arkhangai*, 1 ♀, 90 km NE of Tsetserleg, 48°03'N, 102°25'E, 27.VII.2005, leg. JH (MHC); *Bayankhongor*, 2 ♂♂, 163 km S of Bayankhongor, 46°13'N, 100°30'E, 2165 m, 10.VII.2004, leg. JH (MHC); *Bulgan*, 1 ♂, 170 km W of Ulaanbaatar, dunes, 1070 m, 16.VIII.2007, leg. MH (MHC); *Khentii*, 1 ♀, 100 km NE of Ondorkhaan, Kerulen River, 970 m, 22.VII.2007, leg. MH (MHC); *Selenge*, 4 ♀♀, 2 ♂♂, 90 km N of Ulaanbaatar, Segnez River, 1450 m, 6–8.VII.2003, leg. JH (MHC); *Tuv*, 2 ♀♀, 4 ♂♂, 50 km E of Ulaanbaatar, Tuul River, 22.VI.2003, leg. JH (MHC); 2 ♂♂, ibid, 12.VII.2003, leg. JH (MHC); 10 ♀♀, 6 ♂♂, 50 km N of Ulaanbaatar, E of Mandal, 1180 m, 8–13.VIII.2007, leg. MH (MHC, PRC); *Zavkhan*, 2 ♂♂, 40 km SW of Uliastay, dunes, 18.VII.2005, leg. JH (MHC).

Distribution. Mongolia (*Arkhangai, *Bayankhongor, *Bulgan, *Khentii, *Selenge, *Zavkhan); Russia (Eastern Siberia) (Rosa et al. 2017a).

Remarks. This species was previously reported from Mongolia (Rosa et al. 2019) without exact locality.

Chrysis equestris Dahlbom, 1854

Chrysis equestris Dahlbom, 1854: 307. Holotype ♀; locality unknown [most likely Sweden] (Stockholm) (examined) (*smaragdula* group).

Material examined. MONGOLIA: *Arkhangai*, 1 ♀, 2 ♂♂, 90 km NE of Tsetserleg, 48°03'N, 102°25'E, 24.VII.2004, leg. JH (MHC); 1 ♀, 1 ♂, ibid, leg. MK (MHC);




Figure 4. *Chrysis fulgida* Linnaeus, habitus dorsal view **A** form A, ♀ **B** form B, ♀ **C** form A, ♂ **D** form B, ♂. Scale bars: 1.0 mm.

1 ♀, 1 ♂, 70 km NE Tsetserleg, 25.VII.2005, leg. JH (MHC); *Selenge*, 1 ♂, 90 km N of Ulaanbaatar, Segnez River, 1450 m, 6–8.VII.2003, leg. JH (MHC).

Distribution. *Mongolia (Arkhangai, Selenge); Asiatic-European, from western Europe to Russia (Rosa et al. 2019).

Chrysis fulgida Linnaeus, 1761

Chrysis fulgida Linnaeus, 1761: 415. Lectotype ♀ (designated by Morgan 1984: 9); Sweden: Uppsala (LSL) (*ignita* group).

Material examined. MONGOLIA: (Form A): Arkhangai, 2 ♀♀, 1 ♂, 70 km NE of Tsetserleg, 25.VII.2005, leg. JH (MHC); *Selenge*, 1 ♂, 90 km N of Ulaanbaatar, Segnez River, 1450 m, 6–8.VII.2003, leg. JH (MHC); *Tuv*, 5 ♀♀, 3 ♂♂, 50 km E of Ulaanbaatar, Tuul River, 22.VI.2003, leg. JH (MHC); (Form B): Arkhangai, 2 ♀♀, 1 ♂, 70 km NE of Tsetserleg, 25.VII.2005, leg. JH (MHC); 1 ♀, 1 ♂, Chuluut Gol River, 47°48'N, 100°19'E, 23.VII.2005, leg. JH (MHC); *Tuv*, 1 ♀, 50 km E of Ulaanbaatar, Tuul River, 22.VI.2003, leg. JH (MHC).

Distribution. *Mongolia (Arkhangai, Selenge, Tuv); Asiatic-European, from Europe to eastern Siberia, Russian Far East and North-East China (Manchuria) (Rosa et al. 2014, 2019).

Remarks. Two distinct colour forms (Fig. 4) are recorded from Mongolia, Siberia and Primorsky Territory (Russia), and Heilongjiang (China). Form A is matching with the typical European *Chrysis fulgida* (Fig. 4A, C). Form B is chromatic different

without the typical blue colouration on male and female metasoma and with non-metallic black areas on head vertex and mesosoma (Fig. 4B, D). Male T1 golden-greenish, with or without a narrow transversal green or bluish stripe or patch; T2 red, with or without a basal, narrow black stripe; female T1 golden-greenish, with green to bluish colour on T1 frontal declivity to petiolar insertion. This colour variation has also been observed in specimens from Russia (Siberia and Primorsky Territory) and China (Heilongjiang). The Chinese form was mentioned by Linsenmaier (1968) as *Chrysis aequicolor* Linsenmaier, 1968, which is anyway an unnecessary replacement name for *Chrysis fulgida* var. *concolor* Mocsáry, 1912 nec Mocsáry, 1892 (actually male and female of the same taxon). Other evident different morphological characteristics are not recognizable. However, these two forms may represent two sister species, genetically separate, but difficult to identify on the basis of morphological characteristics, as in other known cases of *Chrysis* of the *ignita* group (Paukkunen et al. 2015; Orlovskyté et al. 2016).

Chrysis ignita (Linnaeus, 1758)

Sphex ignita Linnaeus, 1758: 571. Lectotype ♀ (designated by Richards 1935); Europe (LSL) (*ignita* group).

Chrysis ignita: Buyanjargal and Abasheev 2015: 31 (biol. host of *Euodynerus dantici*, central Mongolia: Khugnu-Khaan Mts, Khugnu-Tarna N.P.).

Material examined. None examined.

Remarks. The identification of *Chrysis ignita* by Buyanjargal and Abasheev (2015) is doubtful and very likely represent another species of the *C. ignita* group or even a member of another species group (e.g., *succincta* group). In fact, the host association with *Euodynerus dantici*, as observed by the two authors, is unusual. *Euodynerus dantici* is known as a possible host for members of the *C. succincta* group (*C. germari* and *C. tristis* (sub *C. succincta succinctula*) Pauli et al. 2019, supplementary file 4). For example, *C. dauriana* Linsenmaier was erroneously identified as *C. ignita* by several authors, including Trautmann (identification label pinned with the type of *C. dauriana*).

Distribution. Mongolia (Bulgan) [doubtful]; West-Palaearctic: from West Europe to central Asia (Linsenmaier 1997b).

Chrysis illecebrosa Semenov, 1967

Chrysis (Tetrachrysis) illecebrosa Semenov-Tian-Shanskij, 1967: 166. Holotype ♂; Bulgaria near Khami, SE from Tian Shan [China, Xinjiang] (ZIN) (examined) (*maculicornis* group).

Material examined. MONGOLIA: Umnugovi, 1 ♂, Deemgin-gobi, 25 km SSO of Khajlastyn-Khuduka, 20.VI.1971, leg. M. Kozlov (ZIN).

Distribution. *Mongolia (Umnugovi); China (Xinjiang) (Rosa et al. 2014).

Chrysis illigeri Wesmael, 1839

Chrysis illigeri Wesmael, 1839: 176. Syntypes ♂♀; Belgium (Bruxelles, MSNG) (examined) (*succincta* group).

Material examined. MONGOLIA: *Selenge*, 1 ♂, 90 km N of Ulaanbaatar, Segnez River, 1450 m, 6–8.VII.2003, leg. JH (MHC); *Tuv*, 1 ♀, 50 km N of Ulaanbaatar, E of Mandal, 1180 m, 8–13.VIII.2007, leg. MH (MHC).

Distribution. *Mongolia (Selenge, Tuv); Asiatic-European, from western Europe to Mongolia (present record).

Chrysis ismaeli Semenov, 1967

Chrysis (Allochrysis) ismaeli Semenov-Tian-Shanskij, 1967: 124. Holotype ♀; Kazakhstan: Balamurun, Karatau Mountain ridge foothills, leg. V. Kozhantschikov (ZIN) (ear group).

Material examined. MONGOLIA: *Dornod*, 1 ♂, 100 km W of Choilbalsan, 820 m, 23.VII.2007, leg. MK (MHC); *Dornogovi*, 5 ♂♂, 65 km SE of Chatan-Bulag, 1020 m, 2.VIII.2007, leg. MH (PRC/MHC); *Umnugovi*, 1 ♀, Gobi, Dalanzadgad, 24–26.VI.2003, leg. JH (MHC); 12 ♀♀, 70 km S of Saynshand, 1100 m, 6.VIII.2007, leg. MH (PRC/MHC).

Distribution. *Mongolia (Dornod, Dornogovi, Umnugovi); Kazakhstan (Rosa 2018).

Notes. As supposed by Rosa (2018), living specimens are red and change to greenish after preparation.

Chrysis jaxartis Semenov, 1910

Chrysis sybarita var. *jaxartis* Semenov-Tian-Shansky, 1910: 222. Lectotype ♂ (designated by Rosa et al. 2017a: 54). Kazakhstan: Djulek (Budapest) (examined) (*graelsii* group).

Material examined. MONGOLIA: *Dornod*, 6 ♀♀, 100 km W of Choilbalsan, 820 m, 23.VII.2007, leg. MH (MHC); 9 ♀♀, 2 ♂♂, 20 km W of Choilbalsan, 800 m, 48°01'N, 114°14'E 24.VII.2007, leg. MH (MHC); *Khentii*, 11 ♀♀, 4 ♂♂, 100 km NE of Ondorkhaan, Kerulen River, 970 m, 22.VII.2007, leg. MH (MHC); *Selenge*, 3 ♀♀, 1 ♂, 90 km N of Ulaanbaatar, Segnez River, 1450 m, 6–8.VII.2003, leg. JH (MHC); *Tuv*, 5 ♀♀, 1 ♂, 50 km N of Ulaanbaatar, E of Mandal, 1180 m, 8–13.VIII.2007, leg. MH (MHC); 1 ♀, same date and locality, and collector (P. Tyrner priv. coll.); *Umnugovi*, 1 ♂, Gobi, Dalanzadgad, 25.VI.2003, leg. JH (MHC).

Distribution. *Mongolia (Dornod, Khentii, Selenge, Tuv, Umnugovi); Asiatic-European, from Greece, Iran, and Turkey to Central Asia (Rosa et al. 2019).

***Chrysis leptomandibularis* Niehuis, 2000**

Chrysis leptomandibularis Niehuis, 2000: 192. Holotype ♀; Germany: Rheinland-Pfalz, Monsheim (Frankfurt) (*ignita* group).

Material examined. MONGOLIA: *Tuv*, 3 ♀♀, 2 ♂♂, 50 km E of Ulaanbaatar, Tuul River, 22.VI.2003, leg. JH, det. J. van der Smissen and MH (MHC).

Distribution. *Mongolia (Tuv); Asiatic-European from Europe to Russia (Rosa et al. 2019).

***Chrysis mane* Semenov, 1912**

Chrysis mane Semenov-Tian-Shanskij, 1912: 192. Lectotype ♂ (designated by Bohart in Kimsey and Bohart 1991: 436); China: Alashan (192 (descr.), depository: ZIN).

Chrysis mane: Kimsey and Bohart 1991: 436 (China [not Mongolia]: Gansu, Qinghai, cat., *ignita* group).

Material examined. Mongolia: *Tuv*, 1 ♂, 50 km N of Ulaanbaatar, E of Mandal, 1180 m, 8–13.VIII.2007, leg. MK (PRC).

Distribution. *Mongolia (Tuv); China (Gansu, Qinghai, Inner Mongolia) (Rosa et al. 2014).

***Chrysis matutina* Semenov, 1967**

Chrysis (Tetrachrysis) matutina Semenov-Tian-Shanskij, 1967: 179. Holotype ♀; China: Gansu (ZIN) (*ignita* group).

Material examined. MONGOLIA: *Arkhangai*, 1 ♂, 90 km NE of Tsetserleg, 48°03'N, 102°25'E, 27.VII.2005, leg. JH (MHC).

Distribution. *Mongolia (Arkhangai); China (Gansu) (Rosa et al. 2014).

***Chrysis mediata* Linsenmaier, 1951**

Chrysis ignita var. *mediata* Linsenmaier, 1951: 76. Lectotype ♀ (designated by Linsenmaier 1959: 154); Switzerland: Wallis (NMLS) (examined) (*ignita* group).

Material examined. MONGOLIA: *Tuv*, 2 ♀♀, Ulaanbaatar Bog Duul, 11.VII.1983, leg. Karl Bleyle, det. Linsenmaier 1992 (NMLS).

Distribution. *Mongolia (Tuv); Palaearctic region excluding Japan (Linsenmaier 1997b).

Chrysis mocsaryi Radoszkowski, 1889

Chrysis (Tetrachrysis) Mocsaryi Radoszkowski, 1889: 29. Holotype ♀; Mongolia: Kobden (Khovd) (ISEA-PAS) (examined) (*comparata* group). Mocsáry 1889: 426 (cat., descr., Mongolia).

Chrysis mocsaryi: Dalla Torre 1892: 78 (cat., Mongolia); Kimsey and Bohart 1991: 440 (cat., Mongolia: Kobden, *comparata-scutellaris* group). Rosa et al. 2015: 41 (cat., type series), 42 (Fig. 4).

Material examined. MONGOLIA: Holotype ♀, golden rounded label, Kansu Kobden-Owatu 12/VIII [handwritten] *Mocsáry* [handwritten by Radoszkowski] // *Chrysis Mocsaryi* Rad. (tres interep.) [?] [handwritten by Mocsáry], label with right flagellum and metasoma, Mus. Pan Krakow [hadwritten by Dylewska].

Distribution. Mongolia (Khovd) (Radoszkowski 1889).

Chrysis mysticalis Linsenmaier, 1959

Chrysis mysticalis Linsenmaier, 1959: 165. Holotype ♀; Spain: Zamora (Luzern) (examined) (*inaequalis* group).

Material examined. MONGOLIA: *Arkhangai*, 1 ♀, 1 ♂, 90 km NE of Tsetserleg, 48°03'N, 102°25'E, 24.VII.2004, leg. MK (MHC); *Bulgan*, 2 ♂♂, Mongol Els Nat. Res., dunes, 47°24'N, 103°39'E, 31.VII.2005, leg. JH (MHC); *Dornogovi*, 1 ♀, 28 km of SE Chatan-Bulag, 3.VIII.2007, leg. MH (MHC); *Tuv*, 1 ♂, 50 km E of Ulaanbaatar, Tuul River, 22.VI.2003, leg. JH (MHC); 2 ♀♀, 50 km N of Ulaanbaatar, E of Mandal, 1180 m, 8–13.VIII.2007, leg. MH (MHC); *Zavkhan*, 1 ♀, 1 ♂, 40 km SW of Uliastay, dunes, 18.VII.2005, leg. JH (MHC).

Distribution. *Mongolia (Arkhangai, Bulgan, Dornogovi, Tuv, Zavkhan); from southern Europe to eastern Siberia (Rosa et al. 2017b).

Chrysis nox Semenov, 1954

Chrysis (Tetrachrysis) nox Semenov in Semenov-Tian-Shanskij and Nikol'skaya 1954: 128. Lectotype ♀ (designated by Bohart in Kimsey and Bohart 1991: 444); Tajikistan [not Mongolia]: Peter the Great Range, Yashil'-Kul' Lake, 7.VIII.1911, leg. Golbek (ZIN) (examined) (*facialis* group). Rosa et al. 2017a: 42 (cat., type series), 158 (plate 97).

Chrysis nox: Kimsey and Bohart 1991: 444 (cat., Mongolia: Yihe Bogdo, Peter the Great Range, *facialis* group).

Material examined. MONGOLIA: *Govi-Altai*, 4 ♀♀, 1 ♂, Ikhe-Bogdo, Gobi Altai, 30.VI–12.VII.1926, leg. P. Kozlov // Paratypes (ZIN); 4 ♀♀, idem, 15–17.VII.1926,

Paratypes (ZIN); 1 ♀, North slope of Ikhe-Bogdo, 30.VI–12.VII.1926, leg. P. Kozlov, Paratypes (ZIN); 1 ♀, Ihe-Bogdo, Gob. Altai, 15–17.VII.1926, leg. P. Kozlov [in Cyrillic], det. M. Nikol'skaya (NMLS); *Tuv*, 1 ♀, Ulaanbaatar Bog Duul, 11.VII.1983, leg. Karl Bleyl, det. Linsenmaier 1990 (NMLS).

Distribution. Mongolia (Govi-Altai, Tuv); Tajikistan (Rosa et al. 2017a).

Chrysis pavesii Rosa, 2017

Chrysis pavesii Rosa in Rosa et al. 2017c: 27. Holotype ♀; Russia: Western Siberia, Altai Rep., 5 km SE of Chagan-Uzun, Tudituyaryk River, 1780 m, 11.VII.2016, leg. M. Proshchalykin & V. Loktionov (ZIN) (examined) (*bihamata* group).

Material examined. MONGOLIA: *Govi-Altai*, 1 ♂, 10 km SSE of Ich-Oba-Ula, 18.VII.1970, leg. E. Narchuk (ZIN).

Distribution. *Mongolia (Govi-Altai, Tuv); Russia (western Siberia) (Rosa et al. 2017c).

Chrysis priapus Rosa, 2018

Chrysis priapus Rosa, 2018: 281. Holotype ♂; Mongolia: Govi-Altai Prov., 8 km SE of Argalant-Ula (ZIN) (examined) (*slava* group).

Material examined. MONGOLIA: *Govi-Altai*, 1 ♂, 8 km SE of Argalant-Ula, 20.VI.1980, leg. G. Medvedev (ZIN).

Distribution. Mongolia (Govi-Altai) (Rosa 2018).

Chrysis pseudobrevitarsis Linsenmaier, 1951

Chrysis ignita var. *pseudobrevitarsis* Linsenmaier, 1951: 79. Lectotype ♀ (designated by Linsenmaier 1959: 158); Switzerland: Wallis (NMLS) (examined) (*ignita* group).

Chrysis (Chrysis) pseudobrevitarsis: Linsenmaier 1997b: 114 (descr., Mongolia, without locality, see in Material examined).

Chrysis pseudobrevitarsis: Rosa et al. 2019: 153 (cat., Mongolia).

Material examined. MONGOLIA: *Arkhangai*, 1 ♂, 70 km NE of Tsetserleg, 25.VII.2005, leg. JH (MHC); *Tuv*, 1 ♀, Tereltz, 8.VII.1983, leg. Karl Bleyl, det. Linsenmaier 1992 (NMLS); 2 ♀♀, 50 km N of Ulaanbaatar, E of Mandal, 1180 m, 8–13.VIII.2007, leg. MH (MHC).

Distribution. Mongolia (Arkhangai, Tuv); Asiatic-European, from western Europe to Mongolia (Linsenmaier 1997a).

Chrysis pupilla Semenov, 1967

Chrysis (Tetrachrysis) pupilla Semenov-Tian-Shanskij, 1967: 174. Holotype ♀; Uzbekistan: Termez (ZIN) (examined) (*varidens* group).

Material examined. MONGOLIA: *Umnugovi*, 1 ♀, “Yuzhno-Gobiyskiy Ajmag, sajri Undyn-Gol, 25 km S of Khan-Bogdo, 7.VII.1971”, leg. M. Kozlov (ZIN).

Distribution. *Mongolia (Umnugovi); Uzbekistan (Semenov-Tian-Shanskij 1967).

Chrysis rutilans Olivier, 1791

Chrysis rutilans Olivier, 1791: 676. Type unknown; France: Angoumois (depository unknown) (*splendidula* group).

Material examined. Mongolia: *Tuv*, 1 ♀, 50 km N of Ulaanbaatar, E of Mandal, 1180 m, 8–13.VIII.2007, leg. MH (MHC).

Distribution. *Mongolia (Tuv); Palaearctic, from western Europe and North Africa to China and Japan (Lisenmaier 1997b).

Chrysis schencki Lisenmaier, 1968

Chrysis (Chrysis) ignita ssp. *schenckiana* Lisenmaier, 1959: 156, nom. praeocc., nec Mocsáry, 1912. Holotype ♀; Switzerland: Graubünden (Luzern) (examined) (*ignita* group).

Chrysis (Chrysis) ignita schencki Lisenmaier, 1968: 99. Replacement name for *C. ignita schenckiana* Lisenmaier, 1959.

Material examined. MONGOLIA: *Arkhangai*, 2 ♀♀, Chuluut Gol River, 47°48'N, 100°19'E, 23.VII.2005, leg. JH, det. J. Van der Smissen (MHC); *Bulgan*, 2 ♀♀, Mongol Els Nat. Res., dunes, 47°24'N, 103°39'E, 31.VII.2005, leg. JH, det. J. Van der Smissen (MHC).

Distribution. *Mongolia (Arkhangai, Bulgan); Asiatic-European, from western Europe to Central Asia, Siberia and Japan (Rosa et al. 2019).

Chrysis sibirica Rosa, 2017

Chrysis sibirica Rosa in Rosa et al., 2017c: 24. Holotype ♀; Russia: Tuva Rep., 31 km NEE of Erzin, Erzin River, 18.vii.2014, leg. A. Lelej, M. Proshchalykin & V. Laktionov (St. Petersburg) (*bihamata* group).

Material examined. MONGOLIA: *Arkhangai*, 1 ♀, Chuluut Gol River, 47°48'N, 100°19'E, 23.VII.2005, leg. JH (MHC).

Distribution. *Mongolia (Arkhangai); Russia (Eastern Siberia) (Rosa et al. 2017c).

***Chrysis solida* Haupt, 1957**

Chrysis ignita solida Haupt, 1957: 115. Lectotype ♀ (designated by Niehuis 2000: 199); Poland: Bellinchen [= Bielinek] (MLU) (*ignita* group).

Chrysis mediata fenniensis Linsenmaier, 1959: Móczár 1967: 189 (cat., Mongolia: 1 ♀, Čojbalsan [= Dornod] aimag: Menengijn valley, 160 km W of Bujr nur Lake, 600 m, Exp. Dr. Z. Kaszab, 1965, nr. 416, 15.VIII.1965).

Material examined. None examined.

Distribution. Mongolia (Dornod); Asiatic-European, from western Europe to Japan (Linsenmaier 1997b).

***Chrysis splendidula unica* Radoszkowski, 1891**

Chrysis splendidula var. *unica* Radoszkowski, 1891: 189. Syntypes ♂, ♀; Turkmenistan: Ashgabad (ISEA-PAS) (examined) (*splendidula* group).

Material examined. MONGOLIA: Arkhangai, 1 ♂, 90 km NE of Tsetserleg, 48°03'N, 102°25'E, 27.VII.2005, leg. JH (MHC).

Distribution. *Mongolia (Arkhangai); Turkmenistan (Radoszkowski 1891).

***Chrysis subcoriacea* Linsenmaier, 1959**

Chrysis (Chrysis) longula ssp. *subcoriacea* Linsenmaier, 1959: 160. Holotype ♀; Finland: Kyrkslätt [= Kirkkonummi] (Luzern) (examined) (*ignita* group).

Material examined. MONGOLIA: Arkhangai, 1 ♂, 70 km NE of Tsetserleg, 25.VII.2005, leg. JH, det. J. Van der Smissen (MHC).

Distribution. *Mongolia (Arkhangai); Asiatic-European, from western Europe to Central Asia, Russia and Japan (Rosa et al. 2019).

***Chrysis viridula* Linnaeus, 1761**

Chrysis viridula Linnaeus, 1761: 415. Type unknown; Sweden (unknown) (*viridula* group).

Material examined. MONGOLIA: Tuv, 1 ♀, 100 km E of Ulaanbaatar, 20 km NE of Tereltz, Tuul River, 15–21.VII.2003, leg. JH (PRC).

Distribution. *Mongolia (Tuv); Asiatic-European, from western Europe to Central Asia, Russia, and Japan (Rosa et al. 2019).

Genus *Chrysura* Dahlbom, 1845

Chrysura Dahlbom, 1845: 6. Type species: *Chrysis austriaca* Fabricius, 1804, by subsequent designation of Bodenstein 1939: 125.

Chrysura dichroa (Dahlbom, 1854)

Chrysis dichroa Dahlbom, 1854: 146. Lectotype ♀ (designated by Rosa and Xu 2015: 17); Hungary: Budapest (MSNT) (*dichroa* group).

Material examined. MONGOLIA: Zavkhan, 1 ♀, 40 km SW of Uliastay, dunes, 18.VII.2005, leg. JH (MHC).

Distribution. *Mongolia (Zavkhan); Asiatic-European, from western Europe to Central Asia and western Siberia (Rosa et al. 2019).

Chrysura ignifrons Brullé, 1833

Chrysis ignifrons Brullé, 1833: 375. Holotype ♂ [not ♀]; Greece: Peloponnese (Paris) (examined) (*austriaca* group).

Material examined. MONGOLIA: Zavkhan, 1 ♂, 40 km SW of Uliastay, dunes, 18.VII.2005, leg. JH (MHC).

Distribution. *Mongolia (Zavkhan); Palaearctic, from southern Europe and northern Africa to Middle East and Central Asia (Rosa et al. 2019).

Genus *Euchroeus* Latreille, 1809

Euchroeus Latreille, 1809: 49. Type species: *Chrysis purpurata* Fabricius, 1787 [= *Euchroeus purpuratus* (Fabricius, 1787)], by monotypy.

Euchroeus mongolicus Tsuneki, 1947

Euchroeus purpuratus f. *mongolicus* Tsuneki, 1947: 54. Holotype ♀; China: Inner Mongolia: Apaka (NIAS).

Euchroeus (*Euchroeus*) *mongolicus*: Linsenmaier 1959: 73 (tax., descr., Mongolia [= Inner Mongolia]), 200 (fig. 213).

Spinolia (*Euchroeus*) *par* Semenov, 1967: 189 (cat., Mongolia: 1 ♀, Uburchangaj aimag: Changaj Mt., 8 km W of Somon Chajrchangulaan, 2000 m, Exp. Dr. Z. Kaszab, 1964, nr. 217, 28.VI.1964; 1 ♀, Southgobi aimag: 60 km E of Somon Bulgan, 1120 m, Exp. Dr. Z. Kaszab 1964, nr. 262, 4.VII.1964).

Brugmoia quadrata f. *mongolica*: Kimsey and Bohart 1991: 296 (cat., China [not Mongolia]: Apaka).

Euchroeus mongolicus: Rosa et al. 2014: 68 (cat.), 111 (Plate 59); Rosa et al. 2019: 326 (Mongolia, figs 83, 84).

Material examined. MONGOLIA: *Govi-Altai*, 25 ♀♀, 1 ♂, 70 km E of Altay city, Guulin, 14.VII.2005, leg. JH (MHC, PRC).

Distribution. Mongolia (**Govi-Altai*, Umnugovi, Uvurkhangai); China (Inner Mongolia, Shanxi) (Rosa et al. 2014).

Euchroeus orientis Semenov, 1910

Pseudochrysis (*Euchroeus*) *purpurata* subsp. *orientis* Semenov-Tian-Shansky, 1910: 214.

Lectotype ♂, designated by Kimsey in Kimsey & Bohart 1991: 296; China: Bugas near Khami, SE of Tian-Shan [China, Xinjiang], 3–5.IX.1895, leg. V. Roborovskij & P. Kozlov (ZIN) (examined).

Spinolia (*Euchroeus*) *orientis*: Móczár 1967: 189 (cat., Mongolia: 1 ♂, Suchebaator [= Sukhbaatar] aimag: Ongon elis, 10 km S of Somon Chongor, 900 m, Exp. Dr. Z. Kaszab, 1965, nr. 356, 3.–4.VIII.1965; 1 ♂, 44 km SSW of Baruun urt, 1050 m, Exp. Dr. Z. Kaszab, 1965, nr. 349, 2.–3.VIII.1965).

Euchroeus (*Euchroeus*) *purpuratus* *orientis*: Linsenmaier 1968: 46 (descr., Mongolia, observed in the collections of HNHM and MHNH, without precise localities).

Material examined. None examined.

Distribution. Mongolia (Sukhbaatar); China (Xinjiang) (Rosa et al. 2014).

Genus *Pentachrysis* Lichtenstein, 1876

Pentachrysis Lichtenstein, 1876: 227. Type species: *Chrysis amoena* Eversmann 1858 [= *Pentachrysis amoena* (Eversmann, 1858)], by subsequent designation of Ashmead 1902: 226.

Pentachrysis amoena (Eversmann, 1858)

Chrysis amoena Eversmann, 1858: 562. Holotype ♀; Russian SFSR: ‘campis transalensibus’ (ISEA-PAS) (examined).

Pentachrysis amoena: Kimsey and Bohart 1991: 521 (Mongolia, without specific locality); Rosa et al. 2019: 197 (cat., Mongolia).

Material examined. None examined.

Distribution. Mongolia (without locality); Asiatic-European, from eastern Europe to Mongolia (Kimsey and Bohart 1991).

Genus *Pseudochrysis* Semenov, 1891

Pseudochrysis Semenov, 1891: 444. Type species: *Chrysura humboldti* Dahlbom, 1845: 6 [= *Pseudochrysis humboldti* (Dahlbom, 1845)], by subsequent designation of Semenov 1892: 485.

Pseudospinolia Linsenmaier, 1951: 65 (as subgenus of *Euchroeus* Latreille, 1809). Type species: *Chrysura uniformis* Dahlbom, 1854: 149, by original designation. Synonymized by Rosa et al. 2017b

Pseudochrysis gengiskhan Rosa, 2017

Pseudochrysis gengiskhan Rosa in Rosa et al. 2017c: 9. Holotype ♀; Mongolia: Övörkhangay [Bulgan], 137 km NE of Aravaykheer, 47°20'N, 103°40.5'E, 1250 m, 26.vii.2004, leg. J. Halada (ZIN) (examined). Rosa et al. 2019: 198 (cat., Mongolia), 328 (fig. 91).

Material examined. MONGOLIA: *Bayankhongor*, 1 ♀, 1 ♂, 129 km NW of Bayankhongor, 47°13'N, 99°55'E, 2590 m, 16.VII.2004, leg. JH (MHC); *Bulgan*, 4 ♀♀, Mongolia, 137 km NE of Aravaykheer, 47°20'N, 103°40.5'E, 1250 m, 26.VII.2004, leg. JH (PRC/ZIN); 1 ♂, Mongol Els Nat. Res., dunes, 47°24'N, 103°39'E, 31.VII.2005, leg. JH (MHC); *Dornod*, 2 ♂♂, 100 km W of Choibalsan, 820 m, 23.VII.2007, leg. MH (MHC); 3 ♂♂, 20 km W of Choibalsan, 48°01'N, 114°14'E, 800 m, 24.VII.2007, leg. MH (MHC); 2 ♂♂, 50 km SW of Choibalsan, 960 m, 25.VII.2007, leg. JH (MHC); *Selenge*, 2 ♂♂, 90 km N of Ulaanbaatar, Segnez River, 1450 m, 6–8.VII. 2003, leg. JH (MHC); *Sukhbaatar*, 1 ♂, 200 km SSE of Baruun-Urt, Moltsoy Els, 1250 m, 27.VII.2007, Allotype, leg. MK (ZIN); 3 ♂♂, ibid, 27.VII.2007, leg. MH (MHC); 2 ♀♀, 7 ♂♂, 100 km SSW of Baruun-Urt, 1100 m, 30.VII.2007, leg. MH (MHC); *Tuv*, 4 ♂♂, Khangay Mts, 5 km N of Khunt, 20.VII.2005, leg. JH (MHC); 4 ♂♂, 75 km W of Ulaanbaatar, dunes, 2.VIII.2005, leg. JH (MHC); 13 ♀♀, 50 km N of Ulaanbaatar, E of Mandal, 1180 m, 8–13.VIII.2007, leg. JH (MHC); 8 ♀♀, ibid, leg. MH (MHC).

Distribution. Mongolia (**Bayankhongor*, **Dornod*, **Selenge*, *Sukhbaatar*, **Tuv*, *Bulgan*); Russia (Siberia) (Rosa et al. 2017c).

Pseudochrysis neglecta (Shuckard, 1837)

Chrysura neglecta Shuckard, 1837: 169. Lectotype ♀ (designated by Morgan 1984: 9); England (LSL).

Material examined. MONGOLIA: *Tuv*, 1 ♂, 50 km E of Ulaanbaatar, Tuul River, 22.VI.2003, leg. JH (MHC).

Distribution. *Mongolia (Tuv); Holarctic: from west Europe to Turkey, Siberia, Manchuria and Russian Far East (Rosa et al. 2019); North America (Bohart and Kimsey 1982).

Genus *Spinolia* Dahlbom, 1854

Spinolia Dahlbom, 1854: 363. Type species: *Spinolia magnifica* Dahlbom, 1854 [= *Spinolia lamprosoma* (Förster, 1853)], by monotypy.

Spinolia spinosa Rosa & Halada, sp. nov.

<http://zoobank.org/A105F4B1-87F4-4005-B1A0-09844A7247B0>

Figures 5A, D, 6A, D

Type material. Holotype: ♀, MONGOLIA: *Bayankhongor*, Edringiyn-Nuru Ridge, 100 km SSW of Bayan-Under, 5.IX.1970, leg. V. Zaitzev (ZIN).

Diagnosis. *Spinolia spinosa* sp. nov. is closely related to Central Asian species of the *unicolor* group, which includes *S. chalcites* (Mocsáry, 1890), *S. rusalka* (Semenov, 1901), *S. hedychroides* (Bingham, 1903) and other small species so far considered synonyms of *S. chalcites* (Kimsey and Bohart 1991). *S. spinosa* sp. nov. female can be easily separated from all these species by: lateral pronotal area and propleuron ventrally V-shaped carinate, displaying two teeth in lateral view (Fig. 5D) (vs. unmodified in other species); mesopleuron with large and deep scrobal sulcus subtended by large projecting subrectangular carina (Fig. 5D) (vs. U-shaped carina); sparse, deep and large punctures on mesosoma (Fig. 6D), and sparse and deep punctures on metasoma (vs. punctuation with dense, shallow and tiny punctures on mesosoma, denser and shallower on metasoma); antennae yellowish, distinctly elongate (Fig. 5A) (vs. black to dark brown, with short to very short flagellomeres); head, in frontal view, transversely subrectangular (Fig. 6A) and not triangular (Fig. 6B); with bulging eyes, similarly to *S. unicolor*. It is additionally separated from *S. unicolor* by punctuation, elongate and yellowish antennae and bronze body colour (entirely blue body in *S. unicolor*, with shortened, blackish flagellomeres).

Description. Female. Body length 6.0 mm. Fore wing length 3.8 mm. OOL = 2.3 MOD; POL = 1.9 MOD; MS = 0.7 MOD; relative length of P:F1:F2:F3 = 1.0:1.4:1.0:0.8; subantennal space: 1.4 MOD. **Head.** Vertex with deep and contiguous punctures, as large as 0.25 MOD; vertex moderately depressed and impunctate in front of anterior ocellus and impunctate laterad of posterior ocelli; median anterior depression developed to upper scapal basin; TFC faint; frons continuous, without two flattened or concave, striate areas; scapal basin almost flat, laterally densely micro-punctate, medially with contiguous punctures forming transverse winkles (Fig. 6A); lower part of scapal basin medially impunctate and sulcate; apex of clypeus discoloured, W-shaped and bent under, medially the folded part measures 0.6 MOD. Malar space very




Figure 5. *Spinolia* species, females **A** *S. spinosa* sp. nov., dorsal view **B** *S. chalcites*, dorsal view **C** *S. unicolor*, dorsal view **D** *Spinolia spinosa* sp. nov., lateral view: arrows pointing at pronotal and propleural spines. Scale bars: 1.0 mm.

short, distinctly less than 1 MOD. Antennae elongate, with flagellomeres as long as $1.5 \times$ their width. Mouth parts elongate (as long as $0.8 \times$ head length) and evidently protruding from oral fossa. **Mesosoma.** Pronotal groove barely visible; anterolateral corner of the pronotum projected to form an acute humeral angle (Fig. 5A); lateral pronotal area ventrally V-shaped carinate forming an acute tooth (Fig. 5D); propleuron ventrally carinate in a large V-shaped tooth (Fig. 5D). Mesosoma punctation dorsally with large, spaced punctures; interspaces medially polished, laterally micro-punctate; notauli incomplete, visible and deep only basally towards the transscutal fissure; parapsidal furrows fully visible; mesopleuron with a large subrectangular area subtended the mesepimeron + mesepisternum; posterior propodeal projections narrow, acute and downward directed. Wing venation unmodified, with long Rs bending slightly away from costal margin, leaving marginal cell broadly open. **Metasoma.** Punctuation on T1 with tiny, sparse punctures (separated by 1–4 PD) (Fig. 6D), laterally micro-punctate on interspaces; T2 with larger and deeper punctures, anterodorsally denser (0.1–2 PD), laterally micro-punctate on interspaces; T3 with coarse to contiguous small punctures;




Figure 6. *Spinolia* species, females, head in frontal view (A–C), and metasoma in dorsal view (D–F). **A** *S. spinosa* sp. nov. **B** *S. chalcites* **C** *S. unicolor*. **D** *S. spinosa* sp. nov. **E** *S. chalcites* **F** *S. unicolor*. Scale bars: 1.0 mm.

T3 pit row barely sunken, with small, round pits, equally spaced; posterior pit row area almost polished, with a few, sparse, tiny punctures; T3 with two lateral angles and fully bordered by hyaline margin. Metasomal invaginated T5, T6, and S5 with several dorsal and lateral lobes. S2 black spots oval, transversally placed and separated 0.5 MOD each other. **Colouration.** Body coppery-bronze, darker to black on median area of mesoscutum; ventrally golden to copper; tegulae golden to non-metallic yellowish on outer margin; tarsi dark brown. Mandible brown, lighter medially. Scape and pedicel coppery, antennomeres yellowish-orange, darker on distal segments. Legs pale coloured, with slight metallic reflections, with non-metallic proximal and distal joints; tarsi yellowish. Forewings hyaline, slightly amber, with light brown veins. **Vestiture.** Whitish, short and sparse setae on head and mesosoma (up to 1.5 MOD long); face with short whitish setae (less than 1.0 MOD); metasoma with short (less than 1.0 MOD) whitish, sparse setae on T3 and ventrally on S2 and S3 and femora.

Male. Unknown.

Etymology. The specific epithet *spinosa* (feminine) is derived from the Latin adjective *spinosus* (thorny) for the long and acute teeth ventrally displayed on pronotum and propleuron and clearly visible in lateral view (Fig. 5D).

Distribution. Mongolia (Bayankhongor).

Spinolia unicolor (Dahlbom, 1831)

Chrysis unicolor Dahlbom, 1831: 32. Syntypes ♂♂; Sweden: Scania: Lomma and Käflinge [= Kävlinge] (ZMUL) (examined).

Spinolia unicolor: Kimsey and Bohart 1991: 552 (cat., Mongolia, without locality).

Material examined. MONGOLIA: *Govi-Altai*, 1 ♂, 70 km E of Altay city, Guulin, 14.VII.2005, leg. JH (MHC).

Distribution. Mongolia (**Govi-Altai*); Asiatic-European: from eastern Europe to Mongolia (Linsenmaier 1959, Kimsey and Bohart 1991).

Genus *Stilbum* Spinola, 1806

Stilbum Spinola, 1806: 9. Type species: *Chrysis calens* Fabricius, 1781, by subsequent designation of Latreille 1810: 437.

Stilbum calens (Fabricius, 1781)

Chrysis calens Fabricius, 1781: 455. Holotype ♀; Russia: Siberia (NHMUK).

Stilbum calens zimmermanni Linsenmaier, 1959: Linsenmaier 1997b: 132 (China, Inner Mongolia [not Mongolia]).

Material examined. MONGOLIA: *Arkhangai*, 9 ♀♀, 25 km NE of Tsetserleg, 47°38'N, 101°45'E, 23.VII.2004, leg. JH (MHC); *Bulgan*, 1 ♀, Mongol Els Nat. Res., dunes, 47°24'N, 103°39'E, 31.VII.2005, leg. JH (MHC); *Dornogovi*, 1 ♂, 28 km SE of Chantan-Bulag, steppe, 3.VIII.2007, leg. MK (PRC); *Tuv*, 2 ♀♀, 1 ♂, Khangayn Mts, 5 km N of Khunt, 20.VII.2005, leg. JH (MHC); 1 ♂, 75 km W of Ulaanbaatar, dunes, 2.VIII.2005, leg. JH (MHC); *Uvurkhangai*, 3 ♀♀, 12 km E of Aravaykheer, 46°22'N, 102°49'E, 1800 m, 3.VII.2004, leg. JH (MHC).

Distribution. *Mongolia (Arkhangai, Bulgan, Dornogovi, Tuv, Uvurkhangai); widely distributed in the Palaearctic Region (Tsuneki 1948, Linsenmaier 1959), Russia (Siberia), China (Liaoning, Beijing, Inner Mongolia, Shanxi) (Rosa et al. 2019). Linsenmaier (1997b) mentioned Mongolia in the distribution range of *Stilbum calens*, yet the specimen examined by the Swiss author was collected in China, Inner Mongolia: 1 ♂, Hujertu Gol [currently Khujirt Gol River, near Bailingmiaozhen monastery, N of Baotou, Inner Mongolia, China] 1927, Sven Hedins Exp. Ctr. Asien Dr. Hummel, det. Linsenmaier 1963 (NMLS).

Genus *Trichrysis* Lichtenstein, 1876

Trichrysis Lichtenstein, 1876: 27. Type species: *Sphex cyanea* Linnaeus, 1758 [= *Trichrysis cyanea* (Linnaeus, 1758)], by monotypy.

Trichrysis cyanea (Linnaeus, 1758)

Sphex cyanea Linnaeus, 1758: 572. Lectotype ♂ (designated by Morgan 1984: 10); Europe (LSL).

Material examined. MONGOLIA: Bayankhongor, 1 ♀, 16 km SW of Bayankhongor, 46°13'N, 100°30'E, 2165 m, 10.VII.2004, leg. JH (MHC); Selenge, 1 ♀, 90 km N of Ulaanbaatar, Segnez River, 1450 m, 6–8.VII.2003, leg. JH (MHC); Tuv, 1 ♀, 100 km E of Ulaanbaatar, 20 km NE Tereltz, Tuul River, 15–21.VII.2003, leg. JH (MHC); 2 ♀♀, 50 km N of Ulaanbaatar, E of Mandal, 1180 m, 8–13.VIII.2007, leg. MH (MHC).

Distribution. *Mongolia (Bayankhongor, Selenge, Tuv); Palaearctic, from western Europe and northern Africa to Central Asia, China and Japan (Rosa et al. 2019).

Trichrysis pellucida (du Buysson, 1887)

Chrysis pellucida du Buysson, 1887: 183. Lectotype ♀ (designated by Rosa et al. 2016: 123); China (MNHN) (examined).

Chrysis (Trichrysis) buyssoni Mocsáry, 1889: 323. Unnecessary replacement name for *Chrysis pellucida* du Buysson, 1887.

Chrysis (Trichrysis) mongolica Mocsáry, 1914: 24. Lectotype ♀ (designated by Bohart in Kimsey and Bohart 1991: 571); Mongolia (HMNH).

Material examined. MONGOLIA: 1 ♀, *mongolica* Mocs. typ. det. Mocsáry, red label, *Chrysis* L. *pellucida* Buyss. Linsenmaier det. 59, Lectotype *Chrysis mongolica* Mocs. ♀ RM Bohart, id nr. 135554 HNHM Hym. coll. Paralectotypes: 4 ♀♀, Mongolia, *mongolica* Mocs. typ. det. Mocsáry, red label, *Chrysis* L. *pellucida* Buyss. Linsenmaier det. 59, Paralectotype *Chrysis mongolica* Mocs. ♀ RM Bohart, id nr. 135555–135558 HNHM Hym. coll.

Distribution. Mongolia (without locality); East-Palaearctic: Russia (Far East), China (Liaoning, Inner Mongolia, Hebei, Beijing, Hunan) (Rosa et al. 2016, 2019).

Trichrysis secernenda (Mocsáry, 1912)

Chrysis (Trichrysis) secernenda Mocsáry, 1912: 376. Lectotype ♂ (designated by Bohart in Bohart and French 1986: 342); Uzbekistan: Gouldsha (HNHM) (examined).

Material examined. MONGOLIA: Selenge, 1 ♂, 90 km N of Ulaanbaatar, Segnez River, 1450 m, 6–8.VII.2003, leg. JH (MHC).

Distribution. *Mongolia (Selenge); Afghanistan, Uzbekistan, China (Xinjiang, Ningxia) (Rosa et al. 2016).

Tribe Elampini

Genus *Colpopyga* Semenov, 1954

Colpopyga Semenov, 1954: 137. Type species: *Hedychrum flavipes* Eversmann, 1858 [= *Colpopyga flavipes* (Eversmann, 1858)], by original designation.

Colpopyga nesterovi Rosa, 2017

Colpopyga nesterovi Rosa, 2017b: 301. Holotype ♀; Kazakhstan: Aktobe Prov., Mughodzhary Mt., Emba River valley, 17.vi.1985, leg. M. Nesterov (ZIN) (examined).

Material examined. MONGOLIA: Dornod, 2 ♀♀, 1 ♂, 20 km W of Choilbalsan, 800 m, 48°01'N, 114°14'E, 24.VII.2007, leg. MH (MHC).

Distribution. *Mongolia (Dornod); Kazakhstan (Rosa 2017b).

Genus *Elampus* Spinola, 1806

Elampus Spinola, 1806: 10. Type species: *Chrysis panzeri* Fabricius, 1804 [= *Elampus panzeri* (Fabricius, 1804)], by subsequent designation of Latreille 1810: 437.

Ellampus Agassiz, 1846: 136. Unjustified emendation of *Elampus* Spinola, 1806 (part.).

Notozus Förster, 1853: 351. Type species: *Notozus frivaldszkii* Förster, 1853 [= *Elampus spina* (Lepeletier, 1806)], by subsequent designation of Ashmead 1902: 228.

Elampus albipennis (Mocsáry, 1889)

Ellampus (Notozus) albipennis Mocsáry, 1889: 80. Lectotype ♂ (designated by Móczár 1964: 447); Russia: Astrakhan (HMNH) (examined).

Material examined. MONGOLIA: Arkhangai, 93 ♀♀, 30 ♂♂, Chuluut Gol River, 47°48'N, 100°19'E, 23.VII.2005, leg. JH (MHC); Dornogovi, 1 ♂, Orgon, 11.VII.2005, leg. JH (MHC).

Distribution. *Mongolia (Arkhangai, Dornogovi); Asiatic-European, from eastern Europe, Saudi Arabia, UAE to Central Asia and eastern Siberia (Rosa et al. 2019).

Elampus coloratus Rosa, 2017

Elampus coloratus Rosa in Rosa et al. 2017d: 2. Holotype ♂; Russia: Tyva Rep., 20 km SSW of Erzin, Tore-Khol' Lake (ZIN) (examined).

Material examined. MONGOLIA: *Sukhbaatar*, 1 ♂, Lun-Ula, 30 km WSW of Dariganga, 1.VII.1971, leg. I. Kerzhner (ZIN).

Distribution. Mongolia (Sukhbaatar); Russia (Tyva Rep.) (Rosa et al. 2017d).

Elampus montanus (Mocsáry, 1890)

Ellampus (Notozus) montanus Mocsáry, 1890: 49. Holotype ♂; Turkey: Buyuk Agri Dagi (Mount Ararat) (ISEA-PAS) (examined).

Material examined. MONGOLIA: *Dornogovi*, 6 ♂♂, Orgon, 11.VII.2005, leg. JH (MHC).

Distribution. *Mongolia (Dornogovi); Turkey (Mocsáry 1890) and Central Asia (unpubl. data).

Elampus panzeri (Fabricius, 1804)

Chrysia scutellaris Panzer, 1798: fig. 51, pl. 11. Type unknown; Germany: Nürnberg (depository unknown), nom. praeocc., nec Fabricius, 1794.

Chrysia panzeri Fabricius, 1804: 172. Replacement name for *Chrysia scutellaris* Panzer, 1798.

Material examined. MONGOLIA: *Arkhangai*, 1 ♂, 25 km NE of Tsetserleg, 47°38'N, 101°45'E, 23.VII.2004, leg MK (MHC); *Zavkhan*, 1 ♀, 2 ♂♂, 40 km SW of Uliastay, dunes, 18.VII.2005, leg. JH (MHC).

Distribution. *Mongolia (Arkhangai, Zavkhan); Asiatic-European, from western Europe to eastern Siberia, Russian Far East, and China (Heilongjiang) (Rosa et al. 2019).

Elampus sanzii Gogorza, 1887

Elampus sanzii Gogorza, 1887: 33. Holotype ♂; Spain: Madrid (MCNM).

Notozus sanzii: Móczár 1967: 183 (cat., Mongolia: Central aimag, Zuun-Chara, 850 m, Exp. Dr. Z. Kaszab, 1964, Nr. 281; 8.VII.1964).

Material examined. None examined.

Distribution. Mongolia (Tuv); Asiatic-European, from Iberian Peninsula to Mongolia and eastern Siberia (Rosa et al. 2019).

Elampus spinifemoris (Móczár, 1967)

Notozus spinifemoris Móczár, 1967: 185. Holotype ♀; Mongolia: Uvurkhangai aimag: Arc Bogd ul, ca. 20 km S of von Sonon Chovd, 1760 m, Exp. Dr. Z. Kaszab,

1964, 22.VI.1964 (HNHM) (examined). Rosa et al. 2017e: 113 (cat., typ., Mongolia: Arc Bogd ul, fig. 87).

Material examined. MONGOLIA: *Uvurkhangai*, 1 ♀, Uburchangaj aimag, Arc Bogd ul, cca 20 km S of von somon Chovd, 1760 m Exp. Dr. Z. Kaszab, 1964, Nr. 170, 22.VI.1964, *Notozus* sp. nov. ♀ det. Móczár 965, ♀ *Omalus* Pz. *Notozus panzeri* F. Linsenmaier det. 1964, Holotype ♀ *Notozus spinifemoris* L. Móczár 1966, Hym. Typ. No. 87 Mus. Budapest, id nr. 134892 HNHM Hym. coll. (HNHM).

Distribution. Mongolia (Uvurkhangai) (Móczár 1967).

Genus *Hedychridium* Latreille, 1802

Hedychridium Abeille de Perrin, 1878: 3. Type species: *Hedychrum minutum* Lepeletier, 1806 [= *Hedychridium ardens* (Coquebert, 1801)], by subsequent designation of Ashmead 1902: 227.

Hedychridium ardens (Coquebert, 1801)

Chrysis ardens Coquebert, 1801: 59. Holotype ♀; France: Bordeaux (MNHN?).

Hedychridium ardens: Móczár 1967: 189 (cat., Mongolia: Uvurkhangai aimag: Baga Bogd ul, between Somon Bogd and Somon Baruun Hajan-ulaan, 1900 m, Exp. Dr. Z. Kaszab, 1964, nr. 176, 23.VI.1964).

Material examined. MONGOLIA: *Arkhangai*, 1 ♀, 90 km NE of Tsetserleg, 48°03'N, 102°25'E, 24.VII.2004, leg. JH (MHC); *Bayankhongor*, 9 ♀♀, 8 ♂♂, 16 km SW of Bayankhongor, 46°13'N, 100°30'E, 2165 m, 10.VII.2004, leg. JH (MHC); *Dornod*, 2 ♀♀, 3 ♂♂, 100 km W of Choilbalsan, 820 m, 23.VII.2007, leg. MH (MHC); 5 ♀♀, 2 ♂♂, 20 km W of Choilbalsan, 800 m, 48°01'N, 114°14'E, 24.VII.2007, leg. MH (MHC); *Khentii*, 1 ♀, 100 km NE of Ondorkhaan, Kerulen River, 970 m, 22.VII.2007, leg. MH (MHC); *Selenge*, 2 ♀♀, 90 km N of Ulaanbaatar, Segnez River, 1450 m, 6–8.VII.2003, leg. JH (MHC); *Sukhbaatar*, 1 ♀, 1 ♂, 100 km SSW of Baruun-Urt, 1100 m, 30.VII.2007, leg. MH (MHC); *Ulaanbaatar*, 7 ♀♀, Ulaanbaatar, Tuul River valley, 12.VII.2003, leg. JH (MHC); *Zavkhan*, 2 ♂♂, 40 km SW of Uliastay, dunes, 18.VII.2005, leg. JH (MHC).

Distribution. Mongolia (**Arkhangai*, **Bayankhongor*, **Dornod*, **Khentii*, **Selenge*, **Sukhbaatar*, **Ulaanbaatar*, Uvurkhangai, **Zavkhan*); Asiatic-European, from Europe and Middle East to Russia (Far East) (Kimsey and Bohart 1991; Kurzenko and Lelej 2007).

Hedychridium asianum Linsenmaier, 1997

Hedychridium integrum ssp. *asianum* Linsenmaier, 1997a: 254. Holotype ♂, Mongolia: Ulan Bator, 1900 m (UKC).

Material examined. MONGOLIA: Arkhangai, 1 ♂, 90 km NE of Tsetserleg, 48°03'N, 102°25'E, 24.VII.2004, leg. JH (MHC); Bayankhongor, 1 ♂, 16 km SW of Bayankhongor, 46°13'N, 100°30'E, 2165 m, 10.VII.2004, leg. JH (MHC); Bulgan, 3 ♂♂, 137 km NE of Aravaykheer, 47°20'N, 103°45.5'E, 1250 m, 2.VII.2004, leg. JS (MHC); 2 ♀♀, 2 ♂♂, 143 km NE of Aravaykheer, 47°24'N, 103°39'E, 1300 m, 26.VII.2004, leg. MH (MHC); Ulaanbaatar, 1 ♂, Ulaanbaatar, 16.VII.1989, 1900 m, leg. Peter Salk, det. Linsenmaier, 1997 (NMLS).

Distribution. Mongolia (*Arkhangai, *Bayankhongor, *Bulgan, Ulaanbaatar); China (Gansu) (Rosa et al. 2014).

Remarks. *Hedychridium asianum* was described as a subspecies of *H. integrum*. As recently pointed out by Paukkunen et al. (2014), *H. integrum* is a synonym of *H. ardens* and *H. integrum* sensu Linsenmaier (1959) is *H. cupreum*.

Hedychridium belokobylskiji Rosa, 2017

Hedychridium belokobylskiji Rosa in Rosa et al. 2017d: 11. Holotype ♀; Russia: Eastern Siberia, Tuva Rep., 12 km SW of Samagaltau, Dyttyg-Khem River, 19.VII.2014, leg. A. Lelej, M. Proshchalykin, V. Loktionov (ZIN) (examined).

Material examined. MONGOLIA: Tuv, 1 ♀, 50 km N of Ulaanbaatar, E of Mandal, 1180 m, 8–13.VIII.2007, leg. MK (PRC).

Distribution. *Mongolia (Tuv); Russia (Eastern Siberia) (Rosa et al. 2019).

Hedychridium cupreum (Dahlbom, 1845)

Hedychrum cupreum Dahlbom, 1845: 3. Lectotype ♀ (designated by Paukkunen et al. 2014: 23); Sweden: Lund (NHMW) (examined).

Material examined. MONGOLIA: Bayankhongor, 2 ♀♀, 16 km SW of Bayankhongor, 46°13'N, 100°30'E, 2165 m, 10.VII.2004, leg. JH (MHC); Dornogovi, 1 ♂, 65 km SE of Chatan-Bulag, 1020 m, 2.VIII.2007, leg. MH (MHC); Govi-Altai, 4 ♀♀, 70 km E of Altay City, Guulin, 14.VII.2005, leg. JH (MHC); Tuv, 2 ♀♀, 1 ♂, 50 km E of Ulaanbaatar, Tuul River, 22.VI.2003, leg. JH (MHC); 1 ♀, Khangaun Mts, 5 km N of Khunt, 20.VII.2005, leg. JH (MHC); Umnugovi, 3 ♀♀, Gobi, 100 km SW of Dalanzadgad, Bayanzag, 1–2.VII.2003, leg. JH (MHC); Uvurkhangai, 1 ♂,

12 km E of Aravaykheer, 46°22'N, 102°49'E, 1800 m, 3.VII.2004, leg. JH (MHC); *Zavkhan*, 2 ♀♀, 1 ♂, 40 km SW of Uliastay, dunes, 18.VII.2005, leg. JH (MHC).

Distribution. *Mongolia (Bayankhongor, Dornogovi, Govi-Altai, Tuv, Umnu-govi, Uvurkhangai, Zavkhan); Asiatic-European, from north-western Europe to Mon-golia and China (Rosa et al. 2014).

Remarks. Specimens from Mongolia display an unusual red colouration.

Hedychridium gabriellae Rosa, 2017

Hedychridium gabriellae Rosa in Rosa et al. 2017d: 19. Holotype ♀; Russia: Eastern Siberia, Tuva Rep., 20 km of SSW Erzin, Tore-Khol' Lake, 30.VI–3.VII.2013, leg. V. Loktionov & M. Proshchalykin (ZIN) (examined).

Material examined. MONGOLIA: *Bayankhongor*, 18 ♂♂, 22 ♀♀, 75 km S of Bay-ankhongor, 45°20'N, 100°48.5'E, 1330 m, 8.VII.2004, leg. JH, JS (MHC); *Dor-nogovi*, 5 ♀♀, 5 ♂♂, 65 km SE of Chatan-Bulag, 1020 m, 2.VIII.2007, leg. MH (MHC); *Tuv*, 1 ♂, 70 km W of Ulaanbaatar, 1070 m, dunes, 16.VIII.2007, leg. MH (MHC).

Distribution. *Mongolia (Bayankhongor, Dornogovi, Tuv); Russia (Eastern Sibe-ria) (Rosa et al. 2019).

Hedychridium longigena Rosa, 2017

Hedychridium longigena Rosa in Rosa et al. 2017d: 21. Holotype ♀; Russia: Irkutsk Prov., 8 km N of Irkutsk, Angara River, sandy slopes, 10.VII.2001, collector un-known (ZIN) (examined).

Material examined. MONGOLIA: *Bayankhongor*, 1 ♀, 1 ♂, 56 km NW of Bay-ankhongor, 46°33'N, 100°12'E, 2200 m, 12.VII.2004, leg. JS (MHC); *Bulgan*, 2 ♀♀, 137 km NE of Aravaykheer, 47°20'N, 103°45.5'E, 1250 m, 2.VII.2004, leg. MK (MHC); 3 ♀♀, ibid, 26.VII.2004, JH (MHC); *Dornod*, 1 ♂, 100 km W of Choibalsan, 820 m, 23.VII.2007, leg. MH (MHC); *Dornogovi*, 1 ♂, 65 km SE of Chatan-Bulag, 1020 m, 2.VIII.2007, leg. MH (MHC); *Khentii*, 2 ♀♀, 2 ♂♂, 100 km NE of Ondorkhaan, Kerulen River, 970 m, 22.VII.2007, leg. MH (MHC); *Selenge*, 1 ♀, 90 km N of Ulaanbaatar, Segnez River, 1450 m, 6–8.VII.2003, leg. JH (MHC); *Tuv*, 10 ♀♀, 9 ♂♂, 50 km N of Ulaanbaatar, E of Mandal, 1180 m, 8–13.VIII.2007, leg. MK (PRC); 4 ♂♂, 70 km W of Ulaanbaatar, 1070 m, dunes, 16.VIII.2007, leg. MH (MHC).

Distribution. *Mongolia (Bayankhongor, Bulgan, Dornod, Dornogovi, Khentii, Selenge, Tuv); Russia (Eastern Siberia) (Rosa et al. 2019).

Hedychridium propodeale Rosa, 2017

Hedychridium propodeale Rosa in Rosa et al. 2017d: 16. Holotype ♀; Russia: Eastern Siberia, Tuva Rep., 20 km SSW of Erzin, Tore-Khol' Lake, 3.VII.2013, leg. V. Loktionov & M. Proshchalykin (ZIN) (examined).

Material examined. MONGOLIA: *Govi-Altai*, 1 ♀, Mongolia W, 70 km E of Altay city, Guulin, 14.VII.2005, leg. JH (MHC).

Distribution. *Mongolia (Govi-Altai); Russia (Eastern Siberia) (Rosa et al. 2019).

Hedychridium roseum (Rossi, 1790)

Chrysis carnea var. *rosea* Rossi, 1790: 75. Syntypes; Italy (Berlin?).

Material examined. Mongolia: *Arkhangai*, 1 ♂, 90 km NE of Tsetserleg, 48°03'N, 102°25'E, 24.7.2004, leg. JH (MHC); *Dornod*, 21 ♀♀, 4 ♂♂, 100 km W of Choilbalsan, 820 m, 23.VII.2007, leg. MH (MHC); 19 ♀♀, 1 ♂, 20 km W of Choilbalsan, 800 m, 48°01'N, 114°14'E, 24.7.2007, leg. MH (MHC); *Dornogovi*, 4 ♂♂, 2 km SE of Khuvgol, 5.VIII.2007, leg. MH (MHC); *Sukhbaatar*, 1 ♀, 1 ♂, 100 km SSW of Baruun-Urt, 1100 m, 30.VII.2007, leg. MH (MHC); 4 ♂♂.

Distribution. *Mongolia (Arkhangai, Dornod, Dornogovi, Sukhbaatar); Asiatic-European, from western Europe to Russia (Far East) (Rosa et al. 2019).

Remarks. The record from Korea (Tsuneki 1953b and Korean checklists) must be referred to *Hedychridium tsunekii* Lisenmaier, 1959. In fact, Lisenmaier (1959) described as *Hedychridium tsunekii* the Korean specimens collected and identified by Tsuneki (1953) as *H. roseum*.

Genus *Hedychrum* Latreille, 1802

Hedychrum Latreille, 1802: 317. Type species: *Chrysis lucidula* Fabricius, 1775 [= *Hedychrum nobile* (Scopoli, 1763)], by monotypy.

Hedychrum chalybaeum Dahlbom, 1854

Hedychrum chalybaeum Dahlbom, 1854: 64. Syntypes ♂♂; Europe: 'Europa media et meridionali', Russia, Prussia, Silesia (MfN, ZMUL) (examined). Móczár 1967: 188 (cat., Mongolia: 1 ♀, Sukhbaatar aimag: 44 km SSW of Baruun urt, 1050 m, Exp. Dr. Z. Kaszab, 1965, nr. 349, 2.-3.VIII.1965; 1 ♀, Chadatin-bulan, 60 km N of Somon Bajanterem, 950 m, Exp. Dr. Z. Kaszab, 1965, nr. 340, 31.VII.1965).

Material examined. MONGOLIA: *Bayankhongor*, 1 ♂, 2 km S of Bayankhongor, 46°12'N, 100°43'E, 1800 m, 10.VII.2004, leg. JH (PRC); 20 ♀♀, 2 ♂♂, 56 km NW of Bayankhongor, 46°33'N, 100°12'E, 2200 m, 12.VII.2004, leg. JH (PRC); *Dornod*, 16 ♀♀, 9 ♂♂, 100 km W of Choibalsan, 820 m, 23.VII.2007, leg. MH (MHC); 16 ♀♀, 6 ♂♂, 20 km W of Choibalsan, 800 m, 48°01'N, 114°14'E 24.VII.2007, leg. MH (MHC); 1 ♂, 15 km W of Choibalsan, Kerulen River, 770 m, 24.VII.2007, leg. MK (PRC); 1 ♂, 50 km SW of Choibalsan, 960 m, 25.VII.2007, leg. JH (MHC); 1 ♀, 50 km SW of Choibalsan, 960 m, 25.VII.2007, leg. MH (MHC); *Govi-Alтай*, 2 ♂♂, 70 km E of Altay city, Guulin, 14.VII.2005, leg. JH (MHC); *Selenge*, 1 ♂, 90 km N of Ulaanbaatar, Segnez River, 1450 m, 6–8.VII.2003, leg. JH (PRC); *Sukhbaatar*, 6 ♀♀, 2 ♂♂, 100 km SSW of Baruun-Urt, 1100 m, 30.VII.2007, leg. MH (MHC); *Tuv*, 1 ♂, 50 km E of Ulaanbaatar, Tuul River, 22.VI.2003, leg. JH (MHC); *Ulaanbaatar*, 1 ♂, Ulaanbaatar, Tuul River valley, 12.VII.2003, leg. JH (PRC); 1 ♀, 7 km E of Ulaanbaatar, Gachuurt, 47°55'N, 107°06'E, 31.VII.2002, 1310 m, leg. JS (PRC).

Distribution. Mongolia (Bayankhongor, *Dornod, *Govi-Alтай, *Selenge, Sukhbaatar, Tuv, *Ulaanbaatar); widely distributed in the Palaearctic Region (Linsenmaier 1959; Kurzenko and Lelej 2007), China (Heilongjiang, Inner Mongolia, Gansu) (Rosa et al. 2014).

Hedychrum gerstaeckeri Chevrier, 1869

Hedychrum gerstaeckeri Chevrier, 1869: 47. Syntypes ♀♀, ♂♂, [not holotype]; Switzerland (Geneva) (examined).

Material examined. MONGOLIA: *Khentii*, 3 ♀♀, 100 km NE of Ondorkhaan, Kerulen River, 970 m, 22.VII.2007, leg. MH (MHC); *Selenge*, 1 ♀, 5 ♂♂, 90 km N of Ulaanbaatar, Segnez River, 1450 m, 6–8.VII.2003, leg. JH (MHC); *Tuv*, 4 ♀♀, 50 km N of Ulaanbaatar, E of Mandal, 1180 m, 8–13.VIII.2007, leg. MH (MHC).

Distribution. *Mongolia (Khentii, Selenge, Tuv); Palaearctic and Oriental region, from western Europe to Russian Far East, Japan, China and Taiwan (Rosa et al. 2014, 2019).

Hedychrum lama du Buysson, 1891

Hedychrum lama du Buysson, 1891: 31. Lectotype ♂ (designated by Kimsey and Bohart 1991: 215); Mongolia: Kansu-Kobden Owatu (MNHN).

Material examined. MONGOLIA: *Khovd*, 1 ♂, Mongolie O. Radoszkowsky, Kansu-Kobden Owatu, 12.8, Mongolie Coll. R. du Buysson 1900, Type, Museum Paris.

Distribution. Mongolia (Khovd) (du Buysson 1891).

Hedychrum longicolle Abeille de Perrin, 1877

Hedychrum longicolle Abeille de Perrin, 1877: 65. Lectotype ♀ (designated by Kimsey 1986: 108); France: Marseille, Toulon (Geneva, Paris) (examined).

Material examined. MONGOLIA: *Bulgan*, 1 ♂, 170 km W of Ulaanbaatar, dunes, 1070 m, 16.VIII.2007, leg. MH (MHC); *Dornod*, 1 ♂, 50 km SW of Choibalsan, 960 m, 25.VII.2007, leg. JH (MHC); *Sukhbaatar*, 1 ♂, 200 km SSE of Baruun-Urt, Moltsoy Els, 1250 m, 27.VII.2007, leg. MH (MHC); 2 ♂♂, 6 ♀♀, 100 km SSW of Baruun-Urt, 1100 m, 30.VII.2007, leg. MH (MHC); *Tuv*, 1 ♀, 75 km W Ulaanbaatar, dunes, 2.VIII.2005, leg. JH (MHC); *Umnugovi*, 1 ♀, Gobi Gurvansaikhan National Park, 44°00'N, 101°50'E, 10.VII.2005, leg. JH (MHC).

Distribution. *Mongolia (Bulgan, Dornod, Sukhbaatar, Tuv, Umnugovi); Palaeoarctic, from southern Europe and northern Africa, to western Asia, Siberia, and China (Rosa et al. 2019).

Hedychrum nobile (Scopoli, 1763)

Sphex nobile Scopoli, 1763: 297. Holotype ♀; Italy [not Austria] (lost).

Material examined. MONGOLIA: *Arkhangai*, 1 ♀, 25 km NE of Tsetserleg, 47°38'N, 101°45'E, 23.VII.2004, leg. JH (MHC); 4 ♂♂, 90 km NE of Tsetserleg, 48°03'N, 102°25'E, 24.VII.2004, leg. JH (MHC); 4 ♀♀, 9 ♂♂, ibid, 27.VII.2005, leg. JH (MHC); *Selenge*, 2 ♀♀, 12 ♂♂, 90 km N of Ulaanbaatar, Segnez River, 1450 m, 6–8.VII.2003, leg. JH (MHC); *Tuv*, 2 ♀♀, 50 km N of Ulaanbaatar, E of Mandal, 1180 m, 8–13.VIII.2007, leg. MH (MHC); *Zavkhan*, 3 ♂♂, 40 km SW of Uliastay, dunes, 18.VII.2005, leg. JH (MHC).

Distribution. *Mongolia (Arkhangai, Selenge, Tuv, Zavkhan); Asiatic-European, from western Europe to Siberia (Rosa et al. 2019).

Hedychrum rutilans ermak Semenov, 1967

Hedychrum intermedium ermak Semenov-Tian-Shanskij, 1967: 142. Holotype ♂; Russia: Siberia, Shira Lake [Khakass Rep.], 24.VII.1897, Yu. Wagner (ZIN) (examined). Móczár 1967: 188 (cat., Mongolia: Sukhbaatar aimag: Ongon elis, 10 km S of Somon Chongor, 900 m, Exp. Dr. Z. Kaszab, 1965, nr. 357, 3.–4.VIII.1965).

Material examined. MONGOLIA: *Arkhangai*, 1 ♀, 90 km NE of Tsetserleg, 48°03'N, 102°25'E, 24.VII.2004, leg. JH (MHC); 1 ♀, 2 ♂♂, 90 km NE of Tsetserleg, 48°03'N, 102°25'E, 27.VII.2005, leg. JH (MHC); *Dornod*, 2 ♂♂, 20 km W of Choibalsan, 48°01'N, 114°14'E, 800 m, 24.VII.2007, leg. MH (MHC); *Selenge*, 1 ♂, 90 km N of Ulaanbaatar, Segnez River, 1450 m, 6–8.VII.2003, leg. JH (MHC); *Tuv*, 1 ♀, 50 km

N of Ulaanbaatar, E of Mandal, 1180 m, 8–13.VIII.2007, leg. MK (PRC); 7 ♀♀, 3 ♂♂, ibid, 8–13.VIII.2007, leg. MH (MHC).

Distribution. Mongolia (*Arkhangai, *Dornod, *Selenge, Sukhbaatar, *Tuv); Russia (Siberia, Far East) (Rosa et al. 2019).

Genus *Holopyga* Dahlbom, 1845

Holopyga Dahlbom, 1845: 4. Type species: *Holopyga amoenula* Dahlbom, 1845, by subsequent designation of Ashmead 1902: 227.

Holopyga generosa asiatica Trautmann, 1926

Holopyga gloriosa var. *asiatica* Trautmann, 1926: 5. Holotype ♀; Turkey: İzmir prov.: Smyrna (MFN) (examined).

Material examined. MONGOLIA: Selenge, 3 ♀♀, 5 ♂♂, 90 km N of Ulaanbaatar, Segnez River, 1450 m, 6–8.VII.2003, leg. JH (MHC).

Distribution. *Mongolia (Selenge); Asiatic-European, from southern Europe to China (Rosa et al. 2019).

Holopyga kaszabi Móczár, 1967

Holopyga kaszabi Móczár, 1967: 187. Holotype ♂; Mongolia: Ostgobi aimag 40 km NW of Chara-Eireg 1150 m Exp. Dr. Z. Kaszab, 1963 (HNHM) (examined). Rosa et al. 2017e: 108 (cat., type series, Mongolia).

Material examined. MONGOLIA: Dornogovi, 1 ♂, Ostgobi aimag 40 km NW of Chara-Eireg, 1150 m Exp. Dr. Z. Kaszab, 1963, Nr. 62, 30.VI.1963, Holotype ♂ *Holopyga kaszabi* n. sp. det. Móczár 1966, Hym. Typ. No. 89, id nr. 134927 HNHM Hym. coll. (HNHM); 1 ♀, Ostgobi aimag 40 km NW of Chara-Eireg 1150 m Exp. Dr. Z. Kaszab, 1963, Nr. 62, 30.VI.1963, Allotype ♀ *Holopyga kaszabi* n. sp. det. Móczár 1966, Hym. Typ. No. 96, id nr. 134933 HNHM Hym. Coll. (HNHM). 1 ♂: Ostgobi aimag 40 km NW of Chara-Eireg 1150 m Exp. Dr. Z. Kaszab, 1963, Nr. 62, 30.VI.1963, *Holopyga* sp. n. ? <handwritten by Móczár>, ♂ Allotype *Holopyga* Dhlb. *diversicolor* Lins. Lisenmaier 1966, Paratype ♂ *Holopyga kaszabi* n. sp. det. Móczár 1966, Hym. Typ. No. 90 (HNHM); 1 ♂, Mongolia, Ostgobi aimag 40 km NW of Chara-Eireg 1150 m Exp. Dr. Z. Kaszab, 1963, Nr. 62, 30.VI.1963, Paratype ♂ *Holopyga kaszabi* n. sp. det. Móczár 1966, Hym. Typ. No. 91, id nr. 134929 HNHM Hym. coll. (HNHM); 1 ♂: Mongolia, Ostgobi aimag 40 km NW of Chara-Eireg 1150 m Exp. Dr. Z. Kaszab, 1963, Nr. 62, 30.VI.1963, Paratype ♂ *Holopyga kaszabi* n. sp. det. Móczár 1966, Hym. Typ. No. 93, id nr. 134930 HNHM Hym. coll. (HNHM); 1

♂, Mongolia, Ostgobi aimag 40 km NW of Chara-Eireg 1150 m Exp. Dr. Z. Kaszab, 1963, Nr. 62, 30.VI.1963, Paratype ♂ *Holopyga kaszabi* n. sp. det. Móczár 1966, Hym. Typ. No. 94, id nr. 134931 HNHM Hym. coll. (HNHM); 1 ♀: Mongolia, Ostgobi aimag 40 km NW of Chara-Eireg 1150 m Exp. Dr. Z. Kaszab, 1963, Nr. 62, 30.VI.63, *Holopyga gloriosa* ? *intermedia* ? det. L. Móczár, Type *Holopyga* Dhlb. *diversicolor* Lins. Linsenmaier 1966, Paratype ♀ *Holopyga kaszabi* n. sp. det. Móczár 1966, Hym. Typ. No. 95 / id nr. 134932 HNHM Hym. coll. (HNHM); 1 ♂, Mongolia, Catgobi aimag 40 Km NW of Chara-Eireg 1150 m, Exp. Dr. Z. Kaszab, 1963, Nr. 62, 30.VI.1963, Paratype (NMLS); 1 ♀, Mongolia, Catgobi aimag 40 Km NW of Chara-Eireg, 1150 m, Exp. Dr. Z. Kaszab, 1963 / Nr. 62, 30.VI.1963, Paratype (NMLS); *Umnugovi*, 1 ♂, Gobi, Dalanzadgad, 25.6.2003, leg. JH (MHC); *Uvurkhangai*, 1 ♂, 139 km SW of Aravaykheer, 45°17'N, 101°41'E, 1430 m, 4.VII.2004, leg. JS (MHC).

Distribution. Mongolia (Dornogovi, *Umnugovi, *Uvurkhangai) (Móczár 1967).

Holopyga minuma Linsenmaier, 1959

Holopyga minuma Linsenmaier, 1959a: 31. Holotype ♀; Turkey: Niğde prov.: Niğde (NMLS) (examined).

Material examined. MONGOLIA: *Dornod*, 22 ♀♀, 14 ♂♂, 100 km W of Choilbalsan, 820 m, 23.VII.2007, leg. MH (MHC); 25 ♀♀, 6 ♂♂, 20 km W of Choilbalsan, 800 m, 48°01'N, 114°14'E 24.VII.2007, leg. MH (MHC); *Sukhbaatar*, 1 ♂, 200 km SSE of Baruun-Urt, Moltsoy Els, 1250 m, 27.VII.2007, leg. MH (MHC).

Distribution. *Mongolia (Dornod, Sukhbaatar); Asiatic-European, from central Europe to eastern Siberia (Rosa et al. 2019).

Genus *Omalus* Panzer, 1801

Omalus Panzer, 1801: 13. Type species: *Chrysis aenea* Fabricius, 1787, by monotypy.

Omalus aeneus (Fabricius, 1787)

Chrysis aenea Fabricius, 1787: 284. Holotype ♀; Germany: Hala Saxonum [= Halle] (Copenhagen) (examined).

Material examined. MONGOLIA: *Tuv*, 1 ♀, 50 km E of Ulaanbaatar, Tuul River, 22.VI.2003, leg. JH (MHC); *Ulaanbaatar*, 1 ♀, Ulaanbaatar, Tuul River valley, 12.VII.2003, leg. JH (MHC).

Distribution. *Mongolia (Tuv, Ulaanbaatar); Holarctic and Oriental: from Europe and North Africa to Japan and Taiwan (Wei et al. 2014). Probably accidentally introduced to North America (Kimsey and Bohart 1991).

Omalus berezovskii (Semenov, 1932)

Ellampus (Dictenulus) berezovskii Semenov-Tian-Shanskij, 1932: 12. Holotype ♀; China: Sichuan (ZIN) (examined).

Material examined. MONGOLIA: *Ulaanbaatar*, 1 ♂, Ulaanbaatar, Tuul River valley, 12.VII.2003, leg. JH (MHC).

Distribution. *Mongolia (Ulaanbaatar); East-Palaearctic: Russia (Eastern Siberia, Far East), China (Ningxia, Sichuan) (Rosa et al. 2019).

Omalus margianus (Semenov, 1932)

Ellampus (Dictenulus) margianus Semenov-Tian-Shanskij, 1932: 15. Lectotype ♀ (designated by Kimsey 1986: 107); Turkmenistan: Imam-baba (ZIN) (examined).

Material examined. MONGOLIA: *Arkhangai*, 1 ♀, 90 km NE of Tsetserleg, 48°03'N, 102°25'E, 27.VII.2005, leg. JH (MHC); *Bayankhongor*, 1 ♀, 75 km S of Bayankhongor, 45°20'N, 100°48.5'E, 1330 m, 8.VII.2004, leg. JS (MHC); *Bulgan*, 2 ♂♂, 137 km NE of Aravaykheer, 47°20'N, 103°45.5'E, 1250 m, 2.VII.2004, leg. JH (MHC); 1 ♀, 1 ♂, 143 km NE of Aravaykheer, 47°24'N, 103°39'E, 1300 m, 26.VII.2004, leg. MH (MHC); 2 ♂♂, Mongol Els Nat. Res., dunes, 47°24'N, 103°39'E, 31.VII.2005, leg. JH (MHC); *Sukhbaatar*, 2 ♀♀, 100 km SSW of Baruun-Urt, 1100 m, 30.VII.2007, leg. MH (MHC); *Tuv*, 1 ♂, 80 km W of Ulaanbaatar, 1230 m, dunes, 17.VIII.2007, leg. MH (MHC).

Distribution. *Mongolia (Arkhangai, Bayankhongor, Bulgan, Sukhbaatar, Tuv); Central Asia (Kimsey and Bohart 1991).

Omalus miramae (Semenov, 1932)

Ellampus (Dictenulus) miramae Semenov-Tian-Shanskij, 1932: 13. Lectotype ♀ (designated by Rosa et al. 2017a: 76); Turkmenistan: Pereval (ZIN) (examined).

Material examined. MONGOLIA: *Bayankhongor*, 1 ♂, 75 km S of Bayankhongor, 45°20'N, 100°48.5'E, 1330 m, 8.VII.2004, leg. JH (MHC); *Dornogovi*, 2 ♀♀, Or-gon, 11.VII.2005, leg. JH (MHC); *Sukhbaatar*, 2 ♀♀, 200 km SSE of Baruun-Urt, Moltsoy Els, 1250 m, 27.VII.2007, leg. MH (MHC).

Distribution. *Mongolia (Bayankhongor, Dornogovi, Sukhbaatar); Central Asia (Kimsey and Bohart 1991).

***Omalus stella* (Semenov, 1932)**

Ellampus (Ellampus) stella Semenov-Tian-Shanskij and Nikol'skaya, 1954: 93. Lectotype ♀ (designated by Kimsey 1986: 107); Tajikistan: Stalinabad (currently Dushambe) (ZIN) (examined).

Material examined. MONGOLIA: Arkhangai, 1 ♀, 90 km NE of Tsetserleg, 48°03'N, 102°25'E, 24.VII.2004, leg. JH (MHC); Tuv, 3 ♂♂, 75 km W of Ulaanbaatar, dunes, 2.VIII.2005, leg. JH (MHC); Uvurkhangai, 1 ♀, 159 km of SW Aravaykheer, 45°11'N, 101°26'E, 1250 m, 5.VII.2004, leg. JH (MHC).

Distribution. *Mongolia (Arkhangai, Tuv, Uvurkhangai); Central Asia (Kimsey and Bohart 1991).

Genus *Philoctetes* Abeille de Perrin, 1879

Philoctetes Abeille de Perrin, 1879: 27. Type species: *Holopyga cicatrix* Abeille de Perrin, 1879 [= *Philoctetes micans* (Klug, 1835)], by subsequent designation of Ashmead 1902: 228.

Ellampus Agassiz, 1846: 136. Unjustified emendation of *Elampus* Spinola, 1806 (part.).

***Philoctetes bogdanovii* (Radoszkowski, 1877)**

Holopyga bogdanovii Radoszkowski, 1877: 5. Holotype ♂; Uzbekistan: Zarafshan (ZMMU) (examined).

Material examined. MONGOLIA: Arkhangai, 1 ♂, 90 km NE of Tsetserleg, 48°03'N, 102°25'E, 27.7.2005, leg. JH (MHC).

Distribution. *Mongolia (Arkhangai); Asiatic-European, from southern Europe, western Asia, Iran, and Turkey to Mongolia (Rosa et al. 2013; present record).

***Philoctetes cynthiae* Rosa, 2017**

Philoctetes cynthiae Rosa in Rosa et al. 2017c: 35. Holotype ♀; Russia: Tyva Rep., 13 km SW of Samagaltai, Dyttyg-Khem River, 9.VII.2013, leg. V. Loktionov & M. Proshchalykin (ZIN) (examined).

Material examined. MONGOLIA: Bayankhongor, 1 ♀, 1 ♂, 75 km S of Bayankhongor, 45°20'N, 100°48.5'E, 1330 m, 8.VII.2004, leg. JH (MHC); 1 ♀, 56 km NW of Bayankhongor, 46°33'N, 100°12'E, 2200 m, 12.VII.2004, leg. JS (MHC); 3 ♀♀,

7 ♂♂, 86 km NW of Bayankhongor, 46°50'N, 100°04'E, 2070 m, 14.VII.2004, leg. JH, MK (MHC); *Sukhbaatar*, 1 ♀, SE Mongolia, 200 km SSE of Baruun-Urt, Moltsoy Els, 1250 m, 27.VII.2007, leg. MK, paratype (PRC); 1 ♂, ibid, leg. MH (MHC); *Ulaanbaatar*, 2 ♂♂, Ulaanbaatar, Tuul River valley, 12.VII.2003, leg. JH (MHC); *Uvurkhangai*, 4 ♂♂, 12 km E of Aravaykheer, 46°22'N, 102°49'E, 1800 m, 3.VII.2004, leg. JH (MHC);

Distribution. Mongolia (*Bayankhongor, Sukhbaatar, *Ulaanbaatar, *Uvurkhangai); Russia (Tyva Rep.) (Rosa et al. 2017c).

Philoctetes diakonovi (Semenov, 1932)

Ellampus (Ellampus) diakonovi Semenov-Tian-Shanskij, 1932: 34. Holotype ♀; Kazakhstan: “Turkestan septentr.: Bajgakum (ZIN) (examined).

Material examined. Mongolia: *Dornogovi*, 1 ♀, 1 ♂, Orgon, 11.VII.2005, leg. JH (MHC).

Distribution. *Mongolia (Dornogovi); Central Asia (Kimsey and Bohart 1991).

Philoctetes lyubae Rosa, 2017

Philoctetes lyubae Rosa in Rosa et al. 2017c: 39. Holotype ♀; Russia: Tuva Rep., 20 km SSW of Erzin, Tore-Khol' Lake, 3.VII.2013, leg. V. Loktionov & M. Proshchalykin (ZIN) (examined).

Material examined. MONGOLIA: *Dornogovi*, 1 ♂, Atayn Mts, Gichigniv Nuruu, 10 km SW of Sain-Shand, 12.VII.2005, leg. JH (MHC).

Distribution. *Mongolia (Dornogovi); Russia (East Siberia) (Rosa et al. 2017c).

Philoctetes mongolicus (du Buysson, 1901)

Ellampus horvathi var. *mongolicus* du Buysson, 1901: 98. Lectotype ♂ (designated by Móczár 1967: 186); N Mongolia (NHMW) (examined). Rosa et al. 2020: 87 (cat., type series), 88 (fig. 55).

Ellampus horwathi (!) var. *mongolicus*: Bischoff 1913: 8 (cat., North Mongolia).

Omalus (Notozus) mongolicus: Lisenmaier 1959: 23 (descr. Mongolia).

Omalus mongolicus: Móczár 1967: 186 (cat., Mongolia: 1 ♂, Uvurkhangai aimag: Arc Bogd ul, ca. 20 km S of Somon Chovd, 1760 m, Exp. Dr. Z. Kaszab, 1964, nr. 170, 22.VI.1964; 1 ♂, Ulan-Baator, Bogdo ul, 1500 m, Exp. Dr. Z. Kaszab, 1963, nr. 4, 16.VI.1963).

Philoctetes horvathi var. *mongolicus*: Kimsey and Bohart 1991: 256 (cat., North Mongolia).

Philoctetes mongolicus: Rosa et al. 2014: 33 (cat., distr.); 2015: 436 (cat., descr., tax., Mongolia: Ulaanbataar, Baruun-Urt Moltsoy Els).

Material examined. MONGOLIA: Arkhangai, 1 ♂, 25 km NE of Tsetserleg, 47°38'N, 101°45'E, 23.VII.2004, leg. JH (MHC); Bayankhongor, 2 ♂♂, 16 km SW of Bayankhongor, 46°13'N, 100°30'E, 2165 m, 10.VII.2004, leg. JH (MHC); 1 ♂, 56 km NW of Bayankhongor, 46°33'N, 100°12'E, 2200 m, 12.VII.2004, leg. JH (MHC); 1 ♂, 86 km NW of Bayankhongor, 46°50'N, 100°04'E, 2070 m, 14.VII.2004, leg. JH (MHC); Khentii, 2 ♀♀, 100 km NE of Ondorkhaan, Kerulen River, 970 m, 22.VII.2007, leg. MH (MHC); Selenge, 1 ♀, 90 km N of Ulaanbaatar, Segnez River, 1450 m, 6–8.VII.2003, leg. JH (MHC); Sukhbaatar, 1 ♀, SE Mongolia, 200 km SSE of Baruun-Urt, Moltsoy Els, 1250 m, 27.VII.2007, MK (PRC); Tuv, 1 ♀, 50 km N of Ulaanbataar, E of Mandal, 1180 m, 8–13.VII.2007, leg. MK (PRC); 1 ♀, 1 ♂, ibid, leg. MH (MHC); Uvurkhangai, 1 ♂, N. Mongolei Leder 92, *Ellampus Horvathi* Mocs. var. *mongolicus* Buyss. var. nov. R. du Buysson det. 1901 ♂, Lectotype v. *mongolicus* Buysson det. L. Móczár <red label> (NHMW). Paralectotypes: 1 ♂ N. Mongolei Leder 92, *Ellampus Horvathi* Mocs. var. *mongolicus* Buyss. var. nov. R. du Buysson det. 1901 ♂, Paralectotype v. *mongolicus* Buysson det. L. Móczár <red label> (NHMW); 1 ♀, N. Mongolei Leder 92, *Ellampus Horvathi* Mocs. var. *mongolicus* Buyss. var. nov. R. du Buysson det. 1901 ♀, *Omalus horvathi* Mocs. det. L. Móczár (NHMW).

Distribution. Mongolia (*Arkhangai, *Bayankhongor, *Khentii, *Sukhbaatar, *Tuv, Ulaanbaatar, Uvurkhangai); widely distributed from Mongolia to Central Asia and southern Russia to Volga (Trautmann 1927), China (Shanxi) (Rosa et al. 2014).

Philoctetes shokalskii (Semenov, 1932)

Ellampus (Dictenulus) shokalskii Semenov-Tian-Shanskij, 1932: 24. Lectotype ♀ (designated by Kimsey 1986: 107); Mongolia: “Mongolia borealis: prope oppid. [um] Urga [Ulaanbaatar], 1–4.VI.1909, leg. P. Kozlov (ZIN) (examined). Rosa et al. 2017a: 78 (cat., type series), 218 (Plate 218). Rosa et al. 2017e: 119 (Mongolia: Chentej aimak 10 km W von Somon Delgerchaan, 1250 m Exp. Dr. Z. Kaszab, 1965 // Nr. 476. 24.VIII.1965).

Omalus shokalskii: Móczár, 1967: 186 (cat., Mongolia: 1 ♂, Ostgobi aimag; 40 km NW of Chara-Eireg, 1150 m, Exp. Dr. Z. Kaszab, 1963, nr. 62, 30.VI.1963; 1 ♂, Ostgobi aimag; 20 km SO of Čojren, 1200 m, Exp. Dr. Z. Kaszab, 1963, nr. 70, 1.VII.1963; 1 ♂, Sukhbaatar aimag; 44 km SSW of Baruun urt, 1050 m, Exp. Dr. Z. Kaszab, 1965, nr. 353, 3.VIII.1965; 1 ♀, Chentej aimag; 10 km W of Somon Delgerchaan, 1250 m, Exp. Dr. Z. Kaszab, 1965, nr. 476, 24.VIII.1965, allotype).

Philoctetes shokalskii: Kimsey and Bohart 1991: 257 (cat., Mongolia: Urga).

Material examined. MONGOLIA: Bayankhongor, 5 ♀♀, 11 ♂♂, 86 km NW of Bayankhongor, 46°50'N, 100°04'E, 2070 m, 14.VII. 2004, leg. JS, MK, JH (MHC); Dornod, 1 ♀, 50 km SW of Choilbalsan, 960 m, 25.VII.2007, leg. JH (MHC); Govi-Altai, 1 ♀, 1 ♂, 70 km of Altay city, Guulin, 14.VII.2005, leg. JH (MHC); Govi-Sümber, 2 ♂♂, 20 km SE of Choyr, 1480 m, 7.VIII.2007, leg. MK (PRC); 1 ♀, 1 ♂, ibid, leg. MH (MHC);

Sukhbaatar, 4 ♂♂, 200 km SSE of Baruun-Urt, Moltsoy Els, 1250 m, 27.VII.2007, leg. MK (PRC); 1 ♂, 210 km SSE of Baruun-Urt, 29.VII.2007, steppe, leg. MK (PRC); 12 ♂♂, 100 km SSW of Baruun-Urt, 1100 m, 30.VII.2007, leg. MH (MHC); *Tuv*, 1 ♀, Teregtin, 1350 m, Exp. Dr. Z. Kaszab, 1963, Nr.73, 2.VII.1963, det. Linsenmaier 1966 (NMLS); *Ulaanbaatar*, ♀ [not ♂], env. Urga [Ulaanbaatar], 1–4.VI.1909, leg. P. Kozlov, *Ellampus shokalskii* m. [mihi] Typ. ♂. A. Semenov-Tian-Shansky det. V.19, Lectotype *Ellampus shokalskii* Sem. design. LS Kimsey <red label> (ZIN); 1 ♀, same data, paralectotype (ZIN); *Umnugovi*, 1 ♀, Gobi Gurvansaikhan National Park, 40 km W of Dalanzadgad, 2000 m, 28–30.VI. 2003, leg. JH (MHC); *Uvurkhangai*, 9 ♂♂, 12 km E of Aravaykheer, 46°22'N, 102°49'E, 1800 m, 3.VII.2004, leg. JH (MHC).

Distribution. Mongolia (*Bayankhongor, *Dornod, Dornogovi, *Govi-Sümber, Khentii, Sukhbaatar, *Tuv, Ulaanbaatar, *Umnugovi, Uvurkhangai) (Kimsey 1986).

Genus *Pseudomalus* Ashmead, 1902

Pseudomalus Ashmead, 1902: 229. Type species: *Omalus semicircularis* Aaron, 1885 [= *Pseudomalus janus* (Haldeman, 1844)], by original designation.

Pseudomalus auratus nigridorsus (Tsuneki, 1953)

Ellampus auratus f. *nigridorsus* Tsuneki, 1953a: 54. Syntypes ♂, ♀; Japan, Korea, Manchuria (NIAS).

Material examined. MONGOLIA: *Bulgan*, 1 ♀, 137 km NE of Aravaykheer, 47°20'N, 103°45.5'E, 1250 m, 2.VII.2004, leg. JS (MHC); 1 ♀, Mongol Els Nat. Res., dunes, 47°24'N, 103°39'E, 31.VII.2005, leg. JH (MHC); *Khentii*, 1 ♀, 100 km NE of On-dorkhaan, Kerulen River, 970 m, 22.VII.2007, leg. MH (MHC); *Tuv*, 2 ♀♀, 18 ♂♂, 50 km E of Ulaanbaatar, Tuul River, 22.VI.2003, leg. JH (MHC); *Zavkhan*, 2 ♀♀, 40 km SW of Uliastay, dunes, 18.VII.2005, leg. JH (MHC).

Distribution. *Mongolia (Bulgan, Khentii, Tuv, Zavkhan); Russia (Eastern Siberia, Far East), China, Korea, Japan (Rosa et al. 2019).

Pseudomalus corensis (Uchida, 1927)

Philoctetes punctatus var. *corensis* Uchida, 1927: 153. Holotype ♂; Korea: Seiryori (descr.) (NIAS).

Omalus joannisi du Buysson, 1908: Móczár 1967: 185. (cat., Mongolia: Central aimag: Zuun-Chara, 850 m, Exp. Dr. Z. Kaszab, 1964, nr. 281; 8.VII.1964) (mis.).

Material examined. MONGOLIA: *Bulgan*, 1 ♀, Mongol Els Nat. Res., dunes, 47°24'N, 103°39'E, 31.VII.2005, leg. JH (MHC); *Dornod*, 1 ♀, 100 km W of Choilbalsan,

820 m, 23.VII.2007, leg. MH (MHC); 2 ♂♂, 20 km W of Choilbalsan, 800 m, 48°01'N, 114°14'E, 24.VII.2007, leg. MH (MHC/PRC); *Khentii*, 1 ♀, 1 ♂, 100 km NE of Ondorkhaan, Kerulen River, 970 m, 22.VII.2007, leg. MH (MHC); *Selenge*, 7 ♀♀, 5 ♂♂, 90 km N of Ulaanbaatar, Segnez River, 1450 m, 6–8.VII.2003, leg. JH (MHC/PRC); *Tuv*, 7 ♀♀, 3 ♂♂, 50 km N of Ulaanbaatar, E of Mandal, 1180 m, 8–13.VIII.2007, leg. MH (MHC, PRC); *Ulaanbaatar*, 3 ♀♀, Ulaanbaatar, Tuul River valley, 12.VII.2003, leg. JH (MHC).

Distribution. Mongolia (*Bulgan, *Dornod, *Khentii, *Selenge, Tuv, *Ulaanbaatar); Russia (Eastern Siberia, Far East); Japan (Hokkaido) (Rosa et al. 2019).

Remarks. The specimen illustrated in the volume of Russian cuckoo wasps (Rosa et al. 2019: fig. 18) is apparently misidentified and currently belonging to an unidentified species. Examination of type material by Uchida is needed for further studies.

Pseudomalus punctatus (Uchida, 1927)

Philoctetes punctatus Uchida, 1927: 152. Syntypes ♂, ♀; Japan: Hokkaido and Honshu (NIAS?).

Omalus punctatus: Móczár, 1967: 185 (cat., Mongolia: 1 ♀, 1 ♂, Čojbalsan [= Dornod] aimag: Chamardavaa ul, 80 km SO of Somon Chalchingol, 600 m, Exp. Dr. Z. Kaszab, 1965, nr. 401, 13.VIII.1963).

Material examined. MONGOLIA: *Bulgan*, 1 ♀, Mongol Els Nat. Res., dunes, 47°24'N, 103°39'E, 31.VII.2005, leg. JH (MHC); *Khentii*, 14 ♀♀, 100 km NE of Ondorkhaan, Kerulen River, 970 m, 22.VII.2007, leg. JH, MH (MHC); *Tuv*, 18 ♂♂, 50 km E of Ulaanbaatar, Tuul River, 22.VI.2003, leg. JH (MHC).

Distribution. Mongolia (*Bulgan, Dornod, *Khentii, *Tuv); Russia (Eastern Siberia, Far East); Korea, Japan (Rosa et al. 2019).

Pseudomalus pusillus (Fabricius, 1804)

Chrysis pusilla Fabricius, 1804: 176. Lectotype ♀ (designated by Rosa et al. 2020: 66); Austria (NHMW) (examined).

Omalus pusillus: Móczár, 1967: 195 (cat., Mongolia: 1 ♀, Čojbalsan [= Dornod] aimag: 50 km SO of Čojbalsan [= Choibalsan], 700 m, Exp. Dr. Z. Kaszab, 1965, nr. 421, 16.VIII.1965; 1 ♂, Čojbalsan [= Dornod] aimag: 44 km NW of Čojbalsan [= Choibalsan], 750 m, Exp. Dr. Z. Kaszab, 1965, nr. 425, 17.VIII.1965).

Material examined. MONGOLIA: *Bayankhongor*, 2 ♀♀, 16 km SW of Bayankhongor, 46°13'N, 100°30'E, 2165 m, 10.VII.2004, leg. JS (MHC); *Bulgan*, 2 ♀♀, 137 km NE of Aravaykheer, 47°20'N, 103°45.5'E, 1250 m, 2.VII.2004, leg. JS (MHC); *Dornod*, 4 ♀♀, 5 ♂♂, 100 km W of Choilbalsan, 820 m, 23.VII.2007, leg. MH (MHC); 4 ♀♀, 20 km W of Choilbalsan, 800 m, 48°01'N, 114°14'E 24.VII.2007,

leg. MH (MHC); 2 ♂♂, 50 km SW of Choilbalsan, 960 m, 25.VII.2007, leg. JH (MHC); *Khentii*, 3 ♀♀, 5 ♂♂, 100 km NE of Ondorkhaan, Kerulen River, 970 m, 22.VII.2007, leg. MH (MHC); *Selenge*, 2 ♂♂, 90 km N of Ulaanbaatar, Segnez River, 1450 m, 6–8.VII.2003, leg. JH (MHC); *Tuv*, 2 ♂♂, 50 km E of Ulaanbaatar, Tuul River, 22.VI.2003, leg. JH (MHC); *Umnugovi*, 3 ♀♀, 12 ♂♂, Gobi Gurvansaikhan National Park, 40 km W of Dalanzadgad, 2000 m, 28–30.VI.2003, leg. JH (MHC); *Uvurkhangai*, 4 ♀♀, 13 ♂♂, 12 km E of Aravaykheer, 46°22'N, 102°49'E, 1800 m, 3.VII.2004, leg. JH (MHC).

Distribution. Mongolia (**Bayankhongor*, **Bulgan*, *Dornod*, **Khentii*, **Selenge*, **Tuv*, **Umnugovi*, **Uvurkhangai*); Palaearctic, from western Europe and northern Africa to Russian Far East (Kurzenko and Lelej 2007).

Tribe *Parnopini*

Genus *Parnopes* Latreille, 1797

Parnopes Latreille, 1797: 126. Type species: *Chrysis carnea* Fabricius, 1775 [= *Parnopes grandior* (Pallas, 1771)], by monotypy.

Parnopes glasunowi Semenov, 1901

Parnopes glasunowi Semenow, 1901: 25. Holotype ♂; Tajikistan: “Turkestan occid. [entalis]: Jagnob: Rovat, 12.VII.1892, leg. D. Glasunow” (ZIN) (examined).

Material examined. MONGOLIA: *Khovd*, 1 ♂, ur. Elkhon, 20 km SE of Altai, Bodoncha, 27.V.1970, leg. E. Narchuk (ZIN).

Distribution. *Mongolia (Khovd); Central Asia, Russia (south of European part) (Rosa et al. 2019).

Parnopes popovii Eversmann, 1858

Parnopes popovii Eversmann, 1858: 567. Holotype ♀; Russia: Siberia “campis orientalibus” (ISEA-PAS) (examined). Kimsey and Bohart 1991: 586 (cat., Mongolia, without locality).

Material examined. MONGOLIA: *Arkhangai*, 1 ♂, 25 km NE of Tsetserleg, 47°38'N, 101°45'E, 23.VII.2004, leg. JH (MHC); 1 ♂, 90 km NE of Tsetserleg, 48°03'N, 102°25'E, 24.VII.2004, leg. JH (MHC); 1 ♀, ibid, 27.VII.2005, leg. JH (MHC); *Bulgan*, 1 ♂, 143 km NE of Arvaykheer, 47°24'N, 103°39'E, 26.VII.2004, 1300 m, sandy dunes, JS (PRC); 4 ♀♀, 3 ♂♂, Mongol Els Nat. Res., 47°24'N, 103°39'E,

dunes, 1320 m, 31.VII.2005, leg. JH (PRC); *Dornod*, 2 ♂♂, 50 km SW of Choibalsan, 960 m, 25.VII.2007, leg. JH (MHC); *Dornogovi*, 1 ♀, 2 ♂♂, 28 km SE of Chatan-Bulag, 3.VIII.2007, leg. MH (MHC); *Sukhbaatar*, 4 ♀♀, 4 ♂♂, 200 km SSE of Baruun-Urt, Moltsoy Els, 1250 m, 27.VII.2007, leg. MH (MHC); 1 ♂, 3 ♀♀, ibid, 27.VII.2007, leg. JH (MHC); 2 ♂♂, 100 km SSW of Baruun-Urt, 1100 m, 30.VII.2007, leg. MH (MHC); *Tuv*, 1 ♀, 75 km W of Ulaanbaatar, dunes, 2.VIII.2005, leg. JH (MHC); *Umnugovi*, 1 ♀, Gobi, Dalanzadgad, 25.VI.2003, leg. JH (MHC).

Distribution. Mongolia (*Arkhangai, *Bulgan, *Dornod, *Dornogovi, *Sukhbaatar, *Tuv, *Umnugovi); China (Heilongjiang, Shanghai, Shandong), Korea, Russia (Rosa et al. 2014).

Species to be excluded from Mongolian fauna

The following 19 taxa were described or listed for Mongolia by Radoszkowski (1887, 1891), Mocsáry (1890), Dalla Torre (1892), du Buysson (1893), Bischoff (1913), Linsenmaier (1959), and Kimsey and Bohart (1991), yet the type localities are situated in Inner Mongolia (China) or Central Asian countries. These species are expected in Mongolia due to the close vicinity of the collecting localities, with the only exception of *Chrysis fouqueti* (du Buysson, 1909), which belongs to the Oriental fauna.

Elampus mocsaryi Radoszkowski, 1887

Elampus mocsari (!) Radoszkowski, 1887: 45. Holotype ♀; Mongolia [= China]: Qinghai: Zaïdam (ISEA-PAS) (examined).

Ellampus (Notozus) mocsaryi: Mocsáry 1889: 80. Justified emendation of *Elampus mocsari* Radoszkowski, 1887. Dalla Torre 1892: 14 (cat., Mongolia [= China]).

Ellampus mocsaryi: Dalla Torre 1892: 14 (cat., Mongolia [= China]); Kimsey and Bohart 1991: 168 (cat., Mongolia [= China]: Zaidam).

Notozus mocsaryi: Bischoff 1913: 6 (cat., Mongolia [= China]).

Omalus (Notozus) mocsaryi: Linsenmaier 1959: 16 (key), 24 (tax., descr., Mongolia [= China]).

Elampus spinipes (Mocsáry, 1890)

Ellampus (Notozus) spinipes Mocsáry, 1890: 49. Holotype ♀; Mongolia [= China, Inner Mongolia]: Mongolia meridionalis (Ta-Wan) (ISEA-PAS) (examined).

Ellampus spinipes: Dalla Torre 1892: 18 (cat., Mongolia [= Inner Mongolia]).

Elampus spinipes: Kimsey and Bohart 1991: 171 (cat., Mongolia [= Inner Mongolia]: Ta-Wan).

Notozus spinipes: Bischoff 1913: 7 (cat., Mongolia [= Inner Mongolia]).

Omalus (Notozus) spinipes: Linsenmaier 1959: 16 (key), 24 (descr., Mongolia [= Inner Mongolia]).

***Hedychridium ardens mongolicum* Tsuneki, 1947**

Hedychridium ardens f. *mongolicum* Tsuneki, 1947: 47. Holotype ♀; China: Inner Mongolia: Apaka (NIAS).

Hedychridium ardens f. *mongolicum*: Kimsey and Bohart 1991: 188 (cat., Mongolia [= China, Inner Mongolia]: Apaka).

***Hedychrum simile* Mocsáry, 1889**

Hedychrum cyaneum Mocsáry in Radoszkowski 1889: 10, nec Brullé, 1846. Lectotype ♀ (designated by French, in Bohart and French 1986: 341); China “Ta-schian-sy” (HNHM) (examined).

Hedychrum simile Mocsáry, 1889: 157. Replacement name for *Hedychrum cyaneum* Radoszkowski, 1889, nec Brullé, 1846.

Hedychrum simile f. *mongolicus* Tsuneki, 1947: 54. Syntypes ♀♀, ♂♂; China: Inner Mongolia: Apaka (NIAS).

Hedychrum simile: Linsenmaier 1959: 39 (descr., key, Mongolia [= China, Inner Mongolia]); Kimsey and Bohart 1991: 220. Mongolia (cat., without locality, related to the record of *H. simile mongolicus* from Inner Mongolia by Tsuneki 1947).

***Philoctetes hypocrita* (du Buysson, 1893)**

Ellampus hypocrita du Buysson, 1893: 246. Syntypes ♀♀; Mongolia [= China, Inner Mongolia], Kansu-Jelisyn-Kuse (ISEA-PAS); Persia (MNHN) (examined). Bischoff 1913: 8 (cat., Mongolia [= China, Inner Mongolia])

Omalus hypocritus: Kimsey and Bohart 1991: 248. Incorrect subsequent spelling.

Pseudomalus hypocrita: Rosa et al. 2015: 77.

Philoctetes hypocrita: Farhad et al. 2018: 199. Lectotype designation: ♂; China: Kansu Jelisyn Kuse (ISEA-PAS).

***Pseudomalus tshingiz* (Semenov, 1954)**

Ellampus tshingiz Semenov in Semenov-Tian-Shanskij and Nikol'skaya, 1954: 93. Holotype ♂; China: Sandzhu [Xinjiang], Gushan Gobi (depository: ZIN) (examined). Rosa et al. 2017a: 80 (cat., type series), 221 (plate 223).

Pseudomalus tshingiz: Kimsey and Bohart 1991: 270 (cat., Mongolia [= China]: Sachow Gobi [= Oasis Sachzhou, Gashunkoe Gobi [= Dunhuang, Gansu]]).

***Chrysis aegle* Semenov, 1967**

Chrysis (Gonodontochrysis) aegle Semenov-Tian-Shanskij, 1967: 160. Holotype ♀; China: Alashan, Maladzhin (ZIN) (examined).

Chrysis aegle: Kimsey and Bohart 1991: 379 (cat., Mongolia [= China, Inner Mongolia]: Alashan, Maladzhin]).

***Chrysis analis altaica* Mocsáry, 1912**

Chrysis (Tetrachrysis) analis var. *altaica* Mocsáry, 1912: 586. Holotype ♀; Kazakhstan: Altai: Semipalatinsk (HNHM) (examined).

Chrysis altaica: Kimsey and Bohart 1991: 381 (cat., Mongolia [= Kazakhstan]: Altai Mts).

***Chrysis fouqueti* (du Buysson, 1909)**

Tetrachrysis fouqueti du Buysson, 1909: 210. Holotype ♀; Viet Nam: Tonkin (MNHN).

Chrysis fouqueti: Kimsey and Bohart 1991: 412 (cat., Mongolia for the erroneous synonymy of *Chrysis csikiana* Mocsáry, 1912).

***Chrysis csikiana* Mocsáry, 1912**

Chrysis (Tetrachrysis) Csikiana Mocsáry, 1912: 406. Lectotype ♂ (designated by Bohart in Bohart and French 1986: 341); Kazakhstan: Semipalatinsk (HMNH) (examined).

Chrysis csikiana: Kimsey and Bohart 1991: 412 (cat., Mongolia [= Kazakhstan]: Altai Mts).

***Chrysis jelisyni* Radoszkowski, 1891**

Chrysis jelisyni Radoszkowski, 1891: 186. Syntypes ♀♀; Mongolia [= China]: Kansu, Jelissyn-Kuce (ISEA-PAS, MfN) (examined).

Chrysis (Tetrachrysis) jelisyni: Bischoff 1913: 54 (cat., Mongolia [= China]: Totau (locality not found)).

Chrysis jelisyni: Kimsey and Bohart 1991: 34 (cat., Mongolia [= China]: Kansu).

***Chrysis keriensis* Radoszkowski, 1887**

Chrysis (Tetrachrysis) keriensis Radoszkowski, 1887: 47. Holotype ♂ [not ♀]; China: Xinjiang, Keria-Daria (ISEA-PAS) (examined).

Chrysis (Tetrachrysis) keriensis: Mocsáry 1889: 516 (tax., descr., Mongolia [= China]); Bischoff 1913: 54 (cat., Mongolia [= China]).

Chrysis keriensis: Dalla Torre 1892: 73 (cat., Mongolia [= China]); Kimsey and Bohart 1991: 427 (cat., Mongolia [= China]: Keria Daria).

***Chrysis kozlovi* Semenov, 1967**

Chrysis (Gonodontochrysis) kozlovi Semenov-Tian-Shanskij, 1967: 160. Holotype ♂; China: Alashan, Uzosto canyon, 14.5.1908, leg. P. Kozlov (ZIN) (examined).

Chrysis kozlovi: Kimsey and Bohart 1991: 429 (cat., Mongolia [= China, Inner Mongolia]: Alashan, Uzosto Canyon).

***Chrysis mongoliana* Bohart, 1991**

Chrysis (Tetrachrysis) mongolica Semenov-Tian-Shanskij, 1967: 178, nec Mocsáry, 1914. Holotype ♀; Russia: Transbaikalia: Ingoda River (ZIN) (examined).

Chrysis mongoliana Bohart in Kimsey and Bohart 1991: 440. Replacement name for *Chrysis (Tetrachrysis) mongolica* Semenov-Tian-Shanskij, 1967, nec Mocsáry, 1914 (cat., Mongolia: Transbaikalia: Ingoda river).

***Chrysis potanini* Radoszkowski, 1891**

Chrysis potanini Radoszkowski, 1891: 186. Holotype ♂; Mongolia [= China]: Tufyn (ISEA-PAS) (examined).

Chrysis (Tetrachrysis) potanini: Bischoff 1913: 57 (cat., Mongolia [= China]).

Chrysis potanini: Kimsey and Bohart 1991: 450 (cat., Mongolia [= China]: Tufyn).

***Chrysis przewalskii* Radoszkowski, 1887**

Chrysis Przewalskii Radoszkowski, 1887: 46. Holotype ♂; Mongolia [= China]: Zaïdam (ISEA-PAS) (examined).

Chrysis (Tetrachrysis) przewalskii: Mocsáry 1889: 504 (tax., descr., Mongolia [= China]); Bischoff 1913: 57 (cat., Mongolia [= China]).

Chrysis przewalskii: Dalla Torre 1892: 86 (cat., Mongolia [= China]); Kimsey and Bohart 1991: 452 (cat., Mongolia [= China]: Zaidam, Keria Mts).

***Chrysis spinidens* Mocsáry, 1887**

Chrysis (Tetrachrysis) spinidens Mocsáry in Radoszkowski, 1887: 48. Holotype ♂; Mongolia [= China]: Zaïdam (ISEA-PAS) (examined). Mocsáry 1889: 516 (cat., descr., Mongolia [= China]); Bischoff 1913: 59 (cat., Mongolia [= China]).

Chrysis spinidens: Dalla Torre 1892: 97 (cat., Mongolia [= China]); Kimsey and Bohart 1991: 464 (cat., Mongolia: Zaidam).

***Chrysura alticola* (Semenov-Tian-Shanskij, 1912)**

Chrysis petri alticola Semenov-Tian-Shanskij, 1912: 190. Lectotype ♀ (designated by Rosa et al. 2017a: 45); Kyrgyzstan: Peter the Great Range, Gardan-Kaftar Pass (ZIN) (examined).

Chrysura alticola: Kimsey and Bohart 1991: 486 (cat., Mongolia [= Kyrgyzstan]).

Conclusions

Approximately 1500 chrysidid specimens were examined for the compilation of this first checklist of the Mongolian Chrysidae. Fifty-seven resulted newly recorded, but still a large number of specimens are laying unidentified in museum and private collections. Nineteen species were excluded from the fauna of Mongolia, because collecting localities are currently included in China territories; however, these species are expected for Mongolia. Based on the available data, distributional records for 90 Mongolian species are listed, representing 18 genera grouped in two subfamilies. In terms of species richness, Cleptinae are represented only by two species so far, and the subfamily Chrysidiinae is the most speciose (88 species, 98%). Among Chrysidiinae, Chrysidiini is the most speciose tribe (47 species, 53.4%), followed by Elampini (39 species, 44.3%), and finally Parnopini (2 species, 2.2%).

Currently eight species (9% of known taxa) are provisionally considered endemic: *Cleptes mongolicus* Rosa, Halada & Agnoli, sp. nov., *Chrysis mocsaryi* Radoszkowski, 1889, *Ch. priapus* Rosa, 2018, *Spinolia spinosa* Rosa & Halada, sp. nov., *Elampus spinifemoris* (Móczár, 1967), *Hedychrum lama* du Buysson, 1891, *Holopyga kaszabi* Móczár, 1967, and *Philoctetes shokalskii* (Semenov, 1932).

From a chorological point of view, one species has a Holarctic distribution (*Pseudochrysis neglecta*), ten have Palaearctic distributions, one has a Holarctic and Oriental distribution (*Omalus aeneus*), one a Palaearctic and Oriental distribution (*Hedychridium gerstaeckeri*), 28 species have an Asiatic-European distribution, 21 have an East Palaearctic distribution, and 19 have a Central Asian distribution.

Another result of the present study is a better knowledge of the distributional limits of some species, and Mongolia represents the easternmost record for seven species: *Chrysis aestiva*, *Ch. illigeri*, *Ch. jaxartis*, *Ch. leptomandibularis*, *Chrysura ignifrons*, *Elampus albipennis* and *Philoctetes bogdanovii*.

The most widespread Mongolian species is the endemic *Philoctetes shokalskii* recorded in ten aimags. *Chrysis consanguinea* and *Hedychridium ardens* were recorded in nine aimags. This is not surprising because *C. consanguinea* resulted in being one of the most common species from Western to Eastern Siberia also (Rosa et al. 2017c, d), whereas *H. ardens* is one of the most common Euro-Siberian species ranging from Central Europe to China (Rosa et al. 2014). *Pseudomalus pusillus* was recorded in eight aimags, whereas *H. cupreum*, *H. chalybaeum* and *Philoctetes mongolicus* were recorded in seven aimags; they are Asiatic-European species, sometimes locally abundant. *Hedychridium longigena* and *Parnopes popovii* were recorded in seven aimags, yet they are East-Palaearctic species.

Although most of the Mongolian aimags are under-represented in the existing data due to inadequacy of surveys, based on the currently available data we can state that the highest number of recorded species was collected in Tuv (42 species), Arkhangai (27 species), and Selenge and Dornod (20 species) aimags (Table 2). The family Chrysidae has not yet been documented from Bayan-Ulgii, Darkhan-Uul,

Table 2. Species diversity of aimags in terms of area size, number of specimens and number of collecting sites.

Aimags	Area, km ²	No. of species	No. of coll. sites	No. of specimens
Arkhangai	55 314	27	4	237
Bayankhongor	115 978	18	10	142
Bulgan	48 733	18	4	61
Dornod	123 597	19	4	228
Dornogovi	109 472	16	4	42
Govi-Alтай	141 448	8	5	49
Govi-Сүмбэр	5 542	1	1	4
Khentii	80 325	12	1	55
Khovd	76 061	4	4	4
Selenge	41 153	20	1	100
Sukhbaatar	82 286	17	4	79
Tuv	74 042	42	10	350
Ulaanbaatar	4 704	10	4	20
Umnugovi	165 381	13	7	34
Uvurkhangai	62 895	12	5	42
Zavkhan	82 457	11	1	21

Comment. Chrysidae are not known in Bayan-Ulgii, Darkhan-Uul, Dundgov, Khuvsulg, Orkhon, and Uvs.

Dundgov, Khuvsulg, Orkhon, and Uvs although it is probable that this cosmopolitan family is present in these aimags and it is only a matter of time before the fauna is sampled and recorded.

Overall, the Mongolian fauna is still too poorly known for a complete analysis of species richness and composition. The faunal richness of Mongolia is doubtless much higher than we currently know, in comparison with the chrysidid fauna of the adjacent countries and considering the geographic position of Mongolian aimags and their different biotopes. For example, at least another 75 species recorded for Siberia (Rosa et al. 2017d) are expected for Mongolia, as well as other five genera known from bordering and central Asian countries (Chrysidiini: *Chrysidea* Bischoff, 1913, *Spintharina* Semenov, 1892; Elampini: *Chrysellampus* Semenov, 1932, *Haba* Semenov, 1954; Parnopini: *Cephaloparnops* Bischoff, 1910). Copious undescribed members of the genus *Prochridium* Linsenmaier, 1968 were collected in Mongolia. The descriptions of Mongolian *Prochridium*, and a revision of this genus, are in preparation. Only a single record of an undescribed *Prochridium* was previously known in literature for Central Asia (Turkmenistan) (Linsenmaier 1994).

Acknowledgements

We are grateful to curators who allowed the study of type materials for this study: Sergey Belokobylskij (Saint Petersburg, Russia), Marco Bernasconi (Luzern, Switzerland), Gian Luca Agnoli (Bologna, Italy), Zoltan Vas (Budapest, Hungary), Daniel Kubisz and Dawid Moroń (ISEA-PAS, Kraków, Poland). We thank Sergey Belokobylskij and Jing-xian Liu (Guangzhou, China) for review of the manuscript.

The reported study for Maxim Proshchalykin and Ulykpan Aibek was funded by RFBR and MECSS, project number 20-54-44014.

References

- Abeille de Perrin E (1877) Diagnoses d'espèces nouvelles et remarques sur des espèces rares. *Feuille Jeunes Naturaliste* 7: 65–68.
- Abeille de Perrin E (1878) Diagnoses de Chrysides nouvelles. Published by the author, Marseille, 6 pp.
- Abeille de Perrin E (1879) Synopsis critique et synonymique des Chrysides de France. *Annales de la Société Linnéenne de Lyon* 26: 1–108. <https://doi.org/10.3406/linly.1879.3979>
- Agassiz A (1846) *Nomenclator zoologicus*, fase. Jent et Gassmann, Soloduri, [12 index, viii +] 393 pp.
- Ashmead WH (1902) Classification of the fossorial, predaceous and parasitic wasps, or the superfamily Vespoidea. *The Canadian Entomologist* 34: 79–88, 131–137, 163–166, 203–210, 219–231, 268–273, 287–291. <https://doi.org/10.4039/Ent34287-11>
- Bischoff H (1913) Hymenoptera. Fam. Chrysididae. In: Wytsman P (Ed.) *Genera insectorum* 151. L. Desmet-Verteneuil, Bruxelles, 86 pp. [5 pls]
- Bodenstein WG (1939) The genotypes of the Chrysididae. *Transactions of the American Entomological Society* 65: 123–133.
- Bohart RM, French LD (1986) Designation of chrysidid lectotypes in the Mocsáry Collection at the Hungarian National Museum, Budapest (Hymenoptera: Chrysididae). *The Pan-Pacific Entomologist* 62(4): 340–343.
- Brullé A (1846) Les Chrysidides. In: Lepeletier de Saint-Fargeau A (Ed.) *Histoire Naturelle des Insectes. Hyménoptères*. Librairie De Ruret Paris, vol. IV, 680 pp.
- Buyanjargal B, Abasheev RYU (2015) Nesting biology and behaviour of *Euodynerus dantici* (Rossi, 1790) (Hymenoptera: Vespidae: Eumeninae) in central Mongolia. *Mongolian Journal of Biological Sciences* 13(1–2): 25–33. <https://doi.org/10.22353/mjbs.2015.13.04>
- Buysson R du (1887) Descriptions de Chrysidides nouvelles. *Revue d'Entomologie* 6: 167–201.
- Buysson R du (1891) Contribution aux Chrysidides du globe. *Revue d'Entomologie* 10: 29–47.
- Buysson R du (1893) Contribution aux Chrysides du Globe – 2^e série. *Revue d'Entomologie de France* 2: 245–252.
- Buysson R du (1901) Sur quelques Chrysidides du Musée de Vienne. *Annalen des K. K. Naturhistorischen Hofmuseums in Wien* 16(1): 97–104.
- Buysson R du (1909 [1908]) Hyménoptères nouveau. *Revue d'Entomologie* 27: 207–219.
- Chevrier F (1869) Description de deux Chrysidides du bassin du Léman. *Mitteilungen der Schweizerischen Entomologischen Gesellschaft* 3: 44–48.
- Coquebert AJ (1801) *Illustratio iconographica insectorum Quae in Musaeis parisiniis observavit et in lucem edidit Joh. Christ. Fabricius. Praemissis ejusdem descriptionibus; Accedunt Species plumariae, vel minus aut nondum cognitae. Tabularum decas secunda. Petri Didot natu majors*, Paris, 43 pp. [48–90 + pls xi–xx].
- Dahlbom AG (1831) *Exercitationes Hymenopterologicae. Monographia Chrysididum Sveciae (Familia Hymenopterorum Octava Latreille)*. Pars II, III (partim). Londini Gothorum, 30 pp. [Pars II pp. 19–32; Pars III, 33–48, Chrysididae 33–36].
- Dahlbom AG (1845) *Dispositio Methodica Specierum Hymenopterorum, secundum Familias Insectorum Naturales. Particula secunda. Chrysis in sensu Linnæano*. Berlingianis, Lund, 20 pp. <https://doi.org/10.5962/bhl.title.66977>

- Dahlbom AG (1854) Hymenoptera Europaæ praecipue borealia, formis typicis nonnullis specierum generumve Exoticorum aut Extraneorum propter nexum systematicum associatis, per familias, genera, species et varietates disposita atque descripta. 2. *Chrysis* in sensu Linnaeano. Friedrich Nicolai, Berlin, [xxiv +] 412 pp. [+ 12 pls]
- Dalla Torre CG de (1892) Catalogus hymenopterorum hucusque descriptorum systematicus et synonymicus. Volumen VI: Chrysidae (Tubulifera). Guilelmi Engelmann, Lipsiae, [viii +] 118 pp. <https://doi.org/10.5962/bhl.title.8794>
- Dathe HH, Proshchalykin MYu (2016) The genus *Hylaeus* Fabricius in Mongolia, an updated species inventory (Hymenoptera: Apoidea: Colletidae). Zootaxa 4121(4): 351–382. <https://doi.org/10.11646/zootaxa.4121.4.1>
- Eversmann E (1858 [1857]) Fauna Hymenopterologica Volgo-Uralensis. Continuatio. Familia Chrysidarum. Bulletin de la Société Imperiale des Naturalistes de Moscou 30: 544–567.
- Fabricius JC (1787) Mantissa insectorum sistens eorum species nuper detectas adiectis characteribus genericis, differentiis specificis, emendationibus, observationibus. Tom. II. Hafniae. Proft. [1], 382 pp. <https://doi.org/10.5962/bhl.title.36471>
- Fabricius JC (1781) Species insectorum, exhibentes eorum differentias specificas, synonyma auctorum, loca natalia, metamorphosin, adiectis observationibus, descriptionibus. Tomo I. C.E. Bohnii, Hambugi et Kiloni [= Hamburg and Kiel], 522 [+ viii] pp. <https://doi.org/10.5962/bhl.title.36509>
- Fabricius JC (1804) Systema Piezatorum secundum ordines, genera, species, adiectis synonymis, locis, observationibus, descriptionibus. Apud Carolum Reichard, Brunsvigae, III–XIV, 439 [+ 30] pp. <https://doi.org/10.5962/bhl.title.10490>
- Farhad A, Rosa P, Talebi AA (2018) Additions to the fauna of Iranian Elampini (Hymenoptera: Chrysidae, Chrysinae), with key to species and taxonomic notes. Journal of Crop Protection 7(2): 191–206.
- Förster A (1853) Eine Centurie neuer Hymenopteren. Beschreibungen neuer Arten aus der Familiae der Chrysididen. Verhandlungen des naturhistorischen Vereins der preussischen Rheinlande und Westphalens 10: 266–362.
- Gogorza J (1887) Crisididos de los alrededores de Madrid. Anales de la Real Sociedad Española de Historia Natural 16: 17–88.
- Hammer K (1936) Schwedisch-chinesische wissenschaftliche Expedition nach den nordwestlichen Provinzen Chinas. 43. Hymenoptera. 8. Scoliiden, Mutilliden und Chrysiden. Arkiv för Zoology 27A(23): 1–3.
- HAO (2020) The Hymenoptera Glossary: Hymenoptera Anatomy Consortium. <http://glossary.hymao.org> [accessed 5 August 2020]
- Haupt H (1957 [1956]) Die unechten und echten Goldwespen Mitteleuropas (*Cleptes* et Chrysidae). Abhandlungen und Berichte aus dem Staatlichen Museum für Tierkunde in Dresden 23: 15–139.
- Kimsey LS (1986) Designation of chrysidid lectotypes. The Pan-Pacific Entomologist 62(2): 105–110.
- Kimsey LS, Bohart RM (1991 [1990]) The Chrysidid wasps of the World. Oxford University Press, New York, 652 pp.

- Kurzenko NV, Lelej AS (2007) Chrysidae. In: Lelej AS, Belokobylskij SA, Kasparyan DR, Kupyanskaya AN, Proshchalkin MYu (Eds) Key to the insects of Russian Far East. Vol. IV. Neuropteroidea, Mecoptera, Hymenoptera. Part 5. Dalnauka, Vladivostok, 998–1006. [In Russian]
- Lanes GO, Kawada R, Azevedo CO, Brothers DJ (2020) Revisited morphology applied for systematics of flat wasps (Hymenoptera, Bethylidae). Zootaxa 4752(1): 1–127. <https://doi.org/10.11646/zootaxa.4752.1.1>
- Latreille PA (1797) Précis des caractères génériques des Insectes, disposés dans un ordre naturel par le Citoyen Latreille. 'An V'. F. Bordeaux, Brive, Prévot, Paris, [xiv +] 201 [+ 7] pp. [+ 1 pl.] <https://doi.org/10.5962/bhl.title.58411>
- Latreille PA (1802) Histoire naturelle générale et particulière des Crustacés et des Insectes. Ouvrage faisant suite à l'Histoire Naturelle générale et particulière, composée par Leclercq de Buffon, et rédigée par C.S. Sonnini, membre de plusieurs Sociétés savantes. Familles naturelles des genres. Tome troisième. F. Dufart, Paris, 455 pp. [xii + 13–467] <https://doi.org/10.5962/bhl.title.15764>
- Latreille PA (1809) Genera Crustaceorum et Insectorum secundum ordinem naturalem in familias disposita, iconibus exemplisque plurimis explicata. Tomus quartus et ultimus. Amand Koenig, Parisiis et Argentorati [= Paris and Strasbourg], 399 pp.
- Latreille PA (1810) Considérations générales de l'ordre des animaux composant les classes des Crustacés, des Arachnides, et des Insectes. F. Schoell, Paris, 444 pp. <https://doi.org/10.5962/bhl.title.39620>
- Lavrenko EM (1979) Vegetation map of Mongolian People's Republic. Scale 1:1,500,000. 4 sheets. GUGK, Moscow. [In Russian and Mongolian]
- Lichtenstein J (1876) Note sur le genre *Chrysis*. Petites Nouvelles Entomologiques 145: 27–27.
- Linnaeus C (1758) Systema Naturae per Regna tria Naturae, secundum Classes, Ordines, Genera, Species, cum characteribus, differentiis, synonymis, locis. Editio Decima, Refurmata, Tomus I. Laurenti Salvii, Holmiae, 824 pp. [+ IV] <https://doi.org/10.5962/bhl.title.542>
- Linnaeus C (1761) Fauna Suecia sistens Animalia Sueciae Regni: Mammalia, Aves, Amphibia, Pisces, Insecta, Vermes. Distributa per Classes et Ordines, enera et Species, cum Differentiis, Specierum, Synonymis, Auctorum, Nominibus Incolarum, Locis natalium, Descriptionibus Insectorum. Editio Altera, Auctior. Laurentius Salvius, Stockholm, 578 pp. [+ 2 pls] <https://doi.org/10.5962/bhl.title.46380>
- Linsenmaier W (1951) Die europäischen Chrysiden (Hymenoptera). Versuch einer natürlichen Ordnung mit Diagnosen. Mitteilungen der Schweizerischen Entomologischen Gesellschaft 24(1): 1–110.
- Linsenmaier W (1959) Revision der Familie Chrysidae (Hymenoptera) mit besonderer Berücksichtigung der europäischen Spezies. Mitteilungen der Schweizerischen Entomologischen Gesellschaft 32(1): 1–232.
- Linsenmaier W (1968) Revision der Familie Chrysidae (Hymenoptera). Zweiter Nachtrag. Mitteilungen der Schweizerischen Entomologischen Gesellschaft 41(1–4): 1–144.
- Linsenmaier W (1994) The Chrysidae (Insecta: Hymenoptera) of the Arabian Peninsula. Fauna of Saudi Arabia 14: 145–206.

- Lisenmaier W (1997a) Altes und Neues von den Chrysididen. (Hymenoptera, Chrysididae). Entomofauna 18(19): 245–300.
- Lisenmaier W (1997b) Die Goldwespen der Schweiz. Veröffentlichungen aus dem Natur-Museum Luzern 9: 1–140.
- Mocsáry A (1889) Monographia Chrysididarum Orbis Terrarum Universi. Hungarian Academy of Science, Budapest, 643 pp.
- Mocsáry A (1890) Additamentum primum ad monographia Chrysididarum Orbis Terrarum Universi. Természetrajzi Füzetek 13(2–3): 45–66.
- Mocsáry A (1912) Species Chrysididarum novae. III. Annales Historico-Naturales Musei Nationalis Hungarici 10(1): 549–592.
- Mocsáry A (1914) Chrysididae plerumque exoticae novae. Annales Historico-Naturales Musei Nationalis Hungarici 12(1): 1–72.
- Móczár L (1964) Über die *Notozus*-Arten Ungarns (Hymenoptera, Chrysididae). Annales Historico-Naturales Musei Nationalis Hungarici 56: 439–447.
- Móczár L (1967) Ergebnisse der zoologischen Forschungen von Dr. Z. Kaszab in der Mongolei. 100. Chrysididae (Hymenoptera). Acta Zoologica Academiae Scientiarum Hungaricae 13(1–2): 183–190.
- Móczár L (1997) Revision of the *Cleptes nitidulus* group of the world (Hymenoptera, Chrysididae, Cleptinae). Entomofauna 18(3): 25–44.
- Morgan D (1984) Cuckoo-Wasps Hymenoptera, Chrysididae. Handbooks for the Identification of British insects. Royal Entomological Society of London 6(5), 37 pp.
- Niehuis O (2000) The European species of the *Chrysia ignita* group: Revision of the *Chrysia angustula* aggregate (Hymenoptera, Chrysididae). Deutsche Entomologische Zeitschrift 47(2): 181–201. <https://doi.org/10.1002/mmnd.4800470207>
- Olivier GA (1791) *Chrysia*. In: Olivier GA (Ed.) Encyclopédie méthodique. Histoire naturelle. Insectes. 5(2). Panckoucke, Paris, 669–678.
- Orlovskyté S, Budrys E, Budrienė A, Radzevičiutė R, Soon V (2016) Sibling species in the *Chrysia ignita* complex: molecular, morphological and trophic differentiation of Baltic species, with a description of two new cryptic species (Hymenoptera: Chrysididae). Systematic Entomology 41: 771–793. <https://doi.org/10.1111/syen.12190>
- Panzer GWF (1798–1806) Faunae Insectorum Germaniae initia oder Deutschlands Insecten, Nürnberg, 49–60 (1798), 80–85 (1801). [Dating after Sherborn, 1923]
- Paukkunen J, Rosa P, Soon V, Johansson N, Ødegaard F (2014) Faunistic review of the cuckoo wasps of Fennoscandia, Denmark and the Baltic countries (Hymenoptera: Chrysididae). Zootaxa 3864(1): 1–67. <https://doi.org/10.11646/zootaxa.3864.1.1>
- Pauli T, Castillo-Cajas RF, Rosa P, Kukowka S, Berg A, van den Berghe E, Fornoff F, Hopfenmüller S, Niehuis M, Peters RS, Staab M, Strumia F, Tischendorf S, Schmitt F, Niehuis O (2018) Phylogenetic analysis of cuckoo wasps (Chrysididae) reveals the partially artificial nature of the current classification at the genus level in this family of Hymenoptera. Systematic Entomology 44(2): 322–335. <https://doi.org/10.1111/syen.12323>
- Proshchalykin MYu (2017) Colletid bees (Hymenoptera, Apoidea: Colletidae) of Mongolia: fauna and zonal distribution. Euroasian Entomological Journal 16(2): 192–200.

- Proshchalykin MYu, Kuhlmann M (2015) The bees of the genus *Colletes* Latreille (Hymenoptera, Colletidae) collected by the Soviet-Mongolian expeditions 1967–1982. Far Eastern Entomologist 296: 1–18.
- Radoszkowski O (1877) Chrysidiformes, Mutillidae et Sphegidae. In: Fedtshenko AP (Ed.) Putieshestvie v Turkestan [Voyage au Turkestan d'Alexis Fedtschenko] (14) 2(5), Sankt-Petersburg, 1–87. [+ 8 pls] [In Russian and Latin]
- Radoszkowski O (1889 [1888]) Révision des armures copulatrices des mâles de la tribu des Chrysides. Hora Socientatis Entomologicae Rossicae 23(1–2): 3–40. [+ 6 pls]
- Radoszkowski O (1891) Descriptions de Chrysides nouvelles. Revue d'Entomologie 10: 183–198.
- Rosa P (2017a) First record of *Cleptes* Latreille for Mongolia, with male description of *Cleptes dauriensis* Móczár (Hymenoptera, Chrysididae). Zoosystematica Rossica 26(2): 287–293. <https://doi.org/10.31610/zsr/2017.26.2.287>
- Rosa P (2017b) Review of the Palaearctic species of the genus *Colpopygia* (Hymenoptera: Chrysididae) with description of a new species. Zoosystematica Rossica 26(2): 294–306. <https://doi.org/10.31610/zsr/2017.26.2.294>
- Rosa P (2018) On the status of the cuckoo wasp genus *Allochrysis* Kimsey & Bohart, 1991, with descriptions of the *Chrysis* ear and *C. slava* species groups (Hymenoptera: Chrysididae). Zoosystematica Rossica 27(2): 268–286. <https://doi.org/10.31610/zsr/2018.27.2.268>
- Rosa P (2019) New species of Chrysididae (Hymenoptera) from Central Asia, Russia and Iran. Part II. Far Eastern Entomologist 377: 1–25. <https://doi.org/10.25221/fee.377.1>
- Rosa P, Belokobylskij SA, Zaytseva LA (2017a) The Chrysididae types described by Semenov-Tian-Shanskij and deposited at the Zoological Institute of the Russian Academy of Sciences, Saint-Petersburg (Insecta: Hymenoptera). Proceedings of the Zoological Institute RAS, Supplement 5: 1–269.
- Rosa P, Lelej AS, Belokobylskij SA, Vinokurov NB, Zaytseva LA (2019) Illustrated and annotated check-list of the Russian cuckoo wasps (Hymenoptera, Chrysididae). Entomofauna, Supplement 23: 1–360. <https://doi.org/10.25221/fee.360.1>
- Rosa P, Lotfalizadeh H, Pourrafei L (2013) First checklist of the chrysidid wasps (Hymenoptera: Chrysididae) of Iran. Zootaxa 3700(1): 1–47. <https://doi.org/10.11646/zootaxa.3700.1.1>
- Rosa P, Madl M, Zettel H, Zimmermann D (2020) Annotated catalogue of the Chrysididae (Insecta, Hymenoptera) types deposited at the Naturhistorisches Museum, Vienna. Annales des Naturhistorischen Museums in Wien, Serie B: 17–140.
- Rosa P, Pavese M, Soon V, Niehuis N (2017b) *Pseudochrysis* Semenov, 1891 is the valid genus name for a group of cuckoo wasps frequently referred to as *Pseudospinolia* Linsenmaier, 1951 (Hymenoptera, Chrysididae). Deutsche Entomologische Zeitschrift 64(1): 69–75. <https://doi.org/10.3897/dez.64.13005>
- Rosa P, Proshchalykin MYu, Lelej AS, Loktionov VM (2017c) Contribution to the Siberian Chrysididae (Hymenoptera). Part 1. Far Eastern Entomologist 341: 1–44. <https://doi.org/10.25221/fee.341.1>
- Rosa P, Proshchalykin MYu, Lelej AS, Loktionov VM (2017d) Contribution to the Siberian Chrysididae (Hymenoptera). Part 2. Far Eastern Entomologist 342: 1–42. <https://doi.org/10.25221/fee.342.1>

- Rosa P, Wei N-s, Feng J, Xu Z-f (2016) Revision of the genus *Trichrysis* Lichtenstein, 1876 from China, with description of three new species (Hymenoptera, Chrysidae). Deutsche Entomologische Zeitschrift 63(1): 109–136. <https://doi.org/10.3897/dez.63.7347>
- Rosa P, Wei N-s, Notton D, Xu Z-f (2015) The genus *Philoctetes* Abeille de Perrin, 1879 from China, with description of two new species (Hymenoptera, Chrysidae). Zootaxa 4040(4): 433–444. <https://doi.org/10.11646/zootaxa.4040.4.3>
- Rosa P, Wei N-s, Xu Z-f (2014) An annotated checklist of the chrysidid wasps (Hymenoptera, Chrysidae) from China. ZooKeys 455: 1–128. <https://doi.org/10.3897/zookeys.455.6557>
- Rossi P (1790) Fauna etrusca sistens Insecta quae in provinciis Florentina et Pisana praesertim collegit. Tomus secundus. Thomae Masi et Sociorum, Liburni, 348 pp. [+ 10 pl.] <https://doi.org/10.5962/bhl.title.15771>
- Schenck ACF (1856) Beschreibung der in Nassau aufgefundenen Goldwespen (Chrysidae) nebst einer Einleitung über die Familie im Allgemeinen und einer kurzen Beschreibung der übrigen deutschen Arten. Jahrbücher des Vereins für Naturkunde im Herzogthum Nassau 11: 13–89.
- Scopoli JA (1763) Entomologia Carniolica exhibens Insecta Carnioliae indigena et distributa in ordines, genera, species, varietates. Methodo Linnaeano. Typis Ioannis Thomae Trattner, Vindobonae, 420 pp. <https://doi.org/10.5962/bhl.title.119976>
- Semenov A (1891) *Pseudochrysis (Spintharis) virgo*, sp. n. Horae Societatis Entomologicae Rossicae 25(1/2): 441–444.
- Semenov A (1892) De genere *Pseudochrysis* m. Horae Societatis Entomologicae Rossicae 26: 480–491.
- Semenov-Tian-Shansky A (1910 [1909]) Chrysidiarum species novae vel parum cognitae (Hymenoptera). IV. Russian Entomological Review 9: 213–226.
- Semenov-Tian-Shanskij A (1912) Chrysidiarum species novae vel parum cognitae (Hymenoptera). V. Russian Entomological Review 12: 177–201.
- Semenov-Tian-Shanskij A (1932) Supplementa ad Chrysidiarum monographias ab A.G. Dahlbom (1854), A. Mocsáry (1889), R. du Buysson (1896) et H. Bischoff (1913) editas. I. Horae Societatis Entomologicae Rossicae 42(3): 1–48.
- Semenov-Tian-Shanskij AP (1954) Classification of the tribe Hedychrini Mocs. (Hymenoptera, Chrysidae) and description of new species. Proceedings of the Zoological Institute, Academy of Sciences of the USSR 15: 138–145. [In Russian]
- Semenov-Tian-Shanskij A (1967) New species of gold wasps (Hymenoptera, Chrysidae). Trudy Zoologicheskogo Instituta Akademii Nauk SSSR 43: 118–184. [In Russian]
- Semenov-Tian-Shanskij A, Nikol'skaya MN (1954) Cuckoo-wasps (Hymenoptera, Chrysidae) of Tajikistan. Trudy Zoologicheskogo Instituta Akademiy Nauk SSSR 15: 89–137. [In Russian]
- Semenow A (1901) Chrysidiarum species novae vel parum cognitae (Hymenoptera). I. Russian Entomological Review 1(1–2): 23–27.
- Shuckard WE (1837) Description of the Genera and Species of the British Chrysidae. Entomologist's Magazine 4: 156–177.
- Spinola M (1806–1808) Insectorum Liguriae species novae aut rariores quas in agro Ligustico nuper detexit, descripsit et iconibus illustravit Maximilianus Spinola, adjecto catalogo

- specierum auctoribus jam enumeratarum, quae in eadem regione passim occurunt. Gravier Yves, Genuae. (1806) 1, [xvii +] 160 pp. [+ 2 pls] (1807) 2, [ii +] 1–82 pp.; (1808) 3, 83–262 pp. [+ 5 pls] [dating after Passerin d'Entrèves 1983] <https://doi.org/10.5962/bhl.title.65985>
- Thomson CG (1870) Öfversigt af de i Sverige funna arter af Slägget *Chrysis* Lin. Opuscula Entomologica 2: 101–108.
- Trautmann W (1926) Untersuchungen an einigen Goldwespenformen. Deutsche Entomologische Zeitschrift 40: 4–12.
- Trautmann W (1927) Die Goldwespen Europas. G. Uschmann, Weimar, 194 pp.
- Tsuneki K (1947) Chrysidae from North China and Inner Mongolia. Mushi 17(9): 43–60.
- Tsuneki K (1948) Chrysidae from Shansi, North China (Hymenoptera). Mushi 18(19): 115–128. [+ 3 pls]
- Tsuneki K (1950) Descriptions of new species and subspecies of the Chrysidae from East Asia (Hymenoptera). Mushi 21(8): 61–81.
- Tsuneki K (1953a) Chrysidae of Manchuria (Hymenoptera). Mushi 25(8): 53–61.
- Tsuneki K (1953b) Chrysidae of Korea (Hymenoptera). Kontyu 20 (1/2): 22–28.
- Uchida T (1927 [1926]) Über die Bethyliden Japans. Zoological Magazine 38: 181–186. [In Japanese]
- Wei NS, Rosa P, Liu JX, Xu ZF (2014) The genus *Omalus* Panzer, 1801 (Hymenoptera, Chrysidae) from China, with descriptions of four new species. ZooKeys 407: 29–54. <https://doi.org/10.3897/zookeys.407.7531>
- Wesmael C (1839) Notice sur les Chrysides de Belgique. Bulletins de l'Académie Royale des Sciences Belles Lettres Bruxelles 6: 167–177.