

The genus *Eriastichus* La Salle (Hymenoptera, Eulophidae, Tetrastichinae) in the Neotropical region, introducing 48 new species

Christer Hansson^{1,2}

1 Scientific Associate Biological Museum (Entomology), Lund University, Sölvegatan 37, SE-22362 Lund, Sweden **2** Natural History Museum, Life Sciences, Cromwell Road, London, UK

Corresponding author: Christer Hansson (christer.hansson@biol.lu.se)

Academic editor: N. Johnson | Received 4 November 2020 | Accepted 14 December 2020 | Published 22 February 2021

<http://zoobank.org/EE1BAF87-5BD7-4E18-9DF9-29E2C9CC7D4C>

Citation: Hansson C (2021) The genus *Eriastichus* La Salle (Hymenoptera, Eulophidae, Tetrastichinae) in the Neotropical region, introducing 48 new species. ZooKeys 1019: 35–91. <https://doi.org/10.3897/zookeys.1019.60364>

Abstract

Our knowledge of the recently described genus *Eriastichus* La Salle is greatly enhanced with the addition of 48 new species: *E. acribis* sp. nov., *E. aphritis* sp. nov., *E. cluridis* sp. nov., *E. coelotis* sp. nov., *E. colenisi* sp. nov., *E. copalensis* sp. nov., *E. daptilis* sp. nov., *E. decoris* sp. nov., *E. denotatis* sp. nov., *E. derilis* sp. nov., *E. diadrys* sp. nov., *E. dotaensis* sp. nov., *E. drupis* sp. nov., *E. ebulis* sp. nov., *E. egestis* sp. nov., *E. eleagnis* sp. nov., *E. ellipsis* sp. nov., *E. eminis* sp. nov., *E. facilis* sp. nov., *E. fenestris* sp. nov., *E. follis* sp. nov., *E. galeatis* sp. nov., *E. geratis* sp. nov., *E. glanis* sp. nov., *E. hilaris* sp. nov., *E. johnlasallei* sp. nov., *E. johnnayosi* sp. nov., *E. maniatis* sp. nov., *E. nebulis* sp. nov., *E. neonis* sp. nov., *E. nexilis* sp. nov., *E. novalis* sp. nov., *E. nugalis* sp. nov., *E. oasis* sp. nov., *E. ononis* sp. nov., *E. orestis* sp. nov., *E. pallidops* sp. nov., *E. parabilis* sp. nov., *E. renodis* sp. nov., *E. rivalis* sp. nov., *E. sannionis* sp. nov., *E. scalaris* sp. nov., *E. sodalis* sp. nov., *E. taraxis* sp. nov., *E. tendrilis* sp. nov., *E. tonioazofeifai* sp. nov., *E. velaminis* sp. nov., and *E. vestis* sp. nov. All species are known only from males, and all material is from Costa Rica. Females show little morphological variation and are not possible to separate to species or link to conspecific males at present. Apart from the diagnostic features for the genus presented in the original description, two new autapomorphies are introduced here: an inflated pleural membrane between Gt_{1-4} and Gs_{1-4} , and tufts of pale and flattened setae laterally on Gt_{4-6} . Both features are found on the gaster and are present in both sexes. The distribution of the genus is firmly established as predominantly neotropical. The biology remains unknown.

Keywords

Autapomorphies, Costa Rica, identification key

Introduction

Taxonomic knowledge of the Tetrastichinae (Chalcidoidea: Eulophidae) fauna in the Neotropical region is very poor, at both the genus and species levels. LaSalle et al. (2006) listed 29 genera in the subfamily from the Neotropical region, and with the addition of a new genus by Hansson (2020) this number is now 30, but hardly any of them have been investigated in this part of the world.

Eriastichus is a genus confined to the Americas, with the great majority of species occurring in tropical parts (La Salle 1994 and present findings). It was described in a paper on the Nearctic genera of Tetrastichinae (LaSalle 1994), including three species that were described in same paper. The species were based on females, and the male was included for only one of the species. With the exception of one specimen from the U.S.A. (Texas), the specimens were from the Neotropical region, ranging from the Caribbean (Dominican Republic) through Central America (Costa Rica, Mexico) to South America (Ecuador). The biology of the species was cited as being unknown.

Materials and methods

Material

Material forming the base of this study includes 63 females and 81 males, and all specimens were collected with a sweep net. The specimens were killed and kept in 80% ethanol until dried with a critical point drier. Thus, the specimens did not shrivel when dried and this should be kept in mind for the measurements/ratios given.

The types are deposited in the following museums with registration numbers pre-fixed by their acronyms:

MZLU Biological Museum (Entomology), Lund University, Lund, Sweden;
MZUCR Museo de Zoología, Universidad de Costa Rica, San José, Costa Rica;
NHMUK Natural History Museum, London, United Kingdom.

Other museum acronyms

CNC Canadian National Collection of Insects, Ottawa, Canada;
USNM United States National Museum of Natural History, Washington, D.C., USA.

Imaging

The colour images of the specimens were made using Canon camera equipment including an EOS 5D Mark IV body, a telezoom lens, 70–300 mm (but using only 135 & 200 mm), with a 10 × Mitutoyo microscope lens attached, and macro twin lite MT-24 EX for illumination. The camera was attached to a Cognisys stackshot macro-

rail system. The picture stacking was done with Helicon Focus version 6, and Adobe Photoshop was used for image processing. The SEM micrographs are from uncoated specimens and were done with a Hitachi SU 3500 microscope, in low vacuum and using a backscatter detector.

Morphological terminology

The terminology used here follows Gibson (1997), except the term mesoscutellum which is used instead of scutellum.

Abbreviations of morphological terms

CC	costal cell in fore wing;	PM	postmarginal vein in fore wing;
F1–4	flagellomeres 1–4;	POL	shortest distance between lateral ocelli;
Gs _{1–7}	gastral sternites 1–7;	SLG	sublateral groove on mesoscutellum (sublateral line in LaSalle 1994);
Gt _{1–7}	gastral tergites 1–7;	SMG	submedian groove on mesoscutellum (submedian line in LaSalle 1994);
HE	height of eye;	ST	stigmal vein in fore wing;
MS	malar space;	WM	width of mouth opening.
MV	marginal vein;		
OOL	shortest distance between lateral ocelli and eyes;		

Species names


To avoid the problems associated with the requirement by the International Code of Zoological Nomenclature (International Commission of Zoological Nomenclature 1999) that the gender of species names must agree with that of the genus, all new names proposed in the present study, except when indicated otherwise, must be treated as arbitrary combinations of letters or nouns in apposition. As a consequence, species names will not change if species are transferred to a genus with a different gender at any time in the future.

Taxonomic accounts


Genus *Eriastichus* La Salle

La Salle 1994: 173–174. Type species *Eriastichus cigdema* La Salle.

Diagnosis. Heavily setose species (Figs 1, 2, 5, 6): head (including eyes), pronotum, mesoscutum, coxae, and lateral parts of propodeum (excl. bare median part – hairy only in *E. masneri* La Salle) ± covered with fine, short setae; malar sulcus curved; base of scape oblique (e.g., Fig. 45); F1–F4 of male antenna with scattered setae of varying length



Figures 1–6. *Eriastichus* spp. **1** *E. eleagnis*, body in lateral view, male paratype **2** *E. sp.*, body in lateral view, female **3** *E. decoris*, gaster in lateral view, male holotype **4** *E. sp.*, right fore wing, female **5** *E. fenestratus*, body in dorsal view, male holotype **6** *E. sp.*, thoracic dorsum, female.



Figures 7–10. *Eriastichus* spp. **7, 8** gaster in lateral view: **7** *E.* sp., female **8** *E. ebulis*, male paratype **9, 10** Gt₆ in lateral view, with brush of flattened setae: **9** *E.* sp., female **10** *E. ebulis*, male paratype.

(Figs 11–58), i.e., without a compact whorl of long setae attached close to base – a feature present in many tetrastichine groups; mesoscutellum transverse (Figs 5–6); dorsellum large, flat to slightly convex and smooth (Fig. 5); fore wing with postmarginal vein 0.3–0.7 × as long as the elongate and slender stigmal vein, wing disc strongly setose, including base (i.e., without speculum) and ventral surface of costal cell (Fig. 4), wings ± uniformly but weakly infuscate (Fig. 4) but in a few species hyaline; gaster with pleural membrane between Gt_{1-4} and Gs_{1-4} inflated (Figs 1–3, 7, 8); both sexes with a tuft of pale and flattened setae laterally on Gt_{3-6} (Figs 7–10), though most species with tufts only on Gt_6 .

Key to males

- 1 Mesoscutellum with parts lateral to submedian grooves strongly setose (Fig. 6)..... 2
- Mesoscutellum with parts lateral to submedian grooves with two or three pairs of strong setae (Fig. 5) 3
- 2 Median propodeum hairy (La Salle 1994: fig. 30) *E. masneri* La Salle (male unknown, this is an anticipated placement; not recorded from Costa Rica)
- Median propodeum bare (as in Fig. 5) *E. nakos* La Salle (male unknown, this is an anticipated placement; not recorded from Costa Rica)
- 3 Scape 1.3–1.4 × as long as ventral plaque (Fig. 50); lower face yellowish brown (Fig. 59)..... *E. pallidops* sp. nov.
- Scape 1.7–8.0 × as long as ventral plaque; colour of lower face variable 4
- 4 Scape 5.3–8.0 × as long as ventral plaque 5
- Scape 1.7–5.0 × as long as ventral plaque 9
- 5 Antenna with dorsobasal setae on F1 0.5 × as long as length of F1 (Fig. 15), pedicel + flagellum 2.0 × as long as width of head *E. copalensis* sp. nov.
- Antenna with dorsobasal setae on F1 at least 1.1 × as long as length of F1, pedicel + flagellum at most 1.6 × as long as width of head 6
- 6 Scape 8.0 × as long as ventral plaque (Fig. 57), and 1.4 × as long as height of eye *E. tonioazofeifai* sp. nov.
- Scape at most 6.7 × as long as ventral plaque, and at most 0.9 × as long as height of eye 7
- 7 Antennal flagellum longer (Fig. 11), e.g., F1 2.5 ×, F4 2.8 ×, clava 7.3 × as long as wide *E. acribis* sp. nov.
- Antennal flagellum shorter, F1 1.5–1.7 ×, F4 2.0 ×, clava 4.8–5.2 × as long as wide 8
- 8 Setae on flagellomeres more erect, and dorsobasal setae on F1 1.5 × as long as length of F1 (Fig. 12) *E. cluridis* sp. nov.
- Setae on flagellomeres less erect, and dorsobasal setae on F1 1.1 × as long as length of F1 (Fig. 16) *E. coelotis* sp. nov.
- 9 Scape 1.3 × as long as height of eye (Fig. 14) *E. colenis* sp. nov.
- Scape at most 1.0 × as long as height of eye 10

10	Scape dark brown	11
—	Scape yellowish brown to pale brown.....	17
11	Dorsobasal setae on F1 0.8–1.1 × as long as length of F1.....	12
—	Dorsobasal setae on F1 0.4–0.6 × as long as length of F1.....	14
12	Ventral plaque on scape situated medially (Fig. 37) <i>E. johnlasallei</i> sp. nov.	
—	Ventral plaque on scape situated below the middle (Figs 17, 20).....	13
13	Each flagellomere with few setae, e.g., F1 with six setae, ventral plaque 0.2 × as long as length of scape (Fig. 17)	<i>E. daptilis</i> sp. nov.
—	Each flagellomere with more setae, e.g., F1 with ca. 15 setae, ventral plaque 0.4 × as long as length of scape (Fig. 20)	<i>E. dotaensis</i> sp. nov.
14	Flagellomeres longer, F1, F3, F4 each 3.7 ×, F2 4.0 × as long as wide (Fig. 18).....	<i>E. decoris</i> sp. nov.
—	Flagellomeres shorter, F1 2.0–3.5, F2 2.4–3.5, F3 2.1–3.4, F4 2.3–3.4 × as long as wide.....	15
15	Fore and mid legs with coxae and femora yellowish brown	
— <i>E. denotatis</i> sp. nov.	
—	Fore and mid legs with coxae and femora dark brown.....	16
16	Ventral plaque 0.5 × as long as length of scape (Fig. 22) .. <i>E. diadrys</i> sp. nov.	
—	Ventral plaque 0.3 × as long as length of scape (Fig. 23).... <i>E. ebulis</i> sp. nov.	
17	Dorsobasal setae on F1 ca. 2 × as long as F1	<i>E. cigdemae</i> La Salle
—	Dorsobasal setae on F1 at most 1.5 × as long as F1	18
18	Dorsobasal setae on F1 at least 1.2 × as long as F1, and setae on flagellomeres erect (Figs 24, 27)	19
—	Dorsobasal setae on F1 at most 1.0 × as long as F1, setae on flagellomeres variable	20
19	Antenna shorter (Fig. 24), pedicel + flagellum 1.1 × as long as width of head.....	<i>E. egestis</i> sp. nov.
—	Antenna longer (Fig. 27), pedicel + flagellum 1.6–1.7 × as long as width of head.....	<i>E. eleagnis</i> sp. nov.
20	Scape 4.5–5.0 × as long as ventral plaque.....	21
—	Scape 1.8–4.2 × as long as ventral plaque.....	24
21	Ventral plaque situated medially (Fig. 25)	<i>E. ellipsis</i> sp. nov.
—	Ventral plaque situated below median part of scape	22
22	Dorsobasal setae on F1 0.6 × as long as length of F1 (Fig. 40)	
— <i>E. johnnnoyesi</i> sp. nov.	
—	Dorsobasal setae on F1 0.9–1.0 × as long as length of F1 (Figs 28, 31)	23
23	Antenna with F1–F4 each 3.3–3.5 × as long as wide, setae on flagellomeres less erect (Fig. 31)	<i>E. eminis</i> sp. nov.
—	Antenna with F1–F4 1.8–2.5 × as long as wide, setae on flagellomeres more erect (Fig. 28)	<i>E. facilis</i> sp. nov.
24	Scape 1.7–2.0 × as long as ventral plaque.....	25
—	Scape 2.2–4.2 × as long as ventral plaque.....	28

25	Setae on flagellomeres more erect, scape dark brown (Fig. 29)	<i>E. fenestris</i> sp. nov.
— Setae on flagellomeres less erect, scape yellowish brown to pale brown 26	
26	Antenna with F1–F4 each 2.0 × and clava 4.3 × as long as wide (Fig. 21)	<i>E. derilis</i> sp. nov.
— Antenna with F1–F4 each 2.7–3.0 × and clava 5.5–6.3 × as long as wide (Figs 13, 30) 27	
27	Flagellomeres with more setae, e.g., F2 with eight setae on dorsal surface (Fig. 13)..... 28	<i>E. aphritis</i> sp. nov.
— Flagellomeres with less setae, e.g., F2 with four setae on dorsal surface (Fig. 30)..... 29	<i>E. follis</i> sp. nov.
28	Frons and legs yellowish brown..... 30	<i>E. sannionis</i> sp. nov.
— Frons partly to completely dark brown, legs yellowish brown to dark brown 31	
29	Coxae and femora predominantly to completely dark brown 32	
— Coxae and femora ±yellowish brown, some coxae or hind femur can be partly dark brown 33	
30	Setae on flagellomeres erect (Figs 34, 58)	31
— Setae on flagellomeres less erect than in alternate (Figs 33, 56)..... 32	
31	Scape narrow at base (Fig. 34)..... 33	<i>E. galeatis</i> sp. nov.
— Scape wide at base (Fig. 58)	<i>E. vestis</i> sp. nov.
32	Flagellomeres with few setae, e.g., F2 with three setae on dorsal surface (Fig. 56)..... 34	<i>E. velaminis</i> sp. nov.
— Flagellomeres with more setae, e.g., F2 with eight setae on dorsal surface (Fig. 33)..... 35	<i>E. geratis</i> sp. nov.
33	Scape 3.5–4.2 × as long as ventral plaque..... 36	
— Scape 2.2–3.1 × as long as ventral plaque..... 37	
34	Dorsobasal setae on F1 0.5–0.6 × as long as F1 (Figs 32, 36)	38
— Dorsobasal setae on F1 0.8–1.0 × as long as F1 (Figs 39, 41)	39
35	Flagellomeres with more setae, e.g., F2 with eight setae on dorsal surface (Fig. 32)..... 40	<i>E. glanis</i> sp. nov.
— Flagellomeres with fewer setae, e.g., F2 with five setae on dorsal surface (Fig. 36)..... 41	<i>E. maniatis</i> sp. nov.
36	Setae on flagellomeres more erect (Fig. 39)	<i>E. neonis</i> sp. nov.
— Setae on flagellomeres less erect (Fig. 41)	<i>E. nexilis</i> sp. nov.
37	Ventral plaque situated below middle of scape	42
— Ventral plaque situated in the middle, or slightly above the middle..... 43	
38	Scape 2.2–2.5 × as long as ventral plaque..... 44	
— Scape 2.8–3.1 × as long as ventral plaque..... 45	
39	Dorsobasal setae on F1 1.0 × as long as F1 (Fig. 44)	<i>E. novalis</i> sp. nov.
— Dorsobasal setae on F1 0.5–0.6 × as long as F1 (Fig. 42)	<i>E. oasis</i> sp. nov.
40	F2–F4 each 2.3 × as long as wide (Fig. 26)	<i>E. drupis</i> sp. nov.
— F2–F4 each at least 3.0 × as long as wide	46

41	Flagellomeres longer, e.g., F2 3.6–3.8 × as long as wide (Figs 38, 43)	42
–	Flagellomeres shorter, F2 3.0–3.2 × as long as wide	43
42	Dorsobasal setae on F1 0.9 × as long as F1 (Fig. 43) <i>E. ononis</i> sp. nov.	
–	Dorsobasal setae on F1 0.6 × as long as F1 (Fig. 38) <i>E. nebulis</i> sp. nov.	
43	Dorsobasal setae on F1 0.5 × as long as F1, scape dark brown (Fig. 46)	
 <i>E. parabilis</i> sp. nov.	
–	Dorsobasal setae on F1 0.9 × as long as F1, scape yellowish brown (Fig. 49)....	
 <i>E. orestis</i> sp. nov.	
44	Dorsobasal setae on F1 0.5 × as long as F1 and pedicel + flagellum 1.9 × as long as width of head.....	45
–	Dorsobasal setae on F1 at least 0.6 × as long as F1 and/or pedicel + flagellum at most 1.7 × as long as width of head	46
45	Setae on flagellomeres more erect (Fig. 47)	<i>E. rivalis</i> sp. nov.
–	Setae on flagellomeres less erect (Fig. 45)	<i>E. nugalis</i> sp. nov.
46	Dorsobasal setae on F1 0.4 × as long as F1 (Fig. 35) and pedicel + flagellum 1.7 × as long as width of head.....	<i>E. hilaris</i> sp. nov.
–	Dorsobasal setae on F1 0.6–1.0 × as long as F1, if dorsobasal setae on F1 is only 0.6 × as long as F1, then pedicel + flagellum are 1.5 × as long as width of head.....	47
47	Antennal flagellum long and slender (Fig. 54), e.g., F4 3.4 × and clava 7.6 × as long as wide	<i>E. renodis</i> sp. nov.
–	F4 2.2–2.8 × and clava 4.8–6.3 × as long as wide.....	48
48	Dorsobasal setae on F1–F4 1.0 × as long as funicular segment it is attached to (Fig. 52).....	<i>E. scalaris</i> sp. nov.
–	Dorsobasal setae on F1–F4 0.6–0.8 × as long as funicular segment it is attached to.....	49
49	Antennal flagellum more slender (Fig. 53), e.g., F3 3.0 × and clava 6.3 × as long as wide	<i>E. taraxis</i> sp. nov.
–	F3 2.3–2.6 × and clava 4.8–5.8 × as long as wide.....	50
50	Pedicel + flagellum 1.9 × as long as width of head	<i>E. sodalis</i> sp. nov.
–	Pedicel + flagellum 1.5–1.7 × as long as width of head	
 <i>E. tendrilis</i> sp. nov.	

Species treatments

Eriastichus acribis sp. nov.

<http://zoobank.org/4CAF1EDF-8C16-4075-B08B-1356397DEC4B>

Figure 11

Type locality. Costa Rica, Heredia, Estación Biológica La Selva, 75 m, 10°26'N, 84°01'W, 24–25.ii.2011, J.S. Noyes leg.

Type specimen. *Holotype* male dried and glued to a paper card. Original labels: "COSTA RICA, Heredia, E.B. La Selva, 75 m, 10°26'N, 84°01'W, 24–25.ii.2011,

J.S. Noyes, NHM (Ent) 2011–93”, “HOLOTYPE Eriastichus acribis Hansson” [red printed label], (NHMUK014431021).

Diagnosis (male). Head brown; antenna (Fig. 11): scape yellowish brown, ventral plaque on scape ca. $0.1 \times$ as long as scape, antenna with dorsobasal setae on F1 $1.1 \times$ as long as F1; gaster with lateral tufts of pale and flattened setae on Gt₆.

Description (male holotype NHMUK014431021). Length of body 1.2 mm. Head brown. Antenna with scape yellowish brown, ventral plaque brown, pedicel and flagellum brown. Mesoscutum and mesoscutellum dark brown; dorsellum and propodeum pale brown. Legs yellowish brown. Gaster dark brown.

Head. Length/width in frontal view 0.8; width/length in dorsal view 2.5; POL/OOL 2.8; WM/MS 3.6; MS/HE 0.4; HE/head length in frontal view 0.6; widths head/mesoscutum 1.2. **Antenna.** Pedicel + flagellum length/mesoscutum width 1.8; pedicel + flagellum length/head width 1.5; lengths scape/ventral plaque 6.7; ventral plaque located in the middle of scape; scape length/width 2.9; lengths scape/head (dorsal view) 0.5; scape length/HE 0.8; length/width F1, F2, F3, F4, clava: 2.5, 2.0, 2.3, 2.8, 7.3; length dorsobasal setae on F1/length F1 1.1. **Mesosoma.** Length/width 1.4; mesoscutum length/width 0.5; mesoscutellum length/width 0.6; widths SMG/SLG 1.0; enclosed space between SMG length/width 2.0; lengths mesoscutum/mesoscutellum 1.2; lengths mesoscutellum/dorsellum 2.3; lengths mesosoma/gaster 0.7. **Wings.** CC length/width 21.9; lengths CC/MV 1.0; lengths MV/ST 2.3; lengths MV/PM 4.9; lengths PM/ST 0.5; submarginal vein with four setae on dorsal surface. **Gaster.** With lateral tufts of pale and flattened setae on Gt₆.

Eriastichus aphritis sp. nov.

<http://zoobank.org/B8A47821-A1EE-4B64-B7C7-5EEE1B059956>

Figure 13

Type locality. Costa Rica, Puntarenas, San Vito, Estación Biológica Las Alturas, 1500 m, 8°57'N, 82°50'W, 17–18.ii.2012, J.S. Noyes leg.

Type specimen. *Holotype* male dried and glued to a paper card. Original labels: “COSTA RICA, Puntarenas, San Vito, E.B. Las Alturas, 1500 m, 8°57'N, 82°50'W, 17–18.ii.2012, J.S. Noyes, NHM (Ent) 2012–91”, “HOLOTYPE Eriastichus aphritis Hansson” [red printed label], (NHMUK014431022).

Diagnosis (male). Head black with metallic tinges with part below antennal toruli yellowish brown, scape and pedicel pale brown, flagellum brown; ventral plaque on scape ca. $0.5 \times$ as long as scape (Fig. 13), antenna with dorsobasal setae on F1 $0.5 \times$ as long as F1; gaster with lateral tufts of pale and flattened setae on Gt₆.

Description (male holotype NHMUK014431022). Length of body 2.2 mm. Head black with metallic tinges with part below antennal toruli yellowish brown, scape and pedicel pale brown, flagellum brown, ventral plaque dark brown. Mesoscutum



Figures 11–17. *Eriastichus* spp. head and antenna in lateral view, male holotypes **11** *E. acribis* **12** *E. cluridis* **13** *E. aphritis* **14** *E. colenis* **15** *E. copalensis* **16** *E. coelotis* **17** *E. daptilis*.

and mesoscutellum black with metallic tinges, dorsellum and propodeum dark brown. Legs yellowish brown. Gaster dark brown.

Head. Length/width in frontal view 0.8; width/length in dorsal view 2.5; POL/OOL 1.9; WM/MS 1.5; MS/HE 0.6; HE/head length in frontal view 0.5; widths head/mesoscutum 1.2. **Antenna.** Pedicel + flagellum length/mesoscutum width 1.9; pedicel + flagellum length/head width 1.8; lengths scape/ventral plaque 2.1; ventral plaque located below the middle of scape; scape length/width 3.0; lengths scape/head (dorsal view) 0.5; scape length/HE 0.9; length/width F1, F2, F3, F4, clava: 3.0, 2.9, 2.8, 2.8, 5.5; length dorsobasal setae on F1/length F1 0.5. **Mesosoma.** Length/width 1.6; mesoscutum length/width 0.6; mesoscutellum length/width 0.8; widths SMG/SLG 1.1; enclosed space between SMG length/width 2.3; lengths mesoscutum/mesoscutellum 1.5; lengths mesoscutellum/dorsellum 3.4; lengths mesosoma/gaster 0.7. **Wings.** CC length/width 17.5; lengths CC/MV 1.1; lengths MV/ST 2.1; lengths MV/PM 3.0; lengths PM/ST 0.7; submarginal vein with ten setae on dorsal surface. **Gaster.** With lateral tufts of pale and flattened setae on Gt₆.

Eriastichus cigdemae La Salle

Eriastichus cigdemae La Salle, 1994: 207. Holotype female in USNM, not examined.

Diagnosis. Mesoscutellum with 2–3 pairs of setae; male with dorsobasal setae on F1 ca. 2× as long as F1.

Description. See La Salle (1994).

Distribution. Costa Rica, Mexico, U.S.A. (Texas) (La Salle 1994).

Eriastichus cluridis sp. nov.

<http://zoobank.org/0C4B7F59-E6A9-4F47-AC10-BA87174C2738>

Figure 12

Type locality. Costa Rica, Alajuela, Parque Nacional Volcan Arenal, 10°26'N, 84°43'W, 600 m, 26.ii.2013, J.S. Noyes leg.

Type specimen. *Holotype* male dried and glued to a paper card. Original labels: "COSTA RICA, Alajuela, P.N. Volcan Arenal, 10°26'N, 84°43'W, 600 m, 26.ii.2013, J.S. Noyes, NHM (Ent) 2013–145AQ", "HOLOTYPE Eriastichus cluridis Hansson" [red printed label], (NHMUK014431023).

Diagnosis (male). Head dark brown with part below antennal toruli yellowish brown; antenna (Fig. 12): scape yellowish brown, ventral plaque on scape ca. 0.2 × as long as scape, antenna with dorsobasal setae on F1 1.5 × as long as F1; gaster with lateral tufts of pale and flattened setae on Gt₆.

Description (male holotype NHMUK014431023). Length of body 1.2 mm. Head dark brown with part below antennal toruli yellowish brown. Antenna with

scape yellowish brown, ventral plaque dark brown, pedicel and flagellum brown. Mesoscutum, mesoscutellum and propodeum dark brown; dorsellum pale brown. Legs yellowish brown, fore and mid coxae brown at base, hind coxa and hind femur brown. Gaster dark brown.

Head. Length/width in frontal view 0.8; width/length in dorsal view 2.1; POL/OOL 2.7; WM/MS 1.5; MS/HE 0.5; HE/head length in frontal view 0.6; widths head/mesoscutum 1.1. **Antenna.** Pedicel + flagellum length/mesoscutum width 1.6; pedicel + flagellum length/head width 1.4; lengths scape/ventral plaque 5.3; ventral plaque located in the middle of scape; scape length/width 3.0; lengths scape/head (dorsal view) 0.5; scape length/HE 0.9; length/width F1, F2, F3, F4, clava: 1.7, 1.7, 2.0, 2.0, 5.2; length dorsobasal setae on F1/length F1 1.5. **Mesosoma.** Length/width 1.4; mesoscutum length/width 0.6; mesoscutellum length/width 0.7; widths SMG/SLG 1.2; enclosed space between SMG length/width 1.7; lengths mesoscutum/mesoscutellum 1.4; lengths mesoscutellum/dorsellum 2.7; lengths mesosoma/gaster 0.8. **Wings.** CC length/width 25.0; lengths CC/MV 1.3; lengths MV/ST 2.0; lengths MV/PM 3.0; lengths PM/ST 0.7; submarginal vein with five setae on dorsal surface. **Gaster.** With lateral tufts of pale and flattened setae on Gt₆.

Eriastichus coelotis sp. nov.

<http://zoobank.org/D18D5E27-7C12-44CC-BBBB-0D18240246CB>

Figure 16

Type locality. Costa Rica, Heredia, Estación Biológica La Selva, 75 m, 10°26'N, 84°01'W, 23–24.ii.2005, J.S. Noyes leg.

Type specimen. *Holotype* male dried and glued to a paper card. Original labels: "COSTA RICA, Heredia, E.B. La Selva, 75 m, 10°26'N, 84°01'W, 23–24. ii.2005, J.S. Noyes", "HOLOTYPE *Eriastichus coelotis* Hansson" [red printed label], (NHMUK014431024).

Diagnosis (male). Head dark brown; antenna (Fig. 16): scape yellowish brown, ventral plaque on scape ca. 0.2 × as long as scape, antenna with dorsobasal setae on F1 1.1 × as long as F1; gaster with lateral tufts of pale and flattened setae on Gt₆.

Description (male holotype NHMUK014431024). Length 1.1 mm. Head dark brown. Antenna with scape yellowish brown, ventral plaque dark brown, pedicel and flagellum pale brown. Mesoscutum, mesoscutellum and propodeum dark brown; dorsellum pale brown. Legs yellowish brown with base of hind coxa brown. Gaster dark brown.

Head. Length/width in frontal view 0.8; width/length in dorsal view 2.3; POL/OOL 2.7; WM/MS 1.5; MS/HE 0.4; HE/head length in frontal view 0.7; widths head/mesoscutum 1.2. **Antenna.** Pedicel + flagellum length/mesoscutum width 1.7; pedicel + flagellum length/head width 1.5; lengths scape/ventral plaque 5.5; ventral plaque located in the middle of scape; scape length/width 3.4; lengths scape/head (dorsal view) 0.6; scape length/HE 0.8; length/width F1, F2, F3, F4, clava: 1.5, 1.8, 1.8,

2.0, 4.8; length dorsobasal setae on F1/length F1 1.1. **Mesosoma.** Length/width 1.5; mesoscutum length/width 0.6; mesoscutellum length/width 0.6; widths SMG/SLG 1.1; enclosed space between SMG length/width 1.8; lengths mesoscutum/mesoscutellum 1.5; lengths mesoscutellum/dorsellum 2.3; lengths mesosoma/gaster 0.8. **Wings.** CC length/width 22.5; lengths CC/MV 1.2; lengths MV/ST 2.1; lengths MV/PM 5.0; lengths PM/ST 0.4; submarginal vein with five setae on dorsal surface. **Gaster.** With lateral tufts of pale and flattened setae on Gt₆.

***Eriastichus colenis* sp. nov.**

<http://zoobank.org/2FA90FB6-94AF-4ABE-B176-ED2970A9177D>

Figure 14

Type locality. Costa Rica, Cartago, Humo, El Copal, 9°47'N, 83°45'W, 1050–1250 m, 29.ii–6.iii.2008, C. Hansson leg.

Type specimen. Holotype male dried and glued to a paper card. Original labels: “COSTA RICA, Cartago, Humo, El Copal, 9°47'N, 83°45'W, 1050–1250 m, 29.ii–6.iii.2008, C. Hansson”, “HOLOTYPE Eriastichus colenis Hansson” [red printed label], (MZLU:7032.1).

Diagnosis (male). Head yellowish brown with vertex and upper ½ of occiput dark brown; antenna (Fig. 14): scape yellowish brown, ventral plaque on scape ca. 0.4 × as long as scape, antenna with dorsobasal setae on F1 0.5 × as long as F1; gaster with lateral tufts of pale and flattened setae on Gt_{5–6}.

Description (male holotype MZLU:7032.1). Length of body 1.6 mm. Head yellowish brown with vertex and upper ½ of occiput dark brown. Antenna with scape yellowish brown, ventral plaque dark brown, pedicel and flagellum pale brown. Mesoscutum and propodeum dark brown; mesoscutellum dark brown with metallic tinges; dorsellum pale brown. Legs yellowish brown. Gaster dark brown.

Head. Length/width in frontal view 0.7; width/length in dorsal view 2.1; POL/OOL 1.9; WM/MS 1.5; MS/HE 0.6; HE/head length in frontal view 0.6; widths head/mesoscutum 1.1. **Antenna.** Pedicel + flagellum length/mesoscutum width 1.7; pedicel + flagellum length/head width 1.5; lengths scape/ventral plaque 2.7; ventral plaque located above the middle of scape; scape length/width 4.2; lengths scape/head (dorsal view) 0.8; scape length/HE 1.3; length/width F1, F2, F3, F4, clava: 2.5, 2.3, 2.2, 2.2, 5.0; length dorsobasal setae on F1/length F1 0.5. **Mesosoma.** Length/width 1.5; mesoscutum length/width 0.5; mesoscutellum length/width 0.8; widths SMG/SLG 1.3; enclosed space between SMG length/width 2.0; lengths mesoscutum/mesoscutellum 1.2; lengths mesoscutellum/dorsellum 3.3; lengths mesosoma/gaster 0.8.

Wings. CC length/width 21.7; lengths CC/MV 1.1; lengths MV/ST 1.9; lengths MV/PM 4.8; lengths PM/ST 0.4; submarginal vein with eight setae on dorsal surface.

Gaster. With lateral tufts of pale and flattened setae on Gt_{5–6}.

***Eriastichus copalensis* sp. nov.**

<http://zoobank.org/55B02464-2EB7-45F9-8862-78EBF38D0631>

Figure 15

Type locality. Costa Rica, Cartago, Humo, El Copal, 9°47'N, 83°45'W, 1050–1250 m, 29.ii–6.iii.2008, C. Hansson leg.

Type specimen. *Holotype* male dried and glued to a paper card. Original labels: "COSTA RICA, Cartago, Humo, El Copal, 9°47'N, 83°45'W, 1050–1250 m, 29.ii–6.iii.2008, C. Hansson", "HOLOTYPE Eriastichus copalensis Hansson" [red printed label], (MZLU:7033.1)

Diagnosis (male). Head dark brown; antenna (Fig. 15): scape brown, ventral plaque on scape ca. 0.2 × as long as scape, antenna with dorsobasal setae on F1 0.5 × as long as F1; gaster with lateral tufts of pale and flattened setae on Gt₆.

Description (male holotype MZLU:7033.1). Length of body 1.5 mm. Head dark brown. Antenna brown, ventral plaque dark brown. Mesoscutum, mesoscutellum and propodeum dark brown; dorsellum pale brown. Legs with coxae and femora dark brown, trochanters, tibiae and tarsi yellowish brown. Gaster dark brown.

Head. Length/width in frontal view 0.8; width/length in dorsal view 2.1; POL/OOL 2.5; WM/MS 1.7; MS/HE 0.5; HE/head length in frontal view 0.6; widths head/mesoscutum 1.2. **Antenna.** Pedicel + flagellum length/mesoscutum width 2.2; pedicel + flagellum length/head width 2.0; lengths scape/ventral plaque 6.0; ventral plaque located slightly above the middle of scape; scape length/width 3.8; lengths scape/head (dorsal view) 0.5; scape length/HE 0.8; length/width F1, F2, F3, F4, clava: 3.0, 3.4, 3.8, 3.4, 8.0; length dorsobasal setae on F1/length F1 0.5. **Mesosoma.** Length/width 1.6; mesoscutum length/width 0.6; mesoscutellum length/width 0.8; widths SMG/SLG 1.2; enclosed space between SMG length/width 2.2; lengths mesoscutum/mesoscutellum 1.4; lengths mesoscutellum/dorsellum 3.0; lengths mesosoma/gaster 0.8. **Wings.** CC length/width 24.0; lengths CC/MV 1.1; lengths MV/ST 1.7; lengths MV/PM 3.1; lengths PM/ST 0.5; submarginal vein with seven setae on dorsal surface. **Gaster.** With lateral tufts of pale and flattened setae on Gt₆.

Etymology. Named after collecting locality.

***Eriastichus daptilis* sp. nov.**

<http://zoobank.org/C60FFDEF-AF6E-434F-AD51-8FA8464C025B>

Figure 17

Type locality. Costa Rica, Heredia, Estación Biológica La Selva, 75 m, 10°26'N, 84°01'W, 28–29.ii.2008, J.S. Noyes leg.

Type specimen. *Holotype* male dried and glued to a paper card. Original labels: "COSTA RICA, Heredia, E.B. La Selva, 75 m, 10°26'N, 84°01'W, 28–29.ii.2008,

J.S. Noyes, NHM (Ent) 2010–21”, “HOLOTYPE Eriastichus daptilis Hansson” [red printed label], (NHMUK014431025).

Diagnosis (male). Head and antenna dark brown; ventral plaque on scape ca. 0.2 × as long as scape (Fig. 17), antenna with dorsobasal setae on F1 1.1 × as long as F1; dorsellum and major parts of legs dark brown; gaster without lateral tufts on tergites.

Description (male holotype NHMUK014431025). Length of body 0.9 mm. Head and antenna dark brown, ventral plaque black. Mesoscutum, mesoscutellum, dorsellum and propodeum dark brown. Legs dark brown, trochanters and tarsi yellowish brown. Gaster dark brown.

Head. Length/width in frontal view 0.8; width/length in dorsal view 2.1; POL/OOL 1.3; WM/MS 4.2; MS/HE 0.7; HE/head length in frontal view 0.6; widths head/mesoscutum 1.0. **Antenna.** Pedicel + flagellum length/mesoscutum width 1.6; pedicel + flagellum length/head width 1.5; lengths scape/ventral plaque 5.0; ventral plaque located below the middle of scape; scape length/width 3.6; lengths scape/head (dorsal view) 0.5; scape length/HE 0.9; length/width F1, F2, F3, F4, clava: 1.6, 1.8, 1.6, 1.6, 5.2; length dorsobasal setae on F1/length F1 1.1. **Mesosoma.** Length/width 1.3; mesoscutum length/width 0.5; mesoscutellum length/width 0.7; widths SMG/SLG 1.1; enclosed space between SMG length/width 1.9; lengths mesoscutum/mesoscutellum 1.3; lengths mesoscutellum/dorsellum 2.5; lengths mesosoma/gaster 0.9.

Wings. CC length/width 16.3; lengths CC/MV 1.2; lengths MV/ST 1.7; lengths MV/PM 2.8; lengths PM/ST 0.6; submarginal vein with four setae on dorsal surface.

Gaster. Without lateral tufts on tergites.

Eriastichus decoris sp. nov.

<http://zoobank.org/A40C4518-95ED-4927-B6E7-5E3A49D80A33>

Figures 3, 18

Type locality. Costa Rica, San José, San Gerardo de Dota, 9°33'N, 83°47'W, 20–21. ii.2013, J.S. Noyes leg.

Type specimen. *Holotype* male dried and glued to a paper card. Original labels: “COSTA RICA, San José, San Gerardo de Dota, 20–21.ii.2013, J.S. Noyes, NHM (Ent) 2012–91”, “HOLOTYPE Eriastichus decoris Hansson” [red printed label], (NHMUK014431026).

Diagnosis (male). Head black and antenna dark brown; ventral plaque on scape ca. 0.3 × as long as scape (Fig. 18), antenna with dorsobasal setae on F1 0.5 × as long as F1; gaster with lateral tufts of pale and flattened setae on Gt_{5-6} .

Description (male holotype NHMUK014431026). Length of body 2.3 mm. Head black, antenna dark brown, ventral plaque black. Mesoscutum and mesoscutellum black with metallic tinges, dorsellum and propodeum black. Legs with coxae, hind femur and 4th tarsomere on all legs dark brown, fore femur pale brown, trochanters, tibiae, mid femur, and tarsomeres 1–3 yellowish brown. Gaster dark brown.

Head. Length/width in frontal view 0.8; width/length in dorsal view 2.1; POL/OOL 2.0; WM/MS 1.5; MS/HE 0.6; HE/head length in frontal view 0.5; widths

head/mesoscutum 1.2. **Antenna.** Pedicel + flagellum length/mesoscutum width 2.1; pedicel + flagellum length/head width 2.1; lengths scape/ventral plaque 3.1; ventral plaque located below the middle of scape; scape length/width 3.4; lengths scape/head (dorsal view) 0.5; scape length/HE 0.9; length/width F1, F2, F3, F4, clava: 3.7, 4.0, 3.7, 3.7, 7.9; length dorsobasal setae on F1/length F1 0.5. **Mesosoma.** Length/width 1.6; mesoscutum length/width 0.7; mesoscutellum length/width 0.8; widths SMG/SLG 1.1; enclosed space between SMG length/width 2.5; lengths mesoscutum/mesoscutellum 1.6; lengths mesoscutellum/dorsellum 2.6; lengths mesosoma/gaster 0.7. **Wings.** CC length/width 18.0; lengths CC/MV 0.9; lengths MV/ST 2.3; lengths MV/PM 3.8; lengths PM/ST 0.6; submarginal vein with nine setae on dorsal surface. **Gaster.** With lateral tufts of pale and flattened setae on Gt₅₋₆.

Eriastichus denotatis sp. nov.

<http://zoobank.org/306B5D9A-CBB9-447E-A4BD-57138BE907CD>

Figures 19, 60

Type locality. Costa Rica, San José, San Gerardo de Dota, 9°33'N, 83°47'W, 20–21. ii.2013, J.S. Noyes leg.


Type specimen. *Holotype* male dried and glued to a paper card. Original labels: "COSTA RICA, San José, San Gerardo de Dota, 20–21.ii.2013, J.S. Noyes, NHM (Ent) 2012–91", "HOLOTYPE Eriastichus denotatis Hansson" [red printed label], (NHMUK014431027).

Additional type material. *Paratype* 1♂ with same label data as holotype (NHMUK014431028).

Diagnosis (male). Head black (Fig. 60) and antenna dark brown (Fig. 19); ventral plaque on scape ca. 0.4 × as long as scape (Fig. 19), antenna with dorsobasal setae on F1 0.6 × as long as F1; gaster with lateral tufts of pale and flattened setae on Gt₆.

Description (male holotype NHMUK014431027). Length of body 2.6 mm (paratype 2.3 mm). Head black, antenna dark brown, ventral plaque black. Mesoscutum and mesoscutellum black with metallic tinges, dorsellum and propodeum black. Legs with hind coxa dark brown, hind femur brown, fore and mid coxae and femora, trochanters, tibiae and tarsi yellowish brown. Gaster dark brown.

Head. Length/width in frontal view 0.8; width/length in dorsal view 2.1; POL/OOL 1.6; WM/MS 1.5; MS/HE 0.6; HE/head length in frontal view 0.5; widths head/mesoscutum 1.1. **Antenna.** Pedicel + flagellum length/mesoscutum width 2.4; pedicel + flagellum length/head width 2.3; lengths scape/ventral plaque 2.4; ventral plaque located in the middle of scape; scape length/width 3.0; lengths scape/head (dorsal view) 0.5; scape length/HE 1.0; length/width F1, F2, F3, F4, clava: 3.1, 3.4, 3.4, 3.4, 7.3; length dorsobasal setae on F1/length F1 0.6. **Mesosoma.** Length/width 1.8; mesoscutum length/width 0.8; mesoscutellum length/width 0.9; widths SMG/SLG 1.2; enclosed space between SMG length/width 2.9; lengths mesoscutum/mesoscutellum 1.6; lengths mesoscutellum/dorsellum 2.7; lengths mesosoma/gaster 0.7. **Wings.** CC length/width 21.7; lengths CC/MV 0.9; lengths MV/ST 2.3; lengths MV/PM



Figures 18–27. *Eriastichus* spp. head and antenna in lateral view, male holotypes **18** *E. decoris* **19** *E. dentotatis* **20** *E. dotaensis* **21** *E. derilis* **22** *E. diadrys* **23** *E. ebulis* **24** *E. egestis* **25** *E. ellipsis* **26** *E. drupis* **27** *E. eleagnis*.

3.9; lengths PM/ST 0.6; submarginal vein with nine setae (ten setae in paratype) on dorsal surface. **Gaster.** With lateral tufts of pale and flattened setae on Gt₆.

***Eriastichus derilis* sp. nov.**

<http://zoobank.org/59EEBC97-92F5-43EB-9205-3816D5D672B0>

Figure 21

Type locality. Costa Rica, Cartago, Humo, El Copal, 9°47'N, 83°45'W, 1050–1250 m, 29.ii–6.iii.2008, C. Hansson leg.

Type specimen. **Holotype** male dried and glued to a paper card. Original labels: "COSTA RICA, Cartago, Humo, El Copal, 9°47'N, 83°45'W, 1050–1250 m, 29.ii–6.iii.2008, C. Hansson", "HOLOTYPE Eriastichus derilis Hansson" [red printed label], (MZLU:7034.1).

Diagnosis (male). Head with lower ½ yellowish brown and upper ½ dark brown, scape yellowish brown, pedicel and flagellum brown; ventral plaque on scape ca. 0.5 × as long as scape (Fig. 21), antenna with dorsobasal setae on F1 0.6 × as long as F1; gaster with lateral tufts of pale and flattened setae on Gt₅₋₆.

Description (male holotype MZLU:7034.1). Length of body 1.2 mm. Head with lower ½ yellowish brown and upper ½ dark brown, scape yellowish brown, pedicel and flagellum brown, ventral plaque dark brown. Mesoscutum, mesoscutellum, dorsellum and propodeum dark brown. Legs yellowish brown. Gaster dark brown.

Head. Length/width in frontal view 0.8; width/length in dorsal view 2.3; POL/OOL 1.9; WM/MS 1.6; MS/HE 0.5; HE/head length in frontal view 0.6; widths head/mesoscutum 1.2. **Antenna.** Pedicel + flagellum length/mesoscutum width 1.7; pedicel + flagellum length/head width 1.5; lengths scape/ventral plaque 2.0; ventral plaque located above the middle of scape; scape length/width 3.0; lengths scape/head (dorsal view) 0.5; scape length/HE 0.9; length/width F1, F2, F3, F4, clava: 2.0, 2.0, 2.0, 2.0, 4.3; length dorsobasal setae on F1/length F1 0.6. **Mesosoma.** Length/width 1.4; mesoscutum length/width 0.6; mesoscutellum length/width 0.9; widths SMG/SLG 1.2; enclosed space between SMG length/width 2.2; lengths mesoscutum/mesoscutellum 1.3; lengths mesoscutellum/dorsellum 4.0; lengths mesosoma/gaster 0.8.

Wings. CC length/width 21.0; lengths CC/MV 1.1; lengths MV/ST 1.8; lengths MV/PM 5.0; lengths PM/ST 0.4; submarginal vein with six setae on dorsal surface. **Gaster.** With lateral tufts of pale and flattened setae on Gt₅₋₆.

***Eriastichus diadrys* sp. nov.**

<http://zoobank.org/08F408C7-5C9C-4BFA-9A92-C11C2A11C881>

Figure 22

Type locality. Costa Rica, San José, San Gerardo de Dota, 9°33'N, 83°47'W, 20–21. ii.2013, J.S. Noyes leg.

Type specimen. *Holotype* male dried and glued to a paper card. Original labels: "COSTA RICA, San José, San Gerardo de Dota, 20–21.ii.2013, J.S. Noyes, NHM (Ent) 2012–91", "HOLOTYPE Eriastichus diadrys Hansson" [red printed label], (NHMUK014431029).

Diagnosis (male). Head black and antenna dark brown; ventral plaque on scape ca. $0.5 \times$ as long as scape (Fig. 22), antenna with dorsobasal setae on F1 $0.5 \times$ as long as F1; gaster with lateral tufts of pale and flattened setae on Gt_{3–6}.

Description (male holotype NHMUK014431029). Length of body 2.1 mm. Head black, antenna dark brown, ventral plaque black. Mesoscutum and mesoscutellum black with metallic tinges, dorsellum and propodeum black. Legs with hind coxa and hind femur dark brown, fore and mid coxae yellowish brown with base dark brown, trochanters, tibiae and tarsi yellowish brown. Gaster dark brown.

Head. Length/width in frontal view 0.7; width/length in dorsal view 2.2; POL/OOL 2.0; WM/MS 1.9; MS/HE 0.5; HE/head length in frontal view 0.6; widths head/mesoscutum 1.1. **Antenna.** Pedicel + flagellum length/mesoscutum width 1.9; pedicel + flagellum length/head width 1.8; lengths scape/ventral plaque 2.1; ventral plaque located in the middle of scape; scape length/width 2.7; lengths scape/head (dorsal view) 0.5; scape length/HE 0.9; length/width F1, F2, F3, F4, clava: 2.7, 2.9, 2.7, 2.6, 5.7; length dorsobasal setae on F1/length F1 0.5. **Mesosoma.** Length/width 1.6; mesoscutum length/width 0.7; mesoscutellum length/width 0.8; widths SMG/SLG 1.5; enclosed space between SMG length/width 2.0; lengths mesoscutum/mesoscutellum 1.6; lengths mesoscutellum/dorsellum 2.9; lengths mesosoma/gaster 0.8. **Wings.** CC length/width 20.6; lengths CC/MV 1.0; lengths MV/ST 2.1; lengths MV/PM 4.8; lengths PM/ST 0.4; submarginal vein with eight setae on dorsal surface. **Gaster.** With lateral tufts of pale and flattened setae on Gt_{3–6}.

Eriastichus dotaensis sp. nov.

<http://zoobank.org/E2228FB1-DE88-47A6-A504-CDEDE886BB50>

Figure 20

Type locality. Costa Rica, San José, San Gerardo de Dota, 9°33'N, 83°47'W, 20–21.ii.2013, J.S. Noyes leg.

Type specimen. *Holotype* male dried and glued to a paper card. Original labels: "COSTA RICA, San José, San Gerardo de Dota, 20–21.ii.2013, J.S. Noyes, NHM (Ent) 2012–91", "HOLOTYPE Eriastichus dotaensis Hansson" [red printed label], (NHMUK014431030).

Diagnosis (male). Head and antenna dark brown; ventral plaque on scape ca. $0.4 \times$ as long as scape (Fig. 20), antenna with dorsobasal setae on F1 $0.8 \times$ as long as F1; gaster with lateral tufts of pale and flattened setae on Gt_{3–6}.

Description (male holotype NHMUK014431030). Length of body 1.6 mm. Head and antenna dark brown, ventral plaque black. Mesoscutum and mesoscutellum black with metallic tinges, dorsellum pale brown, propodeum dark brown. Legs with

coxae and femora dark brown, trochanters, tibiae and tarsi yellowish brown. Gaster dark brown.

Head. Length/width in frontal view 0.7; width/length in dorsal view 2.2; POL/OOL 1.7; WM/MS 1.5; MS/HE 0.6; HE/head length in frontal view 0.6; widths head/mesoscutum 1.2. **Antenna.** Pedicel + flagellum length/mesoscutum width 2.1; pedicel + flagellum length/head width 1.9; lengths scape/ventral plaque 2.7; ventral plaque located below the middle of scape; scape length/width 3.0; lengths scape/head (dorsal view) 0.6; scape length/HE 1.0; length/width F1, F2, F3, F4, clava: 2.5, 2.8, 2.7, 2.7, 6.4; length dorsobasal setae on F1/length F1 0.8. **Mesosoma.** Length/width 1.5; mesoscutum length/width 0.7; mesoscutellum length/width 0.8; widths SMG/SLG 1.1; enclosed space between SMG length/width 2.2; lengths mesoscutum/mesoscutellum 1.6; lengths mesoscutellum/dorsellum 2.6; lengths mesosoma/gaster 0.8. **Wings.** CC length/width 26.0; lengths CC/MV 1.0; lengths MV/ST 2.2; lengths MV/PM 3.6; lengths PM/ST 0.6; submarginal vein with six setae on dorsal surface. **Gaster.** With lateral tufts of pale and flattened setae on Gt₆.

Etymology. Named after collecting locality.

Eriastichus drupis sp. nov.

<http://zoobank.org/4CF0E18A-2BE0-4E3F-BB19-378178073525>

Figure 26

Type locality. Costa Rica, Puntarenas, San Vito, Estación Biológica Las Alturas, 1500 m, 8°57'N, 82°50'W, 17–18.ii.2012, J.S. Noyes leg.

Type specimen. *Holotype* male dried and glued to a paper card. Original labels: "COSTA RICA, Puntarenas, San Vito, E.B. Las Alturas, 1500 m, 8°57'N, 82°50'W, 17–18.ii.2012, J.S. Noyes, NHM (Ent) 2012–91", "HOLOTYPE *Eriastichus drupis* Hansson" [red printed label], (NHMUK014431031).

Diagnosis (male). Head dark brown, scape and pedicel pale brown, flagellum brown; ventral plaque on scape ca. 0.4 × as long as scape (Fig. 26), antenna with dorsobasal setae on F1 0.8 × as long as F1; gaster with lateral tufts of pale and flattened setae on Gt_{5–6}.

Description (male holotype NHMUK014431031). Length of body 1.4 mm. Head dark brown, scape and pedicel pale brown, flagellum brown, ventral plaque dark brown. Mesoscutum and mesoscutellum black with metallic tinges, dorsellum pale brown, propodeum dark brown. Legs with hind coxa and hind femur dark brown, fore and mid coxae yellowish brown with base brown, remaining parts of legs yellowish brown. Gaster dark brown.

Head. Length/width in frontal view 0.8; width/length in dorsal view 2.2; POL/OOL 1.8; WM/MS 1.6; MS/HE 0.5; HE/head length in frontal view 0.6; widths head/mesoscutum 1.3. **Antenna.** Pedicel + flagellum length/mesoscutum width 1.7; pedicel + flagellum length/head width 1.6; lengths scape/ventral plaque 2.8; ventral plaque located below the middle of scape; scape length/width 3.1; lengths scape/head

(dorsal view) 0.5; scape length/HE 0.9; length/width F1, F2, F3, F4, clava: 2.7, 2.3, 2.3, 2.2, 5.3; length dorsobasal setae on F1/length F1 0.8. **Mesosoma.** Length/width 1.5; mesoscutum length/width 0.5; mesoscutellum length/width 0.7; widths SMG/SLG 1.1; enclosed space between SMG length/width 2.1; lengths mesoscutum/mesoscutellum 1.3; lengths mesoscutellum/dorsellum 3.1; lengths mesosoma/gaster 0.9. **Wings.** CC length/width 22.5; lengths CC/MV 1.1; lengths MV/ST 1.9; lengths MV/PM 4.8; lengths PM/ST 0.4; submarginal vein with six setae on dorsal surface. **Gaster.** With lateral tufts of pale and flattened setae on Gt₅₋₆.

Eriastichus ebulis sp. nov.

<http://zoobank.org/44D3C6E8-F596-4AA4-BFD7-DCA4E8D2EC18>

Figures 8, 10, 23

Type locality. Costa Rica, San José, San Gerardo de Dota, 9°33'N, 83°47'W, 20–21. ii.2013, J.S. Noyes leg.

Type specimen. *Holotype* male dried and glued to a paper card. Original labels: "COSTA RICA, San José, San Gerardo de Dota, 20–21.ii.2013, J.S. Noyes, NHM (Ent) 2012–91", "HOLOTYPE Eriastichus ebulis Hansson" [red printed label], (NHMUK014431032).

Additional type material. *Paratypes* (4♂♂): 2♂♂ with same label data as holotype (NHMUK014431033, NHMUK014431034); 2♂♂ "COSTA RICA, Cartago, Humo, El Copal, 9°47'N, 83°45'W, 1050–1250 m, 29.ii–6.iii.2008, C. Hansson" (MZLU:7035.2, MZUCR:01693).

Diagnosis (male). Head dark brown, antenna brown; ventral plaque on scape ca. 0.3 × as long as scape (Fig. 23), antenna with dorsobasal setae on F1 0.5 × as long as F1; gaster with lateral tufts of pale and flattened setae on Gt₅₋₆.

Description (male holotype NHMUK014431032). Length of body 1.5 mm (paratypes 1.6–2.0 mm). Head dark brown, antenna brown, ventral plaque black. Mesoscutum and mesoscutellum black with metallic tinges, dorsellum and propodeum black. Legs with coxae, femora and 4th tarsomere on all legs dark brown, trochanters, tibiae and tarsomeres 1–4 yellowish brown. Gaster dark brown.

Head. Length/width in frontal view 0.8; width/length in dorsal view 2.0; POL/OOL 1.5; WM/MS 1.4; MS/HE 0.6; HE/head length in frontal view 0.6; widths head/mesoscutum 1.0. **Antenna.** Pedicel + flagellum length/mesoscutum width 2.3; pedicel + flagellum length/head width 1.9; lengths scape/ventral plaque 3.0; ventral plaque located in the middle of scape; scape length/width 3.0; lengths scape/head (dorsal view) 0.5; scape length/HE 0.9; length/width F1, F2, F3, F4, clava: 3.2, 3.0, 3.0, 3.0, 7.8; length dorsobasal setae on F1/length F1 0.5. **Mesosoma.** Length/width 1.8; mesoscutum length/width 0.8; mesoscutellum length/width 0.8; widths SMG/SLG 1.6; enclosed space between SMG length/width 2.2; lengths mesoscutum/mesoscutellum 1.6; lengths mesoscutellum/dorsellum 2.7; lengths mesosoma/gaster 0.8. **Wings.** CC length/width 26.0; lengths CC/MV 1.0; lengths MV/ST 1.9; lengths MV/PM

7.4; lengths PM/ST 0.3; submarginal vein with seven setae (5–9 setae in paratypes) on dorsal surface. **Gaster.** With lateral tufts of pale and flattened setae on Gt_{5–6}.

***Eriastichus egrestis* sp. nov.**

<http://zoobank.org/B875A5F8-F524-407E-8D2F-6A874BE758FA>

Figure 24

Type locality. Costa Rica, Puntarenas, San Vito, Las Cruces, 8°46'N, 82°57'W, 1300 m, 15–16.ii.2006, J.S. Noyes leg.

Type specimen. *Holotype* male dried and glued to a paper card. Original labels: "COSTA RICA, Puntarenas, San Vito, Las Cruces, 8°46'N, 82°57'W, 1300 m, 15–16. ii.2006, J.S. Noyes", "HOLOTYPE Eriastichus egrestis Hansson" [red printed label], (NHMUK014431035).

Diagnosis (male). Head dark brown, antenna brown; ventral plaque on scape ca. 0.2 × as long as scape (Fig. 24), antenna with dorsobasal setae on F1 1.3 × as long as F1; gaster with lateral tufts of pale and flattened setae on Gt₆.

Description (male holotype NHMUK014431035). Length of body 1.0 mm. Head dark brown, antenna brown, ventral plaque dark brown. Mesoscutum and mesoscutellum dark brown with metallic tinges, dorsellum and propodeum dark brown. Legs with coxae and femora brown, trochanters, tibiae and tarsi yellowish brown. Gaster dark brown.

Head. Length/width in frontal view 0.8; width/length in dorsal view 2.5; POL/OOL 2.4; WM/MS 1.5; MS/HE 0.6; HE/head length in frontal view 0.5; widths head/mesoscutum 1.3. **Antenna.** Pedicel + flagellum length/mesoscutum width 1.4; pedicel + flagellum length/head width 1.1; lengths scape/ventral plaque 4.5; ventral plaque located below the middle of scape; scape length/width 3.6; lengths scape/head (dorsal view) 0.5; scape length/HE 0.9; length/width F1, F2, F3, F4, clava: 1.4, 1.6, 1.6, 1.6, 4.3; length dorsobasal setae on F1/length F1 1.3. **Mesosoma.** Length/width 1.5; mesoscutum length/width 0.6; mesoscutellum length/width 0.6; widths SMG/SLG 1.4; enclosed space between SMG length/width 1.5; lengths mesoscutum/mesoscutellum 1.5; lengths mesoscutellum/dorsellum 2.5; lengths mesosoma/gaster 0.8. **Wings.** CC length/width 14.5; lengths costal cell/MV 1.0; lengths MV/ST 2.0; lengths MV/PM 3.5; lengths PM/ST 0.6; submarginal vein with five setae on dorsal surface. **Gaster.** With lateral tufts of pale and flattened setae on Gt₆.

***Eriastichus eleagnis* sp. nov.**

<http://zoobank.org/81F12226-5418-40DF-8BD8-D872B5B814B4>

Figures 1, 27

Type locality. Costa Rica, Puntarenas, Estación Biológica Monteverde, 10°20'N, 84°49'W, 1540 m, 18–25.ii.2004, C. Hansson leg.

Type specimen. *Holotype* male dried and glued to a paper card. Original labels: "COSTA RICA, Puntarenas, E.B. Monteverde, 10°20'N, 84°49'W, 1540 m, 18–25. ii.2004, C. Hansson", "HOLOTYPE Eriastichus eleagnis Hansson" [red printed label], (MZLU:7036.1).

Additional type material. *Paratypes* (4♂♂): 1♂ with same label data as holotype (MZLU:7036.2); 1♂ from same locality as holotype but collected 26.ii.2007 (NHMUK014431036); 1♂ "COSTA RICA, Puntarenas, San Vito, E.B. Las Alturas, 1500 m, 8°57'N, 82°50'W, 17–18.ii.2012, J.S. Noyes, NHM (Ent) 2012–91" (NHMUK014431037); 1♂ "COSTA RICA, San José, San Gerardo de Dota, 20–21. ii.2013, J.S. Noyes, NHM (Ent) 2012–91" (NHMUK014431038).

Diagnosis (male). Head dark brown, scape yellowish brown, pedicel and flagellum brown; ventral plaque on scape ca. 0.4 × as long as scape (Fig. 27), antenna with dorsobasal setae on F1 1.5 × as long as F1; gaster with lateral tufts of pale and flattened setae on Gt₆.

Description (male holotype MZLU:7036.1). Length of body 1.4 mm (paratypes 1.3–1.5 mm). Head dark brown, scape yellowish brown, pedicel and flagellum brown, ventral plaque dark brown. Mesoscutum, mesoscutellum and propodeum dark brown, dorsellum pale brown. Legs with coxae and femora brown, trochanters, tibiae and tarsi yellowish brown. Gaster dark brown.

Head. Length/width in frontal view 0.8; width/length in dorsal view 2.4; POL/OOL 2.4; WM/MS 1.5; MS/HE 0.5; HE/head length in frontal view 0.6; widths head/mesoscutum 1.1. **Antenna.** Pedicel + flagellum length/mesoscutum width 1.8; pedicel + flagellum length/head width 1.7; lengths scape/ventral plaque 2.8; ventral plaque located below the middle of scape; scape length/width 3.1; lengths scape/head (dorsal view) 0.6; scape length/HE 1.0; length/width F1, F2, F3, F4, clava: 2.6, 3.0, 3.0, 2.8, 7.6; length dorsobasal setae on F1/length F1 1.5. **Mesosoma.** Length/width 1.4; mesoscutum length/width 0.6; mesoscutellum length/width 0.8; widths SMG/SLG 1.1; enclosed space between SMG length/width 2.1; lengths mesoscutum/mesoscutellum 1.6; lengths mesoscutellum/dorsellum 2.9; lengths mesosoma/gaster 0.8.

Wings. CC length/width 24.0; lengths CC/MV 1.1; lengths MV/ST 1.9; lengths MV/PM 3.8; lengths PM/ST 0.5; submarginal vein with seven setae (5–7 setae in paratypes) on dorsal surface. **Gaster.** With lateral tufts of pale and flattened setae on Gt₆.

Eriastichus ellipsis sp. nov.

<http://zoobank.org/B8933986-9084-433F-B5BC-3BB531D04D8E>

Figure 25

Type locality. Costa Rica, Heredia, Estación Biológica La Selva, 75 m, 10°26'N, 84°01'W, 24–25.ii.2011, J.S. Noyes leg.

Type specimen. *Holotype* male dried and glued to a paper card. Original labels: "COSTA RICA, Heredia, E.B. La Selva, 75 m, 10°26'N, 84°01'W, 24–25.ii.2011,

J.S. Noyes, NHM (Ent) 2011–93”, “HOLOTYPE *Eriastichus ellipsis* Hansson” [red printed label], (NHMUK014431039).

Additional type material. *Paratype* 1♂ with same label data as holotype (NHMUK014431040).

Diagnosis (male). Head dark brown with part below antennal toruli yellowish brown, scape yellowish brown, pedicel and flagellum brown; ventral plaque on scape ca. 0.2 × as long as scape (Fig. 25), antenna with dorsobasal setae on F1 0.6 × as long as F1; gaster with lateral tufts of pale and flattened setae on Gt₆.

Description (male holotype NHMUK014431039). Length of body 1.1 mm (paratype 1.1 mm). Head dark brown with part below antennal toruli yellowish brown, scape yellowish brown, pedicel and flagellum brown, ventral plaque dark brown. Mesoscutum and mesoscutellum black with metallic tinges, dorsellum pale brown, propodeum dark brown. Legs yellowish brown. Gaster dark brown.

Head. Length/width in frontal view 0.7; width/length in dorsal view 2.3; POL/OOL 3.5; WM/MS 1.3; MS/HE 0.5; HE/head length in frontal view 0.6; widths head/mesoscutum 1.2. **Antenna.** Pedicel + flagellum length/mesoscutum width 1.8; pedicel + flagellum length/head width 1.5; lengths scape/ventral plaque 4.8; ventral plaque located in the middle of scape; scape length/width 3.2; lengths scape/head (dorsal view) 0.5; scape length/HE 0.8; length/width F1, F2, F3, F4, clava: 1.5, 2.0, 2.2, 2.2, 6.0; length dorsobasal setae on F1/length F1 0.6. **Mesosoma.** Length/width 1.3; mesoscutum length/width 0.6; mesoscutellum length/width 0.7; widths SMG/SLG 1.4; enclosed space between SMG length/width 1.7; lengths mesoscutum/mesoscutellum 1.5; lengths mesoscutellum/dorsellum 2.8; lengths mesosoma/gaster 0.8. **Wings.** CC length/width 21.3; lengths CC/MV 1.1; lengths MV/ST 2.0; lengths MV/PM 5.0; lengths PM/ST 0.4; submarginal vein with six setae (five setae in paratype) on dorsal surface. **Gaster.** With lateral tufts of pale and flattened setae on Gt₆.

Eriastichus eminis sp. nov.

<http://zoobank.org/91B5BBE0-7453-4F7D-BB97-FD1DFC02224F>

Figure 31

Type locality. Costa Rica, Guanacaste, Estación Pitilla, Área Conservación de Guanacaste, 10°59'N, 85°26'W, 21.ii.2011, J.S. Noyes leg.

Type specimen. *Holotype* male dried and glued to a paper card. Original labels: “COSTA RICA, Guanacaste, Est. Pitilla, ACG, 21.ii.2011, J.S. Noyes, NHM (Ent) 2011–93”, “HOLOTYPE *Eriastichus eminis* Hansson” [red printed label], (NHMUK014431041).

Diagnosis (male). Head dark brown with part below antennal toruli dark yellowish brown, scape yellowish brown, pedicel and flagellum brown; ventral plaque on scape ca. 0.2 × as long as scape (Fig. 31), antenna with dorsobasal setae on F1 1.1 × as long as F1; gaster with lateral tufts of pale and flattened setae on Gt₆.

Description (male holotype NHMUK014431041). Length of body 1.2 mm. Head dark brown with part below antennal toruli dark yellowish brown, scape yellowish brown, pedicel and flagellum brown, ventral plaque dark brown. Mesoscutum, mesoscutellum and propodeum dark brown, dorsellum pale brown. Legs yellowish brown. Gaster dark brown.

Head. Length/width in frontal view 0.8; width/length in dorsal view 2.2; POL/OOL 2.2; WM/MS 1.5; MS/HE 0.5; HE/head length in frontal view 0.6; widths head/mesoscutum 1.2. **Antenna.** Pedicel + flagellum length/mesoscutum width 2.4; pedicel + flagellum length/head width 1.9; lengths scape/ventral plaque 4.5; ventral plaque located below the middle of scape; scape length/width 2.9; lengths scape/head (dorsal view) 0.5; scape length/HE 0.8; length/width F1, F2, F3, F4, clava: 3.5, 3.5, 3.3, 3.5, 8.0; length dorsobasal setae on F1/length F1 1.1. **Mesosoma.** Length/width 1.7; mesoscutum length/width 0.7; mesoscutellum length/width 0.9; widths SMG/SLG 1.3; enclosed space between SMG length/width 2.2; lengths mesoscutum/mesoscutellum 1.5; lengths mesoscutellum/dorsellum 2.9; lengths mesosoma/gaster 0.8.

Wings. CC length/width 26.3; lengths CC/MV 1.1; lengths MV/ST 2.1; lengths MV/PM 5.7; lengths PM/ST 0.4; submarginal vein with six setae on dorsal surface. **Gaster.** With lateral tufts of pale and flattened setae on Gt₆.

Eriastichus facilis sp. nov.

<http://zoobank.org/1759534A-5BB1-4670-8391-9AD918D65A24>

Figure 28

Type locality. Costa Rica, Puntarenas, Estación Biológica Monteverde, 10°20'N, 84°49'W, 1540 m, 26.ii.2007, J.S. Noyes leg.

Type specimen. *Holotype* male dried and glued to a paper card. Original labels: "COSTA RICA, Puntarenas, E.B. Monteverde, 10°20'N, 84°49'W, 1540 m, 26.ii.2007, J.S. Noyes", "HOLOTYPE Eriastichus facilis Hansson" [red printed label], (NHMUK014431042).

Additional type material. *Paratypes* (2♂♂): 1♂ "COSTA RICA, Cartago, Humo, El Copal, 9°47'N, 83°45'W, 1050–1250 m, 29.ii–6.iii.2008, C. Hansson" (MZLU:7037.2); 1♂ "COSTA RICA, Alajuela, Est. Pilón, 10°43'N, 85°59'W, 700 m, 12–18.ii.2003, C. Hansson & J.A. Azofeifa" (MZLU:7037.3).

Diagnosis (male). Head dark brown, scape yellowish brown, pedicel and flagellum brown; ventral plaque on scape ca. 0.2 × as long as scape (Fig. 28), antenna with dorsobasal setae on F1 0.9 × as long as F1; gaster with lateral tufts of pale and flattened setae on Gt₆.

Description (male holotype NHMUK014431042). Length of body 1.5 mm (paratypes 1.0–1.1 mm). Head dark brown, scape yellowish brown, pedicel and flagellum brown, ventral plaque dark brown. Mesoscutum, mesoscutellum and propodeum dark brown, dorsellum pale brown. Legs with hind coxa and hind femur dark brown, remaining parts of legs yellowish brown. Gaster dark brown.

Head. Length/width in frontal view 0.8; width/length in dorsal view 2.4; POL/OOL 2.6; WM/MS 1.4; MS/HE 0.5; HE/head length in frontal view 0.6; widths



Figures 28–34. *Eriastichus* spp. head and antenna in lateral view, male holotypes **28** *E. facilis* **29** *E. fenestris* **30** *E. follis* **31** *E. eminis* **32** *E. glanis* **33** *E. geratis* **34** *E. galeatis*.

head/mesoscutum 1.0. **Antenna.** Pedicel + flagellum length/mesoscutum width 1.7; pedicel + flagellum length/head width 1.6; lengths scape/ventral plaque 4.7; ventral plaque located below the middle of scape; scape length/width 3.5; lengths scape/head

(dorsal view) 0.5; scape length/HE 0.9; length/width F1, F2, F3, F4, clava: 2.2, 2.2, 2.5, 2.5, 6.3; length dorsobasal setae on F1/length F1 0.9. **Mesosoma.** Length/width 1.4; mesoscutum length/width 0.6; mesoscutellum length/width 0.7; widths SMG/SLG 1.2; enclosed space between SMG length/width 2.1; lengths mesoscutum/mesoscutellum 1.4; lengths mesoscutellum/dorsellum 2.7; lengths mesosoma/gaster 0.9. **Wings.** CC length/width 18.0; lengths CC/MV 1.2; lengths MV/ST 2.0; lengths MV/PM 3.8; lengths PM/ST 0.5; submarginal vein with five setae on dorsal surface, in all types. **Gaster.** With lateral tufts of pale and flattened setae on Gt₄₋₆.

Eriastichus fenestratus sp. nov.

<http://zoobank.org/68B54220-6BD6-40CE-9CC0-CEE6962E631>

Figures 5, 29

Type locality. Costa Rica, San José, San Gerardo de Dota, 9°33'N, 83°47'W, 20–21. ii.2013, J.S. Noyes leg.

Type specimen. *Holotype* male dried and glued to a paper card. Original labels: "COSTA RICA, San José, San Gerardo de Dota, 20–21.ii.2013, J.S. Noyes, NHM (Ent) 2012–91", "HOLOTYPE Eriastichus fenestratus Hansson" [red printed label], (NHMUK014431043).

Diagnosis (male). Head black, scape brown, pedicel and flagellum dark brown; ventral plaque on scape ca. 0.6 × as long as scape (Fig. 29), antenna with dorsobasal setae on F1 0.7 × as long as F1; gaster with lateral tufts of pale and flattened setae on Gt₄₋₆.

Description (male holotype NHMUK014431043). Length of body 2.1 mm. Head black, scape brown, pedicel, flagellum and ventral plaque dark brown. Mesoscutum, mesoscutellum and dorsellum black with metallic tinges, propodeum black. Legs with fore and mid coxae yellowish brown with base dark brown, hind coxa black, remaining parts of legs yellowish brown. Gaster dark brown.

Head. Length/width in frontal view 0.7; width/length in dorsal view 2.2; POL/OOL 1.8; WM/MS 1.6; MS/HE 0.6; HE/head length in frontal view 0.6; widths head/mesoscutum 1.0. **Antenna.** Pedicel + flagellum length/mesoscutum width 2.1; pedicel + flagellum length/head width 2.1; lengths scape/ventral plaque 1.7; ventral plaque located in the middle of scape; scape length/width 2.6; lengths scape/head (dorsal view) 0.5; scape length/HE 0.9; length/width F1, F2, F3, F4, clava: 3.2, 3.0, 2.9, 3.0, 5.8; length dorsobasal setae on F1/length F1 0.7. **Mesosoma.** Length/width 1.6; mesoscutum length/width 0.6; mesoscutellum length/width 0.8; widths SMG/SLG 1.2; enclosed space between SMG length/width 2.2; lengths mesoscutum/mesoscutellum 1.5; lengths mesoscutellum/dorsellum 2.3; lengths mesosoma/gaster 1.0. **Wings.** CC length/width 19.0; lengths CC/MV 0.9; lengths MV/ST 2.1; lengths MV/PM 5.5; lengths PM/ST 0.4; submarginal vein with ten setae on dorsal surface. **Gaster.** With lateral tufts of pale and flattened setae on Gt₄₋₆.

***Eriastichus follis* sp. nov.**

<http://zoobank.org/F9EFCF3E-87E4-4157-9AE2-ADDA7AF0E7D3>

Figure 30

Type locality. Costa Rica, Puntarenas, Estación Biológica Monteverde, 10°20'N, 84°49'W, 1540 m, 26.ii.2007, J.S. Noyes leg.

Type specimen. *Holotype* male dried and glued to a paper card. Original labels: "COSTA RICA, Puntarenas, E.B. Monteverde, 10°20'N, 84°49'W, 1540 m, 26.ii.2007, J.S. Noyes", "HOLOTYPE *Eriastichus follis* Hansson" [red printed label], (NHMUK014431044).

Diagnosis (male). Head dark brown with part below antennal toruli pale brown, scape and pedicel pale brown, flagellum brown; ventral plaque on scape ca. 0.6 × as long as scape (Fig. 30), antenna with dorsobasal setae on F1 0.6 × as long as F1; gaster with lateral tufts of pale and flattened setae on Gt₆.

Description (male holotype NHMUK014431044). Length of body 1.5 mm. Head dark brown with part below antennal toruli pale brown, scape and pedicel pale brown, flagellum brown, ventral plaque dark brown. Mesoscutum and mesoscutellum dark brown with metallic tinges, dorsellum and propodeum dark brown. Legs with hind coxa pale brown, remaining parts of legs yellowish brown. Gaster dark brown.

Head. Length/width in frontal view 0.8; width/length in dorsal view 2.1; POL/OOL 1.9; WM/MS 1.2; MS/HE 0.5; HE/head length in frontal view 0.5; widths head/mesoscutum 1.2. **Antenna.** Pedicel + flagellum length/mesoscutum width 2.1; pedicel + flagellum length/head width 1.8; lengths scape/ventral plaque 1.7; ventral plaque located below the middle of scape; scape length/width 2.9; lengths scape/head (dorsal view) 0.5; scape length/HE 0.8; length/width F1, F2, F3, F4, clava: 2.8, 2.7, 2.7, 2.7, 6.3; length dorsobasal setae on F1/length F1 0.6. **Mesosoma.** Length/width 1.7; mesoscutum length/width 0.7; mesoscutellum length/width 0.8; widths SMG/SLG 1.2; enclosed space between SMG length/width 2.3; lengths mesoscutum/mesoscutellum 1.5; lengths mesoscutellum/dorsellum 2.8; lengths mesosoma/gaster 0.8.

Wings. CC length/width 16.7; lengths CC/MV 0.9; lengths MV/ST 2.3; lengths MV/PM 4.9; lengths PM/ST 0.5; submarginal vein with nine setae on dorsal surface.

Gaster. With lateral tufts of pale and flattened setae on Gt₆.

***Eriastichus galeatus* sp. nov.**

<http://zoobank.org/04AE8D37-56D4-411C-88A0-04FDC46979A3>

Figure 34

Type locality. Costa Rica, Puntarenas, Estación Biológica Monteverde, 10°20'N, 84°49'W, 1540 m, 26.ii.2007, J.S. Noyes leg.

Type specimen. *Holotype* male dried and glued to a paper card. Original labels: "COSTA RICA, Puntarenas, E.B. Monteverde, 10°20'N, 84°49'W, 1540 m,

26.ii.2007, J.S. Noyes”, “HOLOTYPE Eriastichus galeatis Hansson” [red printed label], (NHMUK014431045).

Diagnosis (male). Head dark brown, scape and pedicel pale brown, flagellum brown; ventral plaque on scape ca. $0.3 \times$ as long as scape (Fig. 34), antenna with dorsobasal setae on F1 $1.0 \times$ as long as F1; gaster with lateral tufts of pale and flattened setae on Gt₆.

Description (male holotype NHMUK014431045). Length of body 1.5 mm. Head dark brown, scape and pedicel pale brown, flagellum brown, ventral plaque dark brown. Mesoscutum, mesoscutellum, dorsellum and propodeum black with metallic tinges. Legs with coxae and femora dark brown, hind tibia pale brown, remaining parts of legs yellowish brown. Gaster dark brown.

Head. Length/width in frontal view 0.7; width/length in dorsal view 2.3; POL/OOL 2.1; WM/MS 1.6; MS/HE 0.5; HE/head length in frontal view 0.6; widths head/mesoscutum 1.1. **Antenna.** Pedicel + flagellum length/mesoscutum width 1.8; pedicel + flagellum length/head width 1.7; lengths scape/ventral plaque 3.1; ventral plaque located below the middle of scape; scape length/width 3.3; lengths scape/head (dorsal view) 0.5; scape length/HE 0.9; length/width F1, F2, F3, F4, clava: 3.0, 2.5, 2.5, 2.5, 6.3; length dorsobasal setae on F1/length F1 1.0. **Mesosoma.** Length/width 1.5; mesoscutum length/width 0.6; mesoscutellum length/width 0.7; widths SMG/SLG 1.3; enclosed space between SMG length/width 1.9; lengths mesoscutum/mesoscutellum 1.5; lengths mesoscutellum/dorsellum 2.7; lengths mesosoma/gaster 0.9. **Wings.** CC length/width 19.2; lengths CC/MV 1.0; lengths MV/ST 2.2; lengths MV/PM 3.8; lengths PM/ST 0.6; submarginal vein with eight setae on dorsal surface. **Gaster.** With lateral tufts of pale and flattened setae on Gt₆.

Eriastichus geratis sp. nov.

<http://zoobank.org/9BF19DD1-E099-457E-AFA4-0D5D82E0BA08>

Figure 33

Type locality. Costa Rica, Puntarenas, San Vito, Estación Biológica Las Alturas, 1500 m, $8^{\circ}57'N$, $82^{\circ}50'W$, 17–18.ii.2012, J.S. Noyes leg.

Type specimen. *Holotype* male dried and glued to a paper card. Original labels: “COSTA RICA, Puntarenas, San Vito, E.B. Las Alturas, 1500 m, $8^{\circ}57'N$, $82^{\circ}50'W$, 17–18.ii.2012, J.S. Noyes, NHM (Ent) 2012–91”, “HOLOTYPE Eriastichus geratis Hansson” [red printed label], (NHMUK014431046).

Additional type material. *Paratype* 1♂ “COSTA RICA, San José, San Gerardo de Dota, 20–21.ii.2013, J.S. Noyes, NHM (Ent) 2012–91” (NHMUK014431047).

Diagnosis (male). Head dark brown, scape and pedicel pale brown, flagellum brown; ventral plaque on scape ca. $0.4 \times$ as long as scape (Fig. 33), antenna with dorsobasal setae on F1 $0.6 \times$ as long as F1; gaster with lateral tufts of pale and flattened setae on Gt₆.

Description (male holotype NHMUK014431046). Length of body 1.8 mm (paratype 1.7 mm). Head dark brown, scape and pedicel pale brown, flagellum brown, ventral plaque dark brown. Mesoscutum and mesoscutellum black with metallic tinges, dorsellum dark brown, propodeum black. Legs with coxae, mid and hind femora dark brown, fore femur pale brown, remaining parts of legs yellowish brown. Gaster dark brown.

Head. Length/width in frontal view 0.8; width/length in dorsal view 2.4; POL/OOL 1.9; WM/MS 1.6; MS/HE 0.6; HE/head length in frontal view 0.5; widths head/mesoscutum 1.0. **Antenna.** Pedicel + flagellum length/mesoscutum width 1.8; pedicel + flagellum length/head width 1.8; lengths scape/ventral plaque 2.5; ventral plaque located in the middle of scape; scape length/width 2.8; lengths scape/head (dorsal view) 0.5; scape length/HE 0.9; length/width F1, F2, F3, F4, clava: 3.2, 2.8, 3.2, 3.0, 7.7; length dorsobasal setae on F1/length F1 0.6. **Mesosoma.** Length/width 1.5; mesoscutum length/width 0.6; mesoscutellum length/width 0.7; widths SMG/SLG 1.2; enclosed space between SMG length/width 1.9; lengths mesoscutum/mesoscutellum 1.5; lengths mesoscutellum/dorsellum 2.5; lengths mesosoma/gaster 0.8. **Wings.** CC length/width 19.3; lengths CC/MV 1.1; lengths MV/ST 2.2; lengths MV/PM 4.3; lengths PM/ST 0.5; submarginal vein with six setae on dorsal surface, in both types. **Gaster.** With lateral tufts of pale and flattened setae on Gt₆.

Eriastichus glanis sp. nov.

<http://zoobank.org/CC8A526B-4C9D-4B48-8503-9CE11D084C54>

Figure 32

Type locality. Costa Rica, Limón, Reserva Biológica Hitoy-Cerere, Headquarters, 100 m, 9°40'N, 83°02'W, 24–26.ii.2008, J.S. Noyes leg.

Type specimen. Holotype male dried and glued to a paper card. Original labels: "COSTA RICA, Limón, R.B. Hitoy-Cerere, HQ, 100 m, 9°40'N, 83°02'W, 24–26. ii.2008, J.S. Noyes, BMNH(Ent) 2010–21", "HOLOTYPE *Eriastichus glanis* Hansson" [red printed label], (NHMUK014431048).

Diagnosis (male). Head dark brown, scape yellowish brown, pedicel pale brown, flagellum brown; ventral plaque on scape ca. 0.3 × as long as scape (Fig. 32), antenna with dorsobasal setae on F1 0.5 × as long as F1; gaster with lateral tufts of pale and flattened setae on Gt₆.

Description (male holotype NHMUK014431048). Length of body 2.0 mm. Head dark brown, scape yellowish brown, pedicel pale brown, flagellum brown, ventral plaque dark brown. Mesoscutum and mesoscutellum black with metallic tinges, dorsellum and propodeum dark brown. Legs yellowish brown, hind coxa with base dark brown. Gaster dark brown.

Head. Length/width in frontal view 0.7; width/length in dorsal view 2.1; POL/OOL 1.3; WM/MS 1.8; MS/HE 0.4; HE/head length in frontal view 0.6; widths

head/mesoscutum 1.0. **Antenna.** Pedicel + flagellum length/mesoscutum width 2.4; pedicel + flagellum length/head width 2.0; lengths scape/ventral plaque 4.0; ventral plaque located in the middle of scape; scape length/width 3.1; lengths scape/head (dorsal view) 0.5; scape length/HE 0.8; length/width F1, F2, F3, F4, clava: 4.0, 4.0, 4.2, 3.8, 8.2; length dorsobasal setae on F1/length F1 0.5. **Mesosoma.** Length/width 1.6; mesoscutum length/width 0.7; mesoscutellum length/width 0.9; widths SMG/SLG 1.7; enclosed space between SMG length/width 2.2; lengths mesoscutum/mesoscutellum 1.4; lengths mesoscutellum/dorsellum 3.3; lengths mesosoma/gaster 0.7. **Wings.** CC length/width 18.7; lengths CC/MV 0.8; lengths MV/ST 2.9; lengths MV/PM 4.7; lengths PM/ST 0.6; submarginal vein with nine setae on dorsal surface. **Gaster.** With lateral tufts of pale and flattened setae on Gt₆.

Eriastichus hilaris sp. nov.

<http://zoobank.org/9FFEC823-0436-45EE-BA8F-54FD1FB5E740>

Figure 35

Type locality. Costa Rica, San José, San Gerardo de Dota, 9°33'N, 83°47'W, 20–21. ii.2013, J.S. Noyes leg.

Type specimen. *Holotype* male dried and glued to a paper card. Original labels: "COSTA RICA, San José, San Gerardo de Dota, 20–21.ii.2013, J.S. Noyes, NHM (Ent) 2012–91", "HOLOTYPE Eriastichus hilaris Hansson" [red printed label], (NHMUK014431049).

Diagnosis (male). Head black with part below antennal toruli pale brown, scape yellowish brown, pedicel pale brown, flagellum dark brown; ventral plaque on scape ca. 0.5 × as long as scape (Fig. 35), antenna with dorsobasal setae on F1 0.4 × as long as F1; gaster with lateral tufts of pale and flattened setae on Gt_{5–6}.

Description (male holotype NHMUK014431049). Length of body 1.7 mm. Head black with part below antennal toruli pale brown, scape yellowish brown, pedicel pale brown, flagellum dark brown, ventral plaque dark brown. Mesoscutum and mesoscutellum black with metallic tinges, dorsellum dark brown, propodeum black. Legs with hind coxa with basal ½ dark brown and apical ½ yellowish brown, remaining parts of legs yellowish brown. Gaster dark brown.

Head. Length/width in frontal view 0.8; width/length in dorsal view 2.0; POL/OOL 2.0; WM/MS 1.4; MS/HE 0.6; HE/head length in frontal view 0.5; widths head/mesoscutum 1.2. **Antenna.** Pedicel + flagellum length/mesoscutum width 1.9; pedicel + flagellum length/head width 1.7; lengths scape/ventral plaque 2.2; ventral plaque located above the middle of scape; scape length/width 2.9; lengths scape/head (dorsal view) 0.5; scape length/HE 0.9; length/width F1, F2, F3, F4, clava: 2.3, 2.4, 2.1, 2.2, 4.9; length dorsobasal setae on F1/length F1 0.4. **Mesosoma.** Length/width 1.6; mesoscutum length/width 0.5; mesoscutellum length/width 0.8; widths SMG/SLG 1.0; enclosed space between SMG length/width 2.4; lengths mesoscutum/mesoscutellum 1.2; lengths mesoscutellum/dorsellum 2.9; lengths mesosoma/gaster 0.8. **Wings.** CC length/width 17.3; lengths CC/MV 1.0; lengths MV/ST 2.2; lengths

MV/PM 4.5; lengths PM/ST 0.5; submarginal vein with nine setae on dorsal surface. **Gaster.** With lateral tufts of pale and flattened setae on Gt₅₋₆.

***Eriastichus johnlasallei* sp. nov.**

<http://zoobank.org/2A25525C-B5A5-471F-9697-A6936C8FB55E>

Figure 37

Type locality. Costa Rica, Cartago, Humo, El Copal, 9°47'N, 83°45'W, 1050–1250 m, 29.ii–6.iii.2008, C. Hansson leg.

Type specimen. **Holotype** male dried and glued to a paper card. Original labels: "COSTA RICA, Cartago, Humo, El Copal, 9°47'N, 83°45'W, 1050–1250 m, 29.ii–6.iii.2008, C. Hansson", "HOLOTYPE Eriastichus johnlasallei Hansson" [red printed label], (MZLU:7038.1).

Diagnosis (male). Head and antenna dark brown; ventral plaque on scape ca. 0.2 × as long as scape (Fig. 37), antenna with dorsobasal setae on F1 0.8 × as long as F1; gaster with lateral tufts of pale and flattened setae on Gt₆.

Description (male holotype MZLU:7038.1). Length of body 1.5 mm. Head and antenna dark brown, ventral plaque black. Mesoscutum, mesoscutellum and propodeum dark brown; dorsellum pale brown. Legs yellowish brown, coxae and hind femur brown. Gaster dark brown.

Head. Length/width in frontal view 0.7; width/length in dorsal view 2.3; POL/OOL 2.0; WM/MS 1.7; MS/HE 0.5; HE/head length in frontal view 0.6; widths head/mesoscutum 1.2. **Antenna.** Pedicel + flagellum length/mesoscutum width 2.0; pedicel + flagellum length/head width 1.7; lengths scape/ventral plaque 4.7; ventral plaque located in the middle of scape; scape length/width 3.5; lengths scape/head (dorsal view) 0.6; scape length/HE 0.9; length/width F1, F2, F3, F4, clava: 2.5, 2.7, 2.8, 2.8, 6.2; length dorsobasal setae on F1/length F1 0.8. **Mesosoma.** Length/width 1.5; mesoscutum length/width 0.5; mesoscutellum length/width 0.8; widths SMG/SLG 1.6; enclosed space between SMG length/width 1.9; lengths mesoscutum/mesoscutellum 1.2; lengths mesoscutellum/dorsellum 3.6; lengths mesosoma/gaster 0.9. **Wings.** CC length/width 24.0; lengths CC/MV 0.9; lengths MV/ST 2.2; lengths MV/PM 4.7; lengths PM/ST 0.5; submarginal vein with seven setae on dorsal surface. **Gaster.** With lateral tufts of pale and flattened setae on Gt₆.


Etymology. Named after John La Salle, fellow euphorid taxonomist who described *Eriastichus*.

***Eriastichus johnnnoyesi* sp. nov.**

<http://zoobank.org/1F78E014-A16C-4D6F-8CFB-FC8EEF163665>

Figure 40

Type locality. Costa Rica, Heredia, Estación Biológica La Selva, 75 m, 10°26'N, 84°01'W, 24–25.ii.2011, J.S. Noyes leg.



Figures 35–40. *Eriastichus* spp. head and antenna in lateral view, male holotypes **35** *E. hilaris* **36** *E. maniatensis* **37** *E. johnlasallei* **38** *E. nebulis* **39** *E. neonis* **40** *E. johnnnoyesi*.

Type specimen. *Holotype* male dried and glued to a paper card. Original labels: "COSTA RICA, Heredia, E.B. La Selva, 75 m, 10°26'N, 84°01'W, 24–25.ii.2011, J.S. Noyes, NHM (Ent) 2011–93", "HOLOTYPE *Eriastichus johnnnoyesi* Hansson" [red printed label], (NHMUK014431050).

Additional type material. *Paratypes* 2♂♂ from same locality as holotype but collected 30–31.iii.2002 (1♂, MZLU:7039.2), 23–24.ii.2005 (1♂, NHMUK014431051).

Diagnosis (male). Head dark brown, scape yellowish brown, pedicel and flagellum brown; ventral plaque on scape ca. 0.2 × as long as scape (Fig. 40), antenna with dorsobasal setae on F1 0.6 × as long as F1; gaster with lateral tufts of pale and flattened setae on Gt₆.

Description (male holotype NHMUK014431050). Length of body 1.1 mm (paratypes 1.0 mm). Head dark brown, scape yellowish brown, pedicel and flagellum brown, ventral plaque dark brown. Mesoscutum and mesoscutellum black, dorsellum and propodeum dark brown. Legs with coxae and hind femur dark brown, remaining parts of legs yellowish brown. Gaster dark brown.

Head. Length/width in frontal view 0.7; width/length in dorsal view 2.2; POL/OOL 3.0; WM/MS 1.3; MS/HE 0.5; HE/head length in frontal view 0.6; widths head/mesoscutum 1.2. **Antenna.** Pedicel + flagellum length/mesoscutum width 1.8; pedicel + flagellum length/head width 1.6; lengths scape/ventral plaque 5.0; ventral plaque located below the middle of scape; scape length/width 3.3; lengths scape/head (dorsal view) 0.5; scape length/HE 0.9; length/width F1, F2, F3, F4, clava: 2.3, 2.5, 2.3, 3.0, 7.8; length dorsobasal setae on F1/length F1 0.6. **Mesosoma.** Length/width 1.4; mesoscutum length/width 0.6; mesoscutellum length/width 0.7; widths SMG/SLG 1.4; enclosed space between SMG length/width 1.6; lengths mesoscutum/mesoscutellum 1.6; lengths mesoscutellum/dorsellum 2.6; lengths mesosoma/gaster 0.8. **Wings.** CC length/width 18.8; lengths CC/MV 0.9; lengths MV/ST 2.0; lengths MV/PM 4.0; lengths PM/ST 0.5; submarginal vein with four setae (four setae in paratypes) on dorsal surface. **Gaster.** With lateral tufts of pale and flattened setae on Gt₆.

Etymology. Named after John S. Noyes (NHMUK), collector of the material of this and several other species described here.

Eriastichus maniatis sp. nov.

<http://zoobank.org/B3BA25D5-CA2F-4B3D-AB17-B8AAF4848F7D>

Figure 36

Type locality. Costa Rica, Puntarenas, San Vito, Las Cruces, 8°46'N, 82°57'W, 1300 m, 15–16.ii.2006, J.S. Noyes leg.

Type specimen. *Holotype* male dried and glued to a paper card. Original labels: "COSTA RICA, Puntarenas, San Vito, Las Cruces, 8°46'N, 82°57'W, 1300 m, 15–16. ii.2006, J.S. Noyes", "HOLOTYPE *Eriastichus maniatis* Hansson" [red printed label], (NHMUK014431052).

Diagnosis (male). Head dark brown with parts above mouth yellowish brown, scape yellowish brown, pedicel and flagellum brown; ventral plaque on scape ca. 0.3 × as long as scape (Fig. 36), antenna with dorsobasal setae on F1 0.6 × as long as F1; gaster with lateral tufts of pale and flattened setae on Gt₆.

Description (male holotype NHMUK014431052). Length of body 1.5 mm. Head dark brown with parts above mouth yellowish brown, scape yellowish brown, pedicel and flagellum brown, ventral plaque dark brown. Mesoscutum and mesoscutellum black with metallic tinges, dorsellum and propodeum dark brown. Legs yellowish brown. Gaster dark brown.

Head. Length/width in frontal view 0.8; width/length in dorsal view 2.2; POL/OOL 2.1; WM/MS 1.4; MS/HE 0.5; HE/head length in frontal view 0.6; widths head/mesoscutum 1.3. **Antenna.** Pedicel + flagellum length/mesoscutum width 2.6; pedicel + flagellum length/head width 2.1; lengths scape/ventral plaque 3.8; ventral plaque located below the middle of scape; scape length/width 3.8; lengths scape/head (dorsal view) 0.5; scape length/HE 0.9; length/width F1, F2, F3, F4, clava: 3.7, 3.8, 3.8, 3.7, 8.3; length dorsobasal setae on F1/length F1 0.6. **Mesosoma.** Length/width 1.6; mesoscutum length/width 0.5; mesoscutellum length/width 0.8; widths SMG/SLG 1.0; enclosed space between SMG length/width 2.3; lengths mesoscutum/mesoscutellum 1.1; lengths mesoscutellum/dorsellum 3.1; lengths mesosoma/gaster 0.9. **Wings.** CC length/width 21.7; lengths CC/MV 0.9; lengths MV/ST 2.5; lengths MV/PM 3.4; lengths PM/ST 0.7; submarginal vein with seven setae on dorsal surface. **Gaster.** With lateral tufts of pale and flattened setae on Gt₆.

Eriastichus masneri La Salle

Eriastichus masneri La Salle, 1994: 207. Holotype female in CNC, not examined.

Diagnosis. Mesoscutellum with parts lateral to submedian grooves strongly setose (as in Fig. 6); median propodeum hairy (fig. 30 in La Salle (1994)).

Description. See La Salle (1994).

Distribution. Dominican Republic (La Salle 1994).

Eriastichus nakos La Salle

Eriastichus nakos La Salle, 1994: 208. Holotype female in CNC, not examined.

Diagnosis. Mesoscutellum with parts lateral to submedian grooves strongly setose (as in Fig. 6); median propodeum bare (as in Fig. 5).

Description. See La Salle (1994).

Distribution. Dominican Republic, Ecuador (La Salle 1994).

***Eriastichus nebulis* sp. nov.**

<http://zoobank.org/5759023B-4DDD-48BC-BC00-CF7B4ACC38F4>

Figure 38

Type locality. Costa Rica, Puntarenas, San Vito, Estación Biológica Las Alturas, 1500 m, 8°57'N, 82°50'W, 17–18.ii.2012, J.S. Noyes leg.

Type specimen. **Holotype** male dried and glued to a paper card. Original labels: "COSTA RICA, Puntarenas, San Vito, E.B. Las Alturas, 1500 m, 8°57'N, 82°50'W, 17–18.ii.2012, J.S. Noyes, NHM (Ent) 2012–91", "HOLOTYPE *Eriastichus nebulis* Hansson" [red printed label], (NHMUK014431053).

Diagnosis (male). Head dark brown with metallic tinges, scape yellowish brown, pedicel and flagellum brown; ventral plaque on scape ca. 0.3 × as long as scape (Fig. 38), antenna with dorsobasal setae on F1 0.6 × as long as F1; gaster with lateral tufts of pale and flattened setae on Gt₆.

Description (male holotype NHMUK014431053). Length of body 1.9 mm. Head dark brown with metallic tinges, scape yellowish brown, pedicel and flagellum brown, ventral plaque dark brown. Mesoscutum and mesoscutellum black with metallic tinges, dorsellum and propodeum dark brown. Legs yellowish brown. Gaster dark brown.

Head. Length/width in frontal view 0.7; width/length in dorsal view 2.2; POL/OOL 2.3; WM/MS 1.6; MS/HE 0.5; HE/head length in frontal view 0.6; widths head/mesoscutum 1.2. **Antenna.** Pedicel + flagellum length/mesoscutum width 2.3;

pedicel + flagellum length/head width 1.9; lengths scape/ventral plaque 3.0; ventral plaque located below the middle of scape; scape length/width 3.0; lengths scape/head (dorsal view) 0.5; scape length/HE 0.9; length/width F1, F2, F3, F4, clava: 3.6, 3.8, 3.3, 3.3, 7.8; length dorsobasal setae on F1/length F1 0.6. **Mesosoma.** Length/width 1.7; mesoscutum length/width 0.7; mesoscutellum length/width 0.8; widths SMG/SLG 1.2; enclosed space between SMG length/width 2.2; lengths mesoscutum/mesoscutellum 1.5; lengths mesoscutellum/dorsellum 2.9; lengths mesosoma/gaster 0.7.

Wings. CC length/width 20.0; lengths CC/MV 1.0; lengths MV/ST 2.0; lengths MV/PM 3.8; lengths PM/ST 0.5; submarginal vein with eight setae on dorsal surface.

Gaster. With lateral tufts of pale and flattened setae on Gt₆.

***Eriastichus neonis* sp. nov.**

<http://zoobank.org/2D31B245-9233-4289-AD36-D0111F016CC2>

Figure 39

Type locality. Costa Rica, Cartago, Humo, El Copal, 9°47'N, 83°45'W, 1050–1250 m, 29.ii–6.iii.2008, C. Hansson leg.

Type specimen. **Holotype** male dried and glued to a paper card. Original labels: "COSTA RICA, Cartago, Humo, El Copal, 9°47'N, 83°45'W, 1050–1250 m, 29.ii–6.iii.2008, C. Hansson", "HOLOTYPE *Eriastichus neonis* Hansson" [red printed label], (MZLU:7040.1).

Additional type material. *Paratypes* (3♂♂): 1♂ with same label data as holotype (MZLU:7040.2); 1♂ “COSTA RICA, Heredia, E.B. La Selva, 75 m, LN 264463/532850, 30–31.iii.2002, swept, J. Azofeifa” (MZUCR:01694); 1♂ “COSTA RICA, Puntarenas, San Vito, E.B. Las Alturas, 1500 m, 8°57'N, 82°50'W, 17–18. ii.2012, J.S. Noyes, NHM (Ent) 2012–91” (NHMUK014431054).

Diagnosis (male). Head dark brown, scape yellowish brown, pedicel and flagellum brown; ventral plaque on scape ca. 0.3 × as long as scape (Fig. 39), antenna with dorsobasal setae on F1 0.7 × as long as F1; gaster with lateral tufts of pale and flattened setae on Gt₆.

Description (male holotype MZLU:7040.1). Length of body 1.5 mm (paratypes 1.3–1.6 mm). Head dark brown, scape yellowish brown, pedicel and flagellum brown, ventral plaque dark brown. Mesoscutum, mesoscutellum and propodeum black, dorsellum dark brown. Legs yellowish brown. Gaster dark brown.

Head. Length/width in frontal view 0.8; width/length in dorsal view 1.9; POL/OOL 2.5; WM/MS 1.6; MS/HE 0.5; HE/head length in frontal view 0.6; widths head/mesoscutum 1.2. **Antenna.** Pedicel + flagellum length/mesoscutum width 2.5; pedicel + flagellum length/head width 2.1; lengths scape/ventral plaque 4.0; ventral plaque located below the middle of scape; scape length/width 3.0; lengths scape/head (dorsal view) 0.5; scape length/HE 0.8; length/width F1, F2, F3, F4, clava: 3.0, 3.0, 3.0, 3.0, 7.0; length dorsobasal setae on F1/length F1 0.7. **Mesosoma.** Length/width 1.8; mesoscutum length/width 0.7; mesoscutellum length/width 0.9; widths SMG/SLG 1.2; enclosed space between SMG length/width 2.1; lengths mesoscutum/mesoscutellum 1.5; lengths mesoscutellum/dorsellum 2.6; lengths mesosoma/gaster 0.8. **Wings.** CC length/width 23.0; lengths CC/MV 0.9; lengths MV/ST 2.1; lengths MV/PM 5.0; lengths PM/ST 0.4; submarginal vein with six setae on dorsal surface (6–7 setae in paratypes). **Gaster.** With lateral tufts of pale and flattened setae on Gt₆.

Eriastichus nexilis sp. nov.

<http://zoobank.org/A8B195EF-FC82-4D9C-86DF-DA156C09D23F>

Figure 41

Type locality. Costa Rica, Heredia, Estación Biológica La Selva, 75 m, 10°26'N, 84°01'W, 28–29.ii.2008, J.S. Noyes leg.

Type specimen. *Holotype* male dried and glued to a paper card. Original labels: “COSTA RICA, Heredia, E.B. La Selva, 75 m, 10°26'N, 84°01'W, 28–29.ii.2008, J.S. Noyes, NHM (Ent) 2010–21”, “HOLOTYPE *Eriastichus nexilis* Hansson” [red printed label], (NHMUK014431055).

Additional type material. *Paratype* 1♂ “COSTA RICA, Cartago, Humo, El Copal, 9°47'N, 83°45'W, 1050–1250 m, 29.ii–6.iii.2008, C. Hansson” (MZLU:7041.2).

Diagnosis (male). Head dark brown, scape pale brown, pedicel and flagellum brown; ventral plaque on scape ca. 0.2 × as long as scape (Fig. 41), antenna with dorsobasal setae on F1 0.9 × as long as F1; gaster with lateral tufts of pale and flattened setae on Gt₆.

Description (male holotype NHMUK014431055). Length of body 1.1 mm (paratype 1.2 mm). Head dark brown, scape pale brown, pedicel and flagellum brown,

ventral plaque dark brown. Mesoscutum and mesoscutellum black with metallic tinges, dorsellum dark brown, propodeum black. Legs yellowish brown. Gaster dark brown.

Head. Length/width in frontal view 0.8; width/length in dorsal view 2.2; POL/OOL 2.0; WM/MS 1.6; MS/HE 0.5; HE/head length in frontal view 0.6; widths head/mesoscutum 1.1. **Antenna.** Pedicel + flagellum length/mesoscutum width 2.3; pedicel + flagellum length/head width 1.9; lengths scape/ventral plaque 4.2; ventral plaque located below the middle of scape; scape length/width 3.0; lengths scape/head (dorsal view) 0.5; scape length/HE 0.9; length/width F1, F2, F3, F4, clava: 3.3, 3.8, 3.5, 3.5, 8.5; length dorsobasal setae on F1/length F1 0.9. **Mesosoma.** Length/width 1.4; mesoscutum length/width 0.6; mesoscutellum length/width 0.8; widths SMG/SLG 1.1; enclosed space between SMG length/width 2.1; lengths mesoscutum/mesoscutellum 1.4; lengths mesoscutellum/dorsellum 3.2; lengths mesosoma/gaster 0.7. **Wings.** CC length/width 22.5; lengths CC/MV 1.1; lengths MV/ST 1.9; lengths MV/PM 4.0; lengths PM/ST 0.5; submarginal vein with six setae on dorsal surface (five setae in paratype). **Gaster.** With lateral tufts of pale and flattened setae on Gt₆.

Eriastichus novalis sp. nov.

<http://zoobank.org/A656E7CC-7D18-4A45-BDB8-F5F05D7C539D>

Figure 44

Type locality. Costa Rica, San José, San Gerardo de Dota, 9°33'N, 83°47'W, 20–21. ii.2013, J.S. Noyes leg.

Type specimen. *Holotype* male dried and glued to a paper card. Original labels: "COSTA RICA, San José, San Gerardo de Dota, 20–21.ii.2013, J.S. Noyes, NHM (Ent) 2012–91", "HOLOTYPE Eriastichus novalis Hansson" [red printed label], (NHMUK014431056).

Diagnosis (male). Head dark brown with part below antennal toruli yellowish brown, scape yellowish brown, pedicel pale brown, flagellum brown; ventral plaque on scape ca. 0.4 × as long as scape (Fig. 44), antenna with dorsobasal setae on F1 0.9 × as long as F1; gaster with lateral tufts of pale and flattened setae on Gt₆.

Description (male holotype NHMUK014431056). Length of body 1.7 mm. Head dark brown with part below antennal toruli yellowish brown, scape yellowish brown, pedicel pale brown, flagellum brown, ventral plaque dark brown. Mesoscutum and mesoscutellum dark brown with metallic tinges, dorsellum pale brown, propodeum dark brown. Legs yellowish brown. Gaster dark brown.

Head. Length/width in frontal view 0.7; width/length in dorsal view 2.3; POL/OOL 1.8; WM/MS 1.6; MS/HE 0.5; HE/head length in frontal view 0.6; widths head/mesoscutum 1.0. **Antenna.** Pedicel + flagellum length/mesoscutum width 2.0; pedicel + flagellum length/head width 1.8; lengths scape/ventral plaque 2.5; ventral plaque located below the middle of scape; scape length/width 2.5; lengths scape/head (dorsal view) 0.5; scape length/HE 0.8; length/width F1, F2, F3, F4, clava: 3.2, 3.0, 2.7, 2.8, 6.5; length dorsobasal setae on F1/length F1 0.9. **Mesosoma.** Length/width 1.4; mesoscutum length/width 0.6; mesoscutellum length/width 1.9; widths SMG/



Figures 41–50. *Eriastichus* spp. head and antenna in lateral view, male holotypes **41** *E. nexilis* **42** *E. oasis* **43** *E. ononis* **44** *E. novalis* **45** *E. nugalis* **46** *E. parabilis* **47** *E. rivalis* **48** *E. sodalis* **49** *E. orestis* **50** *E. pallidops*.

SLG 1.0; enclosed space between SMG length/width 2.0; lengths mesoscutum/mesoscutellum 1.6; lengths mesoscutellum/dorsellum 2.6; lengths mesosoma/gaster 0.7. **Wings.** CC length/width 23.3; lengths CC/MV 1.0; lengths MV/ST 2.1; lengths MV/PM 5.4; lengths PM/ST 0.4; submarginal vein with ten setae on dorsal surface. **Gaster.** With lateral tufts of pale and flattened setae on Gt₆.

***Eriastichus nugalis* sp. nov.**

<http://zoobank.org/07A69A24-BEB7-4C6C-BA98-953B99362A62>

Figure 45

Type locality. Costa Rica, Cartago, Humo, El Copal, 9°47'N, 83°45'W, 1050–1250 m, 29.ii–6.iii.2008, C. Hansson leg.

Type specimen. *Holotype* male dried and glued to a paper card. Original labels: "COSTA RICA, Cartago, Humo, El Copal, 9°47'N, 83°45'W, 1050–1250 m, 29.ii–6.iii.2008, C. Hansson", "HOLOTYPE Eriastichus nugalis Hansson" [red printed label], (MZLU:7042.1).

Additional type material. *Paratypes* (2♂♂): 1♂ with same label data as holotype (MZLU:7042.2); 1♂ "COSTA RICA, Puntarenas, San Vito, E.B. Las Alturas, 1500 m, 8°57'N, 82°50'W, 17–18.ii.2012, J.S. Noyes, NHM (Ent) 2012–91" (NHMUK014431057).

Diagnosis (male). Head dark brown, scape yellowish brown, pedicel pale brown, flagellum dark brown; ventral plaque on scape ca. 0.4 × as long as scape (Fig. 45), antenna with dorsobasal setae on F1 0.5 × as long as F1; gaster with lateral tufts of pale and flattened setae on Gt₆.

Description (male holotype MZLU:7042.1). Length of body 1.8 mm (paratypes 1.5–1.6 mm). Head dark brown, scape yellowish brown, pedicel pale brown, flagellum dark brown, ventral plaque dark brown. Mesoscutum, mesoscutellum and propodeum dark brown, dorsellum pale brown. Legs with fore and mid coxae yellowish brown with base brown, hind coxa, remaining parts of legs yellowish brown. Gaster dark brown.

Head. Length/width in frontal view 0.8; width/length in dorsal view 2.1; POL/OOL 1.8; WM/MS 1.9; MS/HE 0.5; HE/head length in frontal view 0.6; widths head/mesoscutum 1.1. **Antenna.** Pedicel + flagellum length/mesoscutum width 2.1; pedicel + flagellum length/head width 1.9; lengths scape/ventral plaque 2.5; ventral plaque located in the middle of scape; scape length/width 3.0; lengths scape/head (dorsal view) 0.6; scape length/HE 0.9; length/width F1, F2, F3, F4, clava: 2.5, 2.5, 2.5, 2.5, 5.5; length dorsobasal setae on F1/length F1 0.5. **Mesosoma.** Length/width 1.6; mesoscutum length/width 0.6; mesoscutellum length/width 0.8; widths SMG/SLG 1.0; enclosed space between SMG length/width 2.3; lengths mesoscutum/mesoscutellum 1.4; lengths mesoscutellum/dorsellum 2.8; lengths mesosoma/gaster 0.7. **Wings.** CC length/width 30.0; lengths CC/MV 1.1; lengths MV/ST 1.9; lengths MV/PM 4.9; lengths PM/ST 0.4; submarginal vein with eight setae on dorsal surface (paratypes with seven setae). **Gaster.** With lateral tufts of pale and flattened setae on Gt₆.

***Eriastichus oasis* sp. nov.**

<http://zoobank.org/3008B625-977E-4985-90B1-81A4783D1BB3>

Figure 42

Type locality. Costa Rica, Heredia, Estación Biológica La Selva, 75 m, 10°26'N, 84°01'W, 22–24.ii.2012, J.S. Noyes leg.

Type specimen. *Holotype* male dried and glued to a paper card. Original labels: "COSTA RICA, Heredia, E.B. La Selva, 75 m, 10°26'N, 84°01'W, 22–24.ii.2012, J.S. Noyes, NHM (Ent) 2012–91", "HOLOTYPE *Eriastichus oasis* Hansson" [red printed label], (NHMUK014431058).

Additional type material. *Paratype* 1♂ "COSTA RICA, Cartago, Humo, El Co-pal, 9°47'N, 83°45'W, 1050–1250 m, 29.ii–6.iii.2008, C. Hansson" (MZLU:7043.2).

Diagnosis (male). Head dark brown, scape yellowish brown, pedicel pale brown, flagellum brown; ventral plaque on scape ca. 0.5 × as long as scape (Fig. 42), antenna with dorsobasal setae on F1 0.5 × as long as F1; gaster with lateral tufts of pale and flattened setae on Gt₆.

Description (male holotype NHMUK014431058). Length of body 1.3 mm (paratype 1.4 mm). Head dark brown, scape yellowish brown, pedicel pale brown, flagellum brown, ventral plaque dark brown. Mesoscutum and mesoscutellum black with metallic tinges, dorsellum pale brown, propodeum dark brown. Legs yellowish brown, hind coxa with base dark brown. Gaster dark brown.

Head. Length/width in frontal view 0.8; width/length in dorsal view 2.3; POL/OOL 2.0; WM/MS 1.5; MS/HE 0.5; HE/head length in frontal view 0.6; widths head/mesoscutum 1.2. **Antenna.** Pedicel + flagellum length/mesoscutum width 2.1; pedicel + flagellum length/head width 1.9; lengths scape/ventral plaque 2.2; ventral plaque located below the middle of scape; scape length/width 3.1; lengths scape/head (dorsal view) 0.5; scape length/HE 0.8; length/width F1, F2, F3, F4, clava: 3.3, 3.3, 3.3, 3.0, 8.5; length dorsobasal setae on F1/length F1 0.5. **Mesosoma.** Length/width 1.5; mesoscutum length/width 0.5; mesoscutellum length/width 0.6; widths SMG/SLG 0.9; enclosed space between SMG length/width 2.0; lengths mesoscutum/mesoscutellum 1.2; lengths mesoscutellum/dorsellum 2.3; lengths mesosoma/gaster 0.7.

Wings. CC length/width 19.0; lengths CC/MV 1.2; lengths MV/ST 2.1; lengths MV/PM 4.1; lengths PM/ST 0.5; submarginal vein with seven setae on dorsal surface (six setae in paratype). **Gaster.** With lateral tufts of pale and flattened setae on Gt₆.

***Eriastichus ononis* sp. nov.**

<http://zoobank.org/A538731B-FECE-4287-9566-65DEC5E3F735>

Figure 43

Type locality. Costa Rica, Puntarenas, San Vito, Las Cruces, 8°46'N, 82°57'W, 1300 m, 15–16.ii.2006, J.S. Noyes leg.

Type specimen. **Holotype** male dried and glued to a paper card. Original labels: "COSTA RICA, Puntarenas, San Vito, Las Cruces, 8°46'N, 82°57'W, 1300 m, 15–16. ii.2006, J.S. Noyes", "HOLOTYPE Eriastichus ononis Hansson" [red printed label], (NHMUK014431059).

Diagnosis (male). Head dark brown, scape yellowish brown, pedicel and flagellum brown; ventral plaque on scape ca. 0.3 × as long as scape (Fig. 43), antenna with dorsobasal setae on F1 0.9 × as long as F1; gaster with lateral tufts of pale and flattened setae on Gt₆.

Description (male holotype NHMUK014431059). Length of body 1.5 mm. Head dark brown, scape yellowish brown, pedicel and flagellum brown, ventral plaque dark brown. Mesoscutum and mesoscutellum black with metallic tinges, dorsellum pale brown, propodeum dark brown. Legs yellowish brown. Gaster dark brown.

Head. Length/width in frontal view 0.8; width/length in dorsal view 2.1; POL/OOL 2.4; WM/MS 1.5; MS/HE 0.5; HE/head length in frontal view 0.6; widths head/mesoscutum 1.2. **Antenna.** Pedicel + flagellum length/mesoscutum width 2.5; pedicel + flagellum length/head width 2.0; lengths scape/ventral plaque 3.1; ventral plaque located below the middle of scape; scape length/width 2.8; lengths scape/head (dorsal view) 0.5; scape length/HE 0.9; length/width F1, F2, F3, F4, clava: 3.6, 3.6, 3.6, 3.4, 7.4; length dorsobasal setae on F1/length F1 0.9. **Mesosoma.** Length/width 1.7; mesoscutum length/width 0.7; mesoscutellum length/width 0.9; widths SMG/SLG 1.0; enclosed space between SMG length/width 2.7; lengths mesoscutum/mesoscutellum 1.4; lengths mesoscutellum/dorsellum 3.4; lengths mesosoma/gaster 0.7. **Wings.** CC length/width 16.0; lengths CC/MV 1.0; lengths MV/ST 2.1; lengths MV/PM 5.0; lengths PM/ST 0.4; submarginal vein with seven setae on dorsal surface. **Gaster.** With lateral tufts of pale and flattened setae on Gt₆.

Eriastichus orestis sp. nov.

<http://zoobank.org/C962EAC3-7329-4811-A87E-EE829A92C069>

Figure 49

Type locality. Costa Rica, Puntarenas, San Vito, Estación Biológica Las Alturas, 1500 m, 8°57'N, 82°50'W, 17–18.ii.2012, J.S. Noyes leg.

Type specimen. **Holotype** male dried and glued to a paper card. Original labels: "COSTA RICA, Puntarenas, San Vito, E.B. Las Alturas, 1500 m, 8°57'N, 82°50'W, 17–18.ii.2012, J.S. Noyes, NHM (Ent) 2012–91", "HOLOTYPE Eriastichus orestis Hansson" [red printed label], (NHMUK014431060).

Additional type material. **Paratypes** (2♂♂): 1♂ with same label data as holotype (NHMUK014431061); 1♂ "COSTA RICA, Cartago, Humo, El Copal, 9°47'N, 83°45'W, 1050–1250 m, 29.ii–6.iii.2008, C. Hansson" (MZLU:7044.2).

Diagnosis (male). Head dark brown with part below antennal toruli dark yellowish brown, scape yellowish brown, pedicel and flagellum brown; ventral plaque on

scape ca. 0.3 × as long as scape (Fig. 49), antenna with dorsobasal setae on F1 0.9 × as long as F1; gaster with lateral tufts of pale and flattened setae on Gt₆.

Description (male holotype NHMUK014431060). Length of body 1.7 mm (paratypes 1.7–1.8 mm). Head dark brown with part below antennal toruli pale brown, scape yellowish brown, pedicel pale brown, flagellum brown, ventral plaque dark brown. Mesoscutum, mesoscutellum and propodeum dark brown, dorsellum pale brown. Legs yellowish brown. Gaster dark brown.

Head. Length/width in frontal view 0.7; width/length in dorsal view 2.3; POL/OOL 1.8; WM/MS 1.5; MS/HE 0.6; HE/head length in frontal view 0.6; widths head/mesoscutum 1.3. **Antenna.** Pedicel + flagellum length/mesoscutum width 2.3; pedicel + flagellum length/head width 1.9; lengths scape/ventral plaque 3.0; ventral plaque located below the middle of scape; scape length/width 3.0; lengths scape/head (dorsal view) 0.5; scape length/HE 0.9; length/width F1, F2, F3, F4, clava: 2.8, 3.0, 3.2, 3.2, 6.8; length dorsobasal setae on F1/length F1 0.9. **Mesosoma.** Length/width 1.6; mesoscutum length/width 0.7; mesoscutellum length/width 0.8; widths SMG/SLG 1.1; enclosed space between SMG length/width 2.4; lengths mesoscutum/mesoscuteum 1.5; lengths mesoscutellum/dorsellum 3.3; lengths mesosoma/gaster 0.7. **Wings.** CC length/width 16.2; lengths CC/MV 0.7; lengths MV/ST 2.4; lengths MV/PM 4.8; lengths PM/ST 0.5; submarginal vein with six setae on dorsal surface (paratypes with eight setae). **Gaster.** With lateral tufts of pale and flattened setae on Gt₆.

Eriastichus pallidops sp. nov.

<http://zoobank.org/A4B4B83F-D622-49B8-965F-CECE0B604217>

Figures 50, 59

Type locality. Costa Rica, Cartago, Humo, El Copal, 9°47'N, 83°45'W, 1050–1250 m, 29.ii–6.iii.2008, C. Hansson leg.

Type specimen. *Holotype* male dried and glued to a paper card. Original labels: "COSTA RICA, Cartago, Humo, El Copal, 9°47'N, 83°45'W, 1050–1250 m, 29.ii–6.iii.2008, C. Hansson", "HOLOTYPE Eriastichus pallidops Hansson" [red printed label], (MZLU:7045.1).

Additional type material. *Paratypes* 2♂ with same label data as holotype (MZU-CR:01695, NHMUK014431062).

Diagnosis (male). Head with lower ½ yellowish brown (Fig. 59); antenna (Fig. 50): scape yellowish brown, ventral plaque on scape 0.7 × as long as scape, antenna with dorsobasal setae on F1 0.4 × as long as F1; gaster with lateral tufts of pale and flattened setae on Gt_{5,6}.

Description (male holotype MZLU:7045.1). Length of body 1.9 mm (paratypes 1.5–1.7 mm). Head yellowish brown with upper ½ of frons pale brown, vertex and upper ½ of occiput dark brown. Antenna with scape yellowish brown with dorsal margin pale brown, ventral plaque black; pedicel and flagellum brown. Mesoscutum and mesoscutellum black with metallic tinges; dorsellum pale brown; propodeum dark

brown. Legs yellowish brown with base of hind coxa and 4th tarsomere dark brown. Gaster dark brown.

Head. Length/width in frontal view 0.7; width/length in dorsal view 2.2; POL/OOL 1.8; WM/MS 1.7; MS/HE 0.5; HE/head length in frontal view 0.6; widths head/mesoscutum 1.1. **Antenna.** Pedicel + flagellum length/mesoscutum width 1.5; pedicel + flagellum length/head width 1.4; lengths scape/ventral plaque 1.4; ventral plaque located slightly above the middle of scape; scape length/width 2.8; lengths scape/head (dorsal view) 0.6; scape length/HE 0.9; length/width F1, F2, F3, F4, clava: 2.3, 2.0, 1.8, 1.8, 4.0; length dorsobasal setae on F1/length F1 0.4. **Mesosoma.** Length/width 1.6; mesoscutum length/width 0.5; mesoscutellum length/width 0.7; widths SMG/SLG 1.2; enclosed space between SMG length/width 1.8; lengths mesoscutum/mesoscutellum 1.2; lengths mesoscutellum/dorsellum 2.7; lengths mesosoma/gaster 0.8. **Wings.** CC length/width 19.3; lengths CC/MV 1.1; lengths MV/ST 2.2; lengths MV/PM 3.7; lengths PM/ST 0.6; submarginal vein with ten setae on dorsal surface (8–9 setae in paratypes). **Gaster.** With lateral tufts of pale and flattened setae on Gt₅₋₆.

Etymology. Named for the yellowish brown lower face.

Eriastichus parabilis sp. nov.

<http://zoobank.org/F33852D4-65AF-4B99-9A79-A5C468443274>

Figure 46

Type locality. Costa Rica, San José, San Gerardo de Dota, 9°33'N, 83°47'W, 20–21. ii.2013, J.S. Noyes leg.

Type specimen. *Holotype* male dried and glued to a paper card. Original labels: "COSTA RICA, San José, San Gerardo de Dota, 20–21.ii.2013, J.S. Noyes, NHM (Ent) 2012–91", "HOLOTYPE Eriastichus parabilis Hansson" [red printed label], (NHMUK014431063).

Additional type material. *Paratype* 1♂ with same label data as holotype (NHMUK014431064).

Diagnosis (male). Head black with lower part of clypeus yellowish brown, scape pale brown, pedicel and flagellum dark brown; ventral plaque on scape ca. 0.4 × as long as scape (Fig. 46), antenna with dorsobasal setae on F1 0.5 × as long as F1; gaster with lateral tufts of pale and flattened setae on Gt₆.

Description (male holotype NHMUK014431063). Length of body 2.3 mm (paratype 2.3 mm). Head black with lower part of clypeus yellowish brown, scape pale brown, pedicel and flagellum dark brown, ventral plaque dark brown. Mesoscutum, mesoscutellum, dorsellum and propodeum black with metallic tinges. Legs with hind coxa black, hind femur pale brown, remaining parts of legs yellowish brown. Wings uniformly but weakly infuscated. Gaster dark brown.

Head. Length/width in frontal view 0.7; width/length in dorsal view 2.3; POL/OOL 1.9; WM/MS 1.5; MS/HE 0.6; HE/head length in frontal view 0.6; widths head/mesoscutum 1.1. **Antenna.** Pedicel + flagellum length/mesoscutum width 2.2;

pedicel + flagellum length/head width 2.0; lengths scape/ventral plaque 2.8; ventral plaque located below the middle of scape; scape length/width 3.1; lengths scape/head (dorsal view) 0.7; scape length/HE 0.9; length/width F1, F2, F3, F4, clava: 3.1, 3.0, 2.9, 2.9, 6.3; length dorsobasal setae on F1/length F1 0.5. **Mesosoma.** Length/width 1.7; mesoscutum length/width 0.7; mesoscutellum length/width 0.8; widths SMG/SLG 1.3; enclosed space between SMG length/width 2.1; lengths mesoscutum/mesoscutellum 1.5; lengths mesoscutellum/dorsellum 3.2; lengths mesosoma/gaster 0.8. **Wings.** CC length/width 17.5; lengths CC/MV 1.0; lengths MV/ST 1.9; lengths MV/PM 5.1; lengths PM/ST 0.4; submarginal vein with ten setae on dorsal surface (nine setae in paratype). **Gaster.** With lateral tufts of pale and flattened setae on Gt₆.

***Eriastichus renodis* sp. nov.**

<http://zoobank.org/1BC71A82-5B85-4E61-BD84-27543158E22C>

Figure 54

Type locality. Costa Rica, Cartago, Humo, El Copal, 9°47'N, 83°45'W, 1050–1250 m, 29.ii–6.iii.2008, C. Hansson leg.

Type specimen. *Holotype* male dried and glued to a paper card. Original labels: "COSTA RICA, Cartago, Humo, El Copal, 9°47'N, 83°45'W, 1050–1250 m, 29.ii–6.iii.2008, C. Hansson", "HOLOTYPE Eriastichus renodis Hansson" [red printed label], (MZLU:7046.1).

Diagnosis (male). Head dark brown, scape yellowish brown, pedicel pale brown, flagellum brown; ventral plaque on scape ca. 0.3 × as long as scape (Fig. 54), antenna with dorsobasal setae on F1 0.9 × as long as F1; gaster with lateral tufts of pale and flattened setae on Gt₆.

Description (male holotype MZLU:7046.1). Length of body 1.5 mm. Head dark brown, scape yellowish brown, pedicel pale brown, flagellum brown, ventral plaque dark brown. Mesoscutum and mesoscutellum dark brown with metallic tinges, dorsellum and propodeum dark brown. Legs yellowish brown. Gaster dark brown.

Head. Length/width in frontal view 0.7; width/length in dorsal view 2.2; POL/OOL 2.3; WM/MS 1.3; MS/HE 0.6; HE/head length in frontal view 0.6; widths head/mesoscutum 1.2. **Antenna.** Pedicel + flagellum length/mesoscutum width 2.4; pedicel + flagellum length/head width 2.0; lengths scape/ventral plaque 3.1; ventral plaque located in the middle of scape; scape length/width 3.1; lengths scape/head (dorsal view) 0.6; scape length/HE 0.9; length/width F1, F2, F3, F4, clava: 3.2, 3.4, 3.4, 3.4, 7.6; length dorsobasal setae on F1/length F1 0.9. **Mesosoma.** Length/width 1.7; mesoscutum length/width 0.6; mesoscutellum length/width 0.9; widths SMG/SLG 1.1; enclosed space between SMG length/width 2.3; lengths mesoscutum/mesoscutellum 1.4; lengths mesoscutellum/dorsellum 2.9; lengths mesosoma/gaster 0.8. **Wings.** CC length/width 20.0; lengths CC/MV 0.8; lengths MV/ST 2.3; lengths MV/PM



Figures 51–56. *Eriastichus* spp. head and antenna in lateral view, male holotypes **51** *E. sannionis* **52** *E. scalaris* **53** *E. taraxis* **54** *E. renodis* **55** *E. tendrilis* **56** *E. velaminis*.

4.2; lengths PM/ST 0.5; submarginal vein with nine setae on dorsal surface. **Gaster.** With lateral tufts of pale and flattened setae on Gt₆.

***Eriastichus rivalis* sp. nov.**

<http://zoobank.org/F78216A7-C07E-4635-A5E5-921D46D5E6AA>

Figure 47

Type locality. Costa Rica, San José, San Gerardo de Dota, 9°33'N, 83°47'W, 20–21. ii.2013, J.S. Noyes leg.

Type specimen. **Holotype** male dried and glued to a paper card. Original labels: "COSTA RICA, San José, San Gerardo de Dota, 20–21.ii.2013, J.S. Noyes, NHM (Ent) 2012–91", "HOLOTYPE Eriastichus rivalis Hansson" [red printed label], (NHMUK014431065).

Diagnosis (male). Head black, scape pale brown, pedicel and flagellum dark brown; ventral plaque on scape ca. 0.4 × as long as scape (Fig. 47), antenna with dorsobasal setae on F1 0.5 × as long as F1; gaster with lateral tufts of pale and flattened setae on Gt₆.

Description (male holotype NHMUK014431065). Length of body 2.0 mm. Head black, scape pale brown, pedicel and flagellum dark brown, ventral plaque dark brown. Mesoscutum, mesoscutellum, dorsellum and propodeum black with metallic tinges. Legs with mid and hind coxae dark brown, hind femur pale brown, remaining parts of legs yellowish brown. Gaster dark brown.

Head. Length/width in frontal view 0.8; width/length in dorsal view 2.3; POL/OOL 2.4; WM/MS 1.4; MS/HE 0.6; HE/head length in frontal view 0.6; widths head/mesoscutum 1.2. **Antenna.** Pedicel + flagellum length/mesoscutum width 2.0; pedicel + flagellum length/head width 1.9; lengths scape/ventral plaque 2.8; ventral plaque located below the middle of scape; scape length/width 3.1; lengths scape/head (dorsal view) 0.6; scape length/HE 1.0; length/width F1, F2, F3, F4, clava: 3.7, 3.2, 3.2, 3.2, 7.3; length dorsobasal setae on F1/length F1 0.5. **Mesosoma.** Length/width 1.6; mesoscutum length/width 0.6; mesoscutellum length/width 0.7; widths SMG/SLG 1.2; enclosed space between SMG length/width 2.0; lengths mesoscutum/mesoscutellum 1.5; lengths mesoscutellum/dorsellum 2.7; lengths mesosoma/gaster 0.7.

Wings. CC length/width 21.3; lengths CC/MV 1.0; lengths MV/ST 2.2; lengths MV/PM 4.1; lengths PM/ST 0.5; submarginal vein with nine setae on dorsal surface.

Gaster. With lateral tufts of pale and flattened setae on Gt₆.

***Eriastichus sannionis* sp. nov.**

<http://zoobank.org/F30DC6E4-DDEE-4240-B603-BD32C6EA5DE2>

Figure 51

Type locality. Costa Rica, Cartago, Humo, El Copal, 9°47'N, 83°45'W, 1050–1250 m, 29.ii–6.iii.2008, C. Hansson leg.

Type specimen. *Holotype* male dried and glued to a paper card. Original labels: "COSTA RICA, Cartago, Humo, El Copal, 9°47'N, 83°45'W, 1050–1250 m, 29.ii–6.iii.2008, C. Hansson", "HOLOTYPE *Eriastichus sannionis* Hansson" [red printed label], (MZLU:7047.1).

Additional type material. *Paratype* 1♂ with same label data as holotype (NHMUK014431066).

Diagnosis (male). Head yellowish brown with vertex and upper ½ of occiput brown, antenna yellowish brown; ventral plaque on scape ca. 0.5 × as long as scape (Fig. 51), antenna with dorsobasal setae on F1 0.6 × as long as F1; gaster with lateral tufts of pale and flattened setae on Gt_{5–6}.

Description (male holotype MZLU:7047.1). Length of body 1.5 mm (paratype 1.3 mm). Head yellowish brown with vertex and upper ½ of occiput brown, antenna yellowish brown, ventral plaque dark brown. Mesoscutum and propodeum dark brown, mesoscutellum black with metallic tinges, dorsellum pale brown. Legs yellowish brown. Gaster dark brown.

Head. Length/width in frontal view 0.7; width/length in dorsal view 2.2; POL/OOL 2.1; WM/MS 1.5; MS/HE 0.5; HE/head length in frontal view 0.7; widths head/mesoscutum 1.2. **Antenna.** Pedicel + flagellum length/mesoscutum width 1.8; pedicel + flagellum length/head width 1.5; lengths scape/ventral plaque 2.2; ventral plaque located above the middle of scape; scape length/width 3.3; lengths scape/head (dorsal view) 0.6; scape length/HE 0.9; length/width F1, F2, F3, F4, clava: 2.3, 2.2, 2.0, 2.0, 5.0; length dorsobasal setae on F1/length F1 0.6. **Mesosoma.** Length/width 1.6; mesoscutum length/width 0.6; mesoscutellum length/width 0.8; widths SMG/SLG 1.2; enclosed space between SMG length/width 2.2; lengths mesoscutum/mesoscutellum 1.3; lengths mesoscutellum/dorsellum 3.0; lengths mesosoma/gaster 0.8.

Wings. CC length/width 30.0; lengths CC/MV 1.0; lengths MV/ST 1.9; lengths MV/PM 5.8; lengths PM/ST 0.3; submarginal vein with eight setae on dorsal surface, in both types. **Gaster.** With lateral tufts of pale and flattened setae on Gt_{5–6}.

Eriastichus scalaris sp. nov.

<http://zoobank.org/F4846532-08CB-4CE2-AB52-5FCE0689B574>

Figure 52

Type locality. Costa Rica, San José, San Gerardo de Dota, 9°33'N, 83°47'W, 20–21. ii.2013, J.S. Noyes leg.

Type specimen. *Holotype* male dried and glued to a paper card. Original labels: "COSTA RICA, San José, San Gerardo de Dota, 20–21.ii.2013, J.S. Noyes, NHM (Ent) 2012–91", "HOLOTYPE *Eriastichus scalaris* Hansson" [red printed label], (NHMUK014431067).

Diagnosis (male). Head dark brown with part below antennal toruli pale brown, scape pale brown, pedicel and flagellum brown; ventral plaque on scape ca. 0.4 × as long as scape (Fig. 52), antenna with dorsobasal setae on F1 1.0 × as long as F1; gaster with lateral tufts of pale and flattened setae on Gt_{5–6}.

Description (male holotype NHMUK014431067). Length of body 1.5 mm. Head dark brown with part below antennal toruli pale brown, scape pale brown, pedicel and flagellum brown, ventral plaque dark brown. Mesoscutum, mesoscutellum and propodeum dark brown, dorsellum pale brown. Legs with fore and mid coxae pale brown, hind coxa and hind femur dark brown, remaining parts of legs yellowish brown. Gaster dark brown.

Head. Length/width in frontal view 0.8; width/length in dorsal view 2.1; POL/OOL 2.0; WM/MS 1.4; MS/HE 0.7; HE/head length in frontal view 0.5; widths head/mesoscutum 1.1. **Antenna.** Pedicel + flagellum length/mesoscutum width 2.0; pedicel + flagellum length/head width 1.7; lengths scape/ventral plaque 2.5; ventral plaque located in the middle of scape; scape length/width 3.1; lengths scape/head (dorsal view) 0.5; scape length/HE 1.0; length/width F1, F2, F3, F4, clava: 2.5, 2.3, 2.5, 2.5, 5.8; length dorsobasal setae on F1/length F1 1.0. **Mesosoma.** Length/width 1.6; mesoscutum length/width 0.7; mesoscutellum length/width 0.5; widths SMG/SLG 0.9; enclosed space between SMG length/width 2.4; lengths mesoscutum/mesoscutellum 1.5; lengths mesoscutellum/dorsellum 3.0; lengths mesosoma/gaster 0.8. **Wings.** CC length/width 16.0; lengths CC/MV 1.0; lengths MV/ST 2.2; lengths MV/PM 7.1; lengths PM/ST 0.3; submarginal vein with seven setae on dorsal surface. **Gaster.** With lateral tufts of pale and flattened setae on Gt₅₋₆.

***Eriastichus sodalis* sp. nov.**

<http://zoobank.org/F0A17B16-B048-4891-AF6C-3184ED9B3A21>

Figure 48

Type locality. Costa Rica, San José, San Gerardo de Dota, 9°33'N, 83°47'W, 20–21. ii.2013, J.S. Noyes leg.

Type specimen. Holotype male dried and glued to a paper card. Original labels: "COSTA RICA, San José, San Gerardo de Dota, 20–21.ii.2013, J.S. Noyes, NHM (Ent) 2012–91", "HOLOTYPE Eriastichus sodalis Hansson" [red printed label], (NHMUK014431068).

Diagnosis (male). Head dark brown with metallic tinges, scape pale brown, pedicel and flagellum dark brown; ventral plaque on scape ca. 0.3 × as long as scape (Fig. 48), antenna with dorsobasal setae on F1 0.8 × as long as F1; gaster with lateral tufts of pale and flattened setae on Gt₆.

Description (male holotype NHMUK014431068). Length of body 1.9 mm. Head dark brown with metallic tinges, scape pale brown, pedicel and flagellum dark brown, ventral plaque dark brown. Mesoscutum and mesoscutellum black with metallic tinges, dorsellum dark brown, propodeum black. Legs with mid coxa and hind femur pale brown, hind coxa dark brown, remaining parts of legs yellowish brown. Gaster dark brown.

Head. Length/width in frontal view 0.8; width/length in dorsal view 2.2; POL/OOL 1.6; WM/MS 1.6; MS/HE 0.5; HE/head length in frontal view 0.6; widths

head/mesoscutum 1.1. **Antenna.** Pedicel + flagellum length/mesoscutum width 2.0; pedicel + flagellum length/head width 1.9; lengths scape/ventral plaque 3.0; ventral plaque located in the middle of scape; scape length/width 2.9; lengths scape/head (dorsal view) 0.5; scape length/HE 0.9; length/width F1, F2, F3, F4, clava: 2.9, 2.6, 2.6, 2.4, 5.7; length dorsobasal setae on F1/length F1 0.8. **Mesosoma.** Length/width 1.7; mesoscutum length/width 0.7; mesoscutellum length/width 0.8; widths SMG/SLG 1.1; enclosed space between SMG length/width 2.2; lengths mesoscutum/mesoscutellum 1.5; lengths mesoscutellum/dorsellum 2.8; lengths mesosoma/gaster 0.7. **Wings.** CC length/width 19.3; lengths CC/MV 1.0; lengths MV/ST 2.2; lengths MV/PM 5.3; lengths PM/ST 0.4; submarginal vein with six setae on dorsal surface. **Gaster.** With lateral tufts of pale and flattened setae on Gt₆.

Eriastichus taraxis sp. nov.

<http://zoobank.org/F6E6E4E6-E788-4C49-BA77-2965F2B87E70>

Figure 53

Type locality. Costa Rica, Puntarenas, Estación Biológica Monteverde, 10°20'N, 84°49'W, 1540 m, 26.ii.2007, J.S. Noyes leg.

Type specimen. Holotype male dried and glued to a paper card. Original labels: "COSTA RICA, Puntarenas, E.B. Monteverde, 10°20'N, 84°49'W, 1540 m, 26.ii.2007, J.S. Noyes", "HOLOTYPE *Eriastichus taraxis* Hansson" [red printed label], (NHMUK014431069).

Diagnosis (male). Head dark brown, scape yellowish brown, pedicel pale brown, flagellum brown; ventral plaque on scape ca. 0.4 × as long as scape (Fig. 53), antenna with dorsobasal setae on F1 0.8 × as long as F1; gaster with lateral tufts of pale and flattened setae on Gt₆.

Description (male holotype NHMUK014431069). Length of body 1.6 mm. Head dark brown, scape yellowish brown, pedicel pale brown, flagellum brown, ventral plaque dark brown. Mesoscutum and mesoscutellum black with metallic tinges, dorsellum and propodeum dark brown. Legs yellowish brown, hind coxa with base dark brown. Gaster dark brown.

Head. Length/width in frontal view 0.8; width/length in dorsal view 2.1; POL/OOL 2.1; WM/MS 1.5; MS/HE 0.5; HE/head length in frontal view 0.6; widths head/mesoscutum 1.1. **Antenna.** Pedicel + flagellum length/mesoscutum width 2.1; pedicel + flagellum length/head width 1.9; lengths scape/ventral plaque 2.7; ventral plaque located in the middle of scape; scape length/width 3.1; lengths scape/head (dorsal view) 0.5; scape length/HE 0.8; length/width F1, F2, F3, F4, clava: 3.0, 3.0, 3.0, 2.8, 6.3; length dorsobasal setae on F1/length F1 0.8. **Mesosoma.** Length/width 1.6; mesoscutum length/width 0.7; mesoscutellum length/width 0.8; widths SMG/SLG 1.1; enclosed space between SMG length/width 2.3; lengths mesoscutum/mesoscutellum 1.6; lengths mesoscutellum/dorsellum 2.8; lengths mesosoma/gaster 0.7. **Wings.** CC length/width 22.5; lengths CC/MV 1.0; lengths MV/ST 2.0; lengths MV/PM

6.5; lengths PM/ST 0.3; submarginal vein with seven setae on dorsal surface. **Gaster.** With lateral tufts of pale and flattened setae on Gt₆.

***Eriastichus tendrilis* sp. nov.**

<http://zoobank.org/1EF86DE6-2C33-48A8-9555-08C10059AC76>

Figure 55

Type locality. Costa Rica, Puntarenas, Estación Biológica Monteverde, 10°20'N, 84°49'W, 1540 m, 18–25.ii.2004, C. Hansson leg.

Type specimen. **Holotype** male dried and glued to a paper card. Original labels: "COSTA RICA, Puntarenas, E.B Monteverde, 10°20'N, 84°49'W, 1540 m, 18–25. ii.2004, C. Hansson", "HOLOTYPE Eriastichus tendrilis Hansson" [red printed label], (MZLU:7048.1).

Additional type material. **Paratypes** (5♂♂): 2♂♂ with same label data as holotype (MZLU:7048.2, MZUCR:01696); 2♂♂ from same locality as holotype but collected 26.ii.2007 (NHMUK014431070, NHMUK014431071); 1♂ "COSTA RICA, San José, San Gerardo de Dota, El Manantial, 18–20.ii.2010, J.S. Noyes, NHM (Ent) 2010–21" (NHMUK014431072).

Diagnosis (male). Head black with metallic tinges, scape yellowish brown, pedicel and flagellum brown; ventral plaque on scape ca. 0.3 × as long as scape (Fig. 55), antenna with dorsobasal setae on F1 0.6 × as long as F1; gaster with lateral tufts of pale and flattened setae on Gt₆.

Description (male holotype MZLU:7048.1). Length of body 1.5 mm (paratypes 1.4–1.7 mm). Head black with metallic tinges, scape yellowish brown, pedicel and flagellum brown, ventral plaque dark brown. Mesoscutum and mesoscutellum black with metallic tinges, dorsellum and propodeum black. Legs with fore and mid coxae yellowish brown with base brown, hind coxa black, hind femur pale brown, remaining parts of legs yellowish brown. Gaster dark brown.

Head. Length/width in frontal view 0.7; width/length in dorsal view 2.5; POL/OOL 1.8; WM/MS 1.6; MS/HE 0.6; HE/head length in frontal view 0.6; widths head/mesoscutum 1.1. **Antenna.** Pedicel + flagellum length/mesoscutum width 1.5; pedicel + flagellum length/head width 1.5; lengths scape/ventral plaque 3.0; ventral plaque located in the middle of scape; scape length/width 3.0; lengths scape/head (dorsal view) 0.6; scape length/HE 0.9; length/width F1, F2, F3, F4, clava: 2.7, 2.7, 2.5, 2.5, 5.5; length dorsobasal setae on F1/length F1 0.6. **Mesosoma.** Length/width 1.5; mesoscutum length/width 0.7; mesoscutellum length/width 0.8; widths SMG/SLG 1.0; enclosed space between SMG length/width 2.0; lengths mesoscutum/mesoscutellum 1.6; lengths mesoscutellum/dorsellum 3.0; lengths mesosoma/gaster 1.0. **Wings.** CC length/width 18.0; lengths CC/MV 1.0; lengths MV/ST 2.1; lengths MV/PM 6.5; lengths PM/ST 0.3; submarginal vein with six setae on dorsal surface (7–8 setae in paratypes). **Gaster.** With lateral tufts of pale and flattened setae on Gt₆.

***Eriastichus tonioazofeifai* sp. nov.**

<http://zoobank.org/4E666BE9-F215-4776-AA27-1E865C9F7A12>

Figure 57

Type locality. Costa Rica, Heredia, Estación Biológica La Selva, 100–200 m, 10°26'N, 84°01'W, 30–31.iii.2002, swept, J.A. Azofeifa leg.

Type specimen. *Holotype* male dried and glued to a paper card. Original labels: "COSTA RICA, Heredia, E.B. La Selva, 100–200 m, 10°26'N, 84°01'W, 30–31. iii.2002, swept, J.A. Azofeifa," "HOLOTYPE *Eriastichus tonioazofeifai* Hansson" [red printed label], (MZLU:7049.1).

Diagnosis (male). Head dark brown with part below antennal toruli yellowish brown; antenna (Fig. 57): scape pale brown, ventral plaque on scape ca. 0.1 × as long as scape, antenna with dorsobasal setae on F1 1.1 × as long as F1; gaster with lateral tufts of pale and flattened setae on Gt₆.

Description (male holotype MZLU:7049.1). Length of body 1.3 mm. Head dark brown with part below antennal toruli yellowish brown. Antenna with scape pale brown, ventral plaque dark brown, pedicel and flagellum brown. Mesoscutum and mesoscutellum dark brown with metallic tinges; dorsellum pale brown; propodeum dark brown. Legs yellowish brown with hind coxa and hind femur pale brown. Wings ±hyaline. Gaster dark brown.

Head. Length/width in frontal view 0.7; width/length in dorsal view 2.3; POL/OOL 2.5; WM/MS 1.5; MS/HE 0.5; HE/head length in frontal view 0.6; widths head/mesoscutum 1.2. **Antenna.** Pedicel + flagellum length/mesoscutum width 2.0; pedicel + flagellum length/head width 1.6; lengths scape/ventral plaque 8.0; ventral plaque located in the middle of scape; scape length/width 5.8; lengths scape/head (dorsal view) 0.8; scape length/HE 1.3; length/width F1, F2, F3, F4, clava: 2.0, 2.0, 2.0, 2.2, 5.3; length dorsobasal setae on F1/length F1 1.1. **Mesosoma.** Length/width 1.5; mesoscutum length/width 0.6; mesoscutellum length/width 0.7; widths SMG/SLG 1.1; enclosed space between SMG length/width 2.0; lengths mesoscutum/mesoscutellum 1.4; lengths mesoscutellum/dorsellum 2.9; lengths mesosoma/gaster 0.7. **Wings.** CC length/width 21.0; lengths CC/MV 1.3; lengths MV/ST 1.9; lengths MV/PM 3.6; lengths PM/ST 0.5; submarginal vein with six setae on dorsal surface. **Gaster.** With lateral tufts of pale and flattened setae on Gt₆.

Etymology. Named after collector, Antonio (Tonio) Azofeifa, parataxonomist in Costa Rica.

***Eriastichus velaminis* sp. nov.**

<http://zoobank.org/1263C4A3-410D-4C60-A610-8F8CF591E7A9>

Figure 56

Type locality. Costa Rica, San José, San Gerardo de Dota, 9°33'N, 83°47'W, 20–21. ii.2013, J.S. Noyes leg.

Type specimen. *Holotype* male dried and glued to a paper card. Original labels: "COSTA RICA, San José, San Gerardo de Dota, 20–21.ii.2013, J.S. Noyes, NHM (Ent) 2012–91", "HOLOTYPE Eriastichus velaminis Hansson" [red printed label], (NHMUK014431073).

Diagnosis (male). Head dark brown with metallic tinges, scape and pedicel pale brown, flagellum brown; ventral plaque on scape ca. $0.4 \times$ as long as scape (Fig. 56), antenna with dorsobasal setae on F1 $0.8 \times$ as long as F1; gaster with lateral tufts of pale and flattened setae on Gt₆.

Description (male holotype NHMUK014431073). Length of body 1.2 mm. Head dark brown with metallic tinges, scape and pedicel pale brown, flagellum brown, ventral plaque dark brown. Mesoscutum and mesoscutellum black with metallic tinges, dorsellum and propodeum dark brown. Legs with coxae and femora on fore and mid legs pale brown, hind coxa and hind femur dark brown, remaining parts of legs yellowish brown. Gaster dark brown.

Head. Length/width in frontal view 0.8; width/length in dorsal view 2.1; POL/OOL 1.9; WM/MS 1.4; MS/HE 0.6; HE/head length in frontal view 0.5; widths head/mesoscutum 1.1. **Antenna.** Pedicel + flagellum length/mesoscutum width 1.8; pedicel + flagellum length/head width 1.5; lengths scape/ventral plaque 2.8; ventral plaque located below the middle of scape; scape length/width 3.1; lengths scape/head (dorsal view) 0.5; scape length/HE 1.0; length/width F1, F2, F3, F4, clava: 2.4, 2.4, 2.2, 2.0, 5.4; length dorsobasal setae on F1/length F1 0.8. **Mesosoma.** Length/width 1.5; mesoscutum length/width 0.6; mesoscutellum length/width 0.8; widths SMG/SLG 1.1; enclosed space between SMG length/width 2.0; lengths mesoscutum/mesoscutellum 1.5; lengths mesoscutellum/dorsellum 2.5; lengths mesosoma/gaster 0.8. **Wings.** CC length/width 22.0; lengths CC/MV 1.0; lengths MV/ST 2.1; lengths MV/PM 4.7; lengths PM/ST 0.5; submarginal vein with five setae on dorsal surface. **Gaster.** With lateral tufts of pale and flattened setae on Gt₆.

Eriastichus vestis sp. nov.


<http://zoobank.org/59E0A806-AAE9-4C16-B373-23BBE4FE3BDC>

Figure 58

Type locality. Costa Rica, Puntarenas, San Vito, Estación Biológica Las Alturas, 1500 m, 8°57'N, 82°50'W, 17–18.ii.2012, J.S. Noyes leg.

Type specimen. *Holotype* male dried and glued to a paper card. Original labels: "COSTA RICA, Puntarenas, San Vito, E.B. Las Alturas, 1500 m, 8°57'N, 82°50'W, 17–18.ii.2012, J.S. Noyes, NHM (Ent) 2012–91", "HOLOTYPE Eriastichus vestis Hansson" [red printed label], (NHMUK014431074).

Diagnosis (male). Head dark brown, scape and pedicel pale brown, flagellum brown; ventral plaque on scape ca. $0.3 \times$ as long as scape (Fig. 58), antenna with dorsobasal setae on F1 $0.9 \times$ as long as F1; gaster with lateral tufts of pale and flattened setae on Gt₆.



Figures 57–60. *Eriastichus* spp. **57, 58** head and antenna in lateral view, male holotypes: **57** *E. tonioazofeifai* **58** *E. vestis*; **59, 60** head in frontal view, male holotypes: **59** *E. pallidops* **60** *E. denotatis*.

Description (male holotype NHMUK014431074). Length of body 1.5 mm. Head dark brown, scape and pedicel pale brown, flagellum brown, ventral plaque dark brown. Mesoscutum, mesoscutellum and propodeum black with metallic tinges, dorsellum dark brown. Legs with coxae and femora dark brown, remaining parts of legs yellowish brown. Gaster dark brown.

Head. Length/width in frontal view 0.8; width/length in dorsal view 2.4; POL/OOL 1.9; WM/MS 1.5; MS/HE 0.6; HE/head length in frontal view 0.5; widths head/mesoscutum 1.0. **Antenna.** Pedicel + flagellum length/mesoscutum width 1.8; pedicel + flagellum length/head width 1.8; lengths scape/ventral plaque 2.9; ventral plaque located above the middle of scape; scape length/width 2.9; lengths scape/head (dorsal view) 0.5; scape length/HE 1.0; length/width F1, F2, F3, F4, clava: 3.0, 3.2, 3.0, 3.0, 7.4; length dorsobasal setae on F1/length F1 0.9. **Mesosoma.** Length/width 1.5; mesoscutum length/width 0.6; mesoscutellum length/width 0.7; widths SMG/SLG 1.5; enclosed space between SMG length/width 1.7; lengths mesoscutum/mesoscutellum 1.4; lengths mesoscutellum/dorsellum 2.5; lengths mesosoma/gaster 0.9.

Wings. CC length/width 16.0; lengths CC/MV 1.0; lengths MV/ST 2.0; lengths MV/PM 5.3; lengths PM/ST 0.4; submarginal vein with five setae on dorsal surface. **Gaster.** With lateral tufts of pale and flattened setae on Gt₆.

Discussion

The original description of *Eriastichus* was based on relatively few specimens and was a morphologically distinct genus. With the addition here of some unique features on the gaster of both sexes, the inflated pleural membrane between Gt_{1-4} and Gs_{1-4} , and the tuft of pale and flattened setae laterally on Gt_{4-6} , as well as the features given by La Salle in the original description, *Eriastichus* forms a morphologically well-defined and quite remarkable genus of the Tetrastichinae.

Species included here can only be separated through morphological characters in males, mainly on the antennae, as demonstrated in the identification key. Apart from the antennae, males of the different species are very similar, with the exception of variation in colour on the head and legs in some of the species. The antennae in females show very little variation, which is also the case for other morphological features. Thus, females cannot be separated to species, and they cannot be linked to conspecific males using morphology. Molecular data will be essential for the association of sexes of conspecific specimens, and for the identification of females.

The three species included in the original description were based mainly on females, and with relatively few specimens. Two of the species had the mesoscutellum with parts lateral to submedian grooves strongly setose, a character state not present in any of the new species described here. The third species included males from Mexico, separated in the key from species described here.

Most of the newly described species are represented by single specimens, thus eliminating the possibility of describing morphological variation within species. However, in species with two or more specimens, frequently from different localities and different collecting events, the antennal characters used to separate species show very little or no variation.

Acknowledgements

I am very grateful to John S. Noyes (NHMUK) for loan of material, and to Paul E. Hanson (University of Costa Rica) for valuable comments on the manuscript. The Microscopy Facility at the Department of Biology, Lund University, is acknowledged for the use of their scanning electron microscope.

References

- Gibson GAP (1997) Morphology and terminology. In: Gibson GAP, Huber JT, Woolley JB (Eds) Annotated keys to the genera of Nearctic Chalcidoidea (Hymenoptera). NRC Research Press, Ottawa, Canada, 16–44. <https://doi.org/10.1093/aesa/91.3.359>
- Hansson C (2020) *Lasalleistichus* a new genus of Tetrastichinae (Hymenoptera: Eulophidae) from the Neotropical region, including four new species. Journal of Natural History 54: 621–633. <https://doi.org/10.1080/00222933.2020.1715500>

- International Commission of Zoological Nomenclature (1999) International code of zoological nomenclature : Code international de nomenclature zoologique (4th ed.). International Trust for Zoological Nomenclature, c/o Natural History Museum, London.
- La Salle J (1994) North American genera of Tetrastichinae (Hymenoptera: Eulophidae). *Journal of Natural History* 28: 109–236. <https://doi.org/10.1080/00222939400770091>
- La Salle J, Schauff ME, Hansson C (2006) Familia Eulophidae. In: Hanson PE, Gauld ID (Eds) Hymenoptera de la Región Neotropical. The American Entomological Institute, Gainesville, Florida, USA, 356–374. [in Spanish]