

Geographic and taxonomic notes, addenda and corrigenda on the subtribe Bembidiina Stephens, 1827 of the 2017 'Catalogue of Palaearctic Coleoptera' (Coleoptera, Carabidae, Bembidiina)

Paolo Neri¹, Luca Toledano²

1 Via Guido Rossa 21 "San Lorenzo", 47121 Forlì (FC), Italy **2** Museo Civico di Storia Naturale di Verona, Lungadige Porta Vittoria 9, 37129 Verona, Italy

Corresponding author: Luca Toledano (lucatole2@libero.it)

Academic editor: Thorsten Assmann | Received 30 December 2020 | Accepted 6 February 2021 | Published 16 June 2021

<http://zoobank.org/71DCAFC2-B2BB-475B-9166-2BCDC83C7F07>

Citation: Neri P, Toledano L (2021) Geographic and taxonomic notes, addenda and corrigenda on the subtribe Bembidiina Stephens, 1827 of the 2017 'Catalogue of Palaearctic Coleoptera' (Coleoptera, Carabidae, Bembidiina). In: Spence J, Casale A, Assmann T, Lieberr JK, Penev L (Eds) Systematic Zoology and Biodiversity Science: A tribute to Terry Erwin (1940–2020). ZooKeys 1044: 563–587. <https://doi.org/10.3897/zookeys.1044.62593>

Abstract

Some corrections to the section of subtribe Bembidiina of the Catalogue of Palaearctic Coleoptera, Vol. 1, together with geographic, systematic, and synonymic updates are reported and commented upon. The following five new synonymies are proposed (with junior synonym listed first): *Bembidion* (*Peryphanes*) *dostali* Kirschenhofer, 1984 = *Bembidion* (*Peryphanes*) *sanatum* Bates, 1883 **syn. nov.**; *Bembidion* (*Terminophanes*) *pseudoconsumatum* Kirschenhofer, 1984 = *Bembidion* (*Terminophanes*) *sjoelanderi* Jedlička, 1965 **syn. nov.**; *Bembidion* (*Asioperphus*) *sapporense* Jedlička, 1951 = *Bembidion* (*Politophanes*) *chloreum* Bates, 1873 **syn. nov.**; *Bembidion* (*Peryphus*) *torosiense* Jedlička, 1961 = *Bembidion* (*Peryphus*) *subcostatum* vau Netolitzky, 1913 **syn. nov.**; *Sinechostictus* (*Sinechostictus*) *multisulcatus* cariniger Korge, 1971 = *Sinechostictus* (*Sinechostictus*) *multisulcatus* (Reitter, 1890) **syn. nov.** Furthermore we confirm the following synonymies: *Asaphidion* *weiratheri* Netolitzky, 1935 = *Asaphidion* *ganglbaueri* J. Müller, 1921; *Sinechostictus* (*Sinechostictus*) *effluviolum* (Peyron, 1858) = *Sinechostictus* (*Sinechostictus*) *tarsicus* (Peyron, 1858). The following nine new combinations are proposed: *Bembidion* (*Euperyphus*) *dimidiatum* Ménétriés, 1832 **comb. nov.**; *Bembidion* (*Peryphus*) *psuchrum* Andrewes, 1922 **comb. nov.**; *Bembidion* (*Plataphus*) *pseudolucillum* Netolitzky, 1938 **comb. nov.**; *Bembidion* (*Politophanes*) *chloreum* Bates, 1873 **comb. nov.**; *Bembidion* (*Politophanes*) *gotoense* Habu, 1973 **comb. nov.**; *Bembidion* (*Politophanes*) *shunichii* Habu, 1973 **comb. nov.**; *Bembidion* (*Politophanes*) *umeyai* Habu, 1959 **comb. nov.**; *Bembidion* (*Politophanes*) *yoshidai* Morita, 2009 **comb.**

nov.; *Bembidion* (*Terminophanes*) *sjoelanderi* Jedlička, 1965 **comb. nov.** The species *Bembidion psuchrum* Andrewes, 1922 and *Bembidion sanatum* Bates, 1883 are here redescribed.

Keywords

Asaphidion, *Bembidion*, *Ocys*, *Sinechostictus*, synonymy

Introduction

After the publication of the Catalogue of Palaearctic Coleoptera we correct some mistakes found in the section of the Catalogue regarding the Carabidae, Bembidiina (Maggi et al. 2017) as identified during our studies or following the suggestions of colleagues. In this contribution we provide and discuss these corrections and add some systematic and geographic updates.

Materials and methods

The systematic treatment of the Bembidiina and the geographic acronyms follow Löbl and Löbl (2017). The acronym YU includes the present Serbia (SB), Montenegro (ME), and Kosovo (KO). In the Current distribution sections, abbreviations are presented as they are in the catalogue. The abbreviations referring to new records are listed in bold. Body length was measured for card-mounted specimens from the front margin of the labrum to the apex of the elytra. The measurement of the aedeagus does not include the portion of endophallus protruding from the basal opening. Dissections were made using standard techniques. Genitalia and small parts were preserved in Euparal on acetate mounts fixed on the same pins as the specimens. The photographs of habitus were made by LT with Nikon DSFi1 and Nikon DS-L2 on a Leica Z6 microscope and those of the male genitalia by Gabriele Fiumi with a Nikon D300 on a Leitz Dialux 20 EB microscope. In the following text “the catalogue” refers to the section Bembidiina of the Catalogue of Palaearctic Coleoptera by Maggi et al. (2017).

The examined material is preserved in the following collections:

AP	coll. Andreas Pütz, Eisenhuttenstadt, Germany;
CR	coll. Christoph Reuter, Hamburg, Germany;
CTVR	coll. Luca Toledano, Verona, Italy;
DW	coll. David Wrase, Gusow Platkow (part of Zoologische Staatssammlung München, Germany);
MHB	Museum für Naturkunde. Berlin, Germany;
MSNT	Museo Civico di Storia Naturale, Trieste, Italy;
NHMB	Naturhistorisches Museum, Basilea, Switzerland;
NHMUK	Natural History Museum, London, England;
NHMD	Natural History Museum, Copenhagen, Denmark;
NHMW	Naturhistorisches Museum, Vienna, Austria;

NHRS	Swedish Museum of Natural History, Stockholm, Sweden;
NMPC	National Museum (Natural History), Prague, Czech Republic;
PN	coll. Paolo Neri, Forlì, Italy;
SMTD	Senckenberg Museum für Tierkunde, Dresden, Germany;
VS	coll. Vladimir Skoupy, Kamenne Zhrovice, Czech Republic;
VZ	coll. Vladimir Zieris, Pardubice, Czech Republic.

Taxonomy

Asaphidion ganglbaueri J. Müller, 1921

Asaphidion weiratheri Netolitzky, 1935

Material examined. 1 ♂, “Kysyldscha Hammam [= Kızılcahamam] / Paphlagonien, Asm. B. / Weirather, Innsbruck [printed] // *ganglbaueri* m. [handwritten] / det. J. Müller [printed]” (MSNT); 1 ♀, “Kysyldscha Hammam [= Kızılcahamam] / Paphlagonien, Asm. B. / Weirather, Innsbruck [printed]” (MSNT).

Notes. *Asaphidion ganglbaueri* J. Müller, 1921 was described based on six specimens collected on Bosdagh, Turkey; it is very similar to *Asaphidion rossi* Schaum, 1857 but with the antennae completely blackish. *Asaphidion weiratheri* Netolitzky, 1935 was described from Turkey, and the type locality is Kysilka-Amam, near Ankara; the species is compared with *Asaphidion caraboides* Schrank, 1781 and its subspecies without any mentions of *A. ganglbaueri*. Müller (1937), examining topotypical specimens of *weiratheri* provided by Weirather himself and, after comparison with *ganglbaueri*, stated that they belonged to the same taxon. The synonymy was accepted by Lorenz (1998) and by Casale and Vignataglianti (1999). Marggi et al. (2003) omitted to take into consideration the synonymy stated by Müller (1937) and listed *weiratheri* as a valid species; this treatment was later followed by Lorenz (2005) and Marggi et al. (2017). We requested loan of the type series of both species from NHMW but they are missing (Harry Schillhammer, pers. comm.).

Conclusions. Considering that the topotypical specimens (Kysilka-Amam, Turkey) sent by Weirather and probably belonging to the type series were identified by Müller as *A. ganglbaueri*, that Netolitzky (1935) did not take into consideration or ignored the description of *A. ganglbaueri* when he described *A. weiratheri*, and that Müller (1937) himself stated the synonymy of both species after examination of both type series, we think that the synonymy stated by Müller (1937) should be retained until proven otherwise. Therefore we confirm that *Asaphidion weiratheri* Netolitzky, 1935 is a synonym of *Asaphidion ganglbaueri* J. Müller, 1921.

Bembidion (Bembidion) quadripustulatum Audinet-Serville, 1821

We report the species from Syria: Syria, Dayr az Zawr, 10 km SE, Euphrates River bank (CTVR). Current distribution in Asia: **A:** CY, ES, FE, IN, IQ, IS, KI, KZ, LE, NMO, **SY**, TD, TM, TR, UZ, WS.

***Bembidion (Bembidionetolitzkya) atrocaeruleum* Stephens, 1828**

The mention for Austria provided by Degasperi et al. (2014) was not included in the catalogue. Current distribution: **E:** AU, BE, CZ, FR, GB, GE, IR, IT, NL, PT, SK, SP, SZ, UK.

***Bembidion (Bembidionetolitzkya) rhodopense* Apfelbeck, 1902**

The mention for Greece by Müller-Motzfeld and Marggi (2011) was not reported in the catalogue. Furthermore we report the species for Montenegro: Montenegro, Plav, Vusanje, torr. Grija, 1000 m (PN). Current distribution: **E:** AL, BU, **GR**, MC, **ME**, YU.

***Bembidion (Emphanes) netolitzkyanum* Schatzmayr, 1940**

We mention the species for Syria. Syria, 20 km S of Kassab (PN); Syria, Al Querdaha (VS). Present distribution: **A:** IS, **SY**.

***Bembidion (Euperyphus) dimidiatum* Ménéttriès, 1832 comb. nov.**

Bembidion (incertae sedis) dimidiatum Ménéttriès, 1832

Notes. Ménéttriès (1832) describes *Bembidion (Peryphus) dimidiatum* from the banks of “Potkouvà” near “Pétigorsk” on “quelques individus”. After the description, *B. dimidiatum* was retained as strictly related to *B. oblongum* Dejean, 1831 or *B. ripicola* Dufour, 1820 (currently all species belonging to the subgenus *Euperyphus* Jeannel, 1941). Netolitzky (1914a) synonymized *B. dimidiatum* with *B. tricolor* (Fabricius, 1801) (= *B. varicolor* (Fabricius, 1803); the synonymy derives from the fact that the author examined Caucasian specimens identified as *B. dimidiatum* actually belonging to *B. (Bembidionetolitzkya) varicolor* and was perhaps unaware that this last species also occurs in the Caucasus and shares the same particular pattern of color of the European *varicolor*. But later (1935) the author himself retained *B. dimidiatum* as closely related to *oblongum*, and in any case not similar to *B. conforme* Dejean, 1831 or *B. tricolor* (both belonging to subgenus *Bembidionetolitzkya* Strand, 1929). Netolitzky (1943a: 112/84, note 83) reported that *B. dimidiatum* belongs to the “Gruppe des *B. ripicola – oblongum*” and retained it as close to *B. testaceum parallelipenne* Chaudoir, 1850; moreover, Netolitzky (1943b: 17/113, note 18) mentioned that, if they were synonyms, *B. dimidiatum* should have priority over *B. parallelipenne*; he did not formally state in any way this synonymy. Later, Kryzhanovskij et al. (1995) and Lorenz (1998) listed *B. dimidiatum* as a synonym of *B. parallelipenne* but expressed doubts. Marggi et al. (2003: 271) completely changed direction, moving *dimidiatum* to “incertae sedis”, explaining the decision as follows: “*Bembidion (Peryphus) parallelipenne* ? syn. *dimidiatum* Ménéttriès, 1832, resurrected as *Bembidion*

(incertae sedis) *dimidiatum* Ménériés, 1832. It is not a member of *Actedium*, as given in Lorenz, 1998” (Marggi et al. 2003: 21).

The same settlement is confirmed by Lorenz (2005) and Marggi et al. (2017). Unfortunately, the sentence of Marggi et al. (2003) above and in particular the reference to *Actedium* probably caused the moving of *B. dimidiatum* to “incertae sedis”, and are unclear; we verified that in Lorenz (1998) neither *B. parallelipenne*, nor *B. dimidiatum* are reported as belonging to the subgenus *Actedium*. From the literature we ascertained that *B. dimidiatum* certainly belongs to the subgenus *Euperyphus* (as currently intended, see above); there still remains uncertainty regarding its precise position, with three (*B. parallelipenne parallelipenne*, *B. parallelipenne exisum* Lutshnik, 1938, *B. parallelipenne pseudoricola* Iablokoff-Khnzorian, 1963) taxa currently assigned to the same group. In our opinion it seems unlikely that all the three Caucasian forms could be valid, and it would be necessary to verify them; in any case we suppose that the subspecies more similar to *B. dimidiatum*, mainly due to the elytral coloration, could be *B. pseudoricola*; in case of possible synonymies the problem of priority already reported by Netolitzky (1943b: 17/113, note 18) mentioned above should be considered. However *B. dimidiatum* must be included in the subgenus *Euperyphus* as currently intended.

***Bembidion (Neja) ambiguum* Dejean, 1831**

Bembidion hesperus Crotch, 1867 (Lindroth 1960) must be added to the list of synonyms of *B. ambiguum* Dejean, 1831.

***Bembidion (Ocydromus) hiekei* Müller-Motzfeld, 1986**

We report the species for Afghanistan: Afghanistan, Cheikehabad (CTVR). Current distribution: **A:** AF, IN, TM.

***Bembidion (Ocydromus) semilotum* Netolitzky, 1911**

We report the species from Iraq: N Iraq, W Kurdistan, SE Al-Amādiya (DW, CTVR). Current distribution: **E:** AR; **A:** IN, **IQ**, TR.

***Bembidion (Ocydromus) siculum durudense* Marggi & Huber, 1999**

We report the species from Iraq: N Iraq, N Choman, Halgurd Mt., 3200–3400 m (CR). Current distribution: **E:** AB, AR, GG, ST; **A:** IN, **IQ**, TR.

***Bembidion (Ocydromus) toledanoianum* Echaroux, 2008**

We report the species for Turkey: Riviera bei Kemer, M5/96, Sieber (DW) and Prov. Mugla, north Dalaman (CTVR). The species was formerly known only from Rhodes, Greece. Current distribution: **E:** GR (Rhodes); **A:** TR.

***Bembidion (Ocyturanus) kurdistanicum* Netolitzky, 1920**

We report the species from Iraq: N Choman, Halgurd Mt., 3200–3400 m (PN, CTVR). Current distribution: **A**: TR, IN, **IQ**.

***Bembidion (Ocyturanus) praeustum* Dejean, 1831**

We report the species from Iraq: Rawandoz, Akolan valley, 1900–2000 m (PN). Current distribution: **E**: AL, BH, BU, CR, FR, GR, IT, MA, MC, RO, SB, SL, ST, TR, UK, YU; **N**: EG, LB; **A**: CY, IN, **IQ**, IS, JO, LE, TR.

***Bembidion (Ocyturanus) ronfelixi* Neri & Toledano, 2017**

We report the species from Tajikistan: Pamir-Alai, Seravahan-Mts., Zavron Valley, 2100–3000 m (DW, PN). Current distribution: **A**: KI, KZ, **TD**, TM, UZ.

***Bembidion (Odontium) striatum* (Fabricius, 1792)**

We report the species from Greece: Macedonia, Lake Kerkini c/o Megalochori (CTVR, PN) and Sidirokastro (CTVR); from Poland: 5 Km NE Lezajsk, Kurilowica env., banks San River (CTVR) and Kozki, Bug River (CTVR, VZ); from Kazakhstan: KZ or., Kurkum distr., 50 km N Zaisan, Chernyi river, 400 m (CTVR, AP). Current distribution: **E**: AL, AU, BH, BU, BY, CR, CT, CZ, DE, EN, FI, FR, GE, **GR**, HU, IT, LA, LT, MC, MD, NL, NT, **PL**, PT, RO, SK, SL, SP, ST, SZ, TR, UK, YU; **A**: **KZ**, TR, WS.

***Bembidion (Omotaphus) mixtum* Schaum, 1863**

In the revision of the subgenus *Omotaphus* Netolitzky, 1914 Bonavita et al. (2016) stated that both *B. madagascariense* Chaudoir, 1876 and *B. picturatum* Fairmaire, 1898 were valid species and not synonyms of *B. mixtum* Schaum, 1863. They also stated that *B. stricticollis* Jeannel, 1955 was a synonym of *B. madagascariense*, and that *B. tumidum* Gemminger & Harold, 1868 and *B. variegatum* Boheman, 1848 were synonyms of *B. mellissi* Wollaston, 1869; all of these African species were formerly listed as synonyms of *B. mixtum* and therefore they must be removed from the list of synonyms of *B. mixtum*.

***Bembidion (Pekinium) chinense* Csiki, 1928**

Bembidion (Pekinium) chinense Csiki, 1928, type species of the monospecific *Bembidion* subgenus *Pekinium* Csiki, 1901, was declared “species inquirenda” by Toledano and Sciaky (1998); this species must be transferred to the group of species “incertae sedis”.

***Bembidion (Peryphanes) dalmatinum dalmatinum* Dejean, 1831 and *Bembidion (Peryphanes) dalmatinum levantinum* Všetečka, 1941**

Bembidion (Peryphus) dalmatinum levantinum Všetečka, 1941, was described from Syria (Damascus), Lebanon (Chtaura, Hamana) and Palestine (Haifa, today Israel); the description is in the keys for the identification of the subspecies of *B. dalmatinum* Dejean, 1831 (Všetečka 1941); the diagnostic character mentioned for *levantinum* is the color of the femora, completely reddish yellow. Netolitzky (1943b) includes Cyprus in the distribution of *B. levantinum* after examination of specimens of the Splichal-Spiller collection. Jeanne (1986) mentioned *B. levantinum*, assigned to the subgenus *Peryphanes*, from several localities in Cyprus. Austin et al. (2008) also mentioned the species for Cyprus. Marggi et al. (2017) mentioned *B. dalmatinum levantinum* for Lebanon, Israel, and Cyprus. We received from NHMW the specimens from Cyprus of the Splichal-Spiller collection seen by Netolitzky and we ascertained that the femora actually are blackish at base or on the lower side. We could also examine some specimens from Cyprus (SMTD, MHB, PN, SDEI) and all of them except for a few immature specimens have femora more or less darkened at the base or on the lower side.

Since the difference between *B. dalmatinum levantinum* and *B. dalmatinum dalmatinum* is only in the darkening of the base of the femora, we suppose that only the nominotypical form is present in Cyprus: therefore Cyprus must be removed from the distribution of the subspecies, while Syria must be added, according to the original description. Current distribution of *B. levantinum*: A: IS, LE, SY. On the other hand Cyprus must be added to the distribution of *B. dalmatinum dalmatinum*.

***Bembidion (Peryphanes) deletum* Audinette-Serville, 1821**

The nominotypical form of *B. deletum* Audinette-Serville, 1821 is not present in Spain, as already (correctly) supposed by Serrano (2003) and Ortuño and Toribio (2005). All records from Spain must be referred to *Bembidion (Peryphanes) deletum schulerianum* Müller-Motzfeld, 1986 and Spain must be removed from the distribution of *B. deletum deletum*.

***Bembidion (Peryphanes) fraxator* Ménéttriés, 1832**

Bembidion (Peryphanes) lucidum (Faldermann, 1836)

Notes. *Bembidion lucidum* Faldermann, 1836 was retained, after a study of the type specimen by Netolitzky (1910) as being very close to *B. dalmatinum* Dejean, 1831. Later Netolitzky (1914b) synonymized *lucidum* with *dalmatinum fraxator* Ménéttriés, 1832 (currently *fraxator* is considered a valid species). Marggi et al. (2003: 271) listed *B. lucidum* as a synonym of *B. dimidiatum* Ménéttriés, 1832 which was moved to “incertae sedis”; while the moving of *B. dimidiatum* was mentioned (p. 21) nothing was stated about *B. lucidum*. This situation remains also in Marggi et al. (2017).

After having carefully analyzed the literature regarding these taxa (e.g., Ménétériés 1832; Faldermann 1836; Schaum 1861; Netolitzky 1910, 1914b; Kryzhanovskij et al. 1995; Belousov and Sokolov 1996), we confirm that *B. lucidum* Faldermann is a synonym of *B. fraxator* Ménétériés and not of *B. dimidiatum* Ménétériés, which, moreover, belongs to another subgenus (see above).

***Bembidion (Peryphanes) sanatum* Bates, 1883**

Figures 1, 5, 6

Bembidion (Peryphanes) dostali Kirschenhofer, 1984 syn. nov.

Material examined. 1 ♂, holotype of *Bembidion (Peryphus) sanatum* Bates, “Type [round, bordered in red] // Japan / G. Lewis / 1910 – 310 [printed] // *B. / sanatum* / Bates [handwritten]” (NHMUK); 1 ♂, “Museum Paris / Nippon Moyen / E. Gallois 1912 [printed] // Japon. Chu- / zenji 17.8.1909 / Edme Gallois [yellow, printed] // coll. Netolitzky [printed] // *sanatum* Bts / dt Netolitzky 1937 [handwritten] // Coll. Netolitzky [printed]” (NHMW); 2 ♂♂, 3 ♀♀, paratypes of *B. dostali*, (NHMW, CTVR), with the same labels as the holotype of *B. dostali*.

The type specimen of *B. sanatum* is rather well preserved, except for the broken antennae (the left one complete but divided in three portions, the right one missing three antennomeres in the middle) and legs (the right median leg and the hind right leg with detached tarsi). All the broken parts are glued on the label.

The specimen from NHMW, identified as *sanatum* by Netolitzky and also used by Kirschenhofer (1984) in the description of *Bembidion (Peryphus) dostali* (at present assigned to the subgenus *Peryphanes* Jeannel, 1941), as an example of the closest species, is in very bad condition: completely immature, missing almost all the appendages of the head and a part of the legs; the pronotum is evidently flattened, the elytra are open and the specimen is so immature that a dissection of the male genitalia (Kirschenhofer, 1984 mistakenly mentions this specimen as a ♀) is not recommended. The general habitus and the comparison with the type confirmed the determination of Netolitzky.

Four of the paratypes of *B. dostali* from NHMW are immature. The only mature male, originally glued to the label on the dorsal side, has been dissected by us and mounted again in order to leave the dorsal side visible; the specimen lacks the three left legs.

We also examined photographs of the holotype of *Bembidion (Peryphus) dostali* Kirschenhofer, 1984 kindly provided by Alexey Solodovnikov and Mikkel Høegh Post (NHMD), and of its labels: “Japan, Unzen / 32°46’N, 130°16’E / 19.VII.1934 / Eigin Suenson leg. [printed] // Holotypus [red, handwritten] // *Bembidion* [printed] / *dostali* n.sp. [handwritten] / det.: Kirschenhofer [printed] 82 [handwritten]”.

Type locality. Niohazan [Giappone], “in June (...) near the snow” (Bates 1883).

Redescription of the holotype ♂ (Fig. 1). Total body length 5.60 mm. **Coloration:** head and pronotum dark brown; elytra brownish olive. Legs and palps orange. Antennae orange, slightly darkened from fifth antennomere. **Head:** maximum width, including eyes, 1.02 mm; interocular distance 0.63 mm; frons, clypeus, and neck




1



2



3



4

Figures 1–4. Habitus of **1** *Bembidion (Peryphanes) sanatum* Bates, 1883, holotype (NHMUK), 5.60 mm **2** *B. (Plataphus) pseudolucillum* Netolitzky, 1938 from Kibune, Kyoto (NHMW), 4.20 mm **3** *B. (Peryphus) psuchrum* Andrewes, 1935 Inde, Himachal Pradesh, Mahari, 3500 m (CTVR), 3.50 mm **4** labels of the holotype of *Bembidion sapporensis* Jedlička (NMPC). Photographs: Luca Toledano.

smooth; evident frontal furrows ending posteriorly between the first and the second supraorbital seta. Eyes weakly protruding, temples oblique towards the neck. Antennae long 3.02 mm. **Pronotum:** length along the midline 1.06 mm; width of anterior margin 0.91 mm, maximum width 1.31 mm, width of base 0.98 mm; pronotal width/pronotal length ratio 1.23; convex, more evidently near the anterior angles; sides entirely rebordered, narrowing with evident sinuate shape before the base, with which they form a slightly acute corner; lateral gutter narrow, of homogeneous width; complete surface glossy; laterobasal carina very long and evident; median line sharp, transverse anterior semilunar impression more evident; basal transverse impression punctured between the deep basal foveae. **Elytra:** length 3.55 mm, maximum overall width, at middle of elytra, 2.25 mm; oval, shoulders slightly rounded; humeral margin reaching stria 5; full microsculpture, so fine and irregularly transverse that it looks like shagreening. Striae with punctures clearly visible almost to apex, even though less impressed at apex.


Male genitalia. Aedeagus (Fig. 5) of medium-large size (1.55 mm), ventral margin with a very faint gibbosity and apex only slightly bent ventrally; central brush completely protruding from basal opening; paracopulatrix lamina and main sclerite extending towards the apex, paracopulatrix lamina narrowing anteriorly; parameres of same length (terminology according to Neri and Vigna Taglianti 2010).

Conclusions. The examination of the holotype of *B. sanatum*, currently listed as *Bembidion* “incertae sedis” in Marggi et al. (2017), shows that the species must be assigned to the subgenus *Peryphanes* Jeannel, 1941, as correctly stated by Sundukov and Makarov (2016), unfortunately unavailable to us before the publication of the catalogue in 2017. The examination of the material of *B. dostali* (Fig. 6) instead revealed that the taxon, formerly considered closely related to *B. sanatum*, is conspecific. The differences between the pronota, mentioned and drawn by Kirschenhofer (1984: 91), are due to the comparison with a completely immature specimen of *B. sanatum* (examined by us, see above); furthermore the examined paratypes of *B. dostali* show a pronotum with characters that are slightly variable. The aedeagi are very similar to one another (Figs 5, 6). According to these observations we can state that *Bembidion* (*Peryphanes*) *dostali* Kirschenhofer, 1984 = *Bembidion* (*Peryphanes*) *sanatum* Bates, 1883 syn. nov. We added the following label to the examined specimens: *Bembidion* (*Peryphanes*) *sanatum* Bates – det. Neri and Toledano 2020.

Bembidion sanatum, a Japanese species, is distinguished from the related species of Japan and China by the following characters: from *B. hykosanum* (Habu & Ueno, 1955) (JA) and *B. parepum* Jedlicka, 1933 (SCH) by the completely microsculptured or shagreened elytra, from *B. hayachinense* (Nakane, 1979) (JA) by the first three antennomeres light and the pronotum not microsculptured, and from *B. lulinense* Habu, 1973 (TAI) by the structure of the endophallus (Fig. 7).

***Bembidion* (*Peryphus*) *asiaeminoris* Netolitzky, 1935**

In Mandl (1963) *Bembidion* (*Peryphus*) *asiaeminoris* Netolitzky, 1935 is reported from “Iran (Kuh-räng, östl. Isfahan)”; its presence in Iran is reported also by Marggi et al.



Figures 5–9. Median lobe of the aedeagi of **5** *Bembidion* (*Peryphanes*) *sanatum* Bates, 1883, holotype (NHMUK), 1.55 mm **6** *B. (P.) dostali* Kirschenhofer, 1984, paratype (NHMW), 1.53 mm **7** *B. (P.) lulinense* Habu, 1973 (drawing from Habu 1973), 1.41 mm **8** *Bembidion* (*Peryphus*) *psuchrum* Andrewes, 1935, holotype (NHMUK), 0.91 mm **9** *Bembidion* (*Plataphus*) *pseudolucillum* Netolitzky, paratype (NHMW), 0.80 mm.

(2017). We received two females of the three specimens mentioned by Mandl from NHMB: the examination of these specimens led to a change of identification, revealing that they belong to *Bembidion* (*Peryphanes*) *morvanianum* Müller-Motzfeld, 1986 (= *Peryphus zagrosensis* Morvan, 1973). We are unaware of any other records of *B. asiaeminoris* from Iran, therefore Iran must be removed from the distribution of *B. asiaeminoris*. Current distribution: **E**: AB, AR, ST; **A**: TR.

***Bembidion* (*Peryphus*) *charon* Andrewes, 1926**

In the catalogue the species is reported from Japan, but we think that it is likely to be a mistake, as we are unaware of any reports of the species from this area: therefore Japan must be removed from the distribution of the species. Current distribution: **A**: NP, UP.

***Bembidion* (*Peryphus*) *cordicolle* du Val, 1852**

The mention for Bulgaria provided by Hieke and Wrase (1988) was omitted in the catalogue. Current distribution: **E**: BU, GR; **A**: TR.

***Bembidion* (*Peryphus*) *ehikoense* Habu, 1984**

The species, described from Japan and originally assigned to the subgenus *Peryphus* Dejean, 1821, was subsequently listed as “incertae sedis” (e.g., Marggi et al. 2003; Lorenz 2005; Marggi et al. 2017). The study of the original description and of the aedeagus confirm that the author correctly assigned the species to *Peryphus*.

***Bembidion* (*Peryphus*) *kuhitangi* Mikhailov & Belousov, 1991**

We noticed that the species name in the catalogue is reported on page 339 with the incorrect spelling “*kughitangi*” (sic!) and must be replaced with “*kuhitangi*”; furthermore, the species has been transferred from “incertae sedis” to the subgenus *Peryphus* Dejean, 1821 by Neri and Toledano (2017).

***Bembidion* (*Peryphus*) *obscurellum* Motschulsky, 1845**

Bembidion (*Peryphus*) *obscurellum turanicum* Csiki, 1928

Bembidion (*Peryphus*) *obscurellum fumipenne* Fassati, 1957

Bembidion (*Peryphus*) *obscurellum thibeticum* Fassati, 1957

Bembidion (*Peryphus*) *obscurellum insperatum* Lutshnik, 1938

Notes. Fassati (1957) describes *Bembidion* (*Peryphus*) *fuscicrum thibeticum* ssp. nov. and *Bembidion* (*Peryphus*) *fuscicrum fumipenne* nat. nov.; they are distinguished from the nominotypical form only by the coloration of the elytra. Lindroth (1963) synonymizes

the following taxa with *B. obscurellum* Motschulsky, 1845: *B. fuscicrum* Motschulsky, 1855, *B. fuscicrum turanicum* Csiki, 1928, *B. fumipenne* and *B. thibeticum*, the last two taxa because they are only color varieties. Kryzhanovskij et al. (1995) listed *B. turanicum* as a subspecies of *B. obscurellum* and retained *B. fumipenne* and *B. thibeticum* as varieties only; moreover Belousov's note (Kryzhanovskij et al. 1995: 85, note no. 158) reports that the subspecific situation of this polymorphic species is provisional. Recently, some authors (e.g., Lorenz 2005; Marggi et al. 2017) still list *B. turanicum*, *B. fumipenne*, and *B. thibeticum* as subspecies of *B. obscurellum*.

In light of our studies we maintain that the synonymies proposed by Lindroth (1963) were correct; moreover, our colleague Liang Hongbin (IZCAS, Beijing) confirms our opinion based on the study of hundreds of specimens from Mongolia that clearly show the variability of the species (Liang Hongbin, pers. comm.).

Regarding *Bembidion (Peryphus) obscurellum insperatum* Lutshnik, 1938, since its discriminating characters are not exclusive but shared by other subspecies, we evaluated the synonymy expressed in Marggi et al. (2017) and add to the distribution of the species the Russian Caucasus (ST), the region mentioned in the description, as already reported in Kryzhanovskij et al. (1995): Ciscaucasia, Western and Central Greater Caucasus. Current distribution: **E**: DE, NT, **ST**, SV; **A**: AF, CH, ES, IN, KA, KI, KZ, MG, TM, TR, WS, XIZ; **NAR**.

***Bembidion (Peryphus) psuchrum* Andrewes, 1922 comb. nov.**

Figures 3, 8

Bembidion (incertae sedis) psuchrum Andrewes, 1922

Examined material. 1 ♂, holotype of *Bembidium psuchrum* Andrewes, “♂ 1995 [no. dissection by Müller-Motzfeld] // Holo / type [round, bordered in red] // Type [round, bordered in red] // *Bembidium / psuchrum* Andr. / Type [handwritten] / H. E. Andrewes det. [printed] // Brit. Mus. / 1923-24 // Bunderdhunga V. / W. Almora Divis. / 8000–12000 feet / June' 19 H.G.C. [printed] // 3120 [printed]” (NHMUK); 1 ♂, “Inde / Himachal Pradesh / Mahari // 3500m / 18.VIII.80 / G. Ledoux” (CTVR) (Fig. 3).

Note. The holotype is in perfect condition and the slide with the aedeagus, on a label mounted on the same pin as the specimen, is clear and well preserved.

Type locality. Kumaon, West Almora, Bunderdhunga, 8000–12000 ft. [India, Uttarakhand] (Andrewes 1922).

Redescription of the holotype. Total body length 3.90 mm. **Coloration:** head, pronotum and elytra black, metallic. Palps dark brown with last palpomere light. Legs with femora black metallic and apex slightly lighter, tibiae reddish and tarsi reddish with blackish metallic reflections. Antennae reddish black with metallic reflections. **Head:** maximum width, including eyes, 0.82 mm; interocular distance 0.55 mm; frons and clypeus smooth with a few transverse lines, neck microsculptured at base;

evident, smooth, deep frontal furrows, ending behind at the second supraorbital seta. Eyes moderately convex, temples slightly convex and oblique towards neck. Antennae long 2.05 mm. **Pronotum:** length along mid line 0.84 mm; width of anterior margin 0.77 mm, maximum width 0.93 mm, width of base 0.61 mm; pronotal width/ pronotal length ratio 1.11; very convex, barely transverse; sides entirely rebordered, narrowing with an evident sinuous curve towards the base with which they form a slightly acute corner; marginal gutter very narrow, of homogeneous width; whole surface glossy, laterobasal carina not very evident; median line evident, slightly widened at base; transverse anterior semilunar impression with a few punctures; basal transverse impression punctured between the subquadrate, not deep, basal foveae. **Elytra:** length 2.37 mm, maximum overall elytral width, slightly beyond the middle, 1.59 mm; oval, moderately convex, evident and rounded shoulders; humeral margin reaching stria 5; totally glossy, faint trace of microsculpture only at the extreme apex. Striae with evident punctation, mainly on disc, barely visible at apex; the very faint punctation is visible also at apex. Evident apical stria.

Male genitalia. Medium sized (0.91 mm) aedeagus (Fig. 8), concave ventral margin with apex bent ventrally; central brush completely inside the median lobe, medium sized main sclerite, moderately wide and sinuate.

Conclusion. The study of the holotype of the species, currently listed as *Bembidion* “incertae sedis” (Marggi et al. 2017), suggests that the species could be assigned to the subgenus *Peryphus* Dejean, 1821.

Bembidion (*Peryphus*) *subcostatum* vau Netolitzky, 1913

Bembidion (*Peryphus*) *torosiense* Jedlička, 1961 syn. nov.

Material examined. **Holotype** ♀ of *Bembidion* (*Peryphus*) *torosiense* Jedlička, 1961, “Anatolia – Toros / Berendi Eregli / 2000m, leg. Muche [printed] // Staatl. Museum für / Tierkunde Dresden [yellow, printed] // Holotypus [red, printed] // Ankauf Muche [yellow, printed] // *Bembidion* / *Peryphus* / *torosiense* sp.n. [pink, handwritten] / det. ING. JEDLICKA [pink, printed] // = *Bembidion* / *subcostatum* vau Net. / det. P. Bonavita, 2012 [printed]” (SMTD).

Notes. In Marggi et al. (2017) the species is listed with the *Bembidion*, species “incertae sedis”. The comparison of the holotype with specimens of *B. subcostatum* vau Netolitzky, 1913 from the same locality, Eregly, Turkey (PN, CTVR) confirmed the synonymy noticed by Bonavita but never published. We wish to thank our colleague Paolo Bonavita for reporting this synonymy and allowing us to publish it here.

Bembidion (*Philochthus*) *judaicum* Sahlberg, 1908

We report the species from Iran: Chahar Mahail va Bachtari, Zagros Mts, 5 Km SW Asad Abad, 2400 m (DW). Current distribution: **E:** AB, BU, GG, GR; **A:** IN, IQ, IS, SY, TR.

***Bembidion (Philochthus) pallidiveste* Carret, 1905**

We report the species from Kazakhstan: Esil (CTVR). Current distribution: **E**: MD, ST, UK; **A**: CT, IN, IQ, IS, **KZ**, SY, TR.

***Bembidion (Plataphus) prasinum* Duftschmid, 1812**

We report the species from Montenegro: Mont., Zaton, V.76, leg. Sama (PN), first mention for the balcanic-mediterranean region. Current distribution: **E**: AU, BE, CZ, FI, FR, GB, GE, HU, IT, **ME**, NL, NR, NT, SK, SL, SP, SV, SZ, UK; **A**: ES, FE, KZ, MG, WS.

***Bembidion (Plataphus) pseudolucillum* Netolitzky, 1938 comb. nov.**

Figures 2, 9

***Bembidion (Peryphus) pseudolucillum* Netolitzky, 1938**

Material examined. 1 ♀, “Japan / Settsu / Katsuoji / 16.VI.24 / J.E.A. Lewis [printed] // 199 [handwritten] // Type [red, printed] // *pseudolucillum* Net. / Type Netolitzky 37 // *Peryphus* // H.E. Andrewes coll. / B.M. 1945-97 [printed]” (NHMUK). 3 ♀♀, on the same pin, “Oku Nikko / Nabagawa / 8.9.1937 / Jano, Jap. [handwritten] // *pseudolucillum* / det. Netolitzky [handwritten] // H.E. Andrewes coll. / B.M. 1945-97 [printed]” (NHMUK). 1 ♀, “Japan / Settsu / Katsuoji / J.E.A. Lewis [handwritten] // [one deleted label] // *Peryphus* sp. / striis mediocr- / punctat [illegible] / dt Netolitzky [handwritten] // mit Type / Brit. Museum / identisch Net. [handwritten] // Coll. / Netolitzky [printed] // *pseudolucillum* Net / Cotypus Netolitzky 37 [handwritten] // CO / TYPUS [red, printed] // Coll. / Netolitzky [printed]” (NHMW). 1 ♂, “Oku Nikko / Nabagawa / 8.9.37 Jano [handwritten] // *pseudolucillum* / dt. Netolitzky [handwritten, contoured in red] // coll. / Netolitzky [printed]” (NHMW); from the handwritten label contoured in red we deduce that this last specimen was considered by Netolitzky as a paratype. 1 ♂, 2 ♀♀, on the same pin, “Oku Nikko / Nabagawa / 8.9.37 Jano [handwritten] // coll. / Netolitzky [printed] // *pseudolucillum* / dt. Netolitzky [handwritten]” (NHMW); we mounted the ♂ on a separate pin, the labels have been photocopied in order to mount the specimens on separate pins, and the ♂, dissected, bears the slide with the aedeagus mounted in euparal on the same pin. 1 ♂, “Kibune, Kyoto / 12-IV-1950 / Leg. S. Uéno [handwritten] // Acqu.-Nr. / 1951-23 [printed] // Coll. / Meyer // *pseudolucillum* / Netolitzky 1938 [handwritten] / Det. S. Uéno [printed]” (NHMW) (Fig. 2). 2 ♀♀, “Kibune, Kyoto / 12-IV-1950 / Leg. S. Uéno [handwritten] // Acqu.-Nr. / 1951-23 [printed] // Coll. / Meyer // *pseudolucillum* [handwritten] / det. Paul Meyer [printed]” (NHMW). 1 ♂, “Japan G. Lewis // Japan. Lewis // Sharp Coll. 1905-313 // *Bembidion* cfr. *pseudolucillum* det. L. Toledano 2008 // CTVR Coll. Toledano Verona – Italy” (CTVR). 1 ♂, “Kujukawa / Aomori Pref. / Honehu

Japan / 10.9.54 [handwritten] // *Bembidion* / *pseudolucillum* / Netolitzky 1938 / det. L. Toledano 2011 [printed] // CTVR / Coll. Toledano / Verona – Italy [printed]” (CTVR). 2 ♀♀, “Numata, Yumma Pref., Honshu Japan” (CTVR, PN). 1 ♀, “Amagi-Pass, Kawazu-cho, Shizuoka pref.” (CTVR). 1 ♀, “Sanjo-valley, Tabayama vill., Yamana-shi-pref.” (CTVR). 1 ♀, “J. Honshu, Shiga-ken, Mikunidakeyama, E slope, 700 m” (CTVR). 2 ♀♀, 3 ♂♂, “Japan, Mt. Hiba, Saljo, Hiroshima pref.” (CTVR, PN).

Notes. Netolitzky (1938) describes *Bembidion* (*Peryphus*) *pseudolucillum* from Settsu (Japan), assigning it to the group of *B. nitidulum* Marsham, 1802, s. l. (= *B. deletum* Serville, 1821); again Netolitzky (1943a, 1943b) found it difficult to find a correct position for the species in his Bestimmungs-Tabellen: he mentions similarities both with species of two subgenera, *Bembidionetolitzya* Strand, 1929 and *Peryphus* Dejean, 1821, but preferred the latter. Jedlička (1965), in his monograph on the Bembidiini of Eastern Asia, includes the species in the keys for subgenus *Plataphus* Motschulsky, 1864 and that of the subgenus *Peryphus*. The species is listed in “incertae sedis” in the last catalogues (e.g., Marggi et al. 2003; Lorenz 2005; Marggi et al. 2017).

Conclusions. The examination of the holotype and of the aedeagus (Fig. 9) shows that the species belongs to subgenus *Plataphus* Motschulsky, 1864. The doubts regarding the systematic position of the species are due to its pronotal convexity, more evident than in other species of the subgenus. We added to all the non-type specimens the following label: *Bembidion* (*Plataphus*) *pseudolucillum* Net. – det. Neri and Toledano 2020. The species length varies from 3.6 to 4.7 mm, the aedeagus from 0.75 to 0.84 mm.

***Bembidion* (*Politophanes*) *chloreum* Bates, 1873 comb. nov.**

Bembidium (*Peryphus*) *chloreum* Bates, 1873

Bembidion (*Asioperypus*) *sapporensis* Jedlička, 1951 syn. nov.

Notes. The subgenus *Politophanes* Müller-Morzfeld, 1998 was restored by Schmidt (2018), after Toledano (2008) downgraded *Politophanes* to species group of subgenus *Ocydromus* Clairville, 1806. The subgenus includes the species that show particular endophallic characters (main sclerite from slightly arcuate to clearly bent, S-shaped, even long and coiled in a ring in the central portion in some species). *Bembidion chloreum*, currently listed as member of subgenus *Peryphus*, shows endophallic characters that clearly include it in the subgenus *Politophanes*.

Bembidion sapporensis Jedlička, 1951, currently assigned to the subgenus *Asioperypus* Vysoký, 1986, was described from a single male specimen: we examined the holotype (NMPC) and, based on the aedeagal characters (long main sclerite, coiled in a ring at middle) we assign it to the subgenus *Politophanes*. The labels of the specimen (Fig. 4) show that our colleague Seiji Morita (Tokyo), a specialist in Japanese Carabidae, particularly in the Bembidiina, identified the specimen as *Bembidion chloreum* Bates in 1991. Later he confirmed this identification and authorized us to state it formally here (Morita, pers. comm.): we are very grateful for that. Therefore we state the

following synonymy: *Bembidion* (*Asioperypus*) *sapporensis* Jedlička, 1951 = *Bembidion* (*Politophanes*) *chloreum* Bates, 1873 syn. nov.

***Bembidion* (*Politophanes*) *gotoense* Habu, 1973 comb. nov.**

Bembidion (*Peryphus*) *gotoense* Habu, 1973

Notes. Habu (1973a) described *Bembidion* (*Peryphus*) *gotoense* from Kyushu, Japan, and in a note reported the closeness of the species to *B. (P.) amurense trajectory* Netolitzky, 1939 (currently assigned to the subgenus *Politophanes* Müller-Motzfeld, 1998); *B. gotoense* must therefore also be transferred to *Politophanes*.

***Bembidion* (*Politophanes*) *shunichii* Habu, 1973 comb. nov.**

Bembidion (*Peryphus*) *shunichii* Habu, 1973

Notes. Habu (1973b) described *Bembidion* (*Peryphus*) *shunichii* from Formosa [Taiwan] from three females. Later Toledano (2009) assigned this species to the subgenus *Ocydromus* Clairville, 1806 s. l., “*lenae* Csiki, 1928 group” (currently referred to the subgenus *Politophanes* Müller-Motzfeld, 1998); the image provided of the aedeagus by Toledano (2009) confirms the attribution to the subgenus *Politophanes* proposed here.

***Bembidion* (*Politophanes*) *umeyai* Habu, 1959 comb. nov.**

Bembidion (*Peryphus*) *umeyai* Habu, 1959

Notes. Habu (1959) describes *Bembidion* (*Peryphus*) *umeyai* from Hokkaido, Japan, reporting that the species is closely related to *Bembidion* (*Peryphus*) *amurense trajectory* Netolitzky, 1939 (currently listed in the subgenus *Politophanes*). Later, the species was listed in Marggi et al. (2003) in two different positions, within the subgenus *Ocydromus* but with the incorrect spelling *umeayi* (sic!) and within *Bembidion* “incertae sedis” with the correct name “*umeyai*”. The same mistake, reported to us by our colleague Katsuyuki Terada (Hiroshima), also occurs in Marggi et al. (2017). In Habu’s (1959) original description and in Morita (2001) the illustration of the aedeagus of *umeyai* shows that the species belongs to the subgenus *Politophanes*.

***Bembidion* (*Politophanes*) *yoshidai* Morita, 2009 comb. nov.**

Bembidion (*Peryphus*) *yoshidai* Morita, 2009

Notes. Morita (2009a) described *Bembidion yoshidai* from SW Japan and in a note reported that the species is closely related to *Bembidion trajectory* Netolitzky, 1939 (currently listed as *Politophanes*); slightly later, Morita himself (2009b) followed Toledano (2008, 2009) and assigned the species to the subgenus *Ocydromus* Clairville, 1806 s. l. Based on the aedeagal characters reported by Müller-Morzfeld (1998) and Schmidt (2018), *yoshidai* must also be transferred to *Politophanes*.

***Bembidion* subg. *Pseudosinocys* Toledano, 2005**

Marggi et al. (2017: 334) reported *Pseudosinyocys* (sic!), while the correct spelling of the subgenus is *Pseudosinocys*.

***Bembidion* (*Terminophanes*) *sjoelanderi* Jedlička, 1965 comb. nov.**

Bembidion (*Peryphus*) *sjoelanderi* Jedlička, 1965

Bembidion (*Terminophanes*) *pseudoconsumatum* Kirschenhofer, 1984 syn. nov.

Notes. Jedlička (1965) described *Bembidion* (*Peryphus*) *sjoelanderi* from China, neglecting to specify the exact locality or mention the number of type specimens. Later, following the further division of the subgenus *Peryphus* Dejean, 1821 into several separate subgenera, the species has always been listed as *Bembidion* species “incertae sedis” (e.g., Toledano 2000; Marggi et al. 2017). We received a photograph of the holotype, female, of *sjoelanderi*, from NHRS and the examination of some exoskeletal characters (pronotum with laterobasal carina rudimental or absent; pronotum with basal punctation absent or reduced; elytral striae 1, 2, and 5 deeply impressed up to the apex) suggests that the species should be assigned to the subgenus *Terminophanes* Müller-Morzfeld, 1998. After a further comparison of *sjoelanderi* with many specimens of *Bembidion* (*Terminophanes*) *pseudoconsumatum* Kirschenhofer, 1984, species showing a marked variability in the pronotal width/pronotal length ratio (1.125 to 1.308) we were convinced that the members of both taxa were conspecific, therefore we state the following synonymy: *Bembidion* (*Terminophanes*) *pseudoconsumatum* Kirschenhofer, 1984 = *Bembidion* (*Terminophanes*) *sjoelanderi* Jedlička, 1965 syn. nov.

***Bembidion* (*Testedium*) *idriae* Mesnigg, 1934**

Bembidion idriae Mesnigg, 1934 was declared a “species inquirenda” (Neri et al. 2013), and should be added to the species “incertae sedis”.

***Bembidion* (*Trepanedoris*) *doris* Panzer, 1797**

The year of description in Marggi et al. (2017) was mistakenly stated as 1796 and should be corrected to 1797.

***Ocys tachysoides* Antoine, 1933**

In Marggi et al. (2017: 340) the abbreviation GB was typed twice and one should be deleted.

***Sinechostictus (Sinechostictus) cribrum cribrum* (du Val, 1851)**

The year of description in Marggi et al. (2017) was erroneously stated as 1852 and should be corrected to 1851.

***Sinechostictus (Sinechostictus) dahlii* (Dejean, 1831)**

Bembidion (Sinechostictus) (sic!) *dahli* ssp. *laevigaster* De Monte, 1949

Bembidion (Sinechostictus) (sic!) *dahli* ssp. *nordaffricanum* De Monte, 1949

Notes. In Marggi et al. (2017: 341) the taxa *Sinechostictus laevigaster* (De Monte, 1949) and *S. nordaffricanus* (De Monte 1949), synonyms of *S. dahlii*, were listed in the same font size as the valid species, instead of the font of the synonyms. The mistake is also recognizable by their order of appearance in the catalogue, in which the species are listed in alphabetical order: *S. laevigaster* and *S. nordaffricanus* are listed after *S. dahlii*, but they are followed by *S. decoratus* Duftschmid 1812. After *laevigaster* and *nordaffricanus* is mistakenly reported the geographic distribution, which is not reported after the synonyms in the rest of the volume: probably this misled the typographers.

Both taxa should be considered synonyms of *S. dahlii*, as stated by Ortuño and Toribio (2005). Moreover, Algeria must be added to the distribution pattern of the species, as reported by De Monte (1949). Current distribution: **E:** FR, IT, SP; **N:** **AG**, LB, MO, TU.

***Sinechostictus (Sinechostictus) ruficornis* (Sturm, 1825)**

In the revision of the Caucasian species of *ruficorne* – *stomoides* group, Fassati (1992) ascertains that *Sinechostictus ruficornis* (Sturm, 1825) was not present in the Caucasian region; therefore the Russian Caucasus (ST) must be removed from its distribution.

***Sinechostictus (Sinechostictus) multisulcatus* (Reitter, 1890)**

Sinechostictus (Sinechostictus) multisulcatus lubricus Lutsnik, 1938

Sinechostictus (Sinechostictus) multisulcatus cariniger Korge, 1971 syn. nov.

Notes. In the revision of the Caucasian species of *ruficorne* – *stomoides* group, after downgrading *S. lubricus* Lutsnik, 1938 and *S. cariniger* Korge, 1971 to subspecies of *S. multisulcatus* Reitter, 1890, Fassati (1992) defined the distribution of the subspecies

(*S. multisulcatus multisulcatus* from high altitudes in Central Caucasus; *S. cariniger* from NE Turkey and Georgia; *S. lubricus* from the Western Caucasus, Lesser Caucasus, and Georgia). He also described some diagnostic exoskeletal characters, but in the text frequently repeated the words “in general” and “often”, so that we suppose that, in his opinion, overlapping of characters are frequent in the subspecies. Kryzhanovskij et al. (1995) list the three subspecies from the Caucasus, but a note by Belousov (Kryzhanovskij et al. 1995: 90, note no. 188) says: “... since the infraspecific structure of *multisulcatum* Rtt. remains obscure”. Later *S. lubricus* was considered by all authors as a synonym of *S. multisulcatus* (Lorenz 1998, 2005; Marggi et al. 2003), while *S. cariniger*, formerly listed as synonym of *S. multisulcatus* in Lorenz (1998), was later considered a valid subspecies (Marggi et al. 2003; Marggi et al. 2017).

We examined several specimens from NE Turkey, Georgia, Azerbaijan (new record: N Azerbaijan, Kusary, Shakh-Nabad-Tshai, 3400 m), and southern Russia (Abchasia, Sochi, Krasnodarski Krai and, confirming *S. lubricus* as a synonym, we think that *S. cariniger* should also be synonymized with the nominotypical form. Therefore we state the following synonymy, with the junior synonym listed first: *Sinechostictus* (*Sinechostictus*) *multisulcatus cariniger* Korge, 1971 = *Sinechostictus* (*Sinechostictus*) *multisulcatus* (Reitter, 1890) syn. nov. Current distribution: **E**: AB, GG, ST; **A**: TR.

***Sinechostictus* (*Sinechostictus*) *tarsicus* (Peyron, 1858)**

Sinechostictus (*Sinechostictus*) *effluviolum* (Peyron, 1858)

Notes. In Neri et al. (2015) *Sinechostictus tarsicus* (Peyron, 1858) is synonymized with *S. effluviolum* (Peyron, 1858). In Paill et al. (2018: 273), based on a personal communication from Wolfgang Lorenz, the name *tarsicus* was restored (Schaum 1861), as first reviewer, and synonymized *effluviolum* with *tarsicus*. We think that this act is correct, and we confirm that *Sinechostictus* (*Sinechostictus*) *effluviolum* (Peyron, 1858) is a junior synonym of *Sinechostictus* (*Sinechostictus*) *tarsicus* (Peyron, 1858). The mistake was probably due to a misunderstanding of the Schaum's (1861) considerations. Furthermore, we report the species for Lebanon (Nabeh Safa, 1000 m (PN) and Marjayoun, env. Litani river, 280 m (CR)). Current distribution: **E**: AL, AU, BH, BU, CR, GR, HU, IT, MC, SL, ST, SZ, TR, YU; **A**: CY, IN, IQ, IS, LE, SY, TR.

Errata in references

Page 1318: the reference datum “Sahlberg R.F. 1844: *In faunam insectorum Rossicam symbola, novas ad Ochotzk lectas carabiorum species continens*. Acta Societatis Scientiarum Fennicae 3: 1–66” should be replaced by: “*In faunam insectorum Rossicam symbola, novas ad Ochotzk lectas carabiorum species continens*. Helsingforsiae: ex off. typ. Frenckelliana, 66 pp.”.

Acknowledgements

The present paper is dedicated to our colleague Terry Erwin, a giant of entomology but also a good friend and real gentlemen, and a hospitable host during a memorable visit of LT and his wife Rebecca to the Smithsonian.

We wish to thank Max Barclay and Beulah Garner (NHMUK), Johannes Bergsten (NHRS), Matthias Borer, Eva Sprecher and Isabelle Zürcher (NHMB), Andrea Colla (MSNT), Jiri Hájek (NMPC), Johannes Frisch and Bernd Jaeger (MNB), Olaf Jäger (SMTD), Hongbin Liang (IZCAS, Beijing, China), Werner Marggi (Thun, Switzerland), Seiji Morita (Tokyo, Japan), Maurizio Pavesi (Milan, Italy), Andreas Pütz (Eisenhuttenstadt, Germany), Christoph Reuter (Hamburg, Germany), Harald Schillhammer (NHMW), Vladimir Skoupy (Kamenne Zhrovice, Czech Republic), Alexey Solodovnikov and Mikkel Høegh Post (NHMD), Katsuyuki Terada (Hiroshima, Japan), David Wrase (Gusow-Platkow, Germany), Vladimir Zieris (Pardubice, Czech Republic) for kindly helping us by providing material in their care for our study or in other ways; Gabriele Fiumi (Forlì, Italy) for some photographs, the computerized editing (Adobe Photoshop CS3) and the layout of the tables; Nando Pederzani (Ravenna, Italy) for the final corrections to the draft of the manuscript; Jon Cooter (Oxford, England) for the linguistic revision; Thorsten Assmann (Lüneburg, Germany), Paolo Bonavita (Roma, Italy) and Pier Mauro Giachino (Torino, Italy) for the critical review of the text.

References

- Andrewes HE (1922) Notes on Oriental Carabidae. IV. Some new species of *Bembidium* from the Himalayas (cont.). Entomologist's Monthly Magazine 58: 238–242.
- Austin K, Small E, Lemaire JM, Jeanne C, Makris C, Georgiou G (2008) A revised catalogue of the Carabidae (Coleoptera) of Cyprus – Annales du Museum d'Histoire Naturelle de Nice – Tome XXIII: 1–199.
- Bates HW (1883) XI. Supplement to the Geodephagous Coleoptera of Japan chiefly from the collection of Mr. George Lewis, made during his second visit, from February, 1880, to September, 1881. Transactions of the Entomological Society of London, 205–290. <https://doi.org/10.1111/j.1365-2311.1883.tb02947.x>
- Belousov IA, Sokolov IM (1996) Review of the Caucasian Species of the Subgenus *Peryphanes* Jeannel (Coleoptera: Carabidae: *Bembidion*). Stuttgarter Beiträge zur Naturkunde, (A, Biologie) 40 (549): 1–40.
- Bonavita P, Toledano L, Maddison DR, Vigna Taglianti A (2016) Bembidiina of the Afro-tropical region: *Bembidion* subgenus *Omotaphus* Netolitzky, 1914 (Coleoptera, Carabidae, Bembidiini). Entomologische Blätter und Coleoptera 112: 67–106.
- Bonavita P, Vigna Taglianti A (2010) *Ocydromus* subg. *Nepha* Motschulsky, 1864: revisione tassonomica, filogenesi e biogeografia (Coleoptera Carabidae). Memorie della Società Entomologica Italiana 89: 7–180. <https://doi.org/10.4081/memorieSEI.2010.7>

- Casale A, Vigna Taglianti A (1999) Caraboid beetles (excl. Cicindelidae) of Anatolia, and their biogeographical significance (Coleoptera, Caraboidea). *Biogeographia, Lavori della Società Italiana di Biogeografia*, (n.s.) 20: 277–406. <https://doi.org/10.21426/B620110002>
- Degasperi G, Ecklet A, Kahlen M, Klarica J, Kopf T, Ledferwasch M, Schatz I, Schied J (2014) Bemerkenswerte Funde aus der Käferwelt Tirols (Coleoptera). *Beiträge zur Entomofaunistik* 14: 61–86.
- De Monte T (1949) Due nuove razze del *Bembidion* (*Synechostictus*) *dahli* Dej. (Col. Carabidae). *Bollettino della Società Entomologica Italiana* 79: 14–15.
- Faldermann F (1836) Coleoptera Persico Armenica. I. Pentamera (citata anche come: “Fauna Entomologica Tanscaucasica. I”). *Nouveaux Mémoires de la Société Imperiale des Naturalistes de Moscou* 4(1836): 1–310.
- Fassati M (1957) Eine neue Art und drei Rassen von Mittelasiatischen Bembidien aus der Untergattung *Peryphus* Steph. *Časopis Československé Společnosti Entomologické* 54: 312–321.
- Fassati M (1992) Revision der kaukasischen *Bembidion*-Arten der Untergattung *Synechostictus* Motsch. Aus der Gruppe *ruficorne-stomoides* (Coleoptera, Carabidae). *Acta Universitatis Carolinae Biologica* 36: 429–444.
- Habu A (1959) One new species of *Bembidion* from Hokkaido (Coleoptera, Carabidae). *Kontyû* 27: 257–259.
- Habu A (1973a) A new *Bembidion* Species from the Gotô Islands, Japan (Coleoptera, Carabidae). *The Entomological Review of Japan* 25: 9–10.
- Habu A (1973b) – Notes and descriptions of Formosan Carabidae taken by Dr. S.I. Uéno in 1961 (Coleoptera: Carabidae). III. *Bembidion*. *Transactions of the Shikoku Entomological Society* 11: 107–116.
- Hieke F, Wrase W (1988) Faunistik der Laufkäfer Bulgariens (Coleoptera, Carabidae). *Deutsche Entomologische Zeitschrift*, N.F., 35: 1–171. <https://doi.org/10.1002/mmnd.4810350102>
- Jeanne C (1986) Contribution a l'Histoire naturelle de l'Île de Chypre. Les Coléoptères carabiques. *Biocosme Méditerranéen*. Nice 3: 1–33.
- Jedlička A (1965) Neue Carabiden aus den Sammlungen des Riksmuseums in Stockholm (Coleoptera – Carabidae). *Entomologisk Tidskrift* 86: 202–208.
- Kirschenhofer E (1984) Neue paläarktische Bembidiinae unter besonderer Berücksichtigung der von Eigin Suenson in Ostasien Durchgeführten auf Sammlungen (Col. Carabidae) 1. Teil, *Bembidion* Latreille. *Koleopterologische Rundschau* 57: 57–92.
- Kryzhanovskiy OL, Belokiy OL, Belousov IA, Kabak II, Kataev BM, Makarov KV, Shilenkov VG (1995) Pensoft Series Faunistica n° 3. A Checklist of the Ground-Beetles of Russia and Adjacent Lands (Insecta, Coleoptera, Carabidae). Pensoft Publishers, Sofia-Mosca, 271 pp.
- Kryzhanovskiy OL, Okhotina MV, Bromblei GF, Lafer GF (1975) A survey of Carabidae (Coleoptera) from Kuril Islands. *Trudy Biol. Poc. Inst. Vladivostok* 28(131): 119–142.
- Lindroth CH (1960) The Ground-Beetles of the Azores (Coleoptera: Carabidae). *Boletim do Museu Municipal do Funchal* 23: 5–37.
- Lindroth CH (1963) The ground-beetles (Carabidae, excl. Cicindelinae) of Canada and Alaska. Part 3. *Opuscula Entomologica, Supplementum* 24: 201–408.

- Löbl I, Löbl D [Eds] (2017) Catalogue of Palaearctic Coleoptera (Vol. 1). Revised and Updated Edition. Archostemata-Myxophaga-Adephaga. Brill, Leiden-Boston, 1443 pp.
- Lorenz W (1998) Systematic List of Extant Ground Beetles of the World. Tutzing, 503 pp.
- Lorenz W (2005) Systematic List of Extant Ground Beetles of the World (2nd edn.). Tutzing, 530 pp.
- Mandl K (1963) Ergebnisse der Österreichischen Iran-Expedition 1949/50. Carabidae (Col.). Coleoptera IX. Teil. Sitzungsberichte der österreichischen Akademie der Wissenschaften 172: 179–192.
- Marggi W, Huber C, Müller-Motzfeld G, Hartmann M (2003) Carabidae: Bembidiini: Bembidiina: 242–273. In: Löbl I, Smetana A (Eds) Catalogue of Palaearctic Coleoptera (Vol. 1). Archostemata – Myxophaga – Adephaga. Apollo Books, Stenstrup, 819 pp.
- Marggi W, Toledano L, Neri P (2017) Carabidae: Bembidiini: Bembidiina: 294–342. In: Löbl I, Löbl D (Eds) Catalogue of Palaearctic Coleoptera (Vol. 1). Archostemata – Myxophaga – Adephaga. Brill, Leiden, 1446 pp.
- Ménétriés E (1832) Catalogue raisonné des objets de zoologie recueillis dans un voyage au Caucase et jusqu'aux frontières actuelles de la Perse. Académie Impériale des Sciences, St. Pétersbourg, 271 pp. <https://doi.org/10.5962/bhl.title.51784>
- Morita S (2001) On *Bembidion* (*Peryphus*) *umeyai* Habu: 293–296. Papers on Coleoptera in honor of Toshifumi Hozumi. The Coleopteran Fauna of Tokai District.
- Morita S (2009a) Notes on the Bembidiinae (Coleoptera, Carabidae) of Japan. XVIII. Two New Species of the Genus *Bembidion* from Southwest Japan. Elytra 37: 21–26.
- Morita S (2009b) Notes on the Bembidiinae (Coleoptera, Carabidae) of Japan. XXII. Systematic position of *Bembidion yoshidai*. Elytra 37: e226.
- Müller G (1937) Note su alcuni Carabidi della Balcania e della regione mediterranea. Atti del Museo Civico di Storia naturale di Trieste 13: 119–134.
- Müller-Motzfeld G, Marggi W (2011) *Bembidion* Latreille, 1802. In: Arndt E, Schnitter P, Sfenthourakis S, Wrase DW (Eds) Ground Beetles (Carabidae) of Greece. Pensoft, Sofia/Moscow, 393 pp.
- Neri P, Bonavita P, Vigna Taglianti A (2015) Note tassonomiche e sinonimiche su alcuni Bembidiina del Genere *Sinechostictus* Motschulsky, 1864 (Insecta Coleoptera Carabidae). Quaderno di Studi e Notizie di Storia Naturale della Romagna 41: 83–98. <http://www.ssnr.it/41-8.pdf>
- Neri P, Bonavita P, Vigna Taglianti A, Gudenzi I (2013) Note tassonomiche e nomenclatoriali (3° contributo) su alcuni Bembidiina della Fauna Italiana (Insecta Coleoptera Carabidae). Quaderno di Studi e Notizie di Storia Naturale della Romagna 36 (2012): 89–95. <http://www.ssnr.it/36-8.pdf>
- Neri P, Toledano L (2017) Notes on genus *Bembidion* Latreille, 1802, subgenus (*Ocyturanus*) Müller-Motzfeld, 1986, with particular reference to the *marginipenne* group and description of seven new species (Insecta: Coleoptera: Carabidae: Bembidiina). Quaderno di Studi e Notizie di Storia Naturale della Romagna 46: 59–120. <http://www.ssnr.it/46-5.pdf>
- Neri P, Vigna Taglianti A (2010) Note su *Ocydromus alticola* e *O. incognitus*, con descrizione di una nuova razza di *O. alticola* dei Monti della Laga, Appennino Centrale (Coleoptera Carabidae). Bollettino della Società Entomologica Italiana 142: 111–120. <http://www.ssnr.it/neri10.pdf>

- Netolitzky F (1910) Über einige wenig bekannte *Bembidion*-Arten. Wiener Entomologische Zeitung 29: 41–50. <https://doi.org/10.5962/bhl.part.23333>
- Netolitzky F (1914a) Die Bembidiini in Winklers Catalogus. Entomologische Blätter 10(1/2): 50–55.
- Netolitzky F (1914b) Die Bembidiini in Winklers Catalogus. Entomologische Blätter 10(7/8): 164–176.
- Netolitzky F (1935) Die *Bembidion*-Arten der Sammlung Motschulsky im Museum der Universität zu Moskau. Vereinsschrift der Gesellschaft Luxemburger Naturfreunde 45: 18–37.
- Netolitzky F (1938) Beiträge zur Kenntnis der Bembidion-Arten des fernen Ostens (Japan, Korea, Ostsibirien) (Coleoptera). I. Mitteilung. Proceedings of the Royal Entomological Society of London – Series B: Taxonomy 7: 37–39. <https://doi.org/10.1111/j.1365-3113.1938.tb01241.x>
- Netolitzky F (1943a) Bestimmungs-Tabellen europäischer Käfer (9. Stück). II. Fam. Carabidae. Subfam. Bembidiinae. 66. Gattung: *Bembidion* Latr. Bestimmungstabelle der Bembidion-Arten des paläarktischen Gebietes. (Mit Hinweisen auf holarktische, äthiopische und orientalische Arten). Koleopterologische Rundschau 28(4/6): 69/41–124/96.
- Netolitzky F (1943b) Bestimmungs-Tabellen europäischer Käfer (9. Stück). II. Fam. Carabidae, Subfam. Bembidiinae. 66. Gattung: *Bembidion* Latr. Bestimmungstabelle der *Bembidion*-Arten des paläarktischen Gebietes. (Mit Hinweisen auf holarktische, äthiopische und orientalische Arten). Koleopterologische Rundschau 29: 1/97–70/166.
- Ortuño VM, Toribio M (2005) Carabidae de la Península Ibérica y Baleares (Vol. I). Trechinae, Bembidiini. Argania editio, Barcelona, 455 pp.
- Paill W, Gunczy J, Hristovski S (2018) The Vjosa-floodplains in Albania as natural habitat for ground beetles: a hotspot of rare and stenotopic species (Coleoptera: Carabidae). Acta Zool Bot Austria 155: 269–306. https://www.zobodat.at/pdf/VZBG_155_1_0269-0306.pdf
- Schaum H (1861) Synonymische Bemerkungen. II. Berliner Entomologische Zeitschrift 5: 199–215.
- Schmidt J (2018) Alpha taxonomy from the Himalaya-Tibet orogen: New species and new synonymies in *Bembidion* Latreille, and a new key to species of the subgenus *Bembidionetolitzkyi* Strand (Coleoptera, Carabidae, Bembidiini). Entomologische Blätter und Coleoptera, 114: 361–406.
- Serrano J (2003) Catálogo de los Carabidae (Coleoptera) de la Península Ibérica. Monografías S.E.A., 9: 1–130.
- Sundukov YN, Makarov KV (2016) New or little-known ground beetles (Coleoptera: Carabidae) of Kunashir Island, Kurile Islands, Russia. Russian Entomological Journal 25(2): 121–160. <https://doi.org/10.15298/rusentj.25.2.01>
- Toledano L (2000) Systematic notes on the palaearctic Bembidiini with particular reference to the fauna of China (Coleoptera Carabidae). Memorie della Società Entomologica Italiana, 78(1999): 5–70. http://www.museostoriatnaturaleverona.it/media/_Musei/_StoriaNaturale/_Allegati/Biblioteca/Altre%20pubblicazioni/Pubblicazione%20collaboratori/Toledano/2000_Systematic_notes_palaearctic_Bembidiini.pdf
- Toledano L (2008) Systematic notes on the Palaearctic *Bembidion* Latreille, 1802 (Coleoptera, Carabidae) with particular reference to the fauna of China. In: Toledano L (Ed.) Con-

- tributions to the knowledge of the eastern Palaearctic Bembidiina (Insecta, Coleoptera, Carabidae). *Memorie del Museo Civico di Storia Naturale di Verona* 2. Serie. Sezione Scienze della Vita 18: 5–46. http://www.museostorianaturaleverona.it/media/_Musei/_StoriaNaturale/_Allegati/Biblioteca/Altre%20pubblicazioni/Pubblicazioni%20collaboratori/Toledano/2008_Systematic_notes_Palaearctic_Bembidion_Latreille.pdf
- Toledano L (2009) Notes on the Bembidiina of Taiwan with description of three new species (Coleoptera: Carabidae). *Acta Entomologica Musei Nationalis Pragae* 49: 577–598. https://www.aemnp.eu/data/article-1247/1228-49_2_577.pdf
- Toledano L, Sciaky R (1998) Three subgenera of *Bembidion* new to China, with description of a new species (Coleoptera Carabidae). *Bollettino del Museo Civico di Storia Naturale di Venezia* 48(1997): 1–18. http://www.museostorianaturaleverona.it/media/_Musei/_StoriaNaturale/_Allegati/Biblioteca/Altre_pubblicazioni/Pubblicazioni_collaboratori/Toledano/1997_Three_subgenera_of_Bembidion_China.pdf
- Všetečka K (1941) Nová Rassa od *Bembidium dalmatinum* Dej. (Col. Carabidae). *Acta Entomologica Musei Nationalis Pragae* 19: 89–90. <https://www.aemnp.eu/acta-entomologica/volume-19/442/nova-rassa-od-bembidium-dalmatinum-dej-col-carabidae-eine-neue-rasse-des-bembidium-dalmatinum-dej-col-carabidae.html>