

# Corrigenda: A new species of *Chromis* damselfish from the tropical western Atlantic (Teleostei, Pomacentridae). ZooKeys 1008: 107–138. <https://doi.org/10.3897/zookeys.1008.58805>

Emily P. McFarland<sup>1,2</sup>, Carole C. Baldwin<sup>3</sup>, David Ross Robertson<sup>4</sup>,  
Luiz A. Rocha<sup>5</sup>, Luke Tornabene<sup>1,2</sup>

**1** School of Aquatic and Fishery Sciences, University of Washington, Seattle, WA 98195-5020, USA **2** Burke Museum of Natural History and Culture, Seattle, WA 98105, USA **3** Department of Vertebrate Zoology, National Museum of Natural History, Smithsonian Institution, Washington, DC, 20560, USA **4** Smithsonian Tropical Research Institute, Balboa, Republic of Panama **5** Department of Ichthyology, California Academy of Sciences, San Francisco, California 94118, USA

Corresponding author: Luke Tornabene ([ltorna1@uw.edu](mailto:ltorna1@uw.edu))

---

Academic editor: Kyle Piller | Received 11 March 2021 | Accepted 19 March 2021 | Published 10 May 2021

<http://zoobank.org/6336309E-41C5-459A-9F44-F20DDBCBE4C>

---

**Citation:** McFarland EP, Baldwin CC, Robertson DR, Rocha LA, Tornabene L (2021) Corrigenda: A new species of *Chromis* damselfish from the tropical western Atlantic (Teleostei, Pomacentridae). ZooKeys 1008: 171–172. <https://doi.org/10.3897/zookeys.1008.58805>. ZooKeys 1036: 171–172. <https://doi.org/10.3897/zookeys.1036.65739>

---

Following the publication of our paper (McFarland et al. 2020), it was brought to our attention by Andrew Bentley (University of Kansas) that Table 3 listed the holotype of the species *Chromis enchrysurus* Jordan & Gilbert, 1882 as KU 27029, and mislabeled the species as *Chromis enchrysurus*. We regret this mistake, as there are three syntypes of *C. enchrysurus*, none of which is the specimen listed. The corrected version of the table is provided here.

**Table 3.** Morphometrics and meristics of *Chromis vanbeberae* and *Chromis enchrysurus* specimens examined. Morphometric values are as percentage of SL.

|                           | <i>Chromis vanbeberae</i>                         |         |           | <i>Chromis enchrysurus</i> |           |
|---------------------------|---|---------|-----------|----------------------------|-----------|
|                           | Holotype of <i>Chromis vanbeberae</i> USNM 446947 | Average | Range     | Average                    | Range     |
| standard length           | 73.9  | 48.2    | 13.9–98.4 | 60.7                       | 80.8–17.7 |
| body depth                | 55.2  | 51.1    | 41.6–57.7 | 50.2                       | 53.9–44.0 |
| body width                | 19.4  | 19.1    | 16.5–21.6 | 17.5                       | 19.2–13.8 |
| head length               | 30.2  | 35.4    | 30.2–41.0 | 31.6                       | 36.0–29.8 |
| snout length              | 7.9   | 8.2     | 5.2–10.3  | 8.2                        | 9.3–5.8   |
| orbit diameter            | 11.8  | 14.6    | 11.5–17.4 | 11.7                       | 14.7–10.0 |
| interorbit width          | 10.7  | 10.6    | 8.6–12.1  | 10.6                       | 14.0–9.2  |
| caudal peduncle depth     | 16.1  | 15.1    | 13.3–16.4 | 14.0                       | 15.6–9.8  |
| upper jaw length          | 9.1   | 10.0    | 6.0–14.4  | 9.7                        | 10.9–8.0  |
| predorsal length          | 33.2  | 34.0    | 28.6–42.0 | 33.7                       | 38.3–28.2 |
| spinous dorsal base       | 48.6  | 44.1    | 35.5–50.2 | 46.7                       | 50.8–36.6 |
| soft dorsal base          | 18.9  | 16.5    | 13.4–18.9 | 14.6                       | 18.0–10.4 |
| 1st dorsal spine          | 8.7   | 9.1     | 7.2–11.9  | 8.3                        | 10.3–6.7  |
| 2nd dorsal spine          | 12.9  | 14.3    | 11.4–17.5 | 12.6                       | 16.2–10.6 |
| 3rd dorsal spine          | 15.7  | 17.9    | 15.3–21.6 | 15.5                       | 19.6–12.3 |
| 4th dorsal spine          | 19.4  | 20.2    | 16.6–24.5 | 17.4                       | 22.4–13.5 |
| 5th dorsal spine          | 20.6  | 20.5    | 16.2–25.9 | 17.4                       | 22.4–13.5 |
| 6th dorsal spine          | 19.8  | 18.6    | 15.5–23.7 | 17.0                       | 21.6–13.3 |
| last dorsal spine         | 16.4  | 13.8    | 10.3–17.4 | 12.3                       | 16.1–9.3  |
| longest dorsal ray        | 23.8  | 23.2    | 21.1–28.5 | 19.1                       | 23.0–16.1 |
| preanal length            | 64.1  | 67.0    | 63.2–69.7 | 66.5                       | 69.9–63.1 |
| 1st anal spine            | 9.3   | 8.7     | 5.8–11.6  | 8.1                        | 9.9–5.5   |
| 2nd anal spine            | 19.9  | 19.2    | 15.1–22.4 | 18.8                       | 21.8–16.0 |
| longest anal ray          | 23.4  | 24.1    | 18.9–28.0 | 19.9                       | 26.3–16.3 |
| caudal length             | 41.0  | 36.8    | 29.7–44.9 | 31.4                       | 35.8–27.3 |
| longest pectoral ray      | 34.2  | 33.8    | 31.1–38.1 | 31.2                       | 33.7–28.6 |
| prepelvic length          | 35.2  | 38.4    | 35.2–43.6 | 37.3                       | 41.7–33.8 |
| pelvic spine length       | 22.2  | 20.3    | 18.7–22.4 | 20.0                       | 31.2–17.2 |
| 1st pelvic soft ray       | 40.9  | 35.4    | 28.8–43.2 | 23.4                       | 36.8–30.8 |
| dorsal rays               | 12  | 12.7    | 12–13     | 12.2                       | 11–15     |
| anal rays                 | 12  | 12.6    | 12–13     | 12.1                       | 11–13     |
| pored lateral line scales | 17  | 16.5    | 15–17     | 17.2                       | 16–18     |
| upper gill rakers         | 7   | 7.3     | 7–8       | 7.5                        | 7–8       |
| lower gill rakers         | 17  | 16.9    | 16–18     | 16.8                       | 16–18     |

## Reference

McFarland EP, Baldwin CC, Robertson DR, Rocha LA, Tornabene L (2020) A new species of *Chromis* damselfish from the tropical western Atlantic (Teleostei, Pomacentridae). ZooKeys 1008: 107–138. <https://doi.org/10.3897/zookeys.1008.58805>