

# New species and records of the subgenus *Libnotes* (*Laosa*) Edwards (Diptera, Limoniidae) from China with a key to world species

Zehui Kang<sup>1</sup>, Xiao Zhang<sup>1</sup>

<sup>1</sup> Key Lab of Integrated Crop Pest Management of Shandong Province, College of Plant Health and Medicine, Qingdao Agricultural University, Qingdao 266109, China

Corresponding author: Xiao Zhang ([xzhang\\_cn@163.com](mailto:xzhang_cn@163.com))

---

Academic editor: N. Dorchin | Received 13 March 2021 | Accepted 9 May 2021 | Published 1 June 2021

---

<http://zoobank.org/21E88AC1-646E-4009-9BDC-DA2721173FAE>

---

**Citation:** Kang Z, Zhang X (2021) New species and records of the subgenus *Libnotes* (*Laosa*) Edwards (Diptera, Limoniidae) from China with a key to world species. ZooKeys 1041: 101–112. <https://doi.org/10.3897/zookeys.1041.65906>

---

## Abstract

Twenty species of *Libnotes* (*Laosa*) Edwards, 1926 are known worldwide and three are known from China so far. Here, two species of *Laosa* are added to the Chinese fauna, of which *L. (L.) baiyunensis* sp. nov. is described and illustrated as new to science, and *L. (L.) fuscineris* Brunetti, 1912 is newly recorded from China. Morphologically, the new species is most similar to *L. (L.) charmosyne* (Alexander, 1958) but can be distinguished by the pleura of the thorax, the relative position of the additional cross veins in cell  $r_3$  and  $r_5$ , and the details of the male genitalia. A key to the world species of *Laosa* is presented.

## Keywords

Chinese fauna, crane flies, Limoniinae, new record, taxonomy

## Introduction

*Libnotes* Westwood, 1876 is a species-rich Limoniidae genus with a total number of 293 species and subspecies, separated into eight subgenera: *Afrolimonia* Alexander, 1965, *Goniodineura* van der Wulp, 1895, *Gressittomyia* Alexander, 1936a, *Laosa* Edwards, 1926, *Libnotes* (s. str.), *Metalibnotes* Alexander, 1972, *Neolibnotes* Alexander,

1972 and *Paralibnotes* Alexander, 1972. The subgenus *Laosa* constitutes a small group within the genus with 20 known species from the Oriental (nine species), Australasian/Oceanian (eight species) and Palaearctic (three species) regions (Oosterbroek 2021), and here an additional new species from China is described and illustrated. It can be easily distinguished from other subgenera by the wing having two additional cross veins in cells  $r_3$  (r-r) and  $r_5$  (r-m, absent in some species) and  $Sc_1$  ending far beyond the fork of Rs. Detailed features for recognition were given by Edwards (1926) and Podenas and Byun (2018).

Three species of the subgenus *Laosa* were previously recorded from China: *L. (L.) diphragma* (Alexander, 1934a), *L. (L.) regalis* Edwards, 1916 and *L. (L.) transversalis* de Meijere, 1916. In this paper, two *Laosa* species are added to the Chinese fauna, of which *L. (L.) baiyunensis* sp. nov. is described and illustrated as new to science and *L. (L.) fuscinervis* Brunetti, 1912, known previously only from India, is newly recorded from China. A key to the world species of *Laosa* based on types and non-type specimens, and on the literature is presented.

## Material and methods

Specimens for this study were collected from several localities in China by different entomologists between 2002–2016. Type specimens are deposited in the Entomological Museum of China Agricultural University, Beijing, China (CAU). Other studied specimens are deposited in Qingdao Agricultural University, Shandong, China (QAU). We also examined specimens from the National Museum of Natural History, Smithsonian Institution, Washington D.C., USA (USNM) and the Natural History Museum, London, UK (NHM) (Table 1). Genitalic preparations of males were made by macerating the apical portion of the abdomen in cold 10% NaOH for 12–15 hours. Observations and illustrations were made using a ZEISS Stemi 2000-C stereomicroscope. Photographs were taken with a Canon EOS 77D digital camera through a macro lens. Details of coloration were examined in specimens immersed in 75%  $C_2H_5OH$ .

**Table 1.** Information of the examined specimens from USNM and NHM.

Species	Specimens examined	Collection
<i>L. (L.) charmosyne</i>	Holotype, male, Japan: Shikoku, Mt. Ishizachi (1800 m), 1956.VI.16, T. Yano.	USNM
<i>L. (L.) fuscinervis</i>	Paratype, male, India: East Himalayas, Dajiling (1829 m), 1908.IX.22, E. Brunetti.	NHM
<i>L. (L.) kariyana</i>	Holotype, male, Japan: Honshu, Ontake (1800 m), 1934.VII.6–10, H. Ise.	USNM
<i>L. (L.) manobo</i>	Holotype, male, Philippines: Mindanao, Mt. Apo (1981 m), 1930.IX.14, C. F. Clagg.	USNM
<i>L. (L.) noctipes</i>	Holotype, female, India: Sikkim, karponang (2469 m), 1959.VIII.22, Schmid.	USNM
<i>L. (L.) regalis</i>	Holotype, male?, China: Taiwan, Taihoku, T. Shiraki. Other material: 1 male, China: Taiwan, Arisan, 1917.IV.20, T. Shiraki.	NHM
<i>L. (L.) rotundifoliae</i>	Paratypes, 2 males 2 females, Indonesia: Sulawesi Utara, Dumoga-Bone National Park (211 m), 1985.VIII. 19–30, Chen W. Young.	NHM
<i>L. (L.) taficola</i>	Holotype, female, Papua New Guinea: Mt Tafa (2591 m), 1934.III, L. E. Cheesman.	NHM
<i>L. (L.) transversalis</i>	1 male, China: Taiwan, Arisan, 1919.IV.25, T. Shiraki.	NHM


The morphological terminology mainly follows McAlpine (1981), and that for venation follows Alexander and Byers (1981). The following abbreviations in figures are used: **tg 9** = ninth tergite, **tg 10** = tenth tergite, **goncx** = gonocoxite, **o gonst** = outer gonostylus, **i gonst** = inner gonostylus, **aed** = aedeagus, **pm** = paramere, **cerc** = cercus, **hyp vlv** = hypogynial valve.

## Taxonomy

### Key to world species of *Laosa*

- 1 Basal 1/4 of wing with complete or broken crossband; m-m shorter than basal section of  $M_3$  (Fig. 1a–c)..... **2**
- Basal 1/4 of wing without conspicuous crossband; m-m significantly longer than basal section of  $M_3$  (Figs 2d, 4d) ..... **9**
- 2 (1) Wing with broad and complete crossband extending from cord to distal end of cell dm (Fig. 1c)..... **3**
- Wing without broad or complete crossband extending from cord to distal end of cell dm (Fig. 1a, b)..... **5**
- 3 (2) Tip of wing narrowly falcate ..... *L. (L.) falcata* (Alexander, 1935)
- Tip of wing round ..... **4**
- 4 (3)  $R_s$  nearly straight or slightly curved, r-r far beyond r-m and distance between them more than twice length of r-r (Fig. 1c) .....  
..... *L. (L.) rotundifoliaeos* (Young, 1990)
- $R_s$  strongly arcuated, r-r beyond r-m and distance between them about length of r-r ..... *L. (L.) innuba* (Alexander, 1941)
- 5 (2) Crossvein r-r situated before r-m, basal section of  $CuA_1$  at fork of M (Fig. 1b)..... *L. (L.) iris* (Alexander, 1950)
- Crossvein r-r situated beyond r-m, basal section of  $CuA_1$  distinctly beyond fork of M (Figs 1a, c, 2d) ..... **6**
- 6 (5) Pleura pale yellow without dark area .... *L. (L.) bipartita* (Alexander, 1936b)
- Pleura with conspicuous dark area ..... **7**
- 7 (6) Basal section of  $CuA_1$  slightly beyond fork of M and at about 1/8 of cell dm ..... *L. (L.) manobo* (Alexander, 1931)
- Basal section of  $CuA_1$  far beyond fork of M and at 1/4–1/2 of cell dm (Figs 2d, 4d)..... **8**
- 8 (7) Coxae yellow;  $R_2$  far before tip of  $Sc_2$  and distance between them about twice length of  $R_2$ , tip of  $A_1$  bent very strongly to wing margin .....  
..... *L. (L.) pavo* (Alexander, 1964)
- Coxae brown;  $R_2$  before tip of  $Sc_2$  and distance between them about length of  $R_2$ , tip of  $A_1$  slightly curved..... *L. (L.) suffalcata* (Alexander, 1964)
- 9 (1) Wing without additional cross vein in cell  $r_5$  (Fig. 4d) ..... **10**
- Wing with additional cross vein in cell  $r_5$  (Figs 1a–c, 2d) ..... **15**

- 10 (9) Wing with stripes along veins broad and extensive, nearly covering wing tip ..... ***L. (L.) noctipes* (Alexander, 1967)**  
 – Wing with stripes along veins not as broad or extensive ..... **11**
- 11 (10) Crossvein m-m about four times or more as long as basal section of  $M_3$ .... **12**  
 – Crossvein m-m less than three times as long as basal section of  $M_3$  (Fig. 4d) ..... **13**
- 12 (11) Wing with many conspicuous spots;  $R_2$  and r-r distinct before distal end of cell dm..... ***L. (L.) taficola* (Alexander, 1948)**  
 – Wing nearly unpatterned except very light brown spots at fork of  $Sc$  and over tip of  $Sc_2$ ;  $R_2$  distinct beyond distal end of cell dm, r-r aligned with distal end of cell dm..... ***L. (L.) transversalis* de Meijere, 1916**
- 13 (11) Anterior scutum and pleura dark brown, without conspicuous pattern .....  
 ..... ***L. (L.) dolonigra* (Alexander, 1956)**  
 – Anterior scutum and pleura with conspicuous stripes (Fig. 4c)..... **14**
- 14 (13) Body length of male more than 13.0 mm; r-r aligned with distal end of cell dm (Alexander 1967)..... ***L. (L.) impensa* (Alexander, 1967)**  
 – Body length of male less than 10.0 mm; r-r distinctly before distal end of cell dm (Fig. 4d) ..... ***L. (L.) fuscinervis* Brunetti, 1912**
- 15 (9)  $R_2$  far before tip of  $Sc_2$  ..... **16**  
 –  $R_2$  close to tip of  $Sc_2$  (Fig. 2d) ..... **17**
- 16 (15) Antennal scape yellow, pedicel and flagellomeres dark brown; anterior scutum with four yellow stripes;  $Sc$  relatively short, end aligned with base of cell dm (Alexander 1959)..... ***L. (L.) jocularator* (Alexander, 1959)**  
 – Antenna black throughout; anterior scutum with three confluent dark brown stripes;  $Sc$  long, end aligned with middle of cell dm.....  
 ..... ***L. (L.) kariyana* (Alexander, 1947)**
- 17 (15) Crossvein r-r close to  $R_2$ ..... **18**  
 – Crossvein r-r far before  $R_2$  and distance between them about or more than length of r-r (Figs 2d, 4d) ..... **19**
- 18 (17) Anterior scutum with indistinct median stripe; wing length of male 10.0–15.0 mm, r-m distinctly before distal end of cell dm, tip of  $A_2$  nearly straight or slightly curved ..... ***L. (L.) charmosyne* (Alexander, 1958)**  
 – Anterior scutum with four ill-defined stripes; wing length of male about 25.0 mm, r-m aligned with distal end of cell dm, tip of  $A_2$  bent very strongly toward margin ..... ***L. (L.) regalis* Edwards, 1916**
- 19 (17) Axillary region of wing without spots (Fig. 2d) .... ***L. (L.) baiyunensis* sp. nov.**  
 – Axillary region of wing darkened (Fig. 4d) ..... **20**
- 20 (19) Tibiae yellow with broad, brown subbasal rings; r-r far before  $R_2$  and distance between them about 1.5 times length of r-r .....  
 ..... ***L. (L.) riedelella* (Alexander, 1934b)**  
 – Tibiae brownish yellow without subbasal ring; r-r before  $R_2$  and distance between them less than length of r-r.... ***L. (L.) diphragma* (Alexander, 1934a)**



**Figure 1.** Wings of *Libnotes (Laosa)* **a** *L. (L.) bipartita* (from Alexander 1936b) **b** *L. (L.) iris* (form Edwards 1926) **c** *L. (L.) rotundifoliae* (paratype, photo by Jinlong Ren).

***Libnotes (Laosa) baiyunensis* sp. nov.**

<http://zoobank.org/EE267965-E009-46B7-B13D-6129B6F06E2A>

Figs 2, 3


**Specimens examined.** *Holotype*, male (CAU), China: Henan, Songxian, Mt. Baiyun, 2002.VII.22, Ding Yang. *Paratypes*: 1 male (CAU), same data as holotype. 1 male 1 female (CAU), China: Henan, Songxian, Mt. Baiyun (1500 m), 2008.VIII.14, Ding Yang.

**Diagnosis.** Anterior scutum brown with side edges brownish black. Pleura brownish yellow with a broad brownish black stripe extending from cervical region to base of wing. Tip of wing round. Wing nearly unpatterned except some pale brown patches around cross veins and portions of longitudinal veins, without conspicuous crossband from top to bottom. Sc long, ending near middle of cell dm. Rs slightly curved.  $R_2$  slightly before tip of  $Sc_2$ . Two additional cross veins in cells  $r_3$  and  $r_4$ , the former (r-r) beyond distal end of cell dm, the latter (r-m) aligned with distal end of cell dm; m-m twice as long as basal section of  $M_3$ . Basal section of  $CuA_1$  far beyond fork of M and at about 1/3 of cell dm. Tip of  $A_2$  nearly straight.

**Description. Male.** Body length 12.0–14.0 mm, wing length 19.0–22.0 mm.


**Head** (Fig. 2b). Brown. Hairs on head brown. Antenna length 2.9 mm, brown. Scape long cylindrical; pedicel oval, nearly as long as first flagellomere; flagellomeres oval, tapering apically, terminal flagellomere 1.5 times as long as preceding segment. Mouthparts brown with white hairs; palpus brown with brown hairs.

**Thorax** (Fig. 2c). Pronotum brown with sides brownish black. Prescutum brown with side edges brownish black; posterior scutum brown. Scutellum pale brown. Mediotergite pale brown with sides brownish black. Pleura (Fig. 2a) brownish yellow with a broad brownish black stripe extending from cervical region to base of wing. Hairs on thorax white. Coxae yellow; trochanters pale yellow; femora yellow to brownish yellow with tips dark brown;



**Figure 2.** *Libnotes (Laosa) baiyunensis* sp. nov. **a** habitus of male, lateral view **b** head, dorsal view **c** thorax, dorsal view **d** wing **e** female ovipositor, lateral view. Scale bars: 5.0 mm (**a**); 3.0 mm (**d**); 1.0 mm (**b**, **c**); 0.2 mm (**e**).

tibiae brown; tarsi brown. Hairs on legs dark brown. Wing (Fig. 2d) tinged with pale brownish yellow. Darkened areas around cross veins, distal end of cell dm and CuA<sub>1</sub>, tip of M<sub>1+2</sub>, CuA and A<sub>2</sub>; three small spots at base of Rs, at fork of Sc, and over R<sub>2</sub> and tip of Sc<sub>2</sub>. Venation: Sc long, ending far beyond fork of Rs and near middle of cell dm. Basal section of Sc<sub>2</sub> very close to tip of Sc<sub>1</sub>. Tip of Sc<sub>2</sub> nearly transverse, indistinct at wing margin. Rs very short, slightly sinuous. R<sub>2</sub> slightly before tip of Sc<sub>2</sub>. Radial and medial veins distinctly curved caudally before wing margin. Two additional cross veins



**Figure 3.** *Libnotes (Laosa) baiyunensis* sp. nov. **a** male hypopygium, dorsal view **b** male hypopygium, ventral view. Scale bars: 0.5 mm.

in cells  $r_3$  and  $r_5$ , the former (r-r) at middle of cell  $r_3$ , the latter (r-m) at basal 2/5 of cell  $r_5$  and aligned with distal end of cell dm. Cell dm elongate, more than 5 times as long as its width; m-m elongate, twice as long as basal section of  $M_3$ . Basal section of  $CuA_1$  far beyond fork of M and at about 1/3 of cell dm.  $A_1$  straight.  $A_2$  slightly sinuous. Halter length 2.6 mm, yellow with knob brown.

**Abdomen.** Tergites brownish yellow with a brown median stripe, lateral borders brown; eighth tergite brown. Sternites brownish yellow with eighth sternite brown. Hairs on abdomen white.

**Hypopygium** (Fig. 3). Ninth tergite with widely rounded posterior margin and small median emargination. Gonocoxite elongate, slender with an elongate, blunt-apexed ventromesal lobe; inside edge with small setose bulge. Outer gonostylus arched at 2/3 length, tip acute. Inner gonostylus short, oval with long, arched rostral prolongation armed with two spines near base from a single tubercle; an elongate lobe arising dorsally near base with a brush of long setae at apex, at right angle to lobe and directed laterally. Paramere wide at base, elongate, triangular distally. Penis long, tip sunken in the middle.

**Female.** Body length 11.5 mm, wing length 17.5 mm. Similar to male, but eighth tergite brownish yellow with a broad brown median stripe. Tenth tergite brown. Cercus (Fig. 2e) brown, tip slightly exceeding tip of hypogynial valve. Hypogynial valve brownish yellow with borders darker, base slightly beyond base of tenth tergite.

**Etymology.** The species is named after the type locality Mt. Baiyun.

**Distribution.** China (Henan).

**Remarks.** This species is somewhat similar to *L. (L.) charmosyne* from South Korea and Japan in having similar spots on the wing, but it can be easily distinguished from the latter by the pleura of the thorax being brownish yellow with a broad brownish black stripe, the crossvein r-r being 1.5 to 2 times its length before  $R_2$ , the crossvein r-m being aligned with the distal end of cell dm, the basal section of  $CuA_1$  being far beyond the fork of M and at about 1/3 of cell dm (Fig. 2d), and the inner gonostylus

being about 2/3 length of the gonocoxite (Fig. 3a), whereas in *L. (L.) charmosyne*, the pleura of the thorax is dark brownish gray, the crossvein r-r is near  $R_2$ , the crossvein r-m is distinctly before the distal end of cell dm, the basal section of  $CuA_1$  is beyond the fork of M and at 1/6–1/5 of cell dm, and the inner gonostylus is half the length of the gonocoxite (Alexander 1958; Podenas and Byun 2018).

### *Libnotes (Laosa) fuscinervis* Brunetti, 1912

Figs 4–5

*Libnotes fuscinervis* Brunetti, 1912: 411. Type locality: Dajiling, East Himalayas (India).


**Specimens examined.** *Paratype*, male (NHM), India: East Himalayas, Dajiling (1829 m), 1908.IX.22, E. Brunetti. **Other material:** 1 male (QAU), China: Yunnan, Lvchun, Yakou (1931 m), 2016.VII.7, Qilemoge.

**Diagnosis.** Anterior scutum brown with a broad, posteriorly subdivided, dark brown median stripe and a spot on each side of it; posterior half of median stripe with a paler division that broadens out across posterior scutum and scutellum. Pleura brownish yellow with a broad, anteriorly indistinct, brownish black stripe extending from cervical region to mediotergite. Tip of wing round. Wing with many conspicuous spots but without conspicuous crossband from top to bottom. Sc long, ending at 1/3 of cell dm. Rs slightly curved.  $R_2$  before tip of  $Sc_2$  and distance between them about length of  $R_2$ . Crossvein r-r before distal end of cell dm. Additional cross vein in cell  $r_5$  absent; m-m twice as long as basal section of  $M_3$ . Basal section of  $CuA_1$  far beyond fork of M and at about 1/4 of cell dm. Tip of  $A_2$  slightly curved.

**Description. Male.** Body length 9.5 mm, wing length 14.5 mm.

**Head** (Fig. 4b). Brownish yellow. Hairs on head brown. Antenna length 2.0 mm, dark brownish yellow. Scape long cylindrical; pedicel oval, nearly as long as first flagellomere; flagellomeres oval, tapering apically, terminal flagellomere 1.5 times as long as preceding segment. Mouthparts brown with white hairs; palpus brown with brown hairs.

**Thorax** (Fig. 4c). Pronotum brownish yellow. Prescutum brown. Anterior scutum brown with a broad, posteriorly subdivided, dark brown median stripe and a spot on each side of it; posterior scutum brownish black with a broad yellow median stripe. Scutellum brownish black with a broad yellow median stripe. Mediotergite brownish black with a narrow yellow median stripe. Pleura (Fig. 4a) brownish yellow with a broad, anteriorly indistinct, brownish black stripe extending from cervical region to mediotergite. Hairs on thorax white. Coxae brownish yellow; trochanters yellow; femora brownish yellow with subtips brownish black; tibiae brownish yellow with tips narrowly brownish black; tarsi brownish yellow with tips brownish black. Hairs on legs dark brown. Wing (Fig. 4d) tinged with pale brownish yellow. Many dark patches around cross veins and portions of longitudinal veins as well as patches in cells as shown in Fig. 4d; four darker spots at base of wing, at base of Rs, at fork of Sc, and over  $R_2$  and tip of  $Sc_2$ , the latter two spots connected by a narrow stripe along  $Sc_2$ . Venation: Sc long, ending far beyond fork of Rs and




**Figure 4.** *Libnotes (Laosa) fuscinervis* Brunetti, 1912 **a** habitus of male, lateral view **b** head, dorsal view **c** thorax, dorsal view **d** wing. Scale bars: 5.0 mm (**a**); 2.0 mm (**d**); 1.0 mm (**b, c**).

at 1/3 of cell dm. Basal section of  $Sc_2$  near tip of  $Sc_1$ . Tip of  $Sc_2$  nearly transverse, indistinct at wing margin. Rs very short, slightly sinuous.  $R_2$  before tip of  $Sc_2$  and distance between them about length of  $R_2$ . Radial and medial veins distinctly curved caudally before wing margin. Crossvein r-r at basal 1/3 of cell  $r_3$ . Cell dm elongate, more than 5 times as long as its width; m-m elongate, twice as long as basal section of  $M_3$ . Basal section of  $CuA_1$  far beyond fork of M and at about 1/4 of cell dm.  $A_1$  straight, slightly curved near tip.  $A_2$  slightly sinuous. Halter length 1.5 mm, pale yellow with knob brownish black.

**Abdomen.** Tergites brownish yellow with lateral borders brownish black. Sternites brownish yellow, middle of first sternite paler. Hairs on abdomen white.

**Hypopygium** (Fig. 5). Ninth tergite with rounded posterior margin and small median emargination. Gonocoxite stubby with an elongate, blunt-apexed ventromesal



**Figure 5.** *Libnotes (Laosa) fuscineris* Brunetti, 1912 **a** male hypopygium, dorsal view **b** male hypopygium, ventral view. Scale bars: 0.5 mm.

lobe; inside edge with a large setose bulge. Outer gonostylus arched at 2/3 length, tip acute. Inner gonostylus short, oval with a long arched rostral prolongation armed with two spines at base from a single tubercle; an elongate lobe arising dorsally near base with a brush of long setae at apex, at right angle to lobe and directed laterally. Paramere wide at base, elongate, triangular distally. Penis long, tip sunken in the middle.

**Female.** Unknown.

**Distribution.** China (Yunnan); India.

**Remarks.** This species was known previously only from India. Now it is recorded from China for the first time.

## Acknowledgements

We express our sincere thanks to Ding Yang for his great help during the study. We are very grateful to Jon K. Gelhaus, Sigitas Podenas, David G. Furth, Yan Li and Qifei Liu for great help during the study of type material from USNM, and to Duncan Sivell and Jinlong Ren for assistance in the study of type material from NHM. We also thank Sigitas Podenas and Ali M. Yasir for revising the manuscript. This work was supported by grants from the National Natural Science Foundation of China (41901061), the Shandong Provincial Natural Science Foundation, China (ZR2019BC034, ZR2018LC006), the High-level Talents Funds of Qingdao Agricultural University, China (663-1119008, 663-1118015) and the National Animal Collection Resource Center, China.

## References

Alexander CP (1931) New or little-known Tipulidae from the Philippines (Diptera). XII. Philippine Journal of Science 46: 447–477.

- Alexander CP (1934a) New or little-known Tipulidae from eastern Asia (Diptera). XIX. Philippine Journal of Science 54: 309–342.
- Alexander CP (1934b) New or little-known Tipulidae from eastern Asia (Diptera). XX. Philippine Journal of Science 54: 433–471.
- Alexander CP (1935) The Diptera of the Territory of New Guinea. II. Family Tipulidae. Proceedings of the Linnaean Society of New South Wales 60: 51–70.
- Alexander CP (1936a) New or little-known Tipulidae from eastern Asia (Diptera). XXXII. Philippine Journal of Science 61: 113–149.
- Alexander CP (1936b) Tipulidae. In: Curran CH (Ed.) The Templeton Crocker expedition to western Polynesian and Melanesian islands, 1933. No. 30. Diptera. Proceedings of the California Academy of Sciences 22(4): 2–11.
- Alexander CP (1941) The Diptera of the Territory of New Guinea. XII. Family Tipulidae. Part IV. Proceedings of the Linnaean Society of New South Wales 66: 138–144.
- Alexander CP (1947) Undescribed species of Japanese crane-flies (Diptera: Tipulidae). Part VI. Annals of the Entomological Society of America 40: 350–371. <https://doi.org/10.1093/aesa/40.2.350>
- Alexander CP (1948) New or little-known Tipulidae (Diptera). LXXIX. Oriental-Australasian species. Annals and Magazine of Natural History 14(11): 388–414. <https://doi.org/10.1080/00222934708654649>
- Alexander CP (1950) Notes on the tropical American species of Tipulidae (Diptera). VI. The tribe Limoniini, genus *Limonia*: subgenera *Limonia*, *Neolimnobia*, *Discobola*, and *Rhipidia*. Revista de Entomologia 21: 161–221.
- Alexander CP (1956) New or little-known Tipulidae (Diptera). XCIX. Oriental-Australasian species. Annals and Magazine of Natural History 8(12): 657–674. <https://doi.org/10.1080/00222935508655682>
- Alexander CP (1958) Undescribed species of Japanese Tipulidae (Diptera). Part I. Transactions of the Shikoku Entomological Society 6: 1–8.
- Alexander CP (1959) New or little-known Tipulidae (Diptera). CVI. Oriental-Australasian species. Annals and Magazine of Natural History 1(13): 657–676. <https://doi.org/10.1080/00222935808650994>
- Alexander CP (1964) New or little-known Tipulidae from eastern Asia (Diptera). LII. Philippine Journal of Science 92: 383–419.
- Alexander CP (1965) New or little-known Tipulidae from Madagascar (Diptera). Transactions of the American Entomological Society 91: 39–83.
- Alexander CP (1967) New or little-known Tipulidae from eastern Asia (Diptera). LIX. Philippine Journal of Science 95: 79–120.
- Alexander CP (1972) Diptera: Tipulidae. Insects of Micronesia 12: 733–863.
- Alexander CP, Byers GW (1981) Tipulidae. In: McAlpine JF, Peterson BV, Shewell GE, Teskey HJ, Vockeroth JR, Wood DM (Eds) Manual of Nearctic Diptera. Vol. I. Biosystematic Research Centre, Ottawa, 153–190.
- Brunetti E (1912) Diptera Nematocera (excluding Chironomidae and Culicidae). Fauna of British India, including Ceylon and Burma 1: 1–581. <https://doi.org/10.5962/bhl.title.8711>

- Edwards FW (1916) New and little-known Tipulidae, chiefly from Formosa. *Annals and Magazine of Natural History* 18(8): 245–269. <https://doi.org/10.1080/00222931608693846>
- Edwards FW (1926) On some crane-flies from French Indo-China. *Encyclopedie Entomologique*, (B II), Diptera 3: 48–55.
- McAlpine JF (1981) Morphology and terminology, Adults. In: McAlpine JF, Peterson BV, Shewell GE, Teskey HJ, Vockeroth JR, Wood DM (Eds) *Manual of Nearctic Diptera*. Vol. I. Biosystematic Research Centre, Ottawa, 9–63.
- Meijere JCH de (1916) Studien uber Sudostasiatische Dipteren, 11. Zur Biologie einiger javanischen Dipteren nebst Beschreibung einiger neuen javanischen Arten. *Tijdschrift voor Entomologie* 59: 184–213.
- Oosterbroek P (2021) Catalogue of the Craneflies of the World (Diptera, Tipuloidea: Pediciidae, Limoniidae, Cylindrotomidae, Tipulidae). <http://ccw.naturalis.nl/> [accessed 10 March 2021]
- Podenas S, Byun HW (2018) *Libnotes* crane flies (Diptera: Limoniidae) from Jeju Island (South Korea). *Zootaxa* 4483: 375–384. <https://doi.org/10.11646/zootaxa.4483.2.9>
- Ribeiro GC (2006) Homology of the gonostylus in crane flies, with emphasis on the families Tipulidae and Limoniidae (Diptera, Tipulomorpha). *Zootaxa* 1110: 47–57. <https://doi.org/10.11646/zootaxa.1110.1.5>
- Westwood JO (1876) *Notae Dipterologicae*. No. 2. – Descriptions of some new exotic species of Tipulidae. *Transactions of the Entomological Society of London* 1876: 501–506. <https://doi.org/10.1111/j.1365-2311.1876.tb01926.x>
- Wulp FM van der (1895) Eenige Javaansche Diptera. *Tijdschrift voor Entomologie* 38: 35–48.
- Young CW (1990) A new *Limonia* (subgenus *Laosa*) species from Sulawesi (Dipt., Tipulidae) and its resting behaviour. *Entomologists Monthly Magazine* 126: 239–244.