

# New species of subgenus *Tipula* (*Sivatipula*) from China, with redescription of *T. (S.) parvauricula* and a key to all known species of the Oriental Region (Diptera, Tipulidae, *Tipula*)

Guo-Xi Xue<sup>1</sup>, Qiu-Lei Men<sup>2</sup>

**1** School of Food and Bioengineering, Zhengzhou University of Light Industry, No. 5 Dongfeng Road, Zhengzhou, Henan 450002, P. R. China **2** School of Life Sciences, the Province Key Laboratory of the Biodiversity Study and Ecology Conservation in Southwest Anhui, Anqing Normal University, Anqing, Anhui 246011, P.R. China

Corresponding author: Qiu-Lei Men ([menqiulei888@126.com](mailto:menqiulei888@126.com))

---

Academic editor: J. Salmela | Received 12 November 2015 | Accepted 13 January 2016 | Published 15 February 2016

---

<http://zoobank.org/1789CC48-3763-49BB-B3AC-31BC78361E25>

---

**Citation:** Xue G-X, Men Q-L (2016) New species of subgenus *Tipula* (*Sivatipula*) from China, with redescription of *T. (S.) parvauricula* and a key to all known species of the Oriental Region (Diptera, Tipulidae, *Tipula*). ZooKeys 563: 33–42. doi: 10.3897/zookeys.563.7176

---

## Abstract

Species of *Tipula* (*Sivatipula*) *biprocessa* sp. n. from Guangxi, China is described and illustrated as new in the subgenus *Tipula* (*Sivatipula*) Alexander, 1964. *T. (S.) parvauricula* Alexander, 1941 is redescribed and illustrated based on additional morphological characters. Semen pump of this subgenus is discussed. A key to all described species in this group is compiled.

## Keywords

China, crane flies, new species, semen pump, *Sivatipula*, *Tipula*, Tipulidae

## Introduction

*Tipula* (*Sivatipula*) Alexander, 1964 is a small subgenus with *Tipula mitocera* Alexander, 1927 from the eastern Himalayas, India as its type species. The other species in this subgenus include *T. (S.) pullimargo* Alexander, 1951 from Myanmar, *T. (S.) alhena* Alexander, 1953 from Thailand, *T. (S.) filicornis* Brunetti, 1918 and *T. (S.) bhisma* Alexander, 1964 from India, *T. (S.) lackschewitziana* Alexander, 1928, *T. (S.) suenoniana* Alexander, 1940 and *T. (S.) parvauricula* Alexander, 1941 from China (Oosterbroek 2015). All these species are restricted to the Oriental region. The Chinese fauna of *Tipula* (*Sivatipula*) is poorly represented with only three known species.

This subgenus is characterized by the following characters: male with antennae very long, slightly shorter, equal to or longer than body length, female with antennae relatively short, not beyond half length of body, flagellomere covered with six or seven long strong verticils; wing with squama naked, outer wing veins scattered with small, abundant macrotrichia,  $R_{1+2}$  entire, Rs longer than m-cu; ninth tergite and sternite fused, median region of sternite extensive, more or less protrudent, forming amembranous extension; inner and outer gonostylus irregularly varied in shape. The species of subgenus *Tipula* (*Sivatipula*) had been placed previously in subgenus *Tipula* (*Acutipula*) Alexander, 1924, but treated subsequently as a distinct group based on the combined structural characters of antennae, hypopygium, and wing (Alexander 1964).

A previously unknown taxon of *Tipula* (*Sivatipula*) was noticed while sorting crane flies specimens collected from Leigongshan Mountain, Guizhou Province, and Cenwanglaoshan Mountain, Guangxi Zhuang Autonomous Region, China. In the present paper, the new species is described and illustrated, and a key is provided for separating all known species. *Tipula* (*S.*) *parvauricula* is redescribed based on newly available morphological characters with detailed illustrations. The original description of this species is insufficient and the illustrations are too simple to reveal necessary characters. The character of semen pump of subgenus *Tipula* (*Sivatipula*) is described for the first time. The current study also demonstrated the new distribution pattern for the subgenus *Tipula* (*Sivatipula*) in both Guangxi Zhuang Autonomous Region and Guizhou Province. Future collecting and investigation would undoubtedly increase the species numbers and range extension of this group in China.

## Material and methods

The genital segments of the specimens were soaked in 10% NaOH overnight and observed or drawn in glycerine using a Leica MZ125 (Leica, Germany) stereomicroscope. The genital segments were then preserved in glycerine in 0.20 ml centrifuge tubes. Photographs of partial body of male were taken by Canon 5D Mark II digital single lens reflex camera (Canon, Japan) with MP-E 65mm f/2.8 1-5X macro lens (Canon, Japan). All measurements are in millimeters (mm), made with the aid of a digital caliper. The terminology and methods of description and illustration follow those of Alex-

ander and Byers (1981) and Frommer (1963). The type specimens of the new species are deposited in the animal specimen room, School of Life Sciences, Anqing Normal University, Anqing, Anhui Province, China.

The key was principally constructed from descriptions in the literature without examination of the type species of most of these species, and should be considered preliminary. The characters used in the key rely primarily on the structures of genitalia and the length of antenna of male specimens.

## Taxonomy

### *Tipula* (*Sivatipula*) *parvauricula* Alexander, 1941

Figs 1–11

*Tipula parvauricula* Alexander, 1941: 400 (original description), Pl. 1, fig. 14, Pl. 4, fig. 44.


**Diagnosis.** Generally orange-yellow in coloration; antenna distinctly longer than body; prescutum orange-yellow with three light brown stripes, sometimes not clear; wings pale brown with a dark brown stigma; abdomen orange with segments six and seven black; hypopygium orange, tergite nine divided by a V-shaped notch, produced into a pair of ear-like processes, sometimes terminated into spinous point, ventral-lateral portions of tergite nine projected into two lobes, densely covered with black setae.

**Redescription.** Male. Length: *Body*: 14.0–18.0 mm (not including antenna,  $n = 5$ ); *Wing*: 18.0–20.0 mm ( $n = 5$ ); *Antenna*: 18.0–20.0 mm ( $n = 5$ ).

*Head* orange. Rostrum orange with distinct orange nasus. Marking of vertex absent (Fig. 1). Eyes black (Fig. 1). Antenna: 12-segmented, distinctly longer than body; scape orange, expanded apically; pedicel orange, short; flagellomeres dark brown, each flagellomere cylindrical and subequal in length, with abundant black verticils, longest one longer than one third of flagellomeres length. Palpi light brown.

*Thorax* with pronotum entirely orange-yellow (Figs 2–3). Prescutum with three light brown stripes, sometimes not clear (Fig. 2). Scutum orange-yellow with two light brown stripes (Fig. 2). Scutellum and postnotum orange-yellow (Fig. 2). Pleura entirely bright yellow (Fig. 3). Legs slender, coxae, trochanters and femora orange-yellow, tibiae orange-yellow at basal half, changing to brown at apical half, tarsi brownish black. Halteres with stem yellow, knob darker. Wings pale brown, cell sc darker than ground color, stigma dark brown (Fig. 4). Sc relatively short, subequal to  $R_3$  in length, petiole of cell m1 as long as or slightly longer than m-m, approximately one-half length of discal cell (Fig. 4).

*Abdomen* orange with segments six and seven black. Hypopygium orange, broad, compressed (Figs 5–6). Tergite nine separated by a V-shaped notch in ventral view; gradually narrowed to apex, produced into a pair of ear-like processes, sometimes terminated into spinous point in lateral view; ventral-lateral portions of tergite nine pro-



**Figures 1–11.** *T. (S.) parvaurecula* **1** head, dorsal view **2** thorax, dorsal view **3** thorax, right lateral view **4** wing **5** hypopygium, right lateral view **6** perspective hypopygium, left lateral view **7** ninth tergite, dorsal view **8** inner gonostylus, left lateral view **9** and **10** semen pump, left lateral view **11** compressor apodeme, dorsal view.

jected into two lobes, densely covered with blackened setae (Figs 5–7). Sternite nine broader than tergite nine, median region of sternite nine protruded to a membranous extension (Fig. 6). Outer gonostylus flattened, widened medially, with a slender rod

on ventral-lateral margin, directed dorsally (Fig. 6). Inner gonostylus broad basally, gradually narrowed to apex, rounded with a process truncated apically, ventral margin with a horn-shaped projection, dorsal margin with a finger-like process, basal region of inner gonostylus with a pyramidal process (Fig. 8).

*Semen pump* with compressor apodeme V-shaped, the arms expanded at apex (Fig. 11). Posterior immovable apodeme with only one arm, distinctly longer than compressor apodeme, gradually narrowed to apex and curved cephalad in lateral view (Fig. 9), sometimes abruptly bent to ventral margin (Fig. 10), the arm deeply grooved in dorsal view. Anterior immovable apodeme flattened, shorter than compressor apodeme, gradually narrowed to apex in lateral view (Figs 9–10). Aedeagus elongated, tubular, at least ten times longer than semen pump (Fig. 6).

**Material examined.** CHINA: Guangxi Zhuang Autonomous Region: 2 males, Dalongping, Cenwanglaoshan Mountain, 24°31'N, 106°17'E, 1300 m, 11 May 2015, Guo-Xi Xue leg.; Guizhou Province: 3 males, Leigongshan Mountain, 26°21'N, 108°13'E, 4 Jun. 2015, Guo-Xi Xue leg.

**Distribution.** China (Fujian, NW. Guangxi, SE. Guizhou).

**Remarks.** In the original description of *T. (S.) parvaauricula* (Alexander 1941), the prescutum is unmarked and the tergite nine is terminated into a pair of spinous points in lateral view. After observing five specimens, we noticed that the prescutum generally has three light brown stripes and the spinous point on tergite nine is not always present.


Three types of semen pumps were defined by Frommer (1963) based on morphological studies of the reproductive system of North American crane flies. Type III is the most common type characterized by the strongly bowed intromittent organ and by posterior immovable apodeme generally with two arms (Frommer 1963). According to the overall morphology, the semen pump of *T. (S.) parvaauricula* should belong to Type III. However, its posterior immovable apodeme has only one arm, which differs from the results in previous works in Chinese species (*Ctenophora fumosa* Men, 2014 in Men and Huang 2014; *Tipula (Vestiplex) coxitalis* Alexander, 1935, *Tipula (Pterelachisus) biaciculifera* Alexander, 1937 and *Tipula (Emodotipula) yaoluopingensis* Men, 2015 in Men 2015; *Tipula (Yamatotipula) nova* Walker, 1848 in Men et al. 2015a; *Nephrotoma liankangensis* Men, Xue & Yang and *N. pseudoliankangensis* Men, Xue & Yang in Men et al. 2015b). This may represent a special form of type III.

***Tipula (Sivatipula) biprocessa* sp. n.**

<http://zoobank.org/0121BA77-9947-486A-91B8-15EAABEEFAB8>

Figs 12–23

**Diagnosis.** Generally straw-yellow; prescutum straw-yellow with three light brown stripes; wings pale brown with a dark brown stigma; abdomen bright yellow on basal three segments, gradually changed to light brown on apical ones, segments six and seven suffused with black; hypopygium straw-yellow, tergite nine rounded at posterior



**Figures 12–23.** *T. (S.) biprocessa* sp. n. **12** head, dorsal view **13** thorax, right lateral view **14** thorax, dorsal view **15** wing **16** hypopygium, right lateral view **17** hypopygium, left lateral view **18** ninth tergite, dorsal view **19** inner gonostylus and outer gonostylus, left view **20** inner gonostylus and outer gonostylus, right view **21** inner gonostylus and outer gonostylus, dorsal view **22** semen pump, dorsal view **23** semen pump, right lateral view.

margin and equipped with two finger-like processes, ventral-lateral portions of tergite nine projected into two lobes, densely covered with black setae.

**Description.** Male. Length: *Body*: 14.0–15.0 mm (not including antenna,  $n = 2$ ); *Wing*: 19.0–20.0 mm ( $n = 2$ ); *Antenna*: 18.0–19.0 mm ( $n = 2$ ).

*Head* straw-yellow (Figs 12–13). Rostrum light brown with a light brown nasus. Vertex without marking (Fig. 12). Eyes black (Fig. 12). Antenna: 12-segmented, distinctly longer than body; scape light yellow, expanded apically; pedicel light yellow, short; flagellomeres light brown, each flagellomere cylindrical, subequal in length, with abundant black verticils. Palpi light brown.


*Thorax* with pronotum entirely orange-yellow (Figs 13–14). Prescutum straw-yellow with three light brown stripes (Fig. 14). Scutum orange-yellow with two light brown stripes (Fig. 14). Scutellum and postnotum orange-yellow (Fig. 14). Pleura entirely bright yellow (Fig. 13). Legs slender, coxae and trochanters straw-yellow, femora straw-yellow with light brown tip, tibiae and tarsi light brown. Halteres with stem yellow, knob darker. Wings pale brown, cell sc darker than ground color, stigma dark brown (Fig. 15). Sc relatively short, subequal to  $R_3$  in length, petiole of cell m1 slightly longer than m-m, approximately one-half length of discal cell (Fig. 15).

*Abdomen* bright yellow on basal three segments, gradually changed to light brown on apical ones, segments six and seven suffused with black (Fig. 16). Hypopygium straw-yellow (Fig. 16). Hypopygium broad, compressed (Figs 16–17). Tergite nine rounded at posterior margin with two finger-like processes, lateral sides of tergite nine with numerous long hairy setae, longest one longer than finger-like process (Fig. 18). Sternite nine broader than tergite nine, median region of sternite nine protruded to a membranous extension (Fig. 17). Outer gonostylus narrow, flattened, apical two-fifths curved caudad (Figs 19–21). Inner gonostylus flattened, narrowed medially, a slender lobe generated from the median region of inner gonostylus (Figs 19–21).

*Semen pump* with compressor apodeme V-shaped, the arms expanded at apex, distinctly broader than that of *T. (S.) parvauricula* (Fig. 22). Posterior immovable apodeme with one arm, distinctly longer than compressor apodeme, gradually narrowed to apex and curved cephalad in lateral view, the arm deeply grooved in dorsal view, basal region distinctly wider than that of *T. (S.) parvauricula* (Figs 22–23). Anterior immovable apodeme flattened, gradually narrowed to apex in lateral view (Fig. 22). Aedeagus elongated, tubular, at least ten times longer than semen pump.

**Material examined.** **Holotype** male. **CHINA:** Guangxi Zhuang Autonomous Region, Dalongping, Cenwanglaoshan Mountain, 24°31'N, 106°17'E, 1300 m, 7 May 2015, Guo-Xi Xue leg. **Paratype.** 1 male, same data as holotype.

**Remarks.** We compared the new species with all known species based on published descriptions and illustrations, and found that it is mostly similar to *T. (S.) parvauricula* by the color of body, the structures of antenna and hypopygium. It can be easily distinguished from the latter by the shape of tergite nine which is rounded at posterior margin with two distinct finger-like processes. The latter species has its tergite nine separated by a V-shaped notch with two short truncated processes and produced into a pair of ear-like processed ventrally. There is also a noticeable difference in the



**Figure 24.** Geographic distribution of *Tipula (Sivatipula)* species: *T. (S.) albena* (●), *T. (S.) bhishma* (▲), *T. (S.) filicornis* (◆), *T. (S.) lackschewitziana* (■), *T. (S.) mitocera* (△), *T. (S.) pullimargo* (○), *T. (S.) suensonia* (◇), *T. (S.) parvauricula* (□), *T. (S.) biprocessa* sp. n. (★).

shape of the outer gonostylus which is simple and narrowed in the new species, but flattened and widened medially with a slender rod on ventral-lateral margin in that of *T. (S.) parvauricula*. Distinct interspecific difference is also found in the shape of inner gonostylus as Figures 8, 19 and 20.

**Etymology.** The specific epithet is a noun derived from the Latin ‘processa’ with Latin prefix ‘bi’, referring to the presence of two finger-shaped processes at posterior margin of tergite nine.

**Distribution.** China (NW. Guangxi).

**Key to species of subgenus *Tipula (Sivatipula)***

- 1 Hind margin of ninth tergite rounded apically with a pair of finger-like processes (see Figs 16–18) ..... ***T. (S.) biprocessa* sp. n.** (China: Guangxi, Cenwanglaoshan Mountain. Fig. 24)
- Hind margin of ninth tergite without such process ..... **2**
- 2 Outer gonostylus with an appressed pubescence on outer surface of apical arm (see Alexander 1928: 459, Pl. 2, fig. 5) ..... ***T. (S.) lackschewitziana* Alexander, 1928** (China: Taiwan, Noko Mountain. Fig. 24)
- Outer gonostylus without such pubescence ..... **3**
- 3 Ninth tergite laterally bearing two spinous projections, lower to them submedially with two short spiculate projections (see Joseph, 1974: 277, figs 137–142) ..... ***T. (S.) filicornis* Brunetti, 1918** (India: West Bengal, Dajeeling. Fig. 24)
- Ninth tergite not as above ..... **4**
- 4 Antenna slightly shorter than body ..... **5**

- Antenna equal to or longer than body..... **6**
- 5 Outer gonostylus club-shaped (see Alexander, 1953: 348, fig. 12d) .....  
***T. (S.) pullimargo* Alexander, 1951** (Myanmar: Adung Valley. Fig. 24)
- Outer gonostylus long-attenuate (see Alexander 1940: 110, fig. 11) .....  
 ..... ***T. (S.) suenso-***  
***niana* Alexander, 1940** (China: Zhejiang, Tianmushan Mountain. Fig. 24)
- 6 Ninth tergite produced into a rounded apex, with numerous black setae  
 (see Alexander 1971: 81, fig. 6) ..... ***T. (S.)***  
***albena* Alexander, 1953** (Thailand: Chiangmai, Dio Suthep. Fig. 24)
- Ninth tergite not produced into a rounded apex..... **7**
- 7 Ninth tergite with powerful lateral arms that are tipped with abundant black-  
 ened pegs (see Alexander 1964: 105, Pl. 4, fig. 40) .....  
 ..... ***T. (S.) bhisma* Alexander, 1964** (India: Assam, Manipur. Fig. 24)
- Ninth tergite without such pegs..... **8**
- 8 Median lobe of ninth sternite bearing terminal brush-like setae (see Alexander  
 1927: 182, fig. 3).....  
 ... ***T. (S.) mitocera* Alexander, 1927** (India: West Bengal, Dajeeling. Fig. 24)
- Median lobe of ninth sternite without such brush-like setae (see Alexander 1941:  
 401, Pl. 4, fig. 44; Fig. 6) ..... ***T. (S.)***  
***parvauricula* Alexander, 1941** (China: Fujian, Wuyishan Mountain; Guangxi,  
 Cenwanglaoshan Mountain; Guizhou, Leigongshan Mountain. Fig. 24)

## Acknowledgments

We wish to express sincere thanks to Dr. Pjotr Oosterbroek, University of Amsterdam, Amsterdam, the Netherlands, for his help with literature and for his valuable web site, the Catalogue of the Craneflies of the World (<http://ccw.naturalis.nl/index.php>), which provides much valuable information about distribution and taxonomy. We also thank Dr. Yan Li, Shenyang Agricultural University, Shenyang, for his help with identification of known species. This study is supported by grants from the National Natural Science Foundation of China (No. 41401067, No. 31300551), the start-up grant of scientific research from Anqing Normal University (No. 044-K05000130005).

## References

- Alexander CP (1927) The Oriental Tipulidae in the collection of the Indian museum. Part I. Records of the Indian Museum 29: 167–214.
- Alexander CP (1928) New or little-known Tipulidae from eastern Asia (Diptera). III. Philippine Journal of Science 36: 455–485.
- Alexander CP (1940) Studies on the Tipulidae of China (Diptera). III. New or little-known crane-flies from Tien-mu-shan, Chekiang. Lingnan Science Journal 19: 105–119.

- Alexander CP (1941) New or little-known Tipulidae from eastern Asia (Diptera). XLIII. Philippine Journal of Science 73: 375–420.
- Alexander CP (1953) The Oriental Tipulidae in the collection of the Indian museum. Part III. Records of the Indian Museum 50: 321–357.
- Alexander CP (1964) New or little-known Tipulidae from eastern Asia (Diptera). LIII. Philippine Journal of Science 93: 77–130.
- Alexander CP (1971) New and little-known Indian craneflies (Diptera: Tipulidae). IV. Oriental Insects 5: 73–82. doi: 10.1080/00305316.1971.10433991
- Alexander CP, Byers GW (1981) Tipulidae. In: McAlpine JF, Peterson BV, Shewell GE, Teskey HJ, Vockeroth JR, Wood DM (Eds) Manual of Nearctic Diptera. Vol. 1. Biosystematics Research Institute, Ottawa, Ontario, 153–190.
- Frommer SI (1963) Gross morphological studies of the reproductive system in representative North American crane flies (Diptera: Tipulidae). Kansas University Science Bulletin 44: 535–625.
- Joseph ANT (1974) The Brunetti types of Tipulidae (Diptera) in the collection of the Zoological Survey of India. Part III. *Tipula* Linnaeus. Oriental Insects 8: 241–280. doi: 10.1080/00305316.1974.10434860
- Men QL (2015) Report on crane flies of the genus *Tipula* (Diptera: Tipulidae: Tipulinae) from Anhui Province, China. Acta Entomologica Musei Nationalis Pragae 55(2): 797–810.
- Men QL, Huang MY (2014) A new species of the genus *Ctenophora* Meigen (Diptera: Tipuloidea: Tipulidae) from China, with a key to the world species. Zootaxa 3841(1): 592–600. doi: 10.11646/zootaxa.3841.4.8
- Men QL, Xue GX, Liu Y (2015a) A morphological study on reproductive system of *Tipula* (*Yamatotipula*) *nova* Walker (Diptera: Tipulidae). Zoological Systematics 40(3): 328–338.
- Men QL, Xue GX, Yang H (2015b) Two new species of the genus *Nephrotoma* (Diptera, Tipuloidea, Tipulidae) from China with a key to species from Mainland China. ZooKeys 532: 117–136. doi: 10.3897/zookeys.532.5970
- Oosterbroek P (2015) Catalogue of the Craneflies of the World, (Diptera, Tipuloidea: Pediciidae, Limoniidae, Cyldrotomidae, Tipulidae). <http://ccw.naturalis.nl/index.php> [accessed 24 September 2015]