

# Revision of the legume-feeding leaf beetle genus *Madurasia* Jacoby, including a new species description (Coleoptera, Chrysomelidae, Galerucinae, Galerucini)

Kaniyarikkal Divakaran Prathapan<sup>1</sup>

<sup>1</sup> Department of Entomology, Kerala Agricultural University, Vellayani P.O., Trivandrum–695 522, Kerala, India

Corresponding author: Kaniyarikkal Divakaran Prathapan ([prathapankd@gmail.com](mailto:prathapankd@gmail.com))

---

Academic editor: M. Schmitt | Received 15 December 2015 | Accepted 21 April 2016 | Published 9 June 2016

<http://zoobank.org/FF526EC2-E8AB-45A8-BDC8-C5878139D858>

---

**Citation:** Prathapan KD (2016) Revision of the legume-feeding leaf beetle genus *Madurasia* Jacoby, including a new species description (Coleoptera, Chrysomelidae, Galerucinae, Galerucini). In: Jolivet P, Santiago-Blay J, Schmitt M (Eds) Research on Chrysomelidae 6. ZooKeys 597: 57–79. doi: 10.3897/zookeys.597.7520

---

## Abstract

*Madurasia* Jacoby is revised and *M. andamanica* sp. n., endemic to the Andaman Islands in the Indian Ocean, is described and illustrated. *Madurasia obscurella* Jacoby, syn. n., is a new junior synonym of *Madurasia undulatovittata* (Motschulsky), comb. n. A lectotype is designated for *M. obscurella*. Literature on the biology and management of *M. undulatovittata* is reviewed.

## Keywords

Asia, Africa, biology, pest, pulses, taxonomy

## Introduction

The monotypic galerucine genus *Madurasia* was described by Jacoby (1886) for a new species, *Madurasia obscurella*, from southern India. Aslam (1972) synonymized *Neorudolphia bedfordi* Laboissière, 1926, the only species in this monotypic genus from Sudan, with *M. obscurella* Jacoby. Examination of the type of *Monolepta undulatovittata* (Motschulsky 1866) (originally described in *Teinodactyla* Chevrolat = *Longitarsus* Latreille) from Sri Lanka has shown that *Madurasia obscurella* is a junior synonym of Motschulsky's species. The genus is here revised and a new species is described from

the Andaman Islands in the Indian Ocean. Information on the biology, pest status and management of *M. undulatovittata* comb. n., which is a significant pest of various legume crops in south-east Asia and Africa, is reviewed.

## Materials and methods

Dissecting techniques and descriptive terminology follow Konstantinov (1998). Label data for holotypes, lectotypes, and paralectotypes has been recorded verbatim, with lines on the same label separated by “/” and labels separated by “;”. Material examined is from the following collections:

<b>BMNH</b>	Natural History Museum, London
<b>INPC</b>	National Pusa Collection, Indian Agricultural Research Institute, New Delhi
<b>JBC</b>	Personal collection of Jan Bezděk, Czech Republic
<b>KAU</b>	Travancore Insect Collection, Kerala Agricultural University, Vellayani
<b>NBAIR</b>	National Bureau of Agricultural Insect Resources, Bangalore
<b>UASB</b>	University of Agricultural Sciences, Bengaluru
<b>USNM</b>	National Museum of Natural History, Smithsonian Institution, Washington D.C.
<b>ZMUH</b>	Zoologisches Institut und Zoologisches Museum, Universität von Hamburg, Hamburg, Germany
<b>ZMUM</b>	Zoological Museum, Moscow State University, Moscow

Determination of the gender of the undissected specimens is provisional as sexually dimorphic characteristics are often not clearly discernible externally.

## Systematics

### *Madurasia* Jacoby, 1886

*Madurasia* Jacoby, 1886: 280 (Type species: *Madurasia obscurella* Jacoby, 1886, southern India, by monotypy)—Maulik 1936: 72—Wilcox 1973: 435—Seeno and Wilcox 1982: 107—Jolivet and Hawkeswood 1995: 101 (host plants)—Medvedev and Sprecher—Uebersax 2005: 316 (key)—Beenen 2010: 481.

*Neorudolphia* Laboissière, 1926: 190 (Type species: *Neorudolphia bedfordi* Laboissière, 1926, Sudan, by monotypy)—Wilcox 1973: 435—Aslam 1972: 500 (= *Madurasia* Jacoby 1886: 280).

**Description.** Body: length 2.0–3.0 mm; 1.8–2.3 times longer than wide. Moderately small, oblong, flattened in lateral view, length 3.1–3.4 times height. General color

straw brown to dark brown with a characteristic, more or less distinct, dark, broad longitudinal stripe on each elytron (Figs 1, 3, 5–7, 21); mesal margin of stripe nearly straight; each stripe nearer to suture than to lateral margin of elytron; stripe narrowing laterally posteriorly of humerus and in distal 2/3 of elytron.

Head (Fig. 8) hypognathous with frontal view slightly longer than wide. In lateral view anterior margin forms a moderately convex line with a notch where vertex meets antennal calli and a second notch at anterior end of frontal ridge. Supraorbital pore represented by a large setaceous pore adjacent to orbital sulcus near eye. Seta in supraorbital pore upcurved. Vertex shiny, indistinctly wrinkled, nearly impunctate. Antennal calli trapezoidal, longer than wide, moderately convex, raised above adjacent border of vertex, separated from each other by a deep midfrontal sulcus; anterior ends acutely triangular, enter into interantennal space, reaching well below midlevel of antennal socket. Orbital sulcus short, deep, represented by supraorbital pore and adjacent area. Supracallinal sulcus represented by punctures arranged in an irregular transverse row, each puncture in supracallinal row bearing a short down-curved seta. Midcranial suture absent. Supraorbital sulcus less distinct than midfrontal sulcus. Suprafrontal and supraantennal sulci well defined. Subgenal suture distinct. Transverse diameter of eye 5.2–8.8 times distance between eye and antennal socket, 2.9–4.4 times distance between antennal sockets, 1.7–1.9 times width of antennal socket, 0.6–0.7 times distance between eyes. Eyes lateral, medium sized, convex, inner margins indistinctly concave, and ventrally divergent. Frontal ridge narrowest between antennal sockets, joins anterofrontal ridge anteriorly. Anterofrontal ridge transverse, gently curved. Frontal ridge together with antero-frontal ridge forms T-shaped ridge. Anterofrontal ridge lower than frontal ridge. Frontolateral area coarsely punctate, each puncture bearing a long seta. Frontoclypeal suture with a row of eight setae. Clypeus narrow. Visible part of labrum much wider than long, with a transverse row of eight pores; all eight pores in *M. andamanica* sp. n. with a well-developed seta; while only six pores, excluding third pore from either end, with seta in *M. undulato vittata*. Labrum (Fig. 13) with anterior margin incised medially; about seven sensillae on either side of incision, arranged along anterior margin of labrum's inner surface; tormae longer than width of labrum. Mandible (Fig. 12) palmate with six sharp denticles. Maxilla (Fig. 11) with four palpomeres: first shortest, second and third subequal, both longer than first, but shorter than apical palpomere, apical longest; lacinia wider than galea. Labial palpi (Fig. 10) with three palpomeres, basal two wider than long, middle widest, apical palpomere longer than wide and longest of three. Antenna (Fig. 9) reaches more or less middle of elytron. First antennomere longest, club shaped; second smallest; third a little longer than second; fourth distinctly longer than third; 4–10 subequal in length; eleventh longer than all except first antennomere; five to six distal antennomeres wider than preceding three or four (Fig. 9).

Dorsum glabrous. Pronotum (Fig. 14) 1.2–1.3 times wider than long; greatest width slightly anterior of middle. Posterior margin 1.1–1.2 times wider than anterior margin, lateral margin gently convex, posterior margin nearly straight in middle, curved laterally, and narrowly margined. Anterolateral callosity longer than wide, setigerous pore posterolaterally situated, not forming denticle at pore; posterolateral

callosity protruding slightly laterally, setigerous pore laterally situated. Disc without impressions, shiny, uniformly punctate, punctures small, smaller than those on elytra. Anterior coxal cavity open behind (Figs 4, 16); intercoxal prosternal process short, acutely pointed, not reaching midlevel of procoxa (Fig. 16); procoxae longer than wide and closely associated; shortest distance from anterior margin of prosternum to procoxal cavity about 1/4–1/5 of longitudinal procoxal diameter. Mesoscutellum triangular, about two times wider than long, flat, impunctate to minutely punctate. Intercoxal mesosternal process short, not reaching midlevel of mesocoxa (Figs 4, 17). Mesepisternum broader than mesepimeron (Fig. 17). Metasternum no longer than first two abdominal ventrites combined (Fig. 4).

Elytra broader than pronotum basally, maximum width posterior of middle. Humeral callus well developed; elytral border narrow, becoming indistinct towards apex; elytral apex broadly rounded; epipleuron (Fig. 4) oblique, maximum width near anterior 1/4 of elytron, maximum width subequal to about 1.5 times maximum width of mid-femur, narrows abruptly before middle and then continues very narrowly, becoming indistinct towards the elytral apex. Hind wings present. Metanotum (Fig. 15) well developed with full complement of internal ridges.

All femora oblong in cross section; all tibiae subcylindrical, subcircular in cross section with a minute apical spur; metatibial spur subequal to claw in length; proportionate length of femur–tibia–tarsomeres 1–4 as follows: 1: 1.0–1.1 : 0.2–0.3 : 0.1–0.2 : 0.1–0.2 : 0.2–0.3 (foreleg); 1: 0.9–1.0 : 0.3 : 0.1–0.3 : 0.1–0.2 : 0.2–0.3 (midleg); 1: 1.1–1.2 : 0.4 : 0.1–0.2 : 0.1 : 0.2 (hindleg); joint where metatibia and first metatarsomere meet, black; third tarsomere always bilobed; claws simple and appendiculate, appendix small and basal. Abdomen (Fig. 4) with five distinct ventrites; ventrites 2–4 becoming progressively slightly shorter; fifth ventrite slightly longer than fourth; intercoxal projection of first abdominal ventrite acute; apical abdominal tergite (Fig. 19, 20) without a median longitudinal groove, posterior margin slightly concave medially in male of *M. undulato vittata* (Fig. 20) and distinctly emarginate in *M. andamanica* sp. n.; posterior margin of apical tergite broadly convex (Fig. 19) in females of both species; posterior margin of apical ventrite more or less lobed medially in male (Figs 22, 23), entire in female.

Female genitalia with receptacle of spermatheca (Figs 28, 29) pot-shaped, wider than long; pump curved, longer than receptacle and enlarged distally, appendix well developed; spermathecal duct shorter than receptacle, glandular duct beyond middle of spermathecal duct. Tignum (Figs 32, 33) gently curved near middle, grooved medially, with long setae near distal margin of broad membranous apex. Vaginal palpi (Figs 30, 31) fused from proximal end to a short distance beyond middle, separate distally, each palpus narrowing towards rounded apex, lateral margin concave preapically, with long distal setae. Median lobe of aedeagus strongly curved in lateral view (Figs 26, 27), acutely pointed. Tegmen with stem much longer than arms.

**Host plants.** Fabaceae.

**Distribution.** Asia, Africa (Sudan).

**Remarks.** *Madurasia* closely resembles *Medythia* Jacoby, 1887, and species of both genera are pests of legumes. The general morphology, including the structure of the

head, female genitalia, and even the presence of elytral stripes in some species of *Medythia*, are similar to those in *Madurasia*, making differentiation of these genera difficult. *Madurasia* can be separated from *Medythia* by the structure of the pronotum. The pronotum in *Medythia* is elongate and narrows posteriorly, whereas the pronotum is transverse and a little wider posteriorly in *Madurasia*. The elytral epipleuron is short in *Madurasia*, hardly extending beyond middle of the elytron. In *Medythia quadrimaculata* Jacoby, type species of the genus, the elytral epipleuron is longer, extending beyond the middle of the elytron. However, the epipleura are identical to those of *Madurasia* in a few Indian *Medythia* species examined. In *Madurasia*, the distal antennomeres are darker, while antennomeres 8–10 are whitish in most *Medythia* species, including the type species.

Adults are attracted to light.

***Madurasia andamanica* sp. n.**

<http://zoobank.org/3D810CFF-3113-43FE-8E67-BD8335505E90>

Figs 21, 22, 24, 26, 28, 30, 32, 33


**Diagnosis.** The new species can be recognized by the following characters: 1) elytral stripe not reaching the elytral apex, narrowing in distal 1/4; 2) labrum with a transverse row of eight well developed setae; 3) posterior margin of apical ventrite in male distinctly lobed medially; 4) apex of aedeagus in lateral view curved like a parrot's beak with an acute tip; 5) ventral side of aedeagus depressed in basal 1/2, then distally raised in the form of a narrow ridge which reaches the apex.

**Description.** Body: length 2.1–2.6 mm; width 1.1–1.2 mm; 1.8–2.1 times longer than wide (Fig. 21). Dorsum straw colored. Head dark brown. Antenna with basal three or four antennomeres a pale straw color, distal antennomeres becoming progressively darker. Mandible, maxilla, and labium paler than labrum and anterior aspect. Pronotum with pale orange hue. Elytra a pale straw color, the dark elytral stripe not reaching the elytral apex (Fig. 21), widest anteriorly of middle, narrowing distinctly posteriorly of humerus as well as in distal 1/4. Thoracic sternites and pronotum concolorous, metathoracic sternite often a slightly darker laterally. Abdominal ventrites pale brown, with lateral margins and apical abdominal ventrite darker in many specimens. Legs straw colored, tibia and first two tarsomeres often a slightly darker than femur. Antenna reaching slightly beyond middle of elytron. Proportionate length of antennomeres 1–11: 1: 0.48: 0.45–0.50: 0.63–0.67: 0.63: 0.61–0.62: 0.66–0.67: 0.62–0.70: 0.63–0.69: 0.62–0.63: 0.75–0.88. Transverse diameter of eye 6.3–8.0 times width of orbit, 3.3–3.8 times width of interantennal space, 1.7–1.9 times width of antennal socket, 0.7 times distance between eyes. Pronotum 1.2–1.3 times wider than long, posterior width 1.1–1.2 times wider than anterior width.

Proportionate length of femur:tibia:tarsomeres 1–4 as follows: 1: 1.0–1.1 : 0.2–0.3 : 0.2 : 0.1–0.2 : 0.2–0.3 (foreleg); 1: 0.9–1.0 : 0.3 : 0.1–0.3 : 0.1–0.2 : 0.2–0.3 (midleg); 1: 1.0–1.1 : 0.4 : 0.1–0.2 : 0.1 : 0.2 (hindleg).



1



2



3



4



5




6




7


**Figures 1–7.** *Madurasia undulatovittata*. 1 Lectotype (specimen on card, photograph edited) 2 labels on lectotype 3 and 5–7 dorsal view, color variation 4 ventral view.



**Figures 8–20.** *Madurasia undulatovittata*. **8** head, frontal view **9** antenna **10** labium **11** maxilla **12** mandible **13** labrum **14** pronotum **15** meso- and metanotum **16** prosternum **17** meso- and metasternum and pleurites **18** metendosternite **19** apical visible tergite, female **20** apical visible tergite, male (all specimens, except head, have been macerated).



**Figures 21–27.** *Madurasia andamanica* sp. n. **21** dorsal habitus **22** apical ventrite of *M. andamanica* sp. n. male **23** apical ventrite of *M. undulatovittata* male **24** median lobe of aedeagus in *M. andamanica* sp. n., ventral view **25** median lobe of aedeagus in *M. undulatovittata*, ventral view (bilaterally symmetrical, specimen tilted) **26** median lobe of aedeagus in *M. andamanica* sp. n., lateral view **27** median lobe of aedeagus in *M. undulatovittata*, lateral view.



**Figures 28–33.** Spermatheca in **28** *M. andamanica* sp. n. **29** *M. undulatovittata*; vaginal palpi of **30** *M. andamanica* sp. n. **31** *M. undulatovittata*; tigna in **32** *M. andamanica* sp. n. **33** *M. undulatovittata*.

Posterior margin of apical ventrite in male distinctly lobed medially (Fig. 22). Receptacle of spermatheca 2.4 times wider than long (Fig. 28). Tignum widened proximally; membranous apex wider towards posterior (Fig. 32).


Aedeagus in lateral view (Fig. 26) with greatest width near middle, narrow in proximal 1/4, apex curved like a parrot's beak with an acute tip. In ventral view (Fig. 24), greatest width at base, narrowing abruptly in apical 1/3; ventral aspect depressed in basal 1/2, then distinctly raised in the form of a narrow ridge which reaches the apex.

**Etymology.** Named after the Andaman Islands, where the new species occurs.

**Material examined. Holotype** ♂ “INDIA: Andaman & Nicobar / North Andaman: Diglipur / 13°14'53.9"N, 92°58'37.5"E, / 15 mts. 24.iv.2014. At light / Yeshwanth H. M.” (white label); “HOLOTYPE / *Madurasia andamanica* / Prathapan sp. nov., 2015” (red label) (BMNH).

Paratypes (104). 5♂, 8♀ same data as holotype; 7♂, 19♀ same data as holotype, but 23.iv.2014; 2♀ INDIA: Andaman & Nicobar / South Andaman: Sippighat / 11°67'26"N, 92°67'12"E, / 44 mts. 18.iv.2014, Light trap / Yeshwanth H. M.; 1♂, 16♀ India: South Andaman / Garacharama / 12.xi.2014 / Bharathimeena Coll. / Ex Redgram; 16♂, 7♀, 1 unsexed same data but 8.I.2015; 2♂, 20♀ same data but 4.XII.2014 and collector Krishnaveni (5 BMNH, 5 USNM, 5 JBC, 5 KAU, 5 UASB, 40 NBAIR, 39 INPC).

**Distribution.** India (Andaman Islands) (Fig. 34).



**Figure 34.** Distribution of *M. andamanica* sp. n. on the Andaman Islands.

**Remarks.** Color pattern in *M. andamanica* sp. n. (Fig. 21) appears to be consistent and less variable compared to that in *M. undulatovittata* (Motschulsky, 1866), where the color of specimens collected on the same host at the same locality on the same day

varied greatly. *Madurasia andamanica* sp. n. resembles *M. undulatovittata* externally. However, it can be distinguished based on the structure of the aedeagus and the number of labral setae, as described under *M. undulatovittata*.

**Host.** *Cajanus cajan* (L.) Millsp. (Fabaceae) (red gram or pigeon pea) (Bharathi-meena T., pers. comm. 2015).

***Madurasia undulatovittata* (Motschulsky), comb. n.**

Figs 1, 3–20, 23, 25, 27, 29, 31, 33, 35

*Teinodactyla undulatovittata* Motschulsky, 1866: 417 [Sri Lanka, Lectotype (ZMUM)]–Wagner and Bieneck 2012: 214–215.

*Longitarsus undulatovittatus*: Gemminger and Harold 1876: 3509–Maulik 1926: 361.

*Monolepta undulatovittata*: Ogloblin 1930: 112.

*Madurasia obscurella* Jacoby, 1886: 381 [“Madura, Madras Presidency”, Southern India– Lectotype (BMNH)]–Maulik 1936: 74–Wilcox 1973: 435–Takizawa 1987: 39–Takizawa and Kimoto 1990: 8–Takizawa 1990: 281–Mohamedsaid 1997: 5–Medvedev and Sprecher 1999: 310 (catalogue)–Mohamedsaid 2000: 370–Kimoto 2005: 58–Beenen 2010: 481–Bezděk 2012: 422, 424. **New synonym.**

*Neorudolphia bedfordi* Laboissière, 1926: 191 [Brit. Sudan, on *Cajanus indicus*, Syntype (ZMUH, Hamburg)]–Aslam 1972: 500 (synonymized with *Madurasia obscurella* Jacoby)–Wilcox 1973: 435–Weidner 1976: 229.

**Description.** Body: length 2.0–3.0 mm; width 1.0–1.3 mm; 2.0–2.3 times longer than wide. General color pattern consistent but highly variable in intensity (Figs 1, 3, 5–7). Head dark brown to pale brown, often darker than pronotum. Basal antennomeres 3–6 pale straw brown, distal antennomeres becoming progressively darker. Pronotum more or less pale brown, generally paler than head. Background color of elytron paler than pronotum. Lateral margin of dark elytral stripe emarginate in anterior 1/3 and posterior 1/3; stripe broadening posteriorly, covering width of elytral apex. In some specimens, elytra darker laterally giving the impression of a pale, medially narrowed line on a dark elytron. Intensity of stripe’s darkness varies from pale straw brown (Fig. 1) to dark brown. In type of *M. undulatovittata*, elytral stripes are hardly visible (Fig. 1). In some specimens, widest region in middle of stripe extends to lateral elytral margin, thus dividing pale colored lateral area into anterior and posterior spots (Figs 6, 7). Ventral aspect (Fig. 4) generally paler than head. Metasternum slightly darker than pro- or mesosternum. Metepisternum darker than metasternum. Abdomen darker laterally and posteriorly in many specimens. In darkest specimens, ventral side dark brown to piceous. Legs pale brown, all femora nearly concolorous with abdominal ventrites; metafemora darker distally in some specimens. All tibiae paler than femora. Metatibia and first metatarsomere whitish in some specimens. Claw tarsomere and bilobed tarsomere often darker than preceding ones.

Antenna (Fig. 9) reaches middle of elytron or a little beyond. Proportionate length of antennomeres 1–11: 1: 0.54–0.57 : 0.44–52 : 0.65–0.69 : 0.59: 0.66: 0.66–0.69: 0.62–0.75: 0.73–0.75: 0.69–0.72: 0.69–0.71: 0.81–0.91. Transverse diameter of eye 5.3–8.8 times width of orbit, 2.9–4.4 times width of interantennal space, 1.7–1.8 times width of antennal socket, 0.6 times distance between eyes. Pronotum (Fig. 14) 1.3 times wider than long, posterior 1.1 times wider than anterior.

Proportionate length of femur–tibia–tarsomeres 1–4 as follows: 1: 1.0–1.1 : 0.3 : 0.1–0.2 : 0.1–0.2 : 0.3 (foreleg); 1: 0.9–1.0 : 0.3 : 0.1–0.2 : 0.1–0.2 : 0.2–0.3 (midleg); 1: 1.1–1.2 : 0.4 : 0.1–0.2 : 0.1 : 0.2 (hindleg). Two visible apical tergites completely exposed in most females, particularly when killed in alcohol.

Posterior margin of apical ventrite in male (Fig. 23) indistinctly lobed medially. Receptacle of spermatheca 1.6 times wider than long (Fig. 29). Tignum not widened proximally (Fig. 33); membranous distal region widest medially.

Aedeagus in lateral view (Fig. 27) strongly curved after basal 1/2, acutely narrowed in proximal 1/3, with weakly curved apex. In ventral view (Fig. 25), widest in proximal 1/3, narrowing sharply towards apex in apical 1/3, lateral margin a little abruptly narrowed preapically. Ventral aspect of aedeagus depressed with a convex portion in middle.

**Material examined. Types.** *Madurasia undulatovittata*: Lectotype ♀. “Teinodactila / undulato / vittata Motch / Ceylon”; “Monolepta / undulatovittata Mots. / 1926 D. Ogloblin det.”; “LECTOTYPUS / des Döberl 2005” (ZMUM).

*Madurasia obscurella*: Lectotype ♀. “Type” (rectangular red label); “Andrewes / Bequest. / B. M. 1922–221.”; “Madura”, “738” (3 in 738 is not legible as pierced by pin); “*Madurasia obscurella* Jac. / Type”; “SYNTYPE” (white circular disc with sky blue margin); “Lectotype / *Madurasia obscurella* Jacoby / des. K. D. Prathapan, 2015” (here designated, specimen on card, right antenna missing) (BMNH).

Paralectotype ♀. “Type / H. T.” (white circular disc with red border); “Madura”; “Jacoby Coll. 1909–28a.”; “*Madurasia / obscurella / Jac. Type*” (Blue label); “SYNTYPE” (white circular disc with sky blue margin); “Paralectotype / *Madurasia obscurella* Jacoby / des. K. D. Prathapan, 2015” (BMNH).

**Non-type material. AFRICA: Sudan:** ♀ British Sudan, S. R. J. Madani, 22.ix.1923, H. M. Bedford, feeding on ‘adis’ (illegible) sudani leaves / Blue Nile A 3024 / Pres by Imp. Bur. Ent. Brit. Mus. 1925–228 / standing as *Neorudolfia* (sic) bedfordi; 1 unsexed R. F. Wadmedanai J. W. Cowland 21/9/32 Shotholing seedlings of *Phaseolus mungo* / Ent. Coll. C 12147 / AFRICA 250,000 55–G Map / Pres. by Imp. Inst. Ent. BM 1933–415 / Standing as *Neorudolfia bedfordi* / SUDAN Govt.; 1 unsexed Blue Nile 5429 / Aenk H. H. & D. King 26.5.13 On boot / Pres. by Imp. Bur. Ent. Brit. Mus. 1927–103 / *Neorudolfia bedfordi* V. Laboissière–Dèt. (all BMNH).

**ASIA: Bangladesh:** ♀ (India) Dacca, 2.vi.1945, D. Leston; ♀ (India) Dacca, 10.v.1945, D. Leston (both BMNH); **India:** *Andhra Pradesh*: 3 unsexed Vizagapatnam Dist., Chipurupalli, B.M. 1924–7; *Gujarat*: 2♀ Baruch, 10.xii.1987, Pigeon pea, CIE A19617; Navasari, 15.iii.1992, Assoc with cowpea, IIE 22432, *Madurasia obscurella* Jac det. M. L. Cox 1992 (all BMNH); *Karnataka*: 1 macerated specimen Belgaum, 1–2.viii. 2008, at light, K. Swamy; 2♀, 1♂ Chikkaballapur, 13°25'48"N,

7°43'12"E 694 mt., 29.viii.2010, Nirmala P., at light (all UASB); *Kerala*: 7♂, 2 ♀ Vellayani, N 08°25'47.5"E, 76°59'8.3", 21.vii.2015, 18 m, Prathapan KD; 19 ♀, same data except for the date 5.vii. 2015 and ex Green gram (NBAIR, JBC, INPC, BMNH); *Maharashtra*: 3♀ Bandra, Jayakumar, 1905–152; 2 unsexed Bombay (Mumbai), 79.15; 1♀ Bombay, G. Bryant, 1919–147; 1 unsexed, Poona (Pune), 27.viii.1944, D. Leston, BM 1946–365; 1♀ 21.x.1944, D. Leston, BM 1945–86 (all BMNH); *Meghalaya*: 1♂, 1 ♀ SW of Cherapunjee, 23°13'15"N/ 91°40'E, 500–900 m, 11–12.v.2004, R. Businsky (all JBC); *New Delhi*: 4♀, 6 unsexed 21.viii.1968, on cowpea, *Phaseolus* and urd (all BMNH); *Rajasthan*: 1♀ Jodhpur N 26°21'4.6"E, 73°2'39"5.VIII.2015 255 m, Prathapan K. D. (KAU); 10 unsexed Banswara 24.ix.2015, S. Ramesh Babu (KAU); *Uttar Pradesh*: Saharanpur Div., Siwalik Hills, 8.iv.1928, H. G. Champion, B.M. 1928–518 (BMNH); *Uttarakhand*: 1 ♀ Dehra Dun, 8.ix.'16, H. G.Champion, BM. 1953– 156; 1 ♀ Ranikhet, 6–8. '16, H. G.Champion, BM. 1953–156 (both BMNH); *West Bengal*: 2 unsexed Sarda, H. G. Champion, B.M. 1953–156; 2 unsexed Sunderbans, H. G. Champion, B.M. 1953–156 (all BMNH); **Sri Lanka**: 2 unsexed, 1♀ Girandurukotte no. 68, 16.xii.86 on cowpea, CIE A18795; 2♀ Maha Illupallama, 1976, R. W. Fellowes, R. W. Fellowes, on *Glycine* & *Vigna*, CIE A9047 (all BMNH); **Yemen**: 2 ♀ Al Hudaydah gov., Jabal Bura Valley forest N. P., (stream valley), 240–350 m, 15°52.4–5'N, 43°24.6–25.2'E, J. Bezděk, 4.xi.2010; 1♂, 2♀ Socotra Island, wadi Ayhaft, 12°36.5'N, 53°58.9'E, 200 m, J. Bezděk, 7–8.xi.2010 (all JBC).

**Distribution.** Africa (Sudan); Asia (Bangladesh, India [Andhra Pradesh, Bihar, Gujarat, Haryana, Karnataka, Kerala, Madhya Pradesh, Maharashtra, Meghalaya, New Delhi, Orissa, Punjab, Rajasthan, Tamil Nadu, Uttar Pradesh, Uttarakhand, West Bengal], Nepal, Sri Lanka, Yemen) (Fig. 35).


**Remarks.** *Madurasia undulatovittata* and *M. andamanica* sp. n. are very similar. However, they can be separated as follows: eight labral setae present in *M. andamanica* sp. n. (only six labral setae visible in *M. undulatovittata*, though eight pores are present); elytral stripes are highly variable in *M. undulatovittata*, even in specimens from the same locality, collected during the same season and on the same host. The elytral pattern in *M. andamanica* sp. n. is rather consistent. The stripe in *M. undulatovittata* is wider apically in specimens where it is well defined, while in *M. andamanica* sp. n., it is narrowed apically. In *M. andamanica* sp. n., the stripe is distinct and well defined against the pale background color. Verma (1995) recorded variation in elytral color pattern. Lobe in the middle of the posterior margin of the apical abdominal ventrite in males distinct in *M. andamanica* sp. n., but poorly distinguishable in *M. undulatovittata*. The two species can easily be separated by the structure of the aedeagus. In lateral view, the apex of aedeagus of *M. andamanica* sp. n. is curved and pointed, like the beak of a parrot (Fig. 26), while the same in *M. undulatovittata* is narrowly rounded, and smoothly curved in apical 1/3 (Fig. 27). The sharply raised ridge on the ventral aspect of the aedeagus in *M. andamanica* sp. n. (Fig. 24) is characteristic, however, this ridge is absent in *M. undulatovittata* (Fig. 25). *Madurasia andamanica* sp. n. is confined to the Andaman Islands and reported to feed on pigeon pea, while *M. undulatovittata* is

transcontinental in distribution and a significant pest of a number of species of pulses in southern Asia and Africa (Sudan).

A photograph of the labels provided by Wagner & Bieneck (Fig. 38a in Wagner and Bieneck 2012) shows three labels, two of which show different information for *M. undulatovittata* (Fig. 2). Labels currently on the specimen indicate that M. Döberl designated the lectotype in 2005. However, no publication by Döberl could be traced in which this specimen is mentioned. According to Wagner and Bieneck (2012), the lectotype was designated by Wagner, and they provide photographs of both the lectotype and its labels. The photograph (Fig. 38b) in Wagner and Bieneck (2012), confirms that the specimen examined by me is the one designated as lectotype by Wagner (Fig. 1). Moreover, Wagner and Bieneck (2012) also mention that the only other specimen, a paralectotype in Motschulsky's collection, is a male from which the aedeagus has been dissected and subsequently lost. Dr Wagner's lectotype designation stands valid as that alone is published (Wagner and Bieneck 2012). Dr Döberl designated the same specimen as lectotype in 2005 as there was a long gap of nearly a decade between the lectotype designation by Dr Wagner and the publication of the same in Wagner and Bieneck 2012 (T. Wagner and M. Döberl, pers. comm., 2016). The specimen collected by Bedford on 22.ix.1923, identified as *Neorudolphia bedfordi* by Laboissière from the BMNH, probably belongs to the type series of *N. bedfordi*. The lectotype for *M. obscurella* is here designated, to have a unique name bearer and standard for its application.

**Host plants.** Fabaceae: *Cajanus cajan* (L.) Millsp. (red gram or pigeon pea); *Glycine max* (L.) Merr. (soybean); *Lablab purpureus* (L.) Sweet (= *Dolichos lablab* L.) (lablab bean); *Vigna aconitifolia* (Jacq.) Marechal (moth bean); *Vigna mungo* (L.) Hepper (= *P. mungo* L. = *P. radiatus* Roxb. non L.) (black gram); *Vigna radiata* (L.) R. Wilczek (= *Phaseolus aureus* Roxb. = *P. radiatus* L.) (green gram or moong); *Vigna radiata* (L.) Wilczek var. *sublobata* (Roxb.) (= *Phaseolus sublobatus* Roxb.); *Vigna umbellata* (Thunb.) Ohwi & Ohashi (rice bean) and *Vigna unguiculata* (L.) Walp. (= *Vigna sinensis* (L.) Savi ex Hausskn.) (cowpea).

**Biology and management.** Information on the host plants and biology of *M. undulatovittata* was generated by agricultural entomologists in India, under the name *M. obscurella*, where it is a widely distributed pest of legume crops across many agro climatic zones. The first record of this species as a pest of pulses is that by Menon and Saxena (1970). According to Naresh and Thakkur (1972), it was reported as a major pest of black gram by Naresh and Nene in 1968. However, there is no mention of this leaf beetle in Naresh and Nene (1968). Saxena et al. (1971) described it as a pest of cowpea, green gram or moong and black gram or urd, indicating that it made holes in the leaf lamina. Other recorded host plants include *Glycine* (CAB International Institute of Entomology 1990), moth bean (Pareek et al. 1983), lablab bean (Gupta and Singh 1984a, b), pigeon pea (Saxena 1977, Mishra and Saxena 1983), rice bean (Satyanarayana et al. 1995a, b) and *Vigna radiata* (L.) Wilczek var. *sublobata* (Roxb.) (= *Phaseolus sublobatus* Roxb.) (Kalaichelvan and Verma 2005).



**Figure 35.** Distribution of *M. undulatovittata* (Motschulsky) in the Afrotropical and Oriental regions (red triangles = literature records).

Gupta and Singh (1984a, b) provided the first account of its life cycle. They recorded the total life cycle as varying between 32 and 44 days and that it completes two generations a year on green gram. A second, more detailed study of the life history was reported by Oza et al. (1996) on cowpea. Eggs were laid singly on soil near the root zone of the plant. The total duration of the life cycle, from egg to death of adult, varied between 35 and 48 days in males and 43 to 58 days in females.

The growth of plants is retarded by severe foliage injury, especially in young plants (Srivastava and Singh 1976). Leaf damage on green gram in summer and rainy season crops ranged between 5–10% and 15–50% respectively (Sinha et al. 1985). Larvae are soil dwelling and feed on root hairs (Srivastava and Singh 1976; Gupta and Singh 1981). Odak and Thakur (1978) reported larval feeding on the root nodules. Gowda and Kaul (1982) recorded adult feeding on leaves, buds and flowers. Gowda et al. (2006) also observed feeding damage by adults on the buds and flowers of pigeon pea. Reddy and Varma (1986) established transmission of southern bean mosaic virus in cowpea by *M. undulatovittata*. The success in transmission varied from 25 to 43%.

The extent of damage on black gram, green gram and cowpea was 20–60% (Srivastava and Singh 1976). This is a common pest of mung bean in the first crop season

(*kharif*) in India, coinciding with the southwest monsoon (June to October) (Tiwari 1978). Singh and Gupta (1982) estimated damage to the leaves of green gram and black gram. Infestation was more pronounced in black gram than in green gram. Infestation starts when the plants are in the two leaf-stage and the insects remain active until flowering (Dhuri and Singh 1983, Nayak et al. 2005).

In Haryana, Yadav and Yadav (1983) recorded it from cowpea and Mrig and Singh (1985) observed maximum damage on *D. lablab* during the third week of September, with the pest disappearing after the first week of November. Feleiro and Singh (1985) carried out yield–infestation studies to fix the critical stages of crops requiring protection. They observed that infestation in summer resulted in heavy yield losses, while the pest attack during the rainy season had no significant effect on yield.

Lal (1985) reviewed information on the biology and control of insect pests of mung bean, including *M. undulato vittata*, in India.

According to Faleiro et al. (1986), *M. undulato vittata* is a sporadic, but major pest of cowpea. A peak population of 10.0–10.25 beetles/10 plants in summer and 29.50–30.25 beetles/10 plants in the rainy season were recorded by Gupta and Singh (1993) in green gram. Sahoo and Patnaik (1994) recorded the incidence of insect pests in green and black gram, and their seasonal activity and the extent of damage in Orissa. *Madurasia undulato vittata* was severe on both the crops in the seedling and vegetative stages, and was the first pest to appear at seedling stage on rice bean, continuing to occur until flowering (Satyanarayana et al. 1995b). Ganapathy and Durairaj (1995) reported it as an important pest on black gram and green gram in drought prone Pudukkottai District, Tamil Nadu. There was more damage in black gram (9.78%) than in green gram (1.45%). However, there are also reports of *M. undulato vittata* only being a minor pest (Devsthali and Joshi 1994, Kumar et al. 1998).

Dhuri et al. (1984) observed population buildup of *M. undulato vittata* under ambient temperature of about 32°C, longer duration of bright sunshine and high relative humidity coupled with intermittent rainfall. Sardana and Verma (1986) showed that maximum temperature and sunshine were negatively, but significantly, correlated with the population of the pest, while rainfall showed a significantly positive correlation. Maximum temperature, minimum temperature, sunshine hours and wind velocity had a significantly negative correlation with damage (Irulandi and Balasubramanian 1999). Nayak et al. (2004) reported a significantly negative correlation with minimum temperature and relative humidity during population buildup on black gram. The population did not show any correlation with maximum temperature, relative humidity and rainfall, but it was highly and significantly correlated with minimum temperature (Kumar et al. 2007).

Various cultivars of green gram (Srivastava et al. 1975, Sahoo et al. 1989, Sahoo and Hota 1991) and black gram (Sahoo et al. 1989) vary significantly in their susceptibility to the pest. Pandey et al. (1995) reported that varieties with thicker leaves were preferred by the pest.

Chemical control remains the most effective option against this pest. Several broad spectrum insecticides have been tried against *M. undulato vittata*, with varying degrees of success (Saxena et al. 1971, Naresh and Thakur 1972, Saxena et al. 1975, Verma and

Pant 1975, Verma and Lal 1976, Vyas and Saxena 1977, Verma and Lal 1978, Yadav et al. 1979, Vyas and Saxena 1981, Chaudhary et al. 1981, Rajendran et al. 1981, Vyas and Saxena 1982, Mishra and Saxena 1983, Singh et al. 1983, Singh and Gupta 1984, Sinha 1985, Gattoria and Singh 1988, Rahman 1988, 1991a, b, Chander and Singh 1989, Sinha and Sharma 1989, Verma and Dikshit 1990, Logiswaran and Gopalan 1993, Uddin et al. 1994, Das 1999, Gowda et al. 2006 and Pandey et al. 2007). Application of neem seed kernel extract had no significant effect to increase yield in mung bean (Yadav et al. 1979). Soundararajan and Chitra (2011) tried biological control agents such as *Pseudomonas fluorescens* and *Beauveria bassiana* and reported that intercropping black gram with sorghum reduced infestation (Soundararajan and Chitra 2012).

## Discussion

A revision of the genus *Medythia* is required to define the boundaries between it and the genus *Madurasia*. *Medythia* and *Madurasia* share the same ecological niche and are often collected together on the same host plants, as well as at light. It is likely that economic entomologists have often misidentified one for the other. *Medythia bukit* and *Medythia marginicollis*, described by Mohamedsaid (1999) from Malaysia, with pronotum broader than wide, as well as a little narrower anteriorly than posteriorly, appear atypical for the genus. Reports of a 6.05mm, ovate beetle as *M. obscurella* from Pakistan (Rizvi et al. 2012, Kamaluddin et al. 2012) are incorrect.

*Madurasia andamanica* sp. n. is a significant pest of red gram or pigeon pea (*C. cajan*) in the Great Andaman Islands (Bharathimeena T., pers. comm. 2015), similar to the pest status of *M. undulatovittata* elsewhere. This endemic pest of the islands, in case of accidental introduction to the mainland India, is likely to become a pest of various pulses and spread far and wide, as in the case of *M. undulatovittata*.

## Acknowledgements

I am deeply indebted to Drs L. N. Medvedev and A. Moseyko (Russian Academy of Sciences) for the loan of the syntype of *M. undulatovittata*; Dr M. Geiser (BMNH) for the loan of the Jacoby types of *M. obscurella* and other specimens from south-east Asia and Sudan, and to Dr Jan Bezděk (Mendel University) who provided specimens from Yemen and north-east India. Jan Bezděk helped with many inaccessible papers, critical suggestions and advices. Specimens of *M. andamanica* sp. n. were provided by T. Bharathimeena and H. M. Yeswanth. I am grateful to Jennie Unnikrishnan, Nisha Rakesh and Tayan Raj Gurung, without whose help, it would have been impossible for me to access many obsolete economic entomology papers. Critical reviews by Drs C. A. Viraktamath, J. Poorani, Jan Bezděk, T. Wagner and E. Grobbelaar greatly improved the manuscript. M. C. Kiran prepared the maps. My work on leaf beetles is funded by the Indian Council of Agricultural Research through the Network Project on Insect Systematics.

## References

- Aslam NA (1972) On the genus *Drasa* Bryant (Coleoptera, Chrysomelidae, Galerucinae) with some nomenclatural notes on the Galerucinae. *Journal of Natural History* 6: 483–501. doi: 10.1080/00222937200770451
- Beenen R (2010) Galerucinae. In: Löbl I, Smetana A (Eds) *Catalogue of Palearctic Coleoptera Vol. 6: Chrysomeloidea*. Apollo Books, Stenstrup, 924 pp.
- Bezděk J (2012) Galerucinae (Coleoptera: Chrysomelidae) of Socotra Island, with a review of taxa recorded from Yemen. *Acta Entomologica Musei Nationalis Pragae* 52, Supplementum 2: 403–428.
- CAB International Institute of Entomology (1990) *Distribution maps of pests. Series A (Agricultural)*. Map no. 517.
- Chander S, Singh Y (1989) Control of galerucid beetle, *Madurasia obscurella* Jacoby infesting green gram, *Vigna radiata* (L.) Wiczek. *Indian Journal of Entomology* 51(2): 185–191.
- Chaudhary JP, Singh AK, Yadav LS (1981) Effect of disulfoton granules on pest complex, crop growth and yield of green gram. *Indian Journal of Entomology* 43(4): 369–372.
- Das SB (1999) Management of flea beetle in green gram, *Vigna radiata* (L.) Wiczek. *Insect Environment* 5(1): 39–40.
- Devsthali S, Joshi M (1994) Infestation and varietal preference of insect pests in green gram. *Indian Agriculturist* 38(4): 263–272.
- Dhuri AV, Singh KM (1983) Pest complex and succession of insect pests in black gram *Vigna mungo* (L.) Hepper. *Indian Journal of Entomology* 45(4): 396–401.
- Dhuri AV, Singh KM, Singh RN (1984) Incidence of insect pests in black gram *Vigna mungo* (L.) Hepper. *Indian Journal of Entomology* 46(3): 270–276.
- Faleiro JR, Singh KM (1985) Yield–infestation studies associated with insects infesting cowpea, *Vigna unguiculata* (L.) Walp. in Delhi. *Indian Journal of Entomology* 47(3): 287–291.
- Faleiro JR, Singh KM, Singh RN (1986) Pest complex and succession of insect pests in cowpea *Vigna unguiculata* (L.) Walp. *Indian Journal of Entomology* 48(1): 54–61.
- Ganapathy N, Durairaj C (1995) Pest status of pulses in Pudukkottai District, Tamil Nadu. *Madras Agricultural Journal* 82(4): 322.
- Gatoria GS, Singh H (1988) Effect of insecticidal and fertilizer applications on chewing insect pests of green gram. *Journal of Research, Punjab Agricultural University* 25(2): 225–232.
- Gemminger M, Harold EV (1876) Chrysomelidae. In: *Catalogus Coleopterorum locusque descriptorum synonymicus et systematicus*. Monachus, Paris & London, 3479–3676.
- Gowda CLL, Kaul AK (1982) *Pulses in Bangladesh*. Bangladesh Agricultural Research Institute, Dhaka, 472 pp.
- Gowda DKS, Yelshetty S, Patil BV (2006) Flea beetle, *Madurasia obscurella* Jacoby on pigeonpea. *Insect Environment* 11(4): 154.
- Gupta PK, Singh J (1981) Important insect pests of cowpea (*Vigna unguiculata* L.) in agroecosystem of eastern Uttar Pradesh. *Indian Journal of Zootomy* 22(2): 91–95.
- Gupta PK, Singh J (1984a) Biology of *Madurasia obscurella* Jacoby, an important pest of rainy-season pulses. *Indian Journal of Agricultural Sciences* 54(10): 931–934.

- Gupta PK, Singh J (1984b) Ecological studies on galerucid beetle (*Madurasia obscurella* Jacoby), an important pest of green gram. Plant Protection Bulletin, India 36(4): 15–18.
- Gupta PK, Singh J (1993) Population studies on insect pests of green gram (*Vigna radiata* (L.) Wilczek). Indian Journal of Entomology 55(1): 45–51.
- Irulandi S, Balasubramanian G (1999) Effect of weather parameters on galerucid beetle, *Madurasia obscurella* (Jacoby) (Galerucidae: Coleoptera) and stemfly, *Ophiomyia phaseoli* (Tryon) (Agromyzidae: Diptera) on green gram. Insect Environment 5(1): 8–9.
- Jacoby M (1886) Description of the new genera and species of phytophagous Coleoptera obtained by Mr. Andrews in India. Part II—Chrysomelidae, Halticinae and Galerucinae. Annales de la Société Entomologique de Belgique 40: 250–304.
- Jacoby M (1887) Description of some new genera and species of phytophagous Coleoptera contained in the Leyden Museum. Notes from the Leyden Museum 9: 220–243.
- Jolivet P, Hawkeswood TJ (1995) Host-plants of Chrysomelidae of the world: An Essay about the Relationships between the Leaf-beetles and their Food-plants. Backhuys Publishers, Leiden, 281 pp.
- Kalaichelvan T, Verma KK (2005) Checklist of leaf beetles (Coleoptera: Chrysomelidae) of Bhilai-Durg. Zoos' Print Journal 20(4): 1838–1842. doi: 10.11609/JoTT.ZPJ.1114.1838-42
- Kamaluddin S, Rizvi SG, Javed T (2012) Cladistic analysis of the Galerucinae Species (Coleoptera: Chrysomelidae: Galerucinae) from Pakistan. Pakistan Journal of Entomology 27(2): 89–98.
- Kimoto S (2005) Systematic catalog of the Chrysomelidae (Coleoptera) from Nepal and Bhutan. Bulletin of the Kitakyushu Museum of Natural History and Human History, Series A 3: 13–114.
- Konstantinov AS (1998) Revision of the Palearctic species of *Aphthona* Chevrolat and cladistic classification of the Aphthonini (Coleoptera: Chrysomelidae: Alticinae). Memoirs on entomology, International 11: 1–429.
- Kumar A, Sinha AK, Yazdani SS (1998) Incidence of insect pests in urd [*Vigna mungo* (Linn.) (Hepper)]. Shashpa 5(1): 113–114.
- Kumar R, Ali S, Chandra U (2007) Seasonal incidence of insect-pests on *Vigna mungo* and its correlation with abiotic factors. Annals of Plant Protection Sciences 15(2): 366–369.
- Laboissière V (1926) Description d'un nouveau genre et de trois nouvelles espèces de *Galerucini africains* (Col.). Bulletin de la Société Entomologique de France 1926: 190–193.
- Lal SS (1985) A review of insect pests of mungbean and their control in India. Tropical Pest Management 31(2): 105–114.
- Logiswaran G, Gopalan M (1993) Effect of seed treatment with insecticides on black gram galerucid beetle. Madras Agricultural Journal 80(2): 39–41.
- Maulik S (1926) The Fauna of British India including Burma and Ceylon. Coleoptera. Chrysomelidae. (Chrysomelinae and Halticinae). Taylor and Francis, London, 442 pp.
- Maulik S (1936) The Fauna of British India, including Ceylon and Burma. Chrysomelidae (Galerucinae). Taylor & Francis, London, 439 pp.
- Medvedev LN, Sprecher-Uebersax E (1999) Katalog der Chrysomelidae von Nepal. Entomologica Basiliensia 21: 261–354.

- Medvedev LN, Sprecher–Uebersax E (2005) A key to the leaf beetle genera of Nepal (Coleoptera: Chrysomelidae). *Entomologica Basiliensia et Collection Frey* 27: 297–336.
- Menon MGR, Saxena HP (1970) The common pulse Galerucid beetle at Delhi. *Labdev Journal of Science and Technology* 8–B(1): 62–63.
- Mishra PN, Saxena HP (1983) Phytotonic effects of insecticides in pigeon pea *Cajanus cajan* (L.) Millsp. *Madras Agricultural Journal* 70(5): 309–311.
- Mohamedsaid MS (1997) The galerucine beetles of Sri Lanka, with descriptions of two new species (Coleoptera: Galerucinae). *Stobaeana* 9(15): 1–7.
- Mohamedsaid MS (1999) Two new species of *Medythia* Jacoby from Malaysia (Coleoptera: Chrysomelidae: Galerucinae). *Genus* 10(3): 415–420.
- Mohamedsaid MS (2000) List of the non–Malaysian Chrysomelidae in the collection of UKM. *Serangga* 5(2): 361–377.
- Motschulsky V (1866) Essai d'un catalogue des insectes de l'île de Ceylan. *Bulletin de la Société Impériale des Naturalistes de Moscou* 39(2): 393–446.
- Mrig KK, Singh R (1985) Incidence of insect-pests on garden bean, *Dolichos lablab* Linn. *Bulletin of Entomology* 26(1): 5–7.
- Naresh JS, Nene YL (1968) Survey and evaluation of economic importance of insect pests of pulses. *Annual Report 1968–69, Uttar Pradesh Agricultural University, Pant Nagar*, 177–178.
- Naresh JS, Thakur RP (1972) Efficacy of systemic granular and spray insecticides for the control of insect pests of black-gram (*Phaseolus mungo* Roxb.). *Indian Journal of Agricultural Sciences* 42(8): 732–735.
- Nayak SK, Ujagir R, Chhibber RC (2004) Effect of abiotic factors on the insect population build up on black gram, *Vigna mungo* L. crop. *Shashpa* 11(1): 31–36.
- Nayak SK, Chhibber RC, Ujagir R (2005) Pest complex and their succession on black gram, *Vigna mungo* L. *Shashpa* 12(1): 60–62.
- Odak SC, Thakur BS (1978) New record of *Madurasia obscurella* Jacoby (Coleoptera–Chrysomelidae; Galerucinae) feeding on root nodules of leguminous crops. *Science and Culture* 44(7): 328–329.
- Ogloblin DA (1930) De quelques espèces de Halticinae (Col. Chrysomelidae) de la collection de V. Motschulsky. *Museo Nacional de Ciencias Naturales Madrid* 10: 83–112.
- Oza RU, Patel MB, Patel CB (1996) Biology of galerucid flea beetle, *Madurasia obscurella* Jacoby (Galerucinae : Chrysomelidae : Coleoptera) on cowpea *Vigna unguiculata* (L.) Walp. *Gujarat Agricultural University Research Journal* 21(2): 55–59.
- Pandey KC, Hasan N, Bhaskar RB, Ahmad ST, Kohli KS (1995) Genetic evaluation of cowpea (*Vigna unguiculata* (L.) Walp.) lines for multiple pest resistance. *Indian Journal of Genetics & Plant Breeding* 55(2): 198–203.
- Pandey A, Pandey A, Srivastava SP (2007) Management of pest complex of summer mung by granular insecticides. *Annals of Plant Protection Sciences* 15(2): 376–380.
- Pareek BL, Sharma RC, Yadav CPS (1983) Record of insect faunal complex on mothbean, *Vigna aconitifolia* (Jacq.) Marechel in semi arid zone of Rajasthan. *Bulletin of Entomology* 24(1): 44–45.
- Rahman MM (1988) Efficacy of some promising insecticides on pest incidence, plant growth and grain yield of cowpea. *Tropical Grain Legume Bulletin* 35: 19–22.

- Rahman MM (1991a) Progress and Prospects of Minor Pulses in Bangladesh. In: Advances in pulses research in Bangladesh: proceedings of the second national workshop on pulses, 6–8 Jun 1989, Joydebpur, Bangladesh. International Crops Research Institute for the Semi-Arid Tropics, Patancheru, 40–47.
- Rahman MM (1991b) Control Measures for Important Insect Pests of Major Pulses. In: Advances in Pulses Research in Bangladesh: proceedings of the Second National Workshop on Pulses, 6–8 Jun 1989, Joydebpur, Bangladesh. International Crops Research Institute for the Semi–Arid Tropics, Patancheru, India, 139–146.
- Rajendran R, Gopalan M, Janarthanan R (1981) A note on the control of pod-borers in black gram and green gram with dust formulations. *Pesticides* 15(10): 35–36.
- Reddy DVR, Varma A (1986) *Madurasia obscurella* Jacoby—a new vector of southern bean mosaic virus. *Current Science* 55(2): 109–110.
- Rizvi SG, Kamaluddin S, Javed T (2012) *Madurasia obscurella* Jacoby: a newly recorded species from Pakistan (Coleoptera: Chrysomelidae: Galerucinae). *Pakistan Journal of Entomology* 27(2): 85–87.
- Sahoo BK, Hota AK (1991) Field screening of green gram germplasm against insect pest and disease complex. *Madras Agricultural Journal* 78(1–4): 84–86.
- Sahoo BK, Patnaik NC (1994) Insect pests in green gram and black gram in the South Coastal Region of Orissa with notes on their seasonal activity. *Orissa Journal of Agricultural Research* 7(suppl.): 74–76.
- Sahoo BK, Sontakke BK, Rath LK (1989) Varietal susceptibility of different green gram and black gram cultivars to the leaf beetles and pod borer complex. *Environment and Ecology* 7(2): 345–347.
- Sardana HR, Verma S (1986) Preliminary studies on the prevalence of insect pests and their natural enemies on cowpea crop in relation to weather factors at Delhi. *Indian Journal of Entomology* 48(4): 448–458.
- Satyanarayana J, Singh KM, Singh RN (1995a) Pest complex of rice bean *Vigna umbellata*. *Bulletin of Entomology* 36(1/2): 28–33.
- Satyanarayana J, Singh KM, Singh RN (1995b) Insect pest succession in rice bean. *Bulletin of Entomology* 36(1/2): 78–83.
- Saxena MC (1977) Water, fertilizer and pest control management in food legumes. Food and Agriculture Organization, Rome, Italy, Technology for increasing food production. Proceedings of the Second FAO/SIDA Seminar on Field Food Crops in Africa and the Near East, Lahore, Pakistan, 18 September to 5 October 1977, 653–658.
- Saxena HP, Kumar S, Prasad SK (1971) Efficacy of some systemic granular insecticides against Galerucid beetle, *Madurasia obscurella* Jacoby infesting kharif pulses. *Indian Journal of Entomology* 33(4): 470–471.
- Saxena HP, Phokela A, Singh Y (1975) Effectiveness of carbofuran as seed treatment for controlling pest complex of ‘moong’ (*Phaseolus aureus*) and ‘urd’ (*Phaseolus mungo*). *Entomologists’ Newsletter* 5(4): 27–28.
- Seeno TN, Wilcox JA (1982) Leaf beetle genera (Coleoptera: Chrysomelidae). *Entomography* 1: 1–221.

- Singh J, Gupta PK (1982) Heavy leaf damage of green gram and black gram by galerucid beetle, *Madurasia obscurella* Jacoby (Coleoptera: Chrysomelidae) at Varanasi (U.P.). *Entomon* 7(1): 51–53.
- Singh J, Gupta PK (1984) Relative efficacy of some granular insecticides against galerucid beetle (*Madurasia obscurella* Jacoby), an important pest of green gram. *Indian Journal of Entomology* 46(2): 221–231.
- Singh AK, Gupta DS, Hameed SF (1983) Translocation and dissipation of disulfoton in mung bean, *Vigna radiata* var. *aureus* Hübn. *Journal of Entomological Research* 7(2): 127–132.
- Sinha SN (1985) Field screening of insecticides for seed treatment of pulses. I. Green gram (*Vigna radiata* (L.) Wilczek). *Seed Research* 13(1): 120–128.
- Sinha MM, Yazdani SS, Kumar A, Singh R (1985) Observations of the seasonal occurrence of insect pests of green gram (*Phaseolus aureus* Roxb.) in north Bihar. *Bulletin of Entomology* 26(2): 210–211.
- Sinha SN, Sharma SP (1989) Effect of application of carbofuran granule along with seed on plant growth and seed yield of cowpea (*Vigna unguiculata*). *Indian Journal of Agricultural Sciences* 59(7): 472–475.
- Soundararajan RP, Chitra N (2011) Effect of bioinoculants on sucking pests and pod borer complex in urdbean. *Journal of Biopesticides* 4(1): 7–11.
- Soundararajan RP, Chitra N (2012) Impact of intercrops on insect pests of black gram, *Vigna mungo* L. *Journal of Entomology* 9(4): 208–219. doi: 10.3923/je.2012.208.219
- Srivastava KM, Singh LN (1976) A review of the pest complex of kharif pulses in Uttar Pradesh. *PANS* 22(3): 333–335. doi: 10.1080/09670877609412068
- Srivastava AS, Srivastava KM, Singh LN, Singh RP, Srivastava RL (1975) Relative susceptibility of some green gram (*Phaseolus aureus* Roxb.) varieties to galerucid beetle, *Madurasia obscurella* Jacoby and pod borer, *Euchrysops cnejus* Fabricius. *Labdev Journal of Science and Technology, B (Life Sciences)* 13(3/4): 221–222.
- Takizawa H (1987) Chrysomelidae collected by the Japan–India cooperative survey in India, 1978. Part IV. *Entomological Review of Japan* 42(1): 37–42.
- Takizawa H (1990) Chrysomelid beetles of Nepal, Northeastern India and Western Sikkim collected by the Himalaya expeditions of the National Science Museum, Tokyo. *Japanese Journal of Entomology* 58(2): 275–291.
- Takizawa H, Kimoto S (1990) Notes on chrysomelid beetles of S. India preserved in the collection of the Staatliches Museum of Munich (Coleoptera: Chrysomelidae). *Entomological Review of Japan* 45(1): 7–10.
- Tiwari AS (1978) Mungbean varietal requirements in relation to cropping seasons in India. The 1st international mungbean symposium, 1978, Office of Information Services, Asian Vegetable Research and Development Center, Taiwan, 129–131.
- Uddin MB, Karim MA, Huq SB, Rahman MM, Alam MZ (1994) Effect of some insecticides on leaf growth and leaf infestation by galerucid beetle and grain yield of cowpea (*Vigna unguiculata* L. Walp.). *Annals of Bangladesh Agriculture* 4(2): 109–115.
- Verma KK (1995) Elytral color pattern in *Madurasia obscurella* Jac. (Chrysomelidae, Galerucinae). *Chrysomela* 30: 7.

- Verma S, Dikshit AK (1990) Efficacy and persistence of some insecticides against pests of cowpea. *Journal of Insect Science* 3(2): 162–166.
- Verma S, Lal R (1976) Evaluation of phorate, monocrotophos and carbaryl against the pest of cowpea crop. *Entomologists' Newsletter* 6(8/9): 52–53.
- Verma S, Lal R (1978) Evaluation of pesticides against the pests of cowpea crop (*Vigna sinensis* Savi). *Indian Journal of Entomology* 40(1): 54–58.
- Verma S, Pant NC (1975) Effect of phorate and endosulfan treatments on growth parameters of 'moong' and 'arhar' crops. *Entomologists' Newsletter* 5(8/9): 41–42.
- Vyas HN, Saxena HP (1977) Observations on the phytotoxicity and plant stand due to disulfoton and phorate seed treatment in *Vigna radiata* ('moong'). *Entomologists' Newsletter* 7(9/10): 44–45.
- Vyas HN, Saxena HP (1981) Some observations on the effectiveness of disulfoton and phorate seed treatment against galerucid beetle, jassid, plant height and yield of green gram. *Pesticides* 15(10): 44.
- Vyas HN, Saxena HP (1982) Comparative efficacy of some insecticides against galerucid beetle, *Madurasia obscurella* Jacoby, infesting green gram. *Indian Journal of Agricultural Sciences* 52(10): 689–693.
- Wagner T, Bieneck S (2012) Galerucinae type material described by Victor Motschulsky in 1858 and 1866 from the Zoological Museum Moscow (Coleoptera: Chrysomelidae: Galerucinae). *Entomologische Zeitschrift Stuttgart* 122(5): 205–215.
- Weidner H (1976) Die Entomologischen Sammlungen des Zoologischen Instituts und des Zoologischen Museums der Universität Hamburg. *Mitteilungen aus dem Hamburgischen Zoologischen Museum und Institut* 73: 87–264.
- Wilcox JA (1973) *Coleopterorum Catalogus Supplementa Pars 78 Fasc. 3 Chrysomelidae: Galerucinae Luperini: Luperina*. W. Junk, Gravenhage, 433–664.
- Yadav LS, Yadav PR (1983) Pest complex of cowpea (*Vigna sinensis* Savi) in Haryana. *Bulletin of Entomology* 24(1): 57–58.
- Yadav LS, Yadav PR, Poonia FS (1979) Effectiveness of certain insecticides against the insect pests of mung bean. *Indian Journal of Plant Protection* 7(2): 165–169.