

A new species and additional records of *Lobrathium* Mulsant & Rey (Coleoptera, Staphylinidae, Paederinae) from South China

Zhong Peng¹, Li-Zhen Li¹, Mei-Jun Zhao¹

¹ Department of Biology, College of Life and Environmental Sciences, Shanghai Normal University, Shanghai, 200234, P. R. China

Corresponding author: Mei-Jun Zhao (mjzhao@shnu.edu.cn)

Academic editor: V. Assing | Received 25 December 2015 | Accepted 28 January 2016 | Published 23 February 2016

<http://zoobank.org/3355C7D9-6F35-44CA-9A65-94D1A6622B3F>

Citation: Peng Z, Li L-Z, Zhao M-J (2016) A new species and additional records of *Lobrathium* Mulsant & Rey (Coleoptera, Staphylinidae, Paederinae) from South China. ZooKeys 568: 51–58. doi: 10.3897/zookeys.568.7622

Abstract

Material of the genus *Lobrathium* Mulsant & Rey, 1878 from the Chinese provinces Fujian, Hunan, Sichuan, Guangdong and Guangxi is examined. Six species are identified, four of them described previously and two undescribed. *Lobrathium kedian* Peng & Li, **sp. n.** (Guangxi: Shiwangda Shan) is described and illustrated. One probably undescribed species remains unnamed. The female sexual characters of *L. flexum* Assing, 2014 are described and illustrated for the first time. The genus is now represented in mainland China by 43 species.

Keywords

Coleoptera, Staphylinidae, *Lobrathium*, new species, new records, China

Introduction

Until today, 42 species of the genus *Lobrathium* Mulsant & Rey have been reported from mainland China and 20 species from Taiwan (Assing 2010, 2012, 2013, 2014; Li et al. 2013; Li et al. 2013a, b, c; Lü and Li 2014). With a total of 14 described species, the *Lobrathium* fauna of Sichuan is currently more diverse than that of any of the other Chinese provinces, followed by Yunnan (8 species), Shaanxi (7 species), Guizhou (6 species) and Zhejiang (6 species) (Assing 2012, 2013, 2014; Li et al. 2013; Li et al. 2013a, b, c; Lü and Li 2014).

A study of *Lobrathium* material from southern China yielded a species new to science and additional records of *L. configens* Assing, 2012, *L. flexum* Assing, 2014, *L. hebeatum* Zheng, 1988 and *L. hongkongense* Bernhauer, 1931.

Material and methods

The following abbreviations are used in the text, with all measurements in millimeters:

Body length (BL) from the anterior margin of the labrum to the abdominal apex; forebody length (FL) from the anterior margin of the labrum to the posterior margin of the elytra; head length (HL) from the anterior clypeal margin to the occipital constriction; head width (HW): maximum width of head; length of antenna (AnL); length of pronotum (PL) along midline; maximum width of pronotum (PW); elytral length (EL) at the suture from the apex of the scutellum to the posterior margin of the elytra (at the sutural angles); maximum width of the elytra (EW); length of aedeagus (AL) from the apex of the dorsal plate to the base of the aedeagal capsule.

The type material is deposited in the Insect Collection of Shanghai Normal University, Shanghai, China (SNUC).

Results

Lobrathium configens Assing, 2012

Fig. 4

Material studied. China: Sichuan: 8 ♂♂, 2 ♀♀, Xiaojin County, Jiajin Shan, 30°48'49"N, 102°42'55"E, 2500 m, 20.VII.2015, Jiang, Peng, Tu & Zhou leg. (SNUC).

Comment. *Lobrathium configens* was previously known from the Chinese provinces Shaanxi, Sichuan, Qinghai, Hubei, Yunnan and Zhejiang (Assing 2012, 2013, 2014; Li et al. 2013a, b). For illustrations of *L. configens* see Assing (2012: figures 153–165) and Li et al. (2013a: figure 4).

Lobrathium flexum Assing, 2014

Figs 1A, 2A–C, 5

Material studied. China: Hunan: 1 ♂, 1 ♀, Yanling County, Nanfengmian, 26°18'N 114°00'E, 1600 m, 06.VI.2015, Peng, Shen, Tu & Zhou leg. (SNUC).

Comment. The original description is based on a male from Jiangxi. The previously unknown female sexual characters are as follows: posterior margin of tergite VIII convex (Fig. 2A); sternite VIII (Fig. 2B) weakly transverse, posteriorly broadly convex;




Figure 1. Habitus: **A** *Lobrathium flexum* **B** *Lobrathium kedian*. Scale bars: 2.0 mm.

tergite IX (Fig. 2C) undivided anteriorly. The above record from Hunan represents a new province record. For illustrations of the habitus and the male sexual characters see Assing (2014: figures 12–16).

Lobrathium hebeatum Zheng, 1988

Fig. 6

Material studied. China: Sichuan: 1 ♂, 3 ♀♀, Dayi County, Xiling Xueshan, 30°41'59"N, 103°12'10"E, 2150 m, 29.VII.2015, Jiang, Peng, Tu & Zhou leg. (SNUC).

Comment. The previously known distribution of *L. hebeatum* included the Chinese provinces Shaanxi, Sichuan, Yunnan, Henan and Ningxia (Assing 2012, 2013, 2014; Li et al. 2013a, b; Zheng 1988). For illustrations of *L. hebeatum* see Assing (2012: figures 142–147) and Li et al. (2013a: figure 9).




Figure 2. *Lobrathium flexum* (A–C) and *Lobrathium kedian* (D–E). **A** female tergite VIII **B** female sternite VIII **C** female tergites IX–X **D** antenna **E** forebody. Scale bars: 0.5 mm (A–C), 1.0 mm (D–E).

Lobrathium hongkongense Bernhauer, 1931

Figs 5, 8

Material studied. China: Fujian: 3 ♂♂, Nanping, Mangdang Shan, 26°41'51"N, 118°07'00"E, 400 m, 10.IX.2015, Yan & Tang leg. (SNUC). Hunan: 2 ♂♂, 1 ♀, Yanling County, Nanfengmian, 26°18'N 114°00'E, 1600 m, 06.VI.2015, Peng, Shen, Tu & Zhou leg. (SNUC). Guangdong: 1 ♂, 1 ♀, Ruyuan County, Nanling Nature Reserve, Qingshuigu, 24°54'57"N, 113°01'55"E, 900 m, 04.V.2015, Peng, Tu & Zhou leg. (SNUC); 1 ♀, Jieyang, Puning, Wufeng Shan, 500 m, 08.VI.2015, Aranyu leg. (SNUC).

Comment. *Lobrathium hongkongense* was previously known from Japan, Hong Kong, Taiwan and the Chinese provinces Fujian, Guizhou, Zhejiang, Jiangsu, Sichuan,

Yunnan, Guangxi, Hubei and Shaanxi (Assing 2012, 2013; Li et al. 2013a, b). The specimens represent the first record from Hunan and Guangdong. For illustrations of *L. hongkongense* see Assing (2012: figures 125–132) and Li et al. (2013a: figure 10).

***Lobrathium kedian* Z. Peng & L.-Z. Li, sp. n.**

<http://zoobank.org/546F9FE5-C3BA-469F-A96A-8FC2117C7634>

Figs 1B, 3, 7

Type material. HOLOTYPE: ♂, labelled 'China: Guangxi Prov., Shangsi County, Shiwanda Shan, 300–500 m, 21°54'N, 107°54'E, 25–IV–2011, Peng & Zhu leg.' (SNUC). Paratypes: 8 ♂♂, 8 ♀♀, same label data as holotype (SNUC).

Description. Measurements (in mm) and ratios: BL 9.88–10.20, FL 5.49–5.62, HL 1.37–1.42, HW 1.36–1.44, AnL 3.13–3.20, PL 1.57–1.63, PW 1.24–1.30, EL 1.35–1.39, EW 1.39–1.46, AL 1.13–1.20, HL/HW 0.97–1.00, HW/PW 1.06–1.10, HL/PL 0.85–0.88, PL/PW 1.25–1.27, EL/PL 0.84–0.87.

Habitus as in Fig. 1B. Coloration: body black, mandibles dark brown, labial palpi light brown; antennae dark brown to light brown; legs with blackish brown profemora and protibiae, basal halves of meso- and metafemora yellowish brown, distal halves gradually infusate.

Head as wide as long, widest behind eyes; punctuation coarse and very dense; interstices without microsculpture. Antenna as in Fig. 2D.

Pronotum distinctly longer than wide, with impunctate midline; punctuation coarse and dense, but distinctly sparser than that of head; interstices glossy.

Elytra distinctly broader than pronotum; punctuation coarse, arranged in irregular series only laterally. Hind wings approximately 1.85–2.02 times as long as elytra.

Abdomen somewhat narrower than elytra; punctuation fine and dense; posterior margin of tergite VII with palisade fringe.

Male. Sternite VII (Fig. 3D) strongly transverse and with shallow median impression posteriorly, without modified setae, posterior margin broadly concave; sternite VIII (Fig. 3E) posteriorly with deep impression, this impression with a cluster of numerous short peg-setae, postero-laterally with a cluster of short black setae; posterior excision large, deep and U-shaped; aedeagus (Figs 3F, G) with apically bifid ventral process in ventral view and broad dorsal plate.

Female. Posterior margin of tergite VIII (Fig. 3A) convex; sternite VIII (Fig. 3B) weakly transverse, posterior margin broadly convex; tergite IX (Fig. 3C) slender and undivided anteriorly.

Distribution and natural history. The type locality is situated in Shiwangda Shan to the south of Shangsi, southern Guangxi. The specimens were sifted from leaf litter in broad-leaved forests at altitudes of 300–500 m (Fig. 7).

Etymology. The specific name is the Chinese noun “kedian” (punctuation) in apposition. It refers to the punctuation of the head of *L. kedian*, which is denser than that of other species known from Guangxi.





Figure 3. *Lobrathium kedian*. **A** female tergite VIII **B** female sternite VIII **C** female tergites IX–X **D** male tergite VIII **E** male sternite VIII **F** aedeagus in ventral view **G** aedeagus in lateral view. Scale bars: 0.5 mm.

Comparative notes. *Lobrathium kedian* shares a bifid ventral process with *L. digitatum* Assing, 2010 from Taiwan, but differs from it in many respects, particularly by larger body size, the shape and chaetotaxy of the male sternite VIII and by the shape of the aedeagus. For illustrations of *L. digitatum* see Assing (2010: figures 203–210).

***Lobrathium* sp.**

Fig. 6

Material studied. China: Sichuan: 1 ♀, Dayi County, Xiling Xueshan, 30°41'59"N, 103°12'10"E, 2150 m, 29.VII.2015, Jiang, Peng, Tu & Zhou leg. (SNUC).



Figures 4–8. Habitats of *Lobrathium*. **4** Jiajin Shan, alt. 2500 m (*L. configans*) **5** Nanfengmian, alt. 1600 m (*L. flexum* and *L. hongkongense*) **6** Xiling Xueshan, alt. 2150 m (*L. hebeatum* and *Lobrathium* sp.) **7** Shiwangda Shan, alt. 300–500 m (*L. kedian* sp. nov.) **8** *Lobrathium hongkongense* walking on the stone.

Comment. This species is similar and probably closely related to *L. daxuense* Assing, 2012. The female represents an undescribed species distinguished from its congeners particularly by the light brown coloration, large body size (8.34 mm), much denser punctuation of the head, a slender pronotum, and the female secondary sexual characters.

Acknowledgements

All the collectors mentioned in the text are acknowledged for their field work. Two anonymous reviewers are thanked for comments on a previous version of the manuscript. We are most grateful to Volker Assing (Hannover) for revising an earlier version of the manuscript. The study is supported by the National Natural Science Foundation of China (No. 31201734 and 31101659), the Foundation of Shanghai Municipal Education Commission (No. 12YZ077) and Shanghai Normal University (SK201234, DZL125 and B-9013-11-003127).

References

- Assing V (2010) On the Lathrobiina of Taiwan (Coleoptera: Staphylinidae: Paederinae). *Beiträge zur Entomologie*, Berlin 60: 301–361.
- Assing V (2012) A revision of East Palaearctic *Lobrathium* (Coleoptera: Staphylinidae: Paederinae). *Bonn Zoological Bulletin* 61: 49–128.
- Assing V (2013) A revision of Palaearctic *Lobrathium*. IV. Three new species from Nepal and China, and additional records (Coleoptera: Staphylinidae: Paederinae). *Linzer Biologische Beiträge* 45: 191–204.
- Assing V (2014) A revision of Palaearctic *Lobrathium*. V. Three new species from the Himalaya and China, a new synonymy, and additional records (Coleoptera: Staphylinidae: Paederinae). *Linzer Biologische Beiträge* 46: 429–441.
- Li X-Y, Solodovnikov A, Zhou H-Z (2013) Four new species of the genus *Lobrathium* Mulsant et Rey from China. *Zootaxa* 3635: 569–578. doi: 10.11646/zootaxa.3635.5.6
- Li W-R, Zhao M-J, Dai C-C, Li L-Z (2013a) New species and records of *Lobrathium* Mulsant & Rey (Coleoptera, Staphylinidae, Paederinae) from China. *ZooKeys* 304: 49–81. doi: 10.3897/zookeys.304.5406
- Li W-R, Dai C-C, Li L-Z (2013b) A new species and additional records of *Lobrathium* Mulsant & Rey (Coleoptera, Staphylinidae, Paederinae) from China. *ZooKeys* 326: 47–53. doi: 10.3897/zookeys.326.5970
- Li W-R, Li L-Z (2013c) Discovery of the male of *Lobrathium rotundiceps* (Koch), and a new species of *Lobrathium* from Jiangxi, East China (Coleoptera, Staphylinidae, Paederinae). *ZooKeys* 348: 89–95. doi: 10.3897/zookeys.348.6299
- Lü Z-K, Li L-Z (2014) Two new species of *Lobrathium* Mulsant & Rey (Coleoptera, Staphylinidae, Paederinae) from China. *ZooKeys* 447: 125–131. doi: 10.3897/zookeys.447.8217
- Zheng F-K (1988) Five new species of the genus *Lobrathium* Mulsant et Rey from China (Coleoptera: Staphylinidae: Paederinae). *Acta Entomologica Sinica* 31: 186–193.