

A taxonomic account of the genus *Stenodynerus* from China, with descriptions of five new species (Hymenoptera, Vespidae, Eumeninae)

Zhen-xia Ma¹, Bin Chen¹, Ting-jing Li¹

¹ Institute of Entomology & Molecular Biology, College of Life Sciences, Chongqing Normal University, Chongqing 401331, China

Corresponding author: Ting-jing Li (ltjing1979@hotmail.com)

Academic editor: A. Köhler | Received 8 January 2016 | Accepted 24 May 2016 | Published 2 June 2016

<http://zoobank.org/97AE8A6F-C102-4D67-8B54-CFECA08B1016>

Citation: Ma Z-x, Chen B, Li T-j (2016) A taxonomic account of the genus *Stenodynerus* from China, with descriptions of five new species (Hymenoptera, Vespidae, Eumeninae). ZooKeys 595: 17–48. doi: 10.3897/zookeys.595.7734

Abstract

In this paper, 20 species of the genus *Stenodynerus* are reviewed and identified from China, including five new species: *S. ninglangensis* Ma & Li, sp. n., *S. reflexus* Ma & Li, sp. n., *Stenodynerus similibaronii* Ma & Li, sp. n., *S. strigatus* Ma & Li, sp. n., and *S. tenuilamellatus* Ma & Li, sp. n., and five new records: *S. baronii* Giordani Soika, *S. bluethgeni* van der Vecht, *S. picticrus* (Thomson), *S. pullus* Gusenleitner and *S. nepalensis* Giordani Soika. The five new species are described and illustrated in detail. Moreover, the diagnostic characters of all new records and known species from China are provided, with a key to the Chinese species of *Stenodynerus*.

Keywords

Hymenoptera, Vespidae, Eumeninae, *Stenodynerus*, new species, China, distribution

Introduction

The genus *Stenodynerus* of potter wasps was established by de Saussure (1863). This genus includes 161 species with 26 subspecies worldwide, and is distributed in the Nearctic, Neotropic, Palearctic and Oriental regions. These known species were described or revised by Bohart (1943, 1944, 1948, 1949, 1966, 1980), Buck (2008), Gusenleitner

(1981, 1985, 2004, 2008, 2001a, 2001b, 2012), Giordani Soika (1972, 1975, 1976a, 1976b, 1979, 1981, 1985, 1986, 1994), Kim and Yamane (1999, 2004), and so on. So far, ten species with two subspecies have been recorded from China (de Saussure 1863, 1867; André 1884; Morawitz 1889; Giordani Soika 1976; Kurzenko 1977; Giordani Soika 1979; Li 1985; Guseleitner 2003, 2012; Kim and Yamane 2004). In this study, a total of 20 species of *Stenodynerus* is recognized, including five new species and five newly recorded species. These five new species are described and illustrated in detail, and the diagnostic characters of new records and all known species from China provided. In addition, a key to the Chinese species of *Stenodynerus* is updated. The diagnostic characters and the key were produced based on specimen examination and the information extracted from the literature.

Materials and methods

Descriptions were made under a stereomicroscope (Olympus SZ61). Measurements were taken as the maximal length of body parts under an image analyzer (LY-M-Tupuwiew), and all figures were taken with a stereomicroscope (LY-WN-YH) attached to a computer. The ratios used throughout the descriptions were measured in the same magnification of the stereomicroscope. Body length was measured from the anterior margin of head to the posterior margin of metasomal tergum II. For the density description of punctures, “sparsely” means that the interspaces are larger than one puncture diameter, “moderately” means equal to the diameter, and “densely” means the interspaces are less than one puncture diameter. Terminology principally follows Carpenter (1982) and Kim and Yamane (2004). Specimens examined are deposited in the Institute of Entomology and Molecular Biology, Chongqing Normal University, Chongqing, China (CQNU).

Taxonomy

Stenodynerus de Saussure, 1863

Stenodynerus de Saussure, 1863: 228; Guseleitner 1981: 221; Giordani Soika 1994: 133.

Nannodynerus Blüthgen, 1938 (1937): 281.

Parhypodynerus Giordani Soika, 1973: 110.

Type species. *Odynerus chinensis* de Saussure, 1863, designated by Bohart, 1939.

Diagnosis. Body generally small and slender (Figs 1, 7, 14–15, 23, 30, 37, 63, 73); anterior surface of pronotum usually with a pair of median foveae, which sometimes contiguously forms U-shaped (Fig. 69), V-shaped (Figs 3, 18, 40, 48, 52, 57, 60, 65, 75) or a transverse fovea; tegula campanulate, broadest in the middle, length

somewhat more than width; parategula just reaching apex of tegula; tergum I generally without a basal transverse carina (Figs 5, 10, 26, 33, 50, 59, 71, 78), but in some Nearctic species present; tergum II without an acarinarium; the terminal segment of male antenna bent backward like a hook, apex usually reaching the base or middle of the segment XI (Fig. 19). This genus is similar in some characters to *Parancistrocerus* Bequaert, which can be distinguished by the presence of an acarinarium on the metasomal tergum II in *Parancistrocerus*.

Distribution. Nearctic, Neotropic, Palearctic and Oriental regions.

***Stenodynerus ninglangensis* Ma & Li, sp. n.**

<http://zoobank.org/E2130F59-2C6F-494A-891D-A0E2597FFFCD>

Figs 1–6


Material examined. Holotype, ♀, China, Yunnan Prov., Lijiang City, Ninglang County, Daxing Town, 27°16'37.68"N, 100°51'03.11"E, 2252 m, 26.VII. 2011, Tingjing Li, No. 1004068 (CQNU). Paratype: 1♀, same data as holotype, No. 1004069 (CQNU).

Description. Female (Figs 1–6): body length 7.2 mm, forewing length 5.6 mm. Black; a basal band on clypeus except median interruption (Fig. 2) and inter-antennal spot yellow; the outer margin of mandible yellow to brown; the spots pale ferruginous: ventral scape, apex of ocular sinus (Fig. 2), transverse post-ocular spot, a band on dorsal surface of pronotum, mesepisternal spot, parategula, metanotum except posterior apex ventrally, apexes of femora, all tibiae and tarsi, and apical bands on metasomal terga I–II and sternum II; tegula brown.

Head. Clypeus strongly punctate, with dense setae, its width much more than length (width 1.25 × length), apex emarginated, apical width: emargination depth = 0.20: 0.04, total width: apical width = 0.86: 0.20 (Fig. 2); frons densely punctate, punctures on vertex somewhat sparser than those on frons; frons and vertex with very sparse and short setae; cephalic fovea obsolete.

Mesosoma. Pronotum, mesoscutum and scutellum densely punctate and reticulate, punctures somewhat larger and sparser than those on the head; punctures on metanotum sparser than those on other parts of the mesosoma. Anterior surface of pronotum shining, almost vertical, with few minute punctures, median foveae contiguous and V-shaped, pronotal carina complete (Fig. 3); scutellum distinct convex; metanotum oblique; dorsal and lateral surfaces of propodeum reticulate-punctate, dorsal surface broad with a weak shelf, posterior surface concave with long and transverse rugae and a median longitudinal carina (Fig. 4).

Metasoma. In dorsal view, tergum I domed, densely punctate, width 1.31 × length and 0.85 × width of tergum II; tergum II with smaller and sparser punctures than tergum I, apex with deeper and denser punctures than other parts of tergum II, and apical margin without a distinct lamella, (Fig. 5); sternum II sparsely punctate, with a long median longitudinal furrow basally, and its anterior surface sloping (Fig. 6).



Figures 1–6. Female of *Stenodynerus ninglangensis* Ma & Li, sp. n. **1** habitus of holotype (dorsal view) **2** clypeus **3** anterior surface of pronotum **4** metanotum and propodeum **5** metasoma (dorsal view) **6** metasomal (ventral view).

Male. Unknown.

Remarks. This species is easily distinguished from all the other members of *Stenodynerus* by the following character combinations: a basal band on clypeus except median interruption yellow (Fig. 2), apex of ocular sinus ferruginous (Fig. 2), apical margin of tergum II without a distinct lamella (Fig. 5), sternum II with a long median longitudinal furrow basally, and its anterior surface sloping (Fig. 6).

Distribution. China (Yunnan).

Etymology. The specific name *ninglangensis* is the Latin adjective of the region from which the type specimens were collected.

***Stenodynerus reflexus* Ma & Li, sp. n.**

<http://zoobank.org/18D6BDFC-DD54-4DAA-8B2D-38E7B0670C63>

Figs 7–13

Material examined. Holotype, ♀, China, Yunnan Prov., Baoshan City, Tengchong County, Beihai Village, 25°16'48.82"N, 98°34'49.16"E, 1783 m, 13.VII.2011, Xin Zhou, No. 1004070 (CQNU).

Description. Female (Figs 7–13): body length 7.4 mm, forewing length 5.8 mm. Black; with the following spots ferruginous: clypeus except median interruption basally (Fig. 8), inter-antennal spot, ventral scape, post-ocular spot, a broad band on the pronotal dorsum, mesepisternal spot, parategula, metanotum except posterior apex ventrally, apical bands on metasomal terga I–II and sternum II, a long band on the dorsal surface of the fore tibia; tegula brown.


Head. Clypeus with sparse punctures, and with setae basally and laterally, its width much more than length (width $1.24 \times$ length), apex moderately emarginated, apical width: emargination depth = 0.25: 0.08, total width: apical width = 0.88: 0.25 (Fig. 8); frons densely punctate, punctures on vertex somewhat weaker than those on frons; frons and vertex with dense setae; cephalic fovea obsolete.

Mesosoma. Masosoma densely punctate and reticulate except metanotum, punctures generally larger than those on the head; punctures on metanotum sparser and shallower than those on other parts of the mesosoma. Anterior surface of pronotum slightly sloping, distinctly punctate, and with a pair of round separated median foveae, the interspace between these two median foveae less than one fovea diameter, pronotal carina complete (Fig. 9); scutellum distinctly convex; metanotum nearly vertical; dorsal and lateral surface of propodeum reticulate-punctate, dorsal surface narrow with a developed shelf, posterior surface concave with long and transverse rugae and a median longitudinal carina (Fig. 11).

Metasoma. In dorsal view, tergum I campanulate, coarsely punctate, width $1.37 \times$ length and $0.85 \times$ width of tergum II, anterior surface vertical, almost impunctate, and with a median longitudinal carina in upper half (Fig. 10); tergum II with smaller and sparser punctures than tergum I, and with a broad reflex apical lamella, dense and deep punctures forming a wide transverse groove on the base of lamella (Fig. 13); sternum II sparsely punctate, basally with a very short median longitudinal furrow, and its anterior surface almost vertical (Fig. 12).

Male. Unknown.

Remarks. This species is similar to *S. pappi* by having the propodeal shelf developed, a median longitudinal carina on propodeal concavity (Fig. 11, 81), anterior vertical surface of tergum I with a longitudinal median carina in upper half (Fig. 10). It



Figures 7–13. Female of *Stenodynerus reflexus* Ma & Li, sp. n. **7** habitus of holotype (dorsal view) **8** clypeus **9** anterior surface of pronotum **10** metasomal tergum I (dorsal view) **11** metanotum and prodeum **12** metasomal (ventral view) **13** metasoma (dorsal view).

is different from *S. pappi* and other members of the genus in the following characters: clypeus basally ferruginous except median interruption (Fig. 8), a broad band on pronotum (Fig. 9), and sternum II with a very short median longitudinal furrow basally, and its anterior surface almost vertical (Fig. 12).

Distribution. China (Yunnan).

Etymology. The specific name is the Latin adjective *reflexus*, which refers to the apical lamella of metasomal tergum II broadly reflexed.

***Stenodynerus similibaronii* Ma & Li, sp. n.**

<http://zoobank.org/3B9079BB-25FC-4436-8C61-9A2AFE5E7CC1>

Figs 14–22, 45


Material examined. Holotype, ♀, China, Yunnan Prov., Baoshan City, Tengchong County, Shangying Village, $25^{\circ}0'54.72"N$, $98^{\circ}39'12.86"E$, 1823 m, 31.VII.2015, Zhenxia Ma & Long Li, No. 1004071 (CQNU). Paratypes: 1♀♀2♂♂, the same data as holotype, Nos. 1004072, 10040743, 1004074 (CQNU).

Description. Female (Figs 14, 16, 18, 21–22, 45): body length 7.0 mm, forewing length 6.5 mm. Black; with the following spots pale ferruginous: a basal band of clypeus (Fig. 16), ventral scape, inter-antennal spot, post-ocular spot, spots on outsides of tegula anteriorly and posteriorly, parategula, apexes of femora to terminal tarsi, and apical bands on metasomal terga I–II and sternum II; a band on dorsal surface of pronotum except median interruption (Fig. 18), and metanotum except posterior apex ventrally dark ferruginous (Fig. 21); tegula brown.

Head. Clypeus with moderate punctures, lateral surface with sparse setae, its width somewhat more than length (width $1.09 \times$ length), apex slightly emarginated, apical width: emargination depth = 0.33: 0.07, total width: apical width = 1.00: 0.33 (Fig. 16); frons and vertex densely punctate and reticulate; frons with sparse and very short setae, setae on vertex denser than those on frons.

Mesosoma. Mesosoma densely punctate and reticulate; punctures generally larger than those on the head; punctures on pronotal dorsum and mesoscutum somewhat denser than those on other parts of the mesosoma (Fig. 14). Anterior surface of pronotum almost vertical with few small punctures, median foveae contiguous and V-shaped, a few short transverse carinae above median foveae, pronotal carina interrupted medially (Fig. 18); scutellum distinctly convex; metanotum oblique; dorsal and lateral surfaces of propodeum reticulate-punctate, dorsal surface broad with a weak shelf, posterior surface concave with long and transverse rugae and a median longitudinal carina (Fig. 21).

Metasoma. In dorsal view, tergum I domed, densely punctate, width $1.35 \times$ length and $0.84 \times$ width of tergum II; tergum II with smaller and sparser punctures than tergum I, apex with deeper and denser punctures than other parts of tergum II, and apical margin without a distinct lamella (Fig. 45); sternum II sparsely punctate, with a long median longitudinal furrow basally, and its anterior surface sloping (Fig. 22).



Figures 14–22. *Stenodynerus similibaronii* Ma & Li, sp. n. **14, 16, 18, 21–22** female, **15, 17, 19–20** male. **14** habitus of holotype (dorsal view) **15** habitus of paratype (dorsal view) **16** clypeus **17** clypeus **18** anterior surface of pronotum **19** antennal segment (lateral view) **20** genitalia (ventral view) **21** metanotum and propodeum **22** metasoma (ventral view).

Male (Figs 15, 17, 19, 20): body length 6.8 mm, forewing length 5.6 mm. Sculpture, punctuation, setae and coloration similar to those of female except the follows: entire clypeus, mandible except apical portion, ventral scape and inter-antennal spot yellow; clypeus strongly, convex medially, with sparse and small punctures, its width equal to length, apex deeply emarginated and U-shaped, apical width: emargination depth = 0.27: 0.12, total width: apical width = 0.8: 0.27 (Fig. 17); punctures on apex of tergum II deeper than those in female; width of tergum I $1.45 \times$ length and $0.79 \times$ width of tergum II; the terminal segment of antenna bent backward like a hook, apex reaching the base of segment XI (Fig. 19). Male genitalia as in Fig. 20, volsella with setae and slightly truncate apically, parallel spines elongate without setae, penis valve rounded apically.

Remarks. This species is similar to *S. baronii* by a basal band of clypeus (Fig. 16), metasomal sternum II with a long median longitudinal furrow basally (Fig. 22), and propodeal concavity with a median longitudinal carina (Fig. 21). It is different from *S. baronii* and other members of the genus in the following characters: anterior surface of pronotum with wider V-shaped median foveae (Fig. 18), punctures on apex of metasomal tergum II sparser (Fig. 45), and male volsella of genitalia narrower and slightly truncate apically than the corresponding parts in *S. baronii* (Figs 20, 41).

Distribution. China (Yunnan).

Etymology. The specific name *similibaronii* is a Latin adjective which refers to the similar species of *S. baronii*.

***Stenodynerus strigatus* Ma & Li, sp. n.**

<http://zoobank.org/45B4C174-8CAA-4972-8642-91F292686599>


Figs 23–29

Material examined. Holotype, ♀, China, Shaanxi Prov., Ankang City, Langao County, Huanli Town, $32^{\circ}12'52.77''$ N, $109^{\circ}0'17.90''$ E, 1808 m, 7.VIII.2015, Yan Peng & Wenkai Zhou, No. 1004075 (CQNU).

Description. Female (Figs 23–29): body length 8.3 mm, forewing length 7.7 mm. Black; the following spots yellow: a basal transverse median spot and two small obscure apical spots on clypeus (Fig. 24), inter-antennal spot, and scape ventrally; small post-ocular spot, pronotal dorsum except posterior apex, mesepisternal spot, parategula, metanotum largely (Fig. 27), apical bands on metasomal terga I–II and sternum II, and a long band on fore and mid tibiae dorsally ferruginous; tegula brown.

Head. Clypeus convex medially, moderately punctate, somewhat reticulate, with sparse and short setae, its width more than length (width $1.25 \times$ length), apex slightly emarginated, apical width: emargination depth = 0.29: 0.06, total width: apical width = 1.02: 0.29 (Fig. 24); frons and vertex densely punctate, with short setae; cephalic fovea obsolete.

Mesosoma. Masosoma densely punctate and reticulate, punctures generally larger than those on the head; punctures on pronotal dorsum denser than those on others



Figures 23–29. Female of *Stenodynerus strigatus* Ma & Li, sp. n. **23** habitus of holotype (dorsal view) **24** clypeus **25** anterior surface of pronotum **26** metasomal tergum I (dorsal view) **27** metanotum and propodeum **28** metasomal (ventral view) **29** metasoma (dorsal view).

parts of the masosoma (Fig. 23). Anterior surface of pronotum sloping, shinning, with few punctures and a pair of round separated median foveae, the interspace between these two median foveae much more than one diameter, pronotal carina complete (Fig. 25); scutellum distinctly convex; metanotum nearly vertical; dorsal and lateral surfaces of propodeum reticulate-punctate, dorsal surface narrow with a weak shelf, posterior surface concave with long and transverse rugae and a median longitudinal carina (Fig. 27).

Metasoma. In dorsal view, tergum I campanulate, coarsely punctate, width 1.58 × length and 0.81 × width of tergum II, anterior surface vertical, almost impunctate, and with a median longitudinal carina and two transverse striations (Fig. 26); tergum II with smaller and sparser punctures than tergum I, and with a broad reflex apical lamella, dense and deep punctures forming a wide transverse groove on the base of lamella (Fig. 29); sternum II sparsely punctate, without a median longitudinal furrow basally, and its anterior surface almost vertical (Fig. 28).

Male. Unknown.

Remarks. This species is similar to *S. pappi* by a median longitudinal carina on propodeal concavity (Fig. 27), anterior vertical surface of tergum I with a longitudinal median carina in upper half (Fig. 26), and tergum II with a broad reflex apical lamella (Fig. 29). It is different from *S. pappi* and other members of the genus in the following characters: a transverse median spot and two obscure apical spots on clypeus basally yellow (Fig. 24), pronotal dorsum mostly ferruginous (Fig. 25), the interspace of pronotal median foveae much more than one fovea diameter (Fig. 25), anterior vertical surface of tergum I with two transverse striations (Fig. 26), and sternum II without a median longitudinal furrow basally, and its anterior surface almost vertical (Fig. 28).

Distribution. China (Shaanxi).

Etymology. The specific name is the Latin adjective *strigatus*, which refers to the anterior vertical surface of tergum I with two transverse striations.


***Stenodynerus tenuilamellatus* Ma & Li, sp. n.**

<http://zoobank.org/3691DE51-0508-4583-913B-C873FC5E66AD>

Figs 30–36

Material examined. Holotype, ♀, China, Yunnan Prov., Baoshan City, Tengchong County, Zhonghe Village, 25°31'55.10"N, 98°23'44.21"E, 1663 m, 29.VII.2015, Zenghui Huang & Siyu Xie, No. 1004076 (CQNU). Paratype: 1♀, China, Yunnan Prov., Baoshan City, Tengchong County, Jietou Village, 25°25'11.18"N, 98°39'42.75"E, 1631 m, 15.VII.2006, Li Ma, No. 1004077 (CQNU).

Description. Female (Figs 30–36): body length 8.0 mm, forewing length 6.7 mm. Black; a minute spot on clypeus basally, inter-antennal spot, and scape ventrally yellow; with the following parategula, metanotum except posterior apex ventrally, apical bands on metasomal terga I–II and sternum II, and the dorsal surface of fore femur; tegula brown.



Figures 30–36. Female of *Stenodynerus tenuilamellatus* Ma & Li, sp. n. **30** habitus of holotype (dorsal view) **31** clypeus **32** anterior surface of pronotum **33** metasomal tergum I (dorsal view) **34** metanotum and propodeum **35** metasomal (ventral view) **36** metasoma (dorsal view).

Head. Clypeus convex medially with sparse punctures and setae, its width somewhat more than length (width $1.08 \times$ length), apex slightly emarginated, apical width: emargination depth = 0.26: 0.07, total width: apical width = 0.96: 0.26 (Fig. 31); frons and vertex densely punctate and reticulate, with short setae; cephalic fovea obsolete.

Mesosoma. Masosoma densely punctate and reticulate, punctures generally larger than those on the head; punctures on pronotal dorsum denser than those on other parts of the masosoma (Fig. 30). Anterior surface of pronotum somewhat sloping, with a few punctures and a pair of round separated median foveae, the interspace between two median foveae almost equal to one fovea diameter, pronotal carina complete (Fig. 32); scutellum distinctly convex; metanotum nearly vertical; dorsal and lateral surfaces of propodeum reticulate-punctate; dorsal surface narrow with a moderate shelf; posterior surface concave with long and transverse rugae and a median longitudinal carina (Fig. 34).

Metasoma. In dorsal view, tergum I campanulate, coarsely punctate, width $1.59 \times$ length and $0.81 \times$ width of tergum II, anterior surface vertical, almost impunctate, and with a median longitudinal carina in upper half (Fig. 33); tergum II with smaller and sparser punctures than tergum I, and with a narrow reflex apical lamella, a row of deep and dense punctures forming a narrow transverse groove on the base of lamella (Fig. 36); sternum II sparsely punctate, without a median longitudinal furrow basally, and its anterior surface vertical (Fig. 35).

Male. Unknown.

Remark. This species is similar to *S. pappi* by a median longitudinal carina on propodeal concavity (Fig. 34), and anterior vertical surface of tergum I with a longitudinal median carina in upper half (Fig. 33). It is different from *S. pappi* and other members of the genus in the following characters: clypeus with a minute spot basally (Fig. 31); propodeal shelf moderately (Fig. 34); apical lamella of metasomal tergum II distinctly narrower than that of *S. pappi* (Figs 36, 82), and sternum II without a median longitudinal furrow basally, and its anterior surface vertical (Fig. 35).

Distribution. China (Yunnan).


Etymology. The specific name *tenuilamellatus* is derived from two Latin words: *tenuis* (= narrow) and *lamellatus* (= lamella), which refers to metasomal tergum II with a narrow apical lamella.

***Stenodynerus baronii* Giordani Soika, 1975, new record**

Figs 37–44

Stenodynerus baronii Giordani Soika, 1975: 387; 1994: 133, 137.

Material examined. 1♀ 2♂, China, Tibet Autonomous Region, Nyingchi City, Medog County, Medog Town, 27.VII.2014, Tingjing Li.



Figures 37–45. *Stenodynerus baronii*, 45 *Stenodynerus similibaronii*. 37, 39–40, 42–45 female, 38, 41 male 37 habitus (dorsal view) 38 genitalia (ventral view) 39 clypeus 40 anterior surface of pronotum 41 volsella (ventral view) 42 metasomal (ventral view) 43 metanotum and propodeum 44 apex of metasomal tergum II 45 apex of metasomal tergum II.

Diagnosis. A basal band of clypeus ferruginous (Fig. 39), with moderate punctures, its width more than length; anterior surface of pronotum almost vertical, with few small punctures, median foveae small and V-shaped, a few short transverse carinae above median foveae, pronotal carina interrupted medially (Fig. 40); propodeal shelf almost obsolete, propodeal concavity with a long median longitudinal carina (Fig. 43); punctures on apex of metasomal tergum II strongly dense, deep and irregular (Fig. 44); and sternum II with a long median longitudinal furrow basally (Fig. 42); male genitalia as in Fig. 38, volsella with rounded apically, penis valve rounded apically.

Distribution. China (new record: Tibet), Bhutan.

***Stenodynerus bluethgeni* van der Vecht, 1971, new record**

Figs 46–50

Odynerus pictus Herrich-Schaeffer, 1839: 13, 32.

Odynerus dentisquama: Morawitz 1895: 462.

Euodynerus dentisquama: Blüthgen 1938 (1937): 281.

Nannodynerus dentisquama: Blüthgen 1952: 2; 1961: 107, 111; Blüthgen and Königs-mann 1969: 928.

Stenodynerus dentisquama: Giordani Soika 1970: 110.

Stenodynerus bluethgeni van der Vecht, 1971: 131; van der Vecht and Fischer 1972: 65; Gusenleitner 1981: 217, 221.

Material examined. 3♀♀1♂, China, Shaanxi Prov., Ankang City, Jianming Town, 16.VIII.2012, Xin Zhou & Cheng Yang; 1♀, China, Jilin Prov., Yanji City, Xiaoying Town, Mingzhu Village, 3.VII.2012, Ju You & Yuan Bai.

Diagnosis. Body with pale yellow spots (Figs 46–50). Cephalic fovea distinct, width less than the distance between posterior ocelli; clypeus black, medially convex (Fig. 46); anterior surface of pronotum almost vertical, with a few small punctures, median foveae V-shaped, pronotal carina complete (Fig. 48); propodeal shelf absent, propodeal concavity with a very short median longitudinal carina (Fig. 47); metasomal tergum II with very small and shallow punctures, without a distinct apical lamella (Fig. 50); sternum II black except lateral surface, without a median longitudinal furrow basally, and its anterior surface sloping (Fig. 49).

Distribution. China (new record: Jilin, Shaanxi), Netherlands, Belgium, Germany, France, Spain, Italy, Switzerland, Austria, Hungary, Czechoslovakia, Belarus, Russia, Serbia, Bulgaria, Albania, Greece, Cyprus, Turkey, Iran.


***Stenodynerus nepalensis* Giordani Soika, 1985, new record**

Figs 51–55

Stenodynerus nepalensis Giordani Soika, 1985: 37, 40; 1994: 135, 143; Gusenleitner 1987: 255; 2001b: 659.

Material examined. 5♀♀5♂♂, China, Yunnan Prov., Diqing Zang Autonomous Prefecture, Deqin County, Fushan Town, 22.VII.2014, Tingjing Li; 3♀♀5♂♂, China, Yunnan Prov., Diqing Zang Autonomous Prefecture, Deqin County, Yunling Town, 21.VII.2014, Tingjing Li.

Diagnosis. Clypeus with strong punctures, width somewhat more than length (Fig. 51); anterior surface of pronotum vertical, with few punctures, median foveae V-shaped, pronotal carina interrupted medially (Fig. 52); propodeal shelf absent (Fig. 54); metasomal tergum II without a distinct apical lamella, punctures on apex deep,



Figures 46–50. Female of *Stenodynerus bluetthgeni*. **46** clypeus **47** metanotum and propodeum **48** anterior surface of pronotum **49** metasomal (ventral view) **50** metasomal (dorsal view).

dense and irregular (Fig. 55); sternum II with a long median longitudinal furrow basally, and its anterior surface sloping (Fig. 53).


Distribution. China (new record: Yunnan), Nepal, India, Thailand.

Stenodynerus pullus Guseinleitner, 1981, new record


Figs 56–59

Stenodynerus pullus Guseinleitner, 1981: 209, 220, 246; Kim 1999: 348, 352; Kim and Yamane 2004: 240, 260–262.

Material examined. 2♀♀, China, Inner Mongolia Autonomous Region, Helan Mountain, Gulabenxiaosong Hills, 30.VII.2010, Zejian Li & Junzhe Xue; 1♂, China, Inner Mongolia Autonomous Region, Helan Mountain, Shuimogou, 27.VII.2010, Fangzhou Ma.



Figures 51–55. Female of *Stenodynerus nepalensis*. **51** clypeus **52** anterior surface of pronotum **53** metasomal (ventral view) **54** metanotum and propodeum **55** apex of metasomal tergum II.



Figures 56–59. Female of *Stenodynerus pullus*. **56** clypeus **57** anterior surface of pronotum **58** metasomal (ventral view) **59** metasomal (dorsal view).

Diagnosis. Cephalic fovea obsolete; clypeus black, medially convex, with small punctures (Fig. 56); pronotal dorsum with a pair of spots, anterior surface almost vertical, with a few small punctures, median foveae V-shaped, pronotal carina complete (Fig. 57); propodeum without shelf, propodeal concavity with a very short median longitudinal carina; metasomal tergum II without a distinct apical lamella (Fig. 59); sternum II with a very short median longitudinal furrow basally, and its anterior surface sloping (Fig. 58).

Distribution. China (new record: Inner Mongolia), Turkey, Russia, Mongolia, Korea.

***Stenodynerus tergitus* Kim, 1999, new record**

Figs 60–62

Stenodynerus tergitus Kim, 1999: 349, 352; Kim and Yamane 2004: 238–239, 250.

Material examined. 1♀3♂♂, China, Shaanxi, Prov., Weinan City, Luonan County, Mantoushan, 8.VIII.2012, Xin Zhou & Cheng Yang.

Diagnosis. Cephalic fovea small and shallow, width less than the distance between posterior ocelli; clypeus black, medially convex, its width somewhat more than length; anterior surface of pronotum sloping, with a few small punctures, median foveae V-shaped, prontoal carina complete (Fig. 60); tergum II with a broad reflex lamella apically, deep and dense punctures forming a broad transverse groove on the base of lamella (Fig. 61); sternum II with a median longitudinal furrow basally, and its anterior surface sloping (Fig. 62).

Distribution. China (new record: Shaanxi), Korea.

***Stenodynerus chinensis chinensis* (de Saussure, 1863)**


Figs 63–68

Odynerus chinensis de Saussure, 1863: 230; von Schulthess 1934: 91; van der Vecht 1967: 32; Opinion 1970: 187, 189–191.

Stenodynerus chinensis: van der Vecht and Fischer 1972: 65; Giordani Soika 1972: 105; Gusenleitner 1981: 220, 289; Li 1985: 138; Yamane and Gusenleitner 1996: 43–44.


Stenodynerus chinensis chinensis: Giordani Soika 1986: 124; Kim and Yamane 2004: 239, 256–257.

Material examined. 2♀♀, China, Henan Prov., Sanmenxia City, Lushi County, Wulichuan Town, 9.VIII.2012, Ju You & Yuan Bai; 4♀♀, China, Henan Prov., NanYang City, Yuanyang County, Sangping Town, Huangsha Village; 5♀♀1♂, China, Shaanxi Prov., Ankang City, Langao County, Huanli Town, 7.VIII.2015, Zhenxia Ma & Yan



Figures 60–62. Female of *Stenodynerus tergitus*. **60** anterior surface of pronotum **61** apex of metasomal tergum II **62** metasomal (ventral view).

Peng; 2♀♀3♂♂, China, Shaanxi Prov., Ankang City, Zhenping County, Zengjia Town, 10.VIII.2015, Zhenxia Ma & Yan Peng; 4♀♀2♂♂, China, Shaanxi Prov., Weinan City, Hua County, Jindui Village 7.VIII.2012, Xin Zhou & Cheng Yang; 3♀♀1♂, China, Shaanxi Prov., Baoji City, Mei County, Huaiya Town, 16.VIII.2015, Zhenxia Ma & Yan Peng; 1♀4♂♂, China, Shannxi Prov., Xian City, Hongqing Town, 19.VIII.2015, Zhenxia Ma & Yan Peng; 7♀♀1♂, China, Shaanxi Prov., Baoji City, Mei County, Jinqu Town, 14.VIII.2015, Zhenxia Ma & Lingquan Zeng; 14♀♀7♂♂, China, Shaanxi Prov., Baoji City, Mei County, Qinghua Town, Jinjiazhuang Village, 15.VIII.2015, Zhenxia Ma & Yan Peng; 8♀♀18♂♂, China, Sichuan Prov., Leshan City, Emeishan County, Huangwan Town, Miaoergang Village, 10.VIII.2011, Tingjing Li; 3♀♀1♂, China, Sichuan Prov., Leshan City, Emeishan County, Longchi Town, Longchi Village, 12.VIII.2011, Tingjing Li & Zhenhu Wu; 3♀♀3♂♂, China, Sichuan Prov., Leshan City, Emeishan County, Zhanggou Village, 8.VIII.2011, Tingjing Li; 12♀♀3♂♂, China, Sichuan Prov., Chongzhou City, Tianshun Village, 17.VIII.2011, Tingjing Li & Zhenhu Wu; 2♀♀1♂, China, Sichuan Prov., Leshan City, Emeishan County, Dawei Town, 13.VIII.2011, Tingjing Li; 4♀♀3♂♂, China, Chongqing Municipality, Youyang County, Banqiao Town, Shuangqiao Village, 26.VIII.2012, Cheng Yang; 6♀♀3♂♂, China, Chongqing Municipality, Jiangjin, Heiwan Village, 23.VI.2012, Xin Zhou; 1♀2♂♂, China, Chongqing Municipality, Liangping County, Bishan Town, Xinyuan Village, 5.IX.2014, Chunfa Chen; 1♀2♂♂, China, Chongqing Municipality, Chengkou County, Bashan Town, Lianmen Village, 7.VIII.2015, Tingjing Li & Chunfa Chen; 1♀2♂♂, China, Chongqing Municipality, Chengkou County, Xianyi Town, Shuangqiao Village, 4.VIII.2015, Tingjing Li & Chunfa Chen; 3♀♀2♂♂, China, Chongqing Municipality, University Town, Xiyongtuanjie, 5.VII.2011, Xin Zhou; 7♀♀11♂♂, China, Chongqing Municipality, Chongqing Normal University, 10.VII.2015, Tingjing Li & Zhenxia Ma; 3♂♂, China, Guizhou Prov., Kaili City, Leishan County, Fangxiang Town, Pingxiang Village, 24.VI.2015, Tingjing Li & Yan Peng.



Figures 63–68. Female of *Stenodynerus chinensis chinensis*. **63** habitus (dorsal view) **64** clypeus **65** anterior surface of pronotum **66** metanotum and propodeum **67** apex of metasomal tergum II **68** metasomal (ventral view).

Diagnosis. Almost whole body covered with large and dense punctures, strongly sculptured (Fig. 63). Cephalic fovea small and shallow, width less than the distance between posterior ocelli; basal half of clypeus yellow, with sparse punctures (Fig. 64); anterior surface of pronotum sloping, with distinct and strong punctures, median foveae V-shaped (Fig. 65); propodeal shelf weak, posterior surface concave, with a median longitudinal carina in lower half (Fig. 66); metasomal tergum II without a distinct apical lamella, apex with deep punctures (Fig. 67); sternum II with a long median longitudinal furrow basally, and its anterior surface sloping (Fig. 68).

Distribution. China (Hebei, Sichuan; Shannxi, Henan, Yunnan, Chongqing, Guizhou, Taiwan).

***Stenodynerus copiosus* Guseinleitner, 2012**

Stenodynerus copiosus Guseinleitner, 2012: 1132–1134.

Material examined. No specimens examined.

Diagnosis. Male: clypeus yellow, with small punctures, its width almost equal to length; pronotal carina interrupted medially; metasomal tergum II without a distinct apical lamella; sternum II with a median longitudinal furrow basally; female: unknown (Guseinleitner, 2012).

Distribution. China (Shanxi, Shaanxi).

***Stenodynerus frauenfeldi* (de Saussure, 1867)**

Figs 69–72


Odynerus frauenfeldi de Saussure, 1867: 15; von Schulthess 1934: 91; Yasumatsu 1935: 225.

Odynerus nigriclypeatus Sonan, 1930: 356; Giordani Soika 1986: 124.

Odynerus apiciornatus: Yano 1932: 309.

Stenodynerus frauenfeldi: van der Vecht and Fischer 1972: 67; Giordani Soika 1972: 105; 1986: 124; 1994: 135, 152; Guseinleitner 1981: 220, 287; Kim 1999: 348, 350; Kim and Yamane 2004: 239, 251–252.

Material examined. 1♀2♂♂, China, Jilin Prov., Yanji City, Xiaoying Town, 4.VI.2104, Ju You & Yuan Bai; 2♀♀, China, Jilin Prov., Baishan City, Linjiang County, Naozhi Town, 7.VII.2012, Ju You & Yuan Bai; 3♀♀4♂♂, China, Shannxi Prov., Huayin City, Hua Mountain, 5.VIII.2012, Xin Zhou & Ju You; 5♀♀5♂♂, China, Shannxi Prov., Ankang City, Jianming Town, 15.VIII.2012, Xin Zhou & Cheng Yang; 2♀♀3♂♂, China, Shannxi Prov., Shangluo City, Luonan County, Mantoushan, 8.VIII.2012, Ju You & Yuan Bai; 3♂♂, China, Shannxi Prov., Hanzhong City, Liuba County, Jiangkou Town, 18.VIII.2012, Xin Zhou & Cheng Yang; 42♀♀11♂♂, China, Shanxi Prov., Baoji City, Mei County, Jinqu Town, 14.VIII.2015, Zhenxia Ma & Yan Peng; 14♀♀11♂♂, China, Shannxi Prov., Baoji City, Mei County, Qinghua Town, Jinjiazhuang Village, 15.VIII.2015, Zhenxia Ma & Yan Peng; 34♀♀3♂♂, China, Shannxi Prov., Baoji City, Mei County, Huaiya Town, 16.VIII.2015, Zhenxia Ma & Yan Peng; 4♀♀1♂♂, China, Shannxi Prov., Ankang City, Zhenping County, Zengjia Town, 10.VIII.2015, Zhenxia Ma & Yan Peng; 4♂♂, China, Chongqing Municipality, Chengkou County, Xianyi Village, Tingjing Li & Chunfa Chen; 1♀1♂,



Figures 69–72. Female of *Stenodynerus frauenfeldi*. **69** anterior surface of pronotum **70** metanotum and propodeum **71** metasomal (dorsal view) **72** metasomal (ventral view).

China, Guizhou Prov., Tongren City, Jiang kou County, Minxiao Town, 29.VI.2015, Tingjing Li & Yan Peng; 1♀1♂, China, Guizhou Prov., Tongren City, Jiang kou County, Heiwan Villiage, 28.VI.2015, Tingjing Li & Yan Peng; 1♀22♂♂, China, Sichuan Prov., Dazhou City, Chengbei Town, Hongqi Village, 7.VI.2013, Ju You; 3♀♀49♂♂, China, Sichuan Prov., Dazhou City, Chengbei Town, Qingfeng Village, 4.VI.2013, Ju You; 7♀♀19♂♂, China, Sichuan Prov., Dazhou City, Maanshan, 10.VI.2013, Ju You; 3♀♀11♂♂, China, Sichuan Prov., Dazhou City, 8.VI.2013, Ju You; 17♂♂, China, Sichuan Prov., Chengdu City, Dujiangyan, Daguan Town, 16.VIII.2011, Tingjing Li & Yuan Bai; 3♀♀, China, Sichuan Prov., Leshan City, Emeishan County, Gaoqiao Town, Zhanggou Village, 8.VIII.2012, Tingjing Li; 3♀♀2♂♂, China, Sichuan Prov., Chongzhou City, Tianshun Village, 17.VIII.2011, Tingjing Li; 4♀♀8♂♂, China, Jiangxi Prov., Fuzhou City, Lichuan County, Hufang Villiage, 21.VI.2014, Tingjing Li.

Diagnosis. Body strongly sculptured, with large and strong punctures, and extensive spots (Figs 69–72). Cephalic fovea almost obsolete; anterior surface of pronotum vertical, median foveae shallow and slightly U-shaped, a few long parallel transverse carinae above median foveae, pronotal carina complete (Fig. 69); propodeal shelf weak (Fig. 70); metasomal tergum II without a distinct apical lamella (Fig. 71); median longitudinal furrow on metasomal sternum II basally weak, even obsolete, and its anterior surface sloping (Fig. 72).

Distribution. China (Sichuan, Jilin, Shaanxi, Chongqing, Guizhou, Jiangxi), Russia, Korea, Japan.

***Stenodynerus funebris* (André, 1884)**

Figs 73–76

Odynerus funebris André, 1884: 729; Blüthgen 1955: 412.

Odynerus limbonatatus: Kostylev, 1940: 28.

Stenodynerus funebris: van der Vecht and Fischer 1972: 67; Guseleitner 1981: 219, 243; Kim 1999: 347, 349, 350; Kim and Yamane 2004: 239, 255.

Material examined. 1♀1♂, China, Jilin Prov., Yanji City, Xiaoying Town, Minzhu Village, 3.VII.2012, Xin Zhou & Ju You.

Diagnosis. Body obviously black (Fig. 73), only with the following spots yellow: mandibular basally, scape ventrally, inter-antennal spot, a minute post-ocular spot, pronotum sometimes with a pair of small spots, apical bands on metasomal terga I–II and sternum II, a long band on fore tibiae dorsally; cephalic fovea small and shallow, width less than the distance between posterior ocelli; clypeus medially convex, apex moderately emarginated (Fig. 74); anterior surface of pronotum almost vertical, with V-shaped median foveae, pronotal carina complete (Fig. 75); propodeal shelf developed, posterior surface concave, with a very short median longitudinal carina in lower portion (Fig. 76); median longitudinal furrow on metasomal sternum II basally weak, even obsolete, and its anterior surface sloping.

Distribution. China (Shanxi, Jilin), Korea, Russia.

***Stenodynerus incurvitus* Guseleitner, 2003**

Stenodynerus incurvitus Guseleitner, 2003: 855.

Material examined. No specimens examined.

Diagnosis. Clypeus yellow basally, with short setae, its width equal to its length; cephalic fovea comparatively larger, width more than the distance between posterior ocelli; propodeal concavity almost smooth; metasomal tergum II without a distinct apical lamella, punctures on apex deep; sternum II with a median longitudinal furrow basally, and its anterior surface sloping (Guseleitner 2003).


Distribution. China (Taiwan).

***Stenodynerus morawitzi* Kurzenko, 1977**

Stenodynerus morawitzi Kurzenko, 1977: 554; Guseleitner 1981: 219, 266.

Material examined. No specimens examined.

Diagnosis. Clypeus yellow, with small punctures, its width somewhat more than length, apex deeply emarginated; pronoal carina interrupted medially; tergum II with



Figures 73–76. Female of *Stenodynerus funebris*. **73** habitus (dorsal view) **74** clypeus **75** anterior surface of pronotum **76** metanotum and propodeum.

a broad reflex lamella apically. In male, metasomal terga I–VII and sterna II–III with yellow apical bands (Gusenleitner 1981).

Distribution. China (Inner Mongolia).

Stenodynerus morbillosus Giordani Soika, 1979

Stenodynerus morbillosus Giordani Soika, 1979: 250; Gusenleitner 1981: 220, 286.

Material examined. No specimens examined.

Diagnosis. Larger species, body length generally more than 10 mm. Clypeus yellow; pronotal carina less developed; metasomal tergum II without a distinct apical lamella; sternum II with a short median longitudinal furrow basally (Giordani Soika 1979; Gusenleitner 1981).

Distribution. China (northeast).

***Stenodynerus nudus* (Morawitz, 1889)**

Lionotus nudus Morawitz, 1889: 164; Morawitz 1895: 463; von Schulthess 1934: 91.

Nannodynerus nudus: Blüthgen 1942: 320.

Stenodynerus nudus: van der Vecht and Fischer 1972: 68; Guseinleitner 1981: 219, 271.

Material examined. No specimens examined.

Diagnosis. Clypeus yellow except margin, with sparse and shallow punctures, its width much more than its length; metasomal tergum II without a distinct apical lamella; anterior surface of metasomal sternum II vertical. Metasomal terga I–V and sterna II–III with yellow apical bands (Guseinleitner 1981).

Distribution. China (Inner Mongolia), Turkmenistan, Kazakhstan, Mongolia.

***Stenodynerus pappi luteifasciatus* Kim & Yamane, 2004**

Stenodynerus pappi luteifasciatus Kim & Yamane, 2004: 235, 238, 245.

Material examined. No specimens examined.

Diagnosis. The species differs from *S. p. pappi* as follows: body with extensive orange-yellow spots; a broad band on pronotal dorsum and almost entire metanotum orange-yellow; punctures on anterior surface of pronotum sparser, apical band on tergum I broader and reflex apical lamella of metasomal tergum II narrower than the corresponding parts in *S. p. pappi* (Kim and Yamane 2004).

Distribution. China (Taiwan).

***Stenodynerus pappi pappi* Giordani Soika, 1976**

Figs 77–82


Stenodynerus pappi Giordani Soika, 1976: 290–291; Guseinleitner 1981: 219, 298;

Kim 1999: 204; Kim and Yamane 2004: 237, 242.

Parancistrocerus ussuriensis Kurzenko, 1982 (1981): 117–122.

Material examined. 2♀♀, China, Shaanxi prov., Baoji City, Taibai County, Taochuan town, 12.VIII.2015, Zhenxia Ma & Yan Peng; 1♀, China, Shaanxi prov., Yanan City, Huanglong County, Shibao Town, 2.VIII.2012, Xin Zhou; 1♀, China, Shaanxi prov., Weinan City, Hua County, Jindui Town, 7.VIII.2012, Ju You & Yuan Bai; 1♀, China, Chongqing, Shizhu County, Huangshui Town, 12.VIII.2008, Bin Chen & Tingjing Li.

Diagnosis. Cephalic fovea small and shallow, width less than the distance between posterior ocelli; clypeus black, sparsely punctate, with sparse setae, its width more than length (Fig. 77); anterior surface of pronotum sloping, with distinct punctures and



Figures 77–82. Female of *Stenodynerus pappi pappi*. **77** clypeus **78** metasomal tergum I (dorsal view) **79** metasomal (ventral view) **80** anterior surface of pronotum **81** metanotum and propodeum **82** apex of metasomal tergum II.

a pair of round contiguous median foveae, the interspace between these two median foveae less than one diameter, pronotal carina obsolete (Fig. 80); propodeal shelf developed, posterior surface concave, with a long median longitudinal carina (Fig. 81); anterior surface of metasomal tergum I vertical, almost impunctate, and with a median longitudinal carina in upper half (Fig. 78); metasomal tergum II with a broad reflex lamella apically, dense and deep punctures forming a wide transverse groove on the base of lamella (Fig. 82); sternum II with a short median longitudinal furrow basally, and its anterior surface sloping (Fig. 79).

Distribution. China (Zhejiang, Jiangxi, Shaanxi, Chongqing, Taiwan), Korea.

Stenodynerus taiwanus Kim & Yamane, 2004

Stenodynerus taiwanus Kim & Yamane, 2004: 235, 237, 241.

Material examined. No specimens examined.

Diagnosis. Anterior surface of pronotum distinctly punctate, with a pair of round contiguous median foveae, pronotal carina weak; propodeal shelf absent, posterior surface

concave, with a median longitudinal carina lost in upper half; anterior surface of metasomal tergum I almost impunctate, dorsal surface with weak and sparse punctures, the interspaces between punctures more than one diameter. Whole body with sparse and long setae (Kim and Yamane, 2004).

Distribution. China (Taiwan).

Key to the Chinese species of the genus *Stenodynerus*

The characters are applicable to both sexes unless the sex is specified.

- 1 Metasomal tergum II with a lamella apically (Figs 13, 29, 36, 61, 82) 2
- Metasomal tergum II without a lamella apically (Figs 5, 44, 45, 50, 55, 59, 67, 71) 8
- 2 Anterior surface of metasomal tergum I vertical, with few punctures and a median longitudinal carina in upper half (Figs 10, 26, 33, 78) 3
- Anterior surface of metasomal tergum I with strong punctures, not vertical and somewhat rounded, without a median longitudinal carina 7
- 3 Metasomal tergum II with a broad reflex lamella apically, the transverse groove on the base of lamella wider (Figs 13, 29, 82) 4
- Metasomal tergum II with a narrow reflex lamella apically, the transverse groove on the base of lamella narrower (Fig. 36) *S. tenuilamellatus* sp. n.
- 4 In female, clypeus black (Fig. 77); in profile, anterior surface of metasomal sternum II sloping (Fig. 79) *S. pappi*, 5
- In female, clypeus with yellow or ferruginous spots (Figs 8, 24); in profile, anterior surface of metasomal sternum II vertical (Figs 12, 28) 6
- 5 Body with yellow spots; pronotum with a pair of small spots dorsally (Fig. 80), metanotum with a yellow band anteriorly (Fig. 81), tergum I with a narrower apical band (Fig. 78) *S. pappi pappi* Giordani Soika
- Body with orange yellow spots; pronotal dorsum with a broad band, metanotum almost entire orange yellow, tergum I with a broader apical band (Kim and Yamane 2004) *S. pappi luteifasciatus* Kim & Yamane
- 6 Anterior surface of pronotum with distinct punctures, the interspace between median foveae less than one fovea diameter (Fig. 9); propodeal shelf developed (Fig. 11); anterior vertical surface of tergum I without a transverse striation (Fig. 10); sternum II with a very short median longitudinal furrow basally (Fig. 12) *S. reflexus* sp. n.
- Anterior surface of pronotum with few punctures, the interspace between median foveae more than one fovea diameter (Fig. 25); propodeal shelf weak (Fig. 27); anterior vertical surface of tergum I with two transverse striations (Fig. 26); sternum II without a median longitudinal furrow basally (Fig. 28) *S. strigatus* sp. n.
- 7 In female, clypeus yellow; in male, metasomal terga I–VII and sterna II–III with yellow apical bands (Giordani Soika 1981) *S. morawitzi* Kurzenko

- In female, clypeus black; in male, metasomal terga I–II and sternum II with yellow apical bands *S. tergitus* Kim (new record) **8**
- 8 Larger species, body length generally more than 10 mm (Giordani Soika 1979; Giordani Soika 1981) *S. morbillosus* Giordani Soika **9**
- Smaller species, body length generally less than 10 mm **9**
- 9 Apical bands of metasomal terga I–V yellow or orange-yellow **10**
- Apical bands of metasomal terga I–II yellow or ferruginous **11**
- 10 In profile, anterior surface of metasomal sternum II sloping (Fig. 53)
..... *S. nepalensis* Giordani Soika (new record) **11**
- In profile, anterior surface of metasomal sternum II vertical (Gusenleitner 1981)..... *S. nudus* (Morawitz) **11**
- 11 Propodeal concavity almost smooth (Gusenleitner 2003)
..... *S. incurvitus* Gusenleitner **11**
- Propodeal concavity with long and transverse rugae..... **12**
- 12 Anterior surface of pronotum with a pair of round contiguous median foveae; anterior surface of metasomal tergum I almost impunctate (Kim and Yamane 2004)..... *S. taiwanus* Kim & Yamane **12**
- Pronotal median foveae contiguous and forming U-shaped or V-shaped; anterior surface of metasomal tergum I with strong punctures (Figs 5, 59, 71) **13**
- 13 Pronotal median foveae shallow and slightly U-shaped (Fig. 69)....
..... *S. frauenfeldi* (de Saussure) **13**
- Pronotal median foveae distinct and V-shaped (Figs 3, 18, 40, 48, 57, 65, 75) **14**
- 14 Clypeus with yellow or ferruginous spots; metasomal sternum II with a long median longitudinal furrow basally..... **15**
- Clypeus black; metasomal sternum II with a very short median longitudinal furrow basally, sometimes obsolete **18**
- 15 Body with large and dense punctures, strongly sculptured (Fig. 63); anterior surface of pronotum sloping, and with distinct and strong punctures (Fig. 65)..... *S. chinensis chinensis* (de Saussure, 1863) **15**
- Body with comparatively smaller punctures, moderately sculptured; anterior surface of pronotum almost vertical, with few punctures (Figs 3, 18, 40)... **16**
- 16 Clypeus strongly punctate (Fig. 2); apex of ocular sinus pale ferruginous; pronotal carina complete *S. ningliangensis* sp. n. **16**
- Clypeus moderately punctate (Figs 16, 39); apex of ocular sinus black; pronotal carina interrupted medially **17**
- 17 Anterior surface of pronotum with wider V-shaped median foveae (Fig. 18); apex of metasomal tergum II with somewhat shallower and sparser punctures (Fig. 45); in male, volsella slightly truncate apically (Fig. 20).... *S. similibaronii* sp. n. **17**
- Anterior surface of pronotum with narrower V-shaped median foveae (Fig. 40); apex of metasomal tergum II with strongly deep and dense punctures (Fig. 44); in male, volsella with rounded apically (Figs 38, 41)
..... *S. baronii* Giordani Soika (new record) **17**

- 18 Body with pale yellow spots; metasomal sternum II black except lateral surface (Fig. 49); metasomal tergum II slightly punctate (Fig. 50); sternum II without a median longitudinal furrow basally (Fig. 49).....
..... *S. bluethgeni* van der Vecht (new record)
- Body with yellow spots; metasomal sternum II with a narrow apical band (Fig. 58); metasomal tergum II strongly punctate (Figs 59, 73); sternum II with a very short median longitudinal furrow basally (Fig. 58) 19
- 19 Anterior surface with a few short transverse carinae above median foveae (Fig. 75); propodeal shelf developed (Fig. 76) *S. funebris* (André)
- Anterior surface without a transverse carina above median foveae (Fig. 57); propodeal shelf obsolete..... *S. pullus* Gusenleitner (new record)

Acknowledgements

We are very grateful to Prof. James M. Carpenter (American Museum of Natural History, New York, USA) for providing copies of many references and important comments and to Dr. P. Girish Kumar (Hymenoptera Section, Zoological Survey of India, Kolkata, India) for copies of many references. We also express our hearty thanks to Prof. Meicai Wei (Central South University of Forestry and Technology, Changsha, China) and Prof. Qiang Li (Yunnan Agricultural University, Kunming, China) for providing the specimens deposited in their Insect Collection. This study was funded by the National Natural Science Foundation of China (Nos: 31372247, 31000976, 31372265), Chongqing Science and Technology Commission (cstc2013jcyjA80015), Young Talent Incubation Program of Chongqing Normal University (14CSDG07), and the Par-Eu Scholars Program.

References

- André E (1884) Family Eumenidae. In species des Hyménoptères d'Europe et d'Algérie 2: 608–810.
- Blüthgen P (1938) Systematisches Verzeichnis der Faltenwespen Mitteleuropas, Skandinaviens und Englands. Konowia, Vienna 16: 270–295.
- Blüthgen P (1942) Neue palaark-tischen Faltenwespen (Hymenoptera: Vespidae: Eumenidinae). Mitteilungen aus dem Zoologischen Museum in Berlin 25: 300–322.
- Blüthgen P (1952) Weitere neue oder bemerkenswerte palaarktische Faltenwespen aus der Zoolo-gischen Staatssammlung in München (Hymenoptera: Eumenidae, Masaridae). Mitteilungen der Muenchener Entomologischen Gesellschaft 42: 1–19.
- Blüthgen P (1955) Weitere Beiträge zur Synonymie der palaarktischen Faltenwespen (Hyme-noptera: Vespidae). Mitteilungen der Münchner Entomologischen Gesellschaft, Munich 44–45: 397–417.
- Blüthgen P (1956) 5. Beitrag zur Synonymie der palaarktischen Faltenwespen (Hymenoptera: Vespidae). Deutsche Entomologische Zeitschrift Berlin (N.F.) 3: 70–73.

- Blüthgen P (1961) Die Faltenwespen Mitteleuropas. Abhandlungen der Deutschen Akademie der Wissenschaften zu Berlin, Klasse für Chemie, Geologie und Biologie 2: 1–252.
- Blüthgen P, Konigsmann E (1969) Ergebnisse der Albanien-Expedition 1961 des Deutschen Entomologischen Institutes. 79. Beitrag, Hymenoptera: Vespidae, Eumenidae, Masaridae. Beiträge zur Entomologie, Berlin 19: 917–935.
- Bohart RM (1943) North American *Stenodynerus* of the *anormis* group (Hymenoptera: Vespidae). Bulletin of the Brooklyn Entomological Society 38: 6–11.
- Bohart RM (1944) *Stenodynerus fundatus* and related species in North America (Hymenoptera: Vespidae). Pan-Pacif Entomologist, San Francisco 20: 69–75.
- Bohart RM (1948) The red *Stenodynerus* of Florida (Hymenoptera: Vespidae). Florida Entomologist 31: 71–80. doi: 10.2307/3492654
- Bohart RM (1949) The genus *Stenodynerus* in southern Arizona (Hymenoptera: Vespidae). Proceedings Biological Society Washington 51: 237–259.
- Bohart RM (1966) New *Stenodynerus* from North America (Hymenoptera: Eumenidae). Proceedings Biological Society Washington 79: 73–81.
- Bohart RM (1980) The middle American species of *Stenodynerus* (Hymenoptera: Eumenidae). Polskie Pismo Entomologiczne 50(1): 71–108.
- Buck M, Marshall SA, Cheung DKB (2008) Identification atlas of the Vespidae (Hymenoptera: Aculeata) of the northeastern Nearctic region. Canadian Journal of Arthropod Identification 5: 1–492. doi: 10.3752/cjai.2008.05
- Carpenter JM (1982) The phylogenetic relationships and natural classification of the Vespoidea (Hymenoptera). Systematic Entomology 7(1): 11–38. doi: 10.1111/j.1365-3113.1982.tb00124.x
- Giordani Soika A (1972) Notulae vespidiologicae 32. Nuovi Eumenidi indomalesi. Bollettino della Società Entomologica Italiana 104: 99–110.
- Giordani Soika A (1973) Descrizione di nuovi Eumenidi. Bollettino del Museo Civico di Storia Naturale di Venezia 24: 97–131.
- Giordani Soika A (1974) Sul genere Gribodia Zav. e descrizione di nuove specie (Hymenoptera: Eumenidae). Bollettino della Società Entomologica Italiana 106(5–7): 107–112.
- Giordani Soika A (1975) Ergebnisse der Bhutan-Expedition 1972 des Naturhistorischen Museums in Basel. Hymenoptera: Fain. Eumenidae. Entomologica Basiliensia 1: 387–393.
- Giordani Soika A (1976a) Nuovi Eumenidi Palearctici. Bollettino del Museo Civico di Storia Naturale di Venezia 28: 153–178.
- Giordani Soika A (1976b) Vespidi ed Eumenidi (Hymenoptera) raccolti in Mongolia dal Dr. Z. Kaszab. Acta Zoologica Hungarica 22(3–4): 271–276.
- Giordani Soika A (1979) Notulae vespidiologicae 40. Eumenidi palearctici nuovi o poco noti. Bollettino del Museo Civico di Storia Naturale di Venezia 30: 247–263.
- Giordani Soika A (1981) Notulae vespidiologicae 41. Nuovi Subancistrocerus del pacifico (Hymenoptera). Bollettino della Società Entomologica Italiana 113(8–10): 169–171.
- Giordani Soika A (1985) Notulae Vespidiologicae 44. Descrizione di un nuovo genere e di due nuove specie di Eumenidi Asiatici (Hymenoptera). Società Veneziana di Scienze Naturali Lavori 10: 37–41.

- Giordani Soika A (1986) Eumenidi Paleartici nuovi o poco noti. Bollettino del Museo Civico di Storia Naturale di Venezia 35: 91–162.
- Giordani Soika A (1994) Ricerche sistematiche su alcuni generi di eumenidi della regione orientale e della Papuasia (Hymenoptera: Vespoidea). Annali del Museo Civico di Storia Naturale “Giacomo Doria” 90: 1–348.
- Gusenleitner J (1981) Revision der palaarktischen *Stenodynerus*-Arten (Hymenoptera: Eumenidae). Polskie Pismo Entomologiczne 51(2): 209–305.
- Gusenleitner J (1985) Bemerkenswertes über Faltenwespen 8. (Hymenoptera: Vespoidea). Nachrichtenblatt der Bayerischen Entomologen 34(4): 105–110.
- Gusenleitner J (1987) Über Eumenidae aus Nepal (Hymenoptera: Vespoidea). Linzer Biologische Beiträge 19(1): 255–270.
- Gusenleitner J (2001a) Contribution to the knowledge of the vespid and eumenid wasps of the Oriental Region (Hymenoptera: Vespidae, Eumenidae). Linzer Biologische Beiträge 33(2): 655–662.
- Gusenleitner J (2001b) The *Stenodynerus* species from Asia (Hymenoptera: Vespoidea: Eumenidae). Linzer Biologische Beiträge 33(2): 651–653.
- Gusenleitner J (2004) A contribution to the knowledge and distribution of Palaearctic Eumenidae (Hymenoptera: Vespoidea). Linzer Biologische Beiträge 36(2): 1093–1105.
- Gusenleitner J (2008) One new species of *Stenodynerus* from Kyrgyzstan. Linzer Biologische Beiträge 40(2): 1491–1493.
- Gusenleitner J (2012) Bemerkenswertes über Faltenwespen XII (Hymenoptera: Vespidae: Eumeninae). Linzer Biologische Beiträge 44(2): 1129–1136.
- Herrick-Schaeffer (1839) Fauna Insectorum Germaniae 173.
- Kim JK (1999) Taxonomic review of the genus *Stenodynerus* Saussure (Hymenoptera: Vespidae: Eumeninae) with description of a new species in Korea. Korean Journal of Biological Sciences 3(4): 347–354. doi: 10.1080/12265071.1999.9647506
- Kim JK, Yamane S (2004) Revision of the genus *Stenodynerus* Saussure (Hymenoptera: Vespidae: Eumeninae) in the Far East, with descriptions of a new species and a new subspecies from Taiwan. Japanese Journal of Systematic Entomology 10(2): 235–264.
- Kostylev G (1940) Espèces nouvelles et peu connues de Vespides, d’Euménides et de Masarides paléarctiques (Hymenoptera). I. Bulletin de la Société des naturalistes de Moscou, Section biologique (NS) 49: 137–154.
- Kurzenko NV (1977) Eumenid wasps (Hymenoptera: Eumenidae) of the Mongolian People’s Republic and adjacent regions of China and southern Siberia. Nasekomye Mongolii = Insects of Mongolia 5: 537–582.
- Kurzenko NV (1981) Survey of some wasp genera of the family Eumenidae (Hymenoptera: Vespoidea) of the USSR fauna. Akademiya Nauk SSSR, Vladivostok: 81–112.
- Li Ts (1985) Economic insect fauna of China. Fasc. 30. Hymenoptera: Vespoidea. Economic Insect Fauna of China, Science Press, Beijing, China, i–vii, 1–159.
- Morawitz (1889) Insecta a Cl. G. N. Potanin in China et in Mongolia Novissime Lecta. IV. Hymenoptera Aculeata. Horae Societatis Entomologicae Rossicae 23: 112–168.
- Morawitz (1895) Materialien zu einer Vespidenfauna des Russischen Reiches. Horae Societatis Entomologicae Rossicae 29: 407–493.

- Opinion (1970) Eumenidae names of Saussure (Hymenoptera): grant of availability to certain names proposed for secondary divisions of genera. Bulletin of Zoological Nomenclature 26: 187–191.
- Saussure HF De (1863) Mélanges Hyménoptérologiques II. Mémoires de la Société de Physique et d'Histoire Naturelle de Genève 17: 1–3.
- Saussure HF De (1867) Reise der Oesterreichischen Fregatte Novara um die Erde. Zoologie 2: 1–156.
- Schlüth A (1934) Zur Kenntnis der *Odynerus*-Arten der Japaischen Subregion. Arbeiten über morphologische und taxonomische Entomologie aus Berlin-Dahlem 1: 66–103.
- Sonan J (1930) Some new species of Hymenoptera in Japanese Empire with two known species. Transactions of the Natural History Society of Formosa 20: 355–366.
- van der Vecht J (1967) The status of certain genus-group names in the Eumenidae (Hymenoptera: Vespoidea) Z.N.(S.) 1689. Bulletin of Zoological Nomenclature 24: 27–33. doi: 10.5962/bhl.part.15358
- van der Vecht J (1971) Notes on palaearctic Eumenidae (Hymenoptera). Entomologische Berichten (Amsterdam) 31: 127–134.
- van der Vecht J, Fischer FCJ (1972) Palaearctic Eumenidae. Hymenopterum Catalogus (nova editio) 8: 1–199.
- Yamane S, Tano T (1987) Studies on Japanese Eumenidae (Hymenoptera: Vespoidea). Transactions of the Shikoku Entomological Society 18: 327–345.
- Yano M (1932) Vespidae. In: Uchida S (Ed.) Iconographia Insectorum Japonicorum. 1st. ed. Hokuryûkan, Tokyo, 306–312.