

Taxonomic notes on the genus *Phrynarachne* from China (Araneae, Thomisidae)

Yejie Lin¹, Long Yu², Peter Koomen³, Xunyou Yan¹, Shuqiang Li⁴

1 Hebei Key Laboratory of Animal Diversity, College of Life Science, Langfang Normal University, Langfang 065000, China **2** State Key Laboratory of Biocatalysis and Enzyme Engineering of China & Centre for Behavioural Ecology & Evolution, School of Life Sciences, Hubei University, Wuhan 430062, China **3** Natuur museum Fryslân, Schoenmakersperk 2, Leeuwarden, 8911 EM, The Netherlands **4** Institute of Zoology, Chinese Academy of Sciences, Beijing 100101, China

Corresponding authors: Xunyou Yan (yanxunyou@163.com), Shuqiang Li (lisq@ioz.ac.cn)

Academic editor: Zhiyuan Yao | Received 13 November 2021 | Accepted 19 January 2022 | Published 4 February 2022

<http://zoobank.org/DA1FA04C-FB0F-4325-BDAB-507FF6DEB967>

Citation: Lin Y, Yu L, Koomen P, Yan X, Li S (2022) Taxonomic notes on the genus *Phrynarachne* from China (Araneae, Thomisidae). ZooKeys 1085: 69–99. <https://doi.org/10.3897/zookeys.1085.77966>

Abstract

Four new species of the genus *Phrynarachne* Thorell, 1869 from China are described: *P. deepy* Lin & S. Li, **sp. nov.** (♂♀), *P. xuxiake* Lin & S. Li, **sp. nov.** (♀), *P. yunhui* Lin & S. Li, **sp. nov.** (♀), and *P. zhengzhongi* Lin & S. Li, **sp. nov.** (♀). The unknown sexes of *P. brevis* Tang & S. Li, 2010 (♂), *P. huangshanensis* Li et al., 1985 (♀), *P. lancea* Tang & S. Li, 2010 (♂), and *P. mammillata* Song, 1990 (♀) are described for the first time. *Phrynarachne sinensis* Peng, Yin & Kim is treated as a nomen dubium.

Keywords

Diagnosis, new species, nomen dubium, type specimens

Introduction

The spider genus *Phrynarachne* Thorell, 1869 currently includes 32 species and subspecies distributed in southern Asia, the Australian region, and sub-Saharan Africa. Only five species are described by both sexes, and 10 species have been studied after their original description. Efforts have been made to find *Phrynarachne* types preserved in well-known European museums, but these endeavors have failed.

Seven *Phrynarachne* species were known from China before the current study; only two species, i.e., *P. ceylonica* (O. Pickard-Cambridge, 1884) and *P. katoi* Chikuni, 1955, are described by both sexes. All endemic Chinese *Phrynarachne* species are only described by few single-sex specimens, and the species in the surrounding areas of China, except Japan, are not well revised and most of them have only initial descriptions (Li et al. 2021; WSC 2021; Yao et al. 2021).

Here, we describe four new and six known *Phrynarachne* species from China. Due to the lost holotype and unknown locality in the original description, we treat *P. sinensis* Peng et al. as *nomen dubium*.

Materials and methods

All specimens were preserved in 80% ethanol. Epigynes were cleared in trypsin enzyme solution to dissolve non-chitinous tissues. Specimens were examined under a LEICA M205C stereomicroscope. Photomicroscopy images were taken with an Olympus C7070 zoom digital camera (7.1 megapixels). Laboratory habitus photographs were taken with a Sony A7RIV digital camera equipped with a Sony FE 90mm Goss lens. Photos were stacked with Helicon Focus (v. 7.6.1) or Zerene Stacker (v. 1.04) and processed in Adobe Photoshop CC2019.

All measurements are in millimeters and were obtained with an Olympus SZX16 stereomicroscope with a Zongyuan CCD industrial camera. Total length is measured without chelicerae. Eye sizes are measured as the maximum diameter from either the dorsal or frontal view. Leg measurements are given as follows: total length (femur, patella, tibia, metatarsus, tarsus). The terminology used in the text and figures follows Ono (1988).

Types of the new species reported here are deposited at the Institute of Zoology, Chinese Academy of Sciences in Beijing.

Abbreviations

ALE	anterior lateral eyes;	Mp	median plate;
AME	anterior median eyes;	PLE	posterior lateral eyes;
E	embolus;	PME	posterior median eyes;
FD	fertilization duct;	RTA	retrolateral tibial apophysis;
H	hood;	S	spermathecae;
ITA	intermediate tibial apophysis;	VTA	ventral tibial apophysis.

Taxonomy

Family Thomisidae Sundevall, 1833

Genus *Phrynarachne* Thorell, 1869

Phrynarachne Thorell, 1869: 37. For the complete list of references see WSC (2021).

Type species. *Thomisus rugosus* Walckenaer, 1805, from Mauritius

Diagnosis. Large or medium-sized, male is much smaller than the female (1:2 or more). Prosoma nearly as long as wide, with granulations. Eyes small, subequal in size. Fovea inconspicuous. Chelicerae with two promarginal and one retromarginal teeth. Labium longer than wide, sternum oval, male palp with VTA, ITA and RTA; tegulum flat, disk-shaped; tegular ridge present; embolus slender. Female epigynum simple, with a media plate, spermathecae strong sclerotized.

Phrynarachne brevis Tang & S. Li, 2010

Figs 1A, 6, 18A, 21

Phrynarachne brevis Tang & Li, 2010: 49, figs 35A–D, 36A, B (♂).

Type material. Holotype: ♂ (IZCASAr18535), **China: Yunnan:** Xishuangbanna, Mengla County, Menglun Town, Menglun Nature Reserve, Bamboo plantation near G213 roadside, 21.8940°N, 101.2823°E, 580 m elev., 3.XII.2009, Guo Tang and Zhiyuan Yao leg., examined.

Other material examined. 1♀ (IZCAS-Ar41642), **China: Yunnan:** Xishuangbanna, Jinghong City, Mengla County, Menglun Town, Menglun Nature Reserve, Bamboo plantation, 21.9008°N, 101.2822°E, 597 m elev., 9.V.2019, Zhigang Chen leg.

Diagnosis. See diagnosis of *P. dreepy* sp. nov.

Description. Female (Figs 1A, 6, 18A). Total length 16.53, carapace 6.62 long, 7.09 wide, yellow brown with brown pattern and granulations dorsally. With large projection between ALE and PLE. Eye sizes and interdistances: ALE 0.26, AME 0.23, PLE 0.25, PME 0.22; ALE–AME 0.51, AME–AME 0.90, PLE–PME 1.62, PME–PME 1.02. Chelicerae brown, with two promarginal and one retromarginal teeth; gnathocoxae, labium dark yellow, labium 1.52 long, 1.18 wide. Sternum yellow. Legs brown, femora I and II with dense, varying-sized tubercles; tibiae and metatarsi I, II with dense asymmetrical ventral spines (I, tibia 12, metatarsus 30; II, tibia 10, metatarsus 30). Leg measurements: I 17.25 (5.74, 6.31, 3.10, 2.10), II 17.61 (5.66, 6.61, 3.30, 2.04), III 12.30 (3.67, 5.74, 1.50, 1.39), IV 10.26 (3.31, 4.12, 1.46, 1.37). Opisthosoma dorsally light yellow, each side with 22 long tubercles, middle with pair of black markings.



Figure 1. *Phrynarachne* spp., live **A** *P. brevis*, adult female **B** *P. xuxiake* sp. nov., juvenile. Photos by Chao Wu (**A**) and Fan Gao (**B**).

Epigyne (Fig. 6) with M-shaped sclerotized margins; median plate obvious, with a posterior hood, anterior edge recurved and posterior edge almost straight, the ratio of length to width is 11:3; copulatory opening obvious; spermathecae kidney-shaped, the ratio of anterior edge to posterior edge length is 1:1. Fertilization duct transverse.

Male. See Tang and Li (2010).

Distribution. China (Yunnan).

Notes. The female is described here for the first time.

***Phrynarachne ceylonica* (O. Pickard-Cambridge, 1884)**

Figs 2, 21

Ornithoscatoides ceylonica O. Pickard-Cambridge, 1884: 201, pl. 15, fig. 3. For the complete list of references see WSC (2021).

Type material. *Syntypes* 2♀, “Ceylon, G.H.K. Thwaites leg.”, Hope Department of Entomology, Oxford, UK, not examined; *O. nigra* O. Pickard-Cambridge, 1884: *Syntypes* 2♂, “Ceylon and India, G.H.K. Thwaites leg.”, Hope Department of Entomology, Oxford, UK, not examined.



Figure 2. *Phrynarachne ceylonica* **A** adult male (left) and female (right) **B** male live **C** female live. Photos by Peter Koomen.

Other material examined. 2♂ (IZCAS), Xishuangbanna, Jinghong City, Mengla County, Menglun Town, Menglun Nature Reserve, primary tropical seasonal rain forest, 21.9598°N, 101.2035°E, 822 m elev., 8.VIII.2007, Guo Zheng leg.; 1♂ (IZCAS), Xishuangbanna, Jinghong City, Mengla County, Menglun Town, Menglun Nature Reserve, *Paramichelia baillonii* plantation (about 20 years old), 21.9129°N, 101.2674°E, 556 m elev., 18.VII.2007, Guo Zheng leg.; 2♂ (IZCAS), Xishuangbanna, Jinghong City, Mengla County, Menglun Town, Menglun Nature Reserve, secondary tropical seasonal moist forest, 21.9120°N, 101.2823°E, 645 m elev., 27.VII.2007, Guo Zheng leg.; 1♂ (IZCAS), Xishuangbanna, Jinghong City, Mengla County, Menglun Town, Menglun Nature Reserve, *Anogeissus acuminata* plantation (about 20 yr.), 21.8999°N, 101.2802°E, 611 m elev., 19.VIII.2007, Guo Zheng leg.; 2♀ (IZCAS), Xishuangbanna, Jinghong City, Mengla County, Menglun Town, Menglun Nature Reserve, secondary tropical seasonal moist forest, 21.9065°N, 101.2802°E, 612 m elev., 10.VIII.2007, Guo Zheng leg.

Distribution. Asia: from India and Sri Lanka to Japan, south to Indonesia. In China is known from Guangxi, Taiwan, and Yunnan.

***Phrynarachne dreepy* Lin & S. Li, sp. nov.**

<http://zoobank.org/E0BC4600-8F1E-403A-BBDD-C37EDE540845>

Figs 3, 7, 8, 18C, D, 21

Type material. Holotype: ♂ (IZCAS-Ar41643), **China: Yunnan:** Xishuangbanna, Jinghong City, Mengla County, Menglun Town, Menglun Nature Reserve, 21.9768°N, 101.2010°E, 814 m elev., 17.VIII.2011, Guo Zheng leg.; **Paratypes:** 2♀ (IZCAS-Ar41644, Ar41645), **China: Yunnan:** Xishuangbanna, Jinghong City, Mengla County, Menglun Town, Menglun Nature Reserve, Xishuangbanna Tropical Botanic Garden, 21.9277°N, 101.2622°E, 552 m elev., VIII.2019, Long Yu leg.; 1♂ (IZCAS-Ar41646), same data as holotype, but 21.9502°N, 101.2010°E, 814 m elev., 18.VIII.2011; 2♂ (IZCAS-Ar41647, Ar41648), Xishuangbanna, Jinghong City, Guanping Town, Shiwudui, 22.2280°N, 100.8894°E, 888 m elev., 20.VII.2012, Qingyuan Zhao and Zhi-gang Chen leg.

Etymology. The species is named after *Dreepy*, a fictional character from *Pokémon Sword and Shield*, who has a triangular head that is reminiscent of the opisthosoma of the new species; noun (name) in apposition.

Diagnosis. *Phrynarachne dreepy* sp. nov. is similar to *P. brevis* in that males have a long RTA; in females the epigyne has sclerotized margins and the posterior edge of the median plate has a depression. However, males of *P. dreepy* sp. nov. can be easily distinguished by the long VTA (vs short VTA in *P. brevis*), the length of embolus to the length of the embolus base (7:1 vs 18:1 in *P. brevis*), and the embolus separate from the tegulum (vs close to the tegulum in *P. brevis*). Females can be separated from *P. brevis* by the short, triangular tubercles on the abdomen (vs long, slender tubercles




Figure 3. *Phrymarachne dreepy* sp. nov. **A, B** male **C** female life. Photos by Peter Koomen (**A, B**) and Chao Wu (**C**).

in *P. brevis*), the straight anterior edge of median plate (vs recurved in *P. brevis*), and the procurved posterior edge of the median plate (vs almost straight in *P. brevis*).

Description. Male (Figs 3A, B, 7, 18C), **holotype**: total length 2.26, carapace 1.04 long, 1.02 wide, yellow-brown, with white tubercles. Eye sizes and interdistances: ALE 0.09, AME 0.07, PLE 0.07, PME 0.06; ALE–AME 0.05, AME–AME 0.07, PLE–PME 0.09, PME–PME 0.11. Chelicerae brown, with two promarginal teeth and one retromarginal tooth; gnathocoxae, yellow-brown, labium brown, 0.23 long, 0.18 wide. Sternum yellow-brown. Legs yellow-brown, femora I and II with dense, varying-sized tubercles; tibiae and metatarsi I, II with pairs of ventral spines (I, tibia

6, metatarsus 8; II, tibia 6, metatarsus 6). Leg measurements: I 3.85 (1.21, 1.38, 0.76, 0.50), II 3.78 (1.22, 1.34, 0.73, 0.49), III 1.72 (0.55, 0.60, 0.27, 0.30), IV 1.52 (0.51, 0.52, 0.23, 0.26). Leg formula: 1234. Opisthosoma dark brown, each side with 18 tubercles, each with a clavate seta.

Male palp (Fig. 3A, B). Tibia brown, VTA club-shaped; RTA long, the length of VTA to the length of RTA is 3:1. Cymbium brown. Tegulum flat, disk-shaped, with a tegular ridge. Embolus spiraled, thin, separated from tegulum; the length of embolus to the length of embolus base 7:1.

Female (Figs 3C, 8, 18D) one *paratype*: total length 8.45, carapace 3.77 long, 4.02 wide, pale yellow, green when alive. Eye sizes and interdistances: ALE 0.22, AME 0.12, PLE 0.20, PME 0.15; ALE–AME 0.18, AME–AME 0.24, PLE–PME 0.28, PME–PME 1.02. Chelicerae brown, with two promarginal teeth and one retromarginal tooth; gnathocoxae, labium yellow, labium 0.86 long, 0.63 wide. Sternum yellow. Legs pale yellow, femora I and II with dense, varying-sized tubercles; tibiae and metatarsi I, II with dense asymmetrical ventral spines (I, tibia 25, metatarsus 102; II, tibia 20, metatarsus 83). Leg measurements: I 13.87 (4.26, 4.94, 3.10, 1.57), II 13.99 (4.40, 4.81, 3.16, 1.62), III 7.17 (2.36, 2.67, 1.12, 1.02), IV 6.74 (2.49, 2.57, 0.82, 0.86). Leg formula: 2134. Opisthosoma pale green, each side with 13 triangular tubercles, each with a clavate seta.

Epigyne (Fig. 8) with sclerotized margins; median plate almost rectangular, hood absent, anterior edge straight, posterior edge slightly recurved, the ratio of length to width is 4:1; copulatory opening inconspicuous; spermathecae kidney-shaped, the ratio of anterior edge to posterior edge length is 2:1. Fertilization duct transverse.

Distribution. Known only from the type locality.

Phrynarachne huangshanensis Li, Chen & Song, 1985

Figs 5, 9, 10, 17A, C, 18B, 21

Phrynarachne huangshanensis Li et al., 1985: 73, figs 1, 2. For the complete list of references see WSC (2021).

Type material. Holotype: ♀ (IZCAS-Ar41649), **China: Anhui:** Huangshan City, Huangshan District, Zhaixi Village, 30.0580°N, 118.1664°E, 423 m elev., 14.VI.1982, Youcai Li, Fayang Chen and Daxiang Song leg., examined.

Other material examined. 1♂(IZCAS-Ar416450), **China: Anhui:** Huangshan City, Tangkou Town, Houyuan, ravine, 30.0735°N, 118.1522°E, 470 m elev., IX.2018, Long Yu leg.; 2♂(IZCAS-Ar41651, Ar41652), **China: Anhui:** Huangshan City, Tangkou town, Fangcunxin Village, ravine, shrub with broad leaves, 30.0457°N, 118.1606°E, 430±8 m elev., 5.IX.2019, Long Yu leg.; 3♂ (IZCAS-Ar41653–Ar41655), **China: Anhui:** Huangshan City, Tangkou town, Fangcun Village, shrub with broad leaves, 30.0302°N, 118.1822°E, 356±6 m elev., 5.IX.2019, Long Yu leg.; 5♀(IZCAS-Ar41656–Ar41660), **China: Anhui:** Huangshan City,

Tangkou Town, Fangcunxin Village, ravine, 30.0501°N, 118.1854°E, 450 m elev., IX.2018, Long Yu leg.

Diagnosis. Males of *Phrynarachne huangshanensis* can be distinguished from those of *P. mammillata* by the ratio of the length of the embolus to the length of the embolus base (7:1 in *P. huangshanensis* vs 10:1 in *P. mammillata*), and the ratio of the length of the RTA to the length of the VTA (3:1 in *P. huangshanensis* vs 2:1 in *P. mammillata*). Females can be differentiated by the length to width ratio of the median plate (3:1 in *P. huangshanensis* vs 5:1 in *P. mammillata*), and the V-shaped median plate (vs M-shaped in *P. mammillata*).

Description. Male (Figs 5A, 9, 18B): total length 2.45, carapace 1.10 long, 1.14 wide, dark brown with long setae. Opisthosoma brown in middle, with some tubercles, each with a clavate seta. A pair of white lines from PLE to fovea. Eye sizes and interdistances: ALE 0.09, AME 0.06, PLE 0.07, PME 0.04; ALE–AME 0.05, AME–AME 0.11, PLE–PME 0.13, PME–PME 0.15. Chelicerae brown, with two promarginal teeth and one retromarginal tooth; gnathocoxae, labium dark brown, labium 0.20 long, 0.21 wide. Sternum black. Legs black, femora I and II with dense, varying-sized tubercles, tibiae and metatarsi I, II with pairs of ventral spines (I, tibia 6, metatarsus 6; II, tibia 6, metatarsus 6); femora III, IV with white stripe. Leg measurements: I 3.54 (1.13, 1.23, 0.66, 0.52), II 3.50 (1.18, 1.22, 0.60, 0.50), III 1.69 (0.55, 0.56, 0.26, 0.32), IV 2.08 (0.73, 0.72, 0.28, 0.35). Leg formula: 1234. Opisthosoma dorsally dark brown, each side with 17 tubercles, each with a clavate seta, center with a pair of yellow markings.

Male palp (Fig. 9). Tibia brown, VTA club-shaped; RTA long, the length ratio of VTA to RTA is 3:1. Cymbium black. Tegulum flat, disk-shaped, with a tegular ridge. Embolus spiral, thin, the length ratio of the embolus to the embolus base is 7:1.

Female. See Li et al. (1985).

Distribution. China (Anhui).

Notes. The male is described for the first time here.

Phrynarachne katoi Chikuni, 1955

Figs 4, 21

Phrynarachne katoi Chikuni, 1955: 35, figs 4A–G, pl. 1. For the complete list of references see WSC (2021).

Type material. Holotype 1♀ (Collection of Kyukichi Kishida, Tokyo), from Tojigami, Daisan-ku, Kawajimura, Iida-shi, Shimoyna-gun, Nagano Pref., 470 m elev., 7.IX.1953, S. Sekigawa leg., not examined.

Other material examined. 1♂ 1♀ (IZCAS), **China: Anhui:** Huangshan City, Xiuning County, Mount Qiyun, 29.8186°N, 118.0294°E, 24.X.2021, Fan Gao leg.

Distribution. China, Korea, and Japan. In China it is known from Anhui, Zhejiang.



Figure 4. *Phrynarachne katoi*, live **A** male **B** female. Photos by Fan Gao.



Figure 5. *Phrynarachne huangshanensis*, live **A** male **B** female. Photos by Ruiyang Wang.




Figure 6. *Phrynarachne brevis*, female **A** epigyne, ventral **B** vulva, dorsal.




Figure 7. *Phrynarachne dreepy* sp. nov., holotype male, left palp **A** ventral **B** retrolateral.

***Phrynarachne lancea* Tang & S. Li, 2010**

Figs 11, 19A, 21

Phrynarachne lancea Tang & Li, 2010: 53, figs 37A–D, 38A, B.

Type material. Holotype: ♂ (IZCAS-Ar18536), **China: Yunnan:** Xishuangbanna, Jinghong City, Mengla County, Menglun Town, Menglun Nature Reserve, Tropical seasonal rainforest, 21.9368°N, 101.2701°E, 558 m elev., 1.XII.2009, Guo Tang and Zhiyuan Yao leg., examined. **Paratype:** 1♂(IZCAS-Ar18537), **China: Yunnan:**

Xishuangbanna, Jinghong City, Mengla County, Menglun Town, Menglun Nature Reserve, *Anogeissus acuminata* plantation (about 20 years old), 21.8970°N, 101.2846°E, 609 m elev., 27.XII.2009, Guo Tang and Zhiyuan Yao leg. examined.

Other material examined. 3♀(IZCAS-Ar41661–Ar41663), **China: Yunnan:** Xishuangbanna, Jinghong City, Mengla County, Xishuangbanna Tropical Botanic Garden, Rainforest Valley, 21.9277°N, 101.2622°E, 552 m elev., III.2018, Yu Long leg.; 2♀ (IZCAS-Ar41664, Ar41665), same data as above, but II.2019; 3♂(IZCAS-Ar41666–Ar41668), same data as above, but V.2019.

Diagnosis. *Phrynarachne lancea* males can be easily distinguished from other species by the wide, spear-shaped RTA. Females of *P. lancea* are similar to *P. mammillata* in having an M-shaped median plate and kidney-shaped spermathecae. However, *P. lancea* can be distinguished by the length to width ratio of the median plate (7:1 in *P. lancea* vs 4:1 in *P. mammillata*), the straight posterior edge of the median plate (vs procurred in *P. mammillata*), the posterior edge of the spermathecae shorter than the anterior edge (vs of equal length in *P. mammillata*), and the longitudinal fertilization ducts (vs transverse in *P. mammillata*).

Description. Female (Figs 11, 19A): total length 16.49, carapace 6.53 long, 6.82 wide, white, posterior edge black. Eye sizes and interdistances: ALE 0.21, AME 0.20, PLE 0.24, PME 0.21; ALE–AME 0.14, AME–AME 0.25, PLE–PME 0.33, PME–PME 0.28. Chelicerae white, with two promarginal teeth and one retromarginal tooth; gnathocoxae white with black pattern, labium black, 0.88 long, 0.83 wide. Sternum white. Legs white with black markings, femora I and II with dense, varying-sized tubercles; tibiae and metatarsi I, II with dense asymmetrical ventral spines (I, tibia 28, metatarsus 75; II, tibia 26, metatarsus 68). Leg measurements: I 12.45 (4.23, 4.49, 2.41, 1.32), II 12.15 (4.15, 4.41, 2.31, 1.28), III 6.25 (2.12, 2.38, 0.98, 0.87), IV 5.90 (2.18, 2.00, 0.92, 0.80). Leg formula: 1234. Opisthosoma white, posterior grey, with four obvious brown tubercles.

Epigyne (Fig. 11) with sclerotized margins inconspicuous, M-shaped; median plate M-shaped, hood absent, anterior and posterior edges recurved, the ratio of length to width is 7:1; copulatory opening inconspicuous; spermathecae kidney-shaped, the ratio of anterior edge to posterior edge length is 3:1. Fertilization duct longitudinal.

Male. See Tang and Li (2010).

Distribution. China (Yunnan).

Notes. The female is reported here for the first time.

Phrynarachne mammillata Song, 1990

Figs 12, 13, 17B, D, 19B, 21

Phrynarachne mammillata Song in Song & Chai, 1990: 364, fig. 1A–D. For the complete list of references see WSC (2021).

Type material. Holotype: ♀ (IZCAS-Ar9358), **China: Guizhou:** Tongren City, Jiangkou County, Fanjing Mountain, 10.VII.1988, examined.




Figure 8. *Phrynarachne dreepy* sp. nov., patatype female **A** epigyne, ventral **B** vulva, dorsal.




Figure 9. *Phrynarachne huangshanensis*, male left palp **A** ventral **B** retrolateral.

Other material examined. 2♀ (IZCAS-Ar41669, Ar41670), **China, Yunnan:** Xishuangbanna, Jinghong City, Mengla County, Xishuangbanna Tropical Botanic Garden, Rainforest Valley, 21.927745°N, 101.262194°E, 552 m elev., 2014/VII, Yu Long leg.; 1♂ (IZCAS-Ar41671), Xishuangbanna, Jinghong City, Guanping Town, Shiwudui, 22.2280°N, 100.8894°E, 888 m elev., 20.VII.2012, Qingyuan Zhao and Zhigang Chen leg.

Diagnosis. See diagnosis of *Phrynarachne huangshanensis*.

Description. Male (Figs 12, 19B): total length 1.82, carapace 0.90 long, 0.97 wide, dark brown, cephalic region yellow-brown. Eye sizes and interdistances: ALE 0.09, AME 0.06, PLE 0.07, PME 0.06; ALE–AME 0.05, AME–AME 0.07, PLE–PME 0.12, PME–PME 0.12. Chelicerae brown, with two promarginal teeth and one retromarginal tooth; gnathocoxae, labium dark brown, labium 0.15 long, 0.19 wide. Sternum brown. Legs black, femora I and II with dense, varying-sized tubercles; tibiae and metatarsi I, II with pairs of ventral spines (I, tibia 6, metatarsus 6; II, tibia 6, metatarsus 6), femora



Figure 10. *Phrynarachne huangshanensis*, holotype female **A** epigyne, ventral **B** vulva, dorsal.

III, IV with broad white pattern. Leg measurements: I 2.51 (0.83, 0.83, 0.51, 0.34), II 2.51 (0.81, 0.86, 0.46, 0.38), III 1.47 (0.47, 0.48, 0.26, 0.26), IV 1.36 (0.48, 0.42, 0.24, 0.22). Leg formula: 1 = 234. Opisthosoma dark brown with yellow-brown spots, each side with 18 tubercles, yellow-brown tubercles, each with a clavate seta.

Male palp (Fig. 12). Tibia brown, VTA club-shaped; RTA long, the length ratio of VTA to RTA is 2:1. Cymbium yellow to brown. Tegulum flat, disk-shaped, with a tegular ridge. Embolus spiraled, thin, the length ratio of embolus to embolus base is 10:1.

Female. See Song and Chai (1990).

Distribution. China (Guizhou, Yunnan).

Notes. The male is reported here for the first time.

Phrynarachne sinensis Peng, Yin & Kim, 2004 *nomen dubium*

Phrynarachne sinensis Peng et al., 2004: 21, figs 1–3; Yin et al. 2012: 1265, fig. 680a–c.

Type material. Holotype: ♀ (College of Life Sciences, Hunan Normal University), China (Wang-101), no detailed data, lost, not examined.

Notes. The lost type specimen, lack of clear figures of the holotype, and the vague distributional information make further study of the taxonomy of this species impossible. We treat it as *nomen dubium*.

Phrynarachne xuxiake Lin & S. Li, sp. nov.

<http://zoobank.org/BB4750BD-1DC5-4B3C-BF29-2F26118EA68D>


Figs 1B, 14, 19C, 21

Type material. Holotype: ♀ (IZCAS-Ar41672), **China: Anhui:** Huangshan City, Tangkou Town, Fangcunxin Village, ravine, 30.0501°N, 118.1854°E, 450 m elev., IX.2018, Long Yu leg.

Etymology. The species is named after Xu Xiake, a Chinese travel writer and geographer of the Ming dynasty; noun (name) in apposition.

Diagnosis. Females of *Phrynarachne xuxiake* sp. nov. are similar to *P. katoi* but can be distinguished by the length to width ratio of the median plate (3:1 in *P. xuxiake* vs 5:1 in *P. katoi*) and by the rectangular median plate with its posterior edge straight (vs dumbbell-shaped with procurred posterior edge in *P. katoi*).

Description. Female (Figs 14, 19C), **holotype:** total length 8.78, carapace 3.84 long, 4.45 wide, dark brown with long setae. Eye sizes and interdistances: ALE 0.22, AME 0.17, PLE 0.18, PME 0.15; ALE–AME 0.18, AME–AME 0.24, PLE–PME 0.28, PME–PME 0.28. Chelicerae white, with two promarginal teeth and one retromarginal tooth; gnathocoxae, labium black, labium 0.84 long, 0.74 wide. Sternum black. Legs yellow, femora I and II with dense, varying-sized tubercles; tibiae and metatarsi of I, II with dense ventral spines



A



B

Figure 11. *Phrynarachne lancea*, female **A** epigyne, ventral **B** vulva, dorsal.




Figure 12. *Phrynarachne mammillata*, male left palp **A** ventral **B** retrolateral.

(I, tibia 26, metatarsus 108; II, tibia 21, metatarsus 89). Leg measurements: I 15.11 (4.17, 4.92, 3.70, 2.32), II 13.93 (4.02, 4.68, 3.13, 2.10), III 6.94 (2.12, 2.71, 1.05, 1.06), IV 6.63 (2.39, 2.54, 0.76, 0.94). Leg formula: 1234. Opisthosoma brown, each side with 19 blunt tubercles, each with a clavate seta, with a pair of black markings medially.

Epigyne (Fig. 14). Sclerotized margins inconspicuous, M-shaped; median plate obvious, with a posterior hood, anterior edge recurved and posterior edge almost straight, the ratio of width to length is 3:1; copulatory opening inconspicuous; Spermathecae kidney-shaped, the ratio of anterior edge to posterior edge length is 3:2. Fertilization duct transverse.

Distribution. Known only from the type locality.




Figure 13. *Phrynarachne mammillata* Song, 1990, female **A** epigyne, ventral **B** vulva, dorsal.




Figure 14. *Phrynarachne xuxiake* sp. nov., holotype female **A** epigyne, ventral **B** vulva, dorsal.




Figure 15. *Phrynarachne yunhui* sp. nov., holotype female **A** epigyne, ventral **B** vulva, dorsal.

***Phrynarachne yunhui* Lin & S. Li, sp. nov.**

<http://zoobank.org/51542955-2CCB-4A0B-881A-FED561B28CB8>

Figs 15, 19D, 21

Type material. Holotype: ♀ (IZCAS-Ar41673), **China: Hainan:** Ledong County, Jianfengling Nature Reserve, Mingfenggu, 18.7417°N, 108.8417°E, 989 m elev., 1.VII.2020, Yunhu Mo leg.

Etymology. The species is named after Mr Yunhu Mo, who collected the holotype; noun (name) in genitive case.

Diagnosis. Females of *Phrynarachne yunhui* sp. nov. are similar to *P. mammillata* in having the anterior edge and posterior edges of the median plate procurved and the posterior edge with a depression, and in having kidney-shaped spermathecae. However, *Phrynarachne yunhui* sp. nov. can be distinguished by the oval median plate (vs M-shaped in *P. mammillata*) and the broad anterior edge of the spermathecae (vs narrow in *P. mammillata*).

Description. Female (Figs 15, 19D): total length 10.04, carapace 3.67 long, 4.44 wide, black. Eye sizes and interdistances: ALE 0.22, AME 0.19, PLE 0.25, PME 0.20; ALE–AME 0.16, AME–AME 0.27, PLE–PME 0.35, PME–PME 0.29. Chelicerae black, with two promarginal teeth and one retromarginal tooth; Gnathocoxae, labium black, labium 0.83 long, 0.75 wide. Sternum black. Legs black, femora I and II with dense, varying-sized tubercles; tibiae and metatarsi I, II with dense asymmetrical ventral spines (I, tibia 17, metatarsus 41; II, tibia 16, metatarsus 36). Leg measurements: I 13.01 (4.38, 4.63, 2.56, 1.44), II 12.74 (4.33, 4.58, 2.43, 1.40), III 6.78 (2.16, 2.42, 1.18, 1.02), IV 6.44 (2.35, 2.19, 1.07, 0.93). Leg formula: 1234. Opisthosoma grey, with dense, varying-sized, red-brown tubercles, each with a clavate seta.

Epigyne (Fig. 15). Sclerotized margins inconspicuous; median plate obvious, with a small posterior hood, anterior and posterior edges recurved, the ratio of width to length is 15:4; copulatory opening inconspicuous; spermathecae kidney-shaped, the ratio of posterior edge to anterior edge length is 1:1. Fertilization duct transverse.

Distribution. Known only from the type locality.

***Phrynarachne zhengzhongi* Lin & S. Li, sp. nov.**

<http://zoobank.org/77674036-F817-4088-A525-2DF0AB763CDC>

Figs 16, 20, 21

Type material. Holotype: ♀ (IZCAS-Ar41674), **China: Yunnan:** Xishuangbanna, Jinghong City, Guanping Town, Shiwudui, 22.2310°N, 100.9172°E, 872 m elev., 27.IV.2018, Zhengzhong Huang leg. **Paratype** ♀ (IZCAS-Ar41675), same data as holotype.

Etymology. The species is named after Mr Zhengzhong Huang, who collected the holotype and paratype; noun (name) in genitive case.





Figure 16. *Phrynarachne zhengzhongi* sp. nov., holotype female **A** epigyne, ventral **B** vulva, dorsal.



Phrynarachne sp. 1♀
李叔才·陳發揚·湖南梵淨山, 1982.6.

乳突瘤蟹蛛

Phrynarachne mammillata
1♀, 贵州江口梵净山, 1982.7.10

Inst. Zool., Chinese Academy of Sciences D

Coll. /, Det. Li Shuqiang

IZCAS-Ar 9358

1♀

Thomisidae

Phrynarachne mammillata

Guizhou(贵州江口县梵净山)

10. VII. 1982 HOLOTYPE

Figure 17. *Phrynarachne* spp., holotype females **A, C** habitus (**A**) and original labels (**C** handwriting by Daxiang Song) of *P. huangshanensis* **B, D** habitus (**B**) and original label (**D** handwriting by Daxiang Song) of *P. mammillata*.




Figure 18. *Phrynarachne* spp., habitus dorsal **A** *P. brevis*, female **B** *P. huangshanensis*, male **C** *P. dreepy* sp. nov., holotype male **D** same, paratype female.




Figure 19. *Phrynarachne* spp., habitus dorsal **A** *P. lancea*, female **B** *P. mammillata*, male **C** *P. xuxiake* sp. nov., holotype female **D** *P. yunhui* sp. nov., holotype female.




Figure 20. *Phrynarachne zhengzhongi* sp. nov., female holotype (A) and paratype (B) habitus **A** dorsal **B** lateral.

Diagnosis. Females of *Phrynarachne zhengzhongi* sp. nov. are similar to *P. brevis* by the shape of the spermathecae; the posterior edge of the spermathecae is as wide as the anterior edge. However, females of *P. zhengzhongi* sp. nov. can be distinguished by the triangular tubercles on the abdomen (vs long, slender apophysis in *P. brevis*), the epigyne with a hood, and the absence of sclerotized margins (vs hood absent, sclerotized margins present in *P. brevis*), and the straight anterior edge of the median plate (vs recurved in *P. brevis*).

Description. Female (Figs 16, 20): total length 10.79, carapace 5.27 long, 5.53 wide, brown, with small projection, ocular tubercle white. Projection present between ALE and PLE. Eye sizes and interdistances: ALE 0.21, AME 0.17, PLE 0.18, PME 0.10; ALE–AME 0.35, AME–AME 0.66, PLE–PME 0.50, PME–PME 0.87. Chelicerae pale yellow, with two promarginal teeth and one retromarginal tooth; gnathocoxae, labium yellow, labium 1.19 long, 0.94 wide. Sternum yellow. Legs brown, femora I and II with dense, varying-sized tubercles; tibiae and metatarsi I, II with dense asymmetrical ventral spines (I, tibia 22, metatarsus 43; II, tibia 17, metatarsus 37). Leg measurements: I 14.43 (4.71, 5.91, 2.35, 1.46), II 14.78 (5.05, 6.31, 2.09, 1.33), III 7.76 (2.24, 3.48, 1.03, 1.01), IV 7.30 (2.44, 3.01, 0.92, 0.93). Leg




Figure 21. Distribution records of *Phrynarachne* species in China **1** *P. brevis* **2** *P. ceylonica* **3** *P. dreepy* sp. nov. **4** *P. huangshanensis* **5** *P. katoi* **6** *P. lancea* **7** *P. mammillata* **8** *P. xuxiake* sp. nov. **9** *P. yunhui* sp. nov. **10** *P. zhengzhongi* sp. nov.

formula: 2134. Opisthosoma dorsally light yellow, each side with 22 tubercles, each with some tubercles.

Epigyne (Fig. 16). Sclerotized margins inconspicuous; median plate obvious, with a posterior hood, anterior and posterior edges almost straight, the ratio of width to length is 4:1; copulatory opening inconspicuous; spermathecae kidney-shaped, the ratio of posterior edge to anterior edge length is 1:1. Fertilization duct transverse.

Distribution. Known only from the type locality.

Acknowledgements

The manuscript benefited greatly from comments by Hirotugu Ono, Yuri Marusik, Zhiyuan Yao. Sarah Crews checked the English. Zhengzhong Huang (IZCAS), Fan Gao (Nanjing, China), and Yunhu Mo (Haikou, China) helped with fieldwork. Fan Gao (Nanjing, China), Chao Wu (Beijing, China), and Ruiyang Wang (Nanjing, China) took the photographs.

References

- Chikuni Y (1955) Five interesting spiders from Japan highlands. *Acta Arachnologica* 14: 29–40.
<https://doi.org/10.2476/asjaa.14.29>
- Li J, Yan X, Lin Y, Li S, Chen H (2021) Challenging Wallacean and Linnean shortfalls: Ectatosticta spiders (Araneae, Hypochilidae) from China. *Zoological Research* 42(6): 791–794. <https://doi.org/10.24272/j.issn.2095-8137.2021.212>
- Li Y, Chen F, Song D (1985) A new species of the genus *Phrynarachne* (Araneae: Thomisidae). *Journal of Anhui Normal University, Natural Sciences* 1985(1): 72–74.
- Ono H (1988) A Revisional Study of the Spider Family Thomisidae (Arachnida, Araneae) of Japan. National Science Museum, Tokyo, 252 pp.
- Peng X, Yin C, Kim J (2004) One new species of the genus *Phrynarachne* (Araneae, Thomisidae) from China. *Korean Arachnology* 20: 21–25.
- Pickard-Cambridge O (1884) On two new genera of spiders. *Proceedings of the Zoological Society of London* 52(2): 196–205. [pl. 40] <https://doi.org/10.1111/j.1096-3642.1884.tb02819.x>
- Song D, Chai J (1990) Notes of some species of the family Thomisidae (Arachnida: Araneae) from Wuling Shan area. In: Zhao E (Ed.) *From Water onto Land*. CSSAR, Beijing, 364–374.
- Tang G, Li S (2010) Crab spiders from Xishuangbanna, Yunnan Province, China (Araneae, Thomisidae). *Zootaxa* 2703: 1–105. <https://doi.org/10.1111/j.1096-3642.1884.tb02819.x>
- Thorell T (1869) On European spiders. Part I. Review of the European genera of spiders, preceded by some observations on zoological nomenclature. *Nova Acta Regiae Societatis Scientiarum Upsaliensis* (3)7: 1–108.
- Yao Z, Wang X, Li S (2021) Tip of the iceberg: species diversity of *Pholcus* spiders (Araneae, Pholcidae) in Changbai Mountains, Northeast China. *Zoological Research* 42(3): 267–271. <https://doi.org/10.24272/j.issn.2095-8137.2020.214>
- Yin CM, Peng XJ, Yan HM, Bao YH, Xu X, Tang G, Zhou QS, Liu P (2012) *Fauna Hunan: Araneae in Hunan, China*. Hunan Science and Technology Press, Changsha, 1590 pp.
- WSC (2021) World Spider Catalog, Version 22.5. Natural History Museum Bern. <http://wsc.nmbe.ch> [Accessed on: 2021–09–11]