

Four new species of the genus *Otacilia* Thorell, 1897 from Hunan Province, China (Araneae, Phrurolithidae)

Chi Jin¹, Lina Fu¹, Xiangchu Yin^{1,2}, Feng Zhang¹

1 Key Laboratory of Invertebrate Systematics and Application, College of Life Sciences, Hebei University, Baoding, Hebei 071002, China **2** Northwest Plateau Institute of Biology, Chinese Academy of Sciences, Xining, 810001, China

Corresponding author: Feng Zhang (dudu06042001@163.com)

Academic editor: S. Li | Received 31 January 2016 | Accepted 7 September 2016 | Published 29 September 2016

<http://zoobank.org/3223A81C-4630-40FA-9967-2A80B4FD2039>

Citation: Jin C, Fu L, Yin X, Zhang F (2016) Four new species of the genus *Otacilia* Thorell, 1897 from Hunan Province, China (Araneae, Phrurolithidae). ZooKeys 620: 33–55. doi: 10.3897/zookeys.620.7982

Abstract

Four new species of the genus *Otacilia* Thorell, 1897 are reported from Hunan Province, China: *Otacilia hippocampa* sp. n., *Otacilia yangmingensis* sp. n., *Otacilia curvata* sp. n., and *Otacilia submicrostoma* sp. n. All new species are described based on both sexes. In addition, the 55 known *Otacilia* species are divided into four species groups.

Keywords

Description, diagnosis, etymology, morphology, taxonomy

Introduction

Phrurolithidae Banks, 1892 was elevated to family rank by Ramírez (2014), consistent with the suggestion of Deeleman-Reinhold (2001). The family is currently represented by 211 species belonging to 14 genera worldwide. Of these, four genera and 65 species are recorded from China (World Spider Catalog 2016, Fu et al. 2015, Fu et al. 2016a, b). The Phrurolithidae are mostly ground-dwelling spiders living in leaf litter, especially bamboo leaves, woody debris or on the forest floor, very few species are found in the canopy (Deeleman-Reinhold 2001; Fu et al. 2014).

Otacilia Thorell, 1897 is one of the species-rich genera of the family comprising 55 species distributed in south-east Asia and east Asia. Among these *Otacilia* species, 35 are reported from China (Fu et al. 2015, Fu et al. 2016a, Fu et al. 2016b). The genus *Otacilia* was established by Thorell (1897) with description of *O. armatissima* based on a single female specimen from Myanmar.

The genus *Otacilia* is closely related to *Phrurolithus* C. L. Koch, 1839, comprising 74 species mostly distributed in the holarctic region. Until now, there is no clear way to differentiate between *Otacilia* and *Phrurolithus*. The diagnostic characters provided by Kamura (2005) were inaccurate (Wang et al. 2015), and the differences listed by Jäger and Wunderlich (2012) were also not distinct with the addition of more new species of these two genera.

Wang et al. (2015) listed ten Chinese *Otacilia* species in two groups. Subsequently, Fu et al. (2016b) reviewed the 31 Chinese *Otacilia* species and agreed with Wang et al.'s (2015) assignment and also established a third species group to accommodate the *Otacilia* species: the *armatissima* group, the *revoluta* group, and the *pseudostella* group.

While examining the collections from Hunan Province, China, some *Otacilia* specimens were found that differed from the currently known *Otacilia* species. They are identified as four new species, *Otacilia hippocampa* sp. n., *Otacilia yangmingensis* sp. n., *Otacilia curvata* sp. n., and *Otacilia submicrostoma* sp. n., and are described and illustrated here.

Material and methods

The terminology used follows Jäger and Wunderlich (2012). All measurements given in the text are in millimeters. Total length is the sum of the carapace and abdomen lengths, regardless of the pedicel. Eye sizes were measured as the maximum diameter in dorsal or frontal view. Leg measurements are shown as: total length (femur, patella, tibia, metatarsus, tarsus). Epigyne were removed and cleared in a warm solution of 10% potassium hydroxide (KOH), transferred to ethanol and temporarily mounted for drawing. All specimens are preserved in 75% alcohol and were examined, drawn, and measured under a Leica M205A stereomicroscope equipped with an Abbe drawing device. Photographs were taken using a Leica M205A stereomicroscope equipped with a DFC450 CCD. The specimens are deposited in the Museum of Hebei University, Baoding, China (MHBU).

Abbreviations

ALE	anterior lateral eyes;
AME	anterior median eyes;
a.s.l.	above sea level;
B	bursa;

C	conductor;
CD	copulatory duct;
CO	copulatory opening;
CT	connecting tube;
DTA	dorsal tibial apophysis;
E	embolus;
FA	femoral apophysis;
FD	fertilization duct;
GA	glandular appendage;
MOA	median ocular area;
MP	median plate;
PLE	posterior lateral eyes;
PME	posterior median eyes;
RTA	retrolateral tibial apophysis;
S	spermatheca;
TA	tegular apophysis.

Taxonomy

Phrurolithidae Banks, 1892

Otacilia Thorell, 1897

Diagnosis. Chelicerae each with two bristles (rarely with one bristle) on anterior side; leg formula: 4123 (rarely 1423); spination: femora I–II d 0–2, III–IV d 0–1, I pl 3–6, II pl 0–3; tibiae I–II usually with 6–8 pairs of ventral spines; tibia I always one more rv than pv spine and tibia II always one more pv than rv spine; metatarsi I–II usually with 3–4 pairs of ventral spines, and always one more pv than rv spine.

Male palp: femur with ventral apophysis or hump; tibia usually with single strong RTA, some species with BTA or DTA; embolus hook-shaped or needle-like, originating antero-prolaterally; tegular apophysis sclerotized or transparent, present or absent, antero-retrolaterally located; conductor membranous, well developed or absent. Female genitalia: epigynal median plate distinct or absent; vulva with pair of transparent bursae anteriorly and pair of strongly sclerotized spermathecae posteriorly.

Species groups of *Otacilia*. After reviewing 59 *Otacilia* species (including the four new Chinese species described in this paper), the grouping was revised and the current species assigned to four groups based on assessment of Fu et al. (2016b). The *revoluta* group was divided into two new groups, the *longituba* group (16 species) and the *ambon* group (two species). The *armatissima* group (29 species) and the *pseudostella* group (nine species) were preserved and updated. Three species were not assigned to any group because of their poor original description and figures or peculiar structure: *Otacilia luzonica* (Simon, 1898) (female is unknown; description and figures are not

diagnostic), *O. papilla* Dankittipakul & Singtripop, 2014 (male is unknown; epigyne medially with lobe and absence of bursae) and *O. paracymbium* Jäger & Wunderlich, 2012 (female is unknown; cymbium with paracymbium).

Here the male and female diagnostic characters are listed for each species group, followed by a list of all of the included *Otacilia* species (Table 1).

***Otacilia hippocampa* sp. n.**

<http://zoobank.org/C336230A-1FD5-435C-BA9C-13657E682F6F>

Figs 1–3

Type material. Holotype ♂, China, *Hunan Province*: Dao County, Qingtang Town, Dajiangyuan Village, Mt. Jiucailing (25°27'37.678"N, 111°21'12.499"E), 448 m a.s.l., 29 September 2015, Chi Jin leg. Paratypes: 2♀2♂, same data as for holotype.

Etymology. The species name is taken from the Latin generic name of the seahorse, “*Hippocampus*”, referring to the seahorse-shaped internal ducts (copulatory duct, connecting tube and spermatheca) in the female epigyne; adjective.

Diagnosis. The male can be distinguished from all other *armatissima* group species, except *O. bicolor* Jäger & Wunderlich, 2012, *O. onoi* Deeleman-Reinhold, 2001 and *O. truncata* Dankittipakul & Singtripop, 2014, by having a long DTA and can be distinguished from these three species by the absence of conductor (Figs 2A–D, 3A–C). The female of the new species can be easily distinguished from all of the other *armatissima* group species by the seahorse-shaped internal ducts (except the bursae), whereas they are S-shaped or crescent-shaped in the other congeners (Figs 2E–G, 3D–E).

Description. Male (Fig. 1A–B). Total length 2.17–2.65 (n = 3). Holotype: body 2.65 long; carapace 1.28 long, 1.12 wide; abdomen 1.37 long, 0.90 wide. Carapace yellowish brown, with black marginal bands; fovea longitudinal, brown. Eye diameters: AME 0.08, ALE 0.09, PME 0.08, PLE 0.09. Eye interdistances: AME–AME 0.02, AME–ALE 0.01, PME–PME 0.09, PME–PLE 0.05, ALE–PLE 0.07. MOA 0.19 long, front 0.17 wide, back 0.25 wide. Clypeus 0.14 high. Chelicerae with two strong anterior bristles; promargin with three well-separated teeth, and retromargin with two teeth close to each other. Labium and sternum dark yellow. Legs light yellowish brown; all femora with distal black distal annulus; patellae I–II black, patellae III–IV with distal black annulus; tibiae I–II almost all black, tibiae III–IV with black distal annulus; metatarsi I–II distal half part black, metatarsi III–IV with black distal annulus. Measurements of legs: leg I 4.95 (1.29, 0.49, 1.39, 1.23, 0.55), II 4.09 (1.11, 0.46, 1.01, 0.98, 0.53), III 3.59 (0.91, 0.44, 0.74, 0.94, 0.56), IV 5.19 (1.36, 0.45, 1.14, 1.49, 0.75). Leg formula: 4123. Femur I with two dorsal spines and three pro-lateral spines, femur II with one dorsal spine and two pro-lateral spines, femur III lacks dorsal spine, femur IV with one dorsal spine; tibia I with six proventral spines and seven retroventral spines, tibia II with six pairs of ventral spines; metatarsus I with four pairs of ventral spines, metatarsus II with four proventral spines and three retroventral spines. Femora I–III lack dorsal spines, femur IV four with one dorsal spine, femur I



Figure 1. *Otacilia hippocampa* sp. n. **A** male habitus, dorsal view **B** same, ventral view **C** Female habitus, dorsal view **D** same, ventral view.

Table 1. Definition of species groups of *Otacilia* species, together with lists of included species (species marked with an asterisk are reported from China).

Species group name	Diagnostic Character	Included Species
<i>armatissima</i>	<p>1) Palpal organ with a distinct sclerotized TA or membranous conductor; embolus hook-shaped.</p> <p>2) Epigyne with a distinct median plate, without concavity; CO located anteriorly or medially, higher than the spermathecae; CD (the left one from the ventral view) anti-clockwise from the CO; connecting tube usually crescent-shaped; spermathecae separated by more than half a spermatheca's diameter.</p>	<p>1) <i>O. armatissima</i> Thorell, 1897</p> <p>2) <i>O. bawangling</i> Fu, Zhang & Zhu, 2010*</p> <p>3) <i>O. biarclata</i> Fu, He & Zhang, 2015*</p> <p>4) <i>O. bicolor</i> Jäger & Wunderlich, 2012</p> <p>5) <i>O. florifera</i> Fu, He & Zhang, 2015*</p> <p>6) <i>O. forcipata</i> Yang, Wang & Yang, 2013*</p> <p>7) <i>O. foveata</i> (Song, 1990)*</p> <p>8) <i>O. fujiana</i> Fu, Jin & Zhang, 2014*</p> <p>9) <i>O. hengshan</i> (Song, 1990)*</p> <p>10) <i>O. jianfengling</i> Fu, Zhang & Zhu, 2010*</p> <p>11) <i>O. kao</i> Jäger & Wunderlich, 2012</p> <p>12) <i>O. komurai</i> (Yaginuma, 1952)*</p> <p>13) <i>O. limushan</i> Fu, Zhang & Zhu, 2010*</p> <p>14) <i>O. liupan</i> Hu & Zhang, 2011*</p> <p>15) <i>O. luna</i> (Kamura, 1994)</p> <p>16) <i>O. lynx</i> (Kamura, 1994)*</p> <p>17) <i>O. macrospora</i> Fu, Zhang & Zhang, 2016*</p> <p>18) <i>O. obesa</i> Fu, Zhang & Zhang, 2016*</p> <p>19) <i>O. onoi</i> Deeleman-Reinhold, 2001</p> <p>20) <i>O. papilion</i> Fu, Zhang & Zhang, 2016*</p> <p>21) <i>O. pyriformis</i> Fu, Zhang & Zhang, 2016*</p> <p>22) <i>O. sinifera</i> Deeleman-Reinhold, 2001</p> <p>23) <i>O. songi</i> Wang et al., 2015*</p> <p>24) <i>O. subliupan</i> Wang et al., 2015*</p> <p>25) <i>O. taiwanica</i> (Hayashi & Yoshida, 1993)*</p> <p>26) <i>O. truncata</i> Dankittipakul & Singtripop, 2014</p> <p>27) <i>O. yangi</i> Zhang, Fu & Zhu, 2009*</p> <p>28) <i>O. hippocampa</i> sp. n.*</p> <p>29) <i>O. yangmingensis</i> sp. n.*</p>
<i>ambon</i>	<p>1) Palpal organ without a distinct TA; the bulb is not pyriform but oval; embolus claw-like.</p> <p>2) Epigyne without median plate and concavity; Cos located posterior, lower than the spermathecae; spermathecae well separated from each other by more than three spermatheca's diameter.</p>	<p>1) <i>O. ambon</i> Deeleman-Reinhold, 2001</p> <p>2) <i>O. revoluta</i> (Yin et al., 2004)*</p>
<i>longituba</i>	<p>1) Palpal organ without a distinct TA; conductor well developed or degenerated; embolus needle-like.</p> <p>2) Epigyne without median plate and concavity; Cos located medially, higher than the spermathecae; CD (the left one from the ventral view) straight or slightly clockwise from the CO; spermathecae close together or separated by less than half a spermatheca's diameter.</p>	<p>1) <i>O. bifurcata</i> Dankittipakul & Singtripop, 2014</p> <p>2) <i>O. christae</i> Jäger & Wunderlich, 2012</p> <p>3) <i>O. flexa</i> Fu, Zhang & Zhang, 2016*</p> <p>4) <i>O. longituba</i> Wang, Zhang & Zhang, 2012*</p> <p>5) <i>O. loriot</i> Jäger & Wunderlich, 2012</p> <p>6) <i>O. microstoma</i> Wang et al., 2015*</p> <p>7) <i>O. mingsheng</i> Yang, Wang & Yang, 2013*</p> <p>8) <i>O. mira</i> Fu, Zhang & Zhang, 2016*</p> <p>9) <i>O. mustela</i> Kamura, 2008</p>

Species group name	Diagnostic Character	Included Species
<i>longituba</i>		10) <i>O. namkhan</i> Jäger & Wunderlich, 2012 11) <i>O. parva</i> Deeleman-Reinhold, 2001 12) <i>O. simianshan</i> Zhou, Wang & Zhang, 2013* 13) <i>O. vangvieng</i> Jäger & Wunderlich, 2012 14) <i>O. zebra</i> Deeleman-Reinhold, 2001 15) <i>O. curvata</i> sp. n.* 16) <i>O. submicrostoma</i> sp. n.*
<i>pseudostella</i>	1) Palpal organ without a distinct TA; an apophysis present near embolic base (PEA); embolus needle-like. 2) Epigyne without indistinct median plate, but with a pair of shallow concavities; Cos located anteriorly or medially, higher than the spermathecae; spermathecae separated by more than one spermatheca's diameter.	1) <i>O. acuta</i> Fu, Zhang & Zhang, 2016* 2) <i>O. aurita</i> Fu, Zhang & Zhang, 2016* 3) <i>O. digitata</i> Fu, Zhang & Zhang, 2016* 4) <i>O. leibo</i> Fu, Zhang & Zhang, 2016* 5) <i>O. ovata</i> Fu, Zhang & Zhang, 2016* 6) <i>O. pseudostella</i> Fu, Jin & Zhang, 2014* 7) <i>O. stella</i> Kamura, 2005 8) <i>O. vulpes</i> (Kamura, 2001) 9) <i>O. zhangii</i> Fu, Jin & Zhang, 2014*
the others		1) <i>O. luzonica</i> (Simon, 1898) 2) <i>O. papilla</i> Dankittipakul & Singtripop, 2014 3) <i>O. paracymbium</i> Jäger & Wunderlich, 2012*

with three prolateral spines; tibia I with six proventral spines and seven retroventral spines, tibia II with six proventral spines and five retroventral spines; metatarsus I with four pairs of ventral spines, metatarsus II with four proventral spines and three retroventral spines. Abdomen oval, dorsum light grey, with several chevron-like black stripes, anterior half with a small dorsal scutum; venter light grey.

Palp (Figs 2A–D, 3A–C). Femur distally with an inflated hump on ventral side and a retrolateral concavity. RTA basally thick, tapering to a sharp apex, bent prolaterally. DTA shaped similarly to RTA, with one spine basally. Embolus short, needle-like. Conductor absent. Tegular apophysis triangular, sclerotized.

Female (Fig. 1C–D). Total length 2.56–2.96 (n = 2). One paratype: body 2.96 long; carapace 1.47 long, 1.22 wide; abdomen 1.49 long, 0.99 wide. Eye diameters: AME 0.07, ALE 0.09, PME 0.08, PLE 0.10. Eye interdistances: AME–AME 0.03, AME–ALE 0.01, PME–PME 0.09, PME–PLE 0.06, ALE–PLE 0.09. MOA 0.22 long, front 0.18 wide, back 0.27 wide. Clypeus 0.13 high. Leg measurements: I 5.40 (1.41, 0.58, 1.54, 1.32, 0.55); II 4.53 (1.21, 0.53, 1.16, 1.04, 0.59); III 3.96 (1.04, 0.46, 0.81, 1.02, 0.63); IV 5.63 (1.46, 0.50, 1.27, 1.60, 0.80). Leg formula: 4123. Leg spination as in male. Abdomen light grey, anterior half lacks dorsal scutum. Other characters as in male.

Epigyne (Figs 2E–F, 3D): median plate narrow, with parallel lateral margin; copulatory openings situated centrally, tiny and pore-like. Vulva (Figs 2G, 3E–F): copulatory ducts short, connected with a pair of slender tubes leading to the large, transparent ovoid bursae; spermathecae located posteriorly, small and ovoid, separated by more than one spermatheca's diameter; connecting tubes curved and sigmoid. Glandular appendages absent.

Distribution. Known only from the type locality, Hunan, China (Fig. 13).




Figure 2. *Otacilia hippocampa* sp. n. **A** left male palp, ventral view **B** same, dorsal view **C** same, prolateral view **D** same, retrolateral view **E** epigyne, ventral view **F** same, cleared by potassium hydroxide, ventral view **G** vulva, dorsal view.




Figure 3. *Otacilia hippocampa* sp. n. **A** left male palp, ventral view **B** same, dorsal view **C** same, retrolateral view **D** epigyne, ventral view **E** vulva, dorsal view. Scale bars equal for **A** and **B**, equal for **D** and **E**.

***Otacilia yangmingensis* sp. n.**

<http://zoobank.org/9FA1C1B9-0F0B-455C-B1D8-E17B2897AA68>

Figs 4–6

Type material. Holotype ♂, China, *Hunan Province*: Shuangpai County, Mt. Yangming, Wanshou Temple (26°06'27.490"N, 111°55'19.186"E), 1375 m a.s.l., 26 September 2015, Chi Jin leg. Paratypes: 5♀4♂, same data as for holotype; 1♂, Shuangpai County, Mt. Yangming, Hongjun Pavilion (26°04'34.924"N, 111°56'19.223"E), 1324 m a.s.l., 27 September 2015, Xiangbo Guo leg.; 1♂, Jiangyong County, Qianjiadong Town, Daboshui (25°24'25.70"N, 111°19'04.33"E), 224 m a.s.l., 3 October 2015, Jingchao He leg.

Etymology. The species name refers to the holotype locality; adjective.

Diagnosis. The male can be distinguished from all other *armatissima* group species, except *O. macrospora* Fu. Zhang & Zhang, 2016, by the RTA base with a triangular




Figure 4. *Otacilia yangmingensis* sp. n. **A** male habitus, dorsal view **B** same, ventral view **C** Female habitus, dorsal view **D** same, ventral view.

process dorsally and by the absent DTA and can be distinguished from it by the long needle-like embolus (embolus stout and hook-shaped in *O. macrospora*) and the thumb-shaped tegular apophysis (tegular apophysis sickle-shaped in *O. macrospora*) (Figs 5A–B, 6A–B; Fu et al. 2016a: figs 16, 18, 22–23). The female of the new species can be distinguished from all other *armatissima* group species, except *O. macrospora* Fu, Zhang & Zhang, 2016, by the copulatory openings connected with a pair of shallow concavities anteriorly, and the concavities have distinct anterior and inner lateral margins, and can be distinguished from it by the median plate (narrower than that of *O. macrospora*) and bursae (long ovoid, whereas they are spherical in *O. macrospora*) (Figs 5E–G, 6D–E; Fu et al. 2016a: figs 20–21, 25–26).

Description. Male (Fig. 4A–B). Total length 3.04–3.16 (n = 7). Holotype: body 3.16 long; carapace 1.58 long, 1.31 wide; abdomen 1.58 long, 0.97 wide. Carapace light yellowish brown, lateral margin black, middle with broad longitudinal black stripe, from ocular area to the posterior margin of carapace; fovea longitudinal, distinct. Eye diameters: AME 0.11, ALE 0.12, PME 0.10, PLE 0.10. Eye interdistances: AME–AME 0.03, AME–ALE 0.01, PME–PME 0.10, PME–PLE 0.05, ALE–PLE 0.09. MOA 0.25 long, front 0.23 wide, back 0.27 wide. Clypeus 0.14 high. Chelicerae with two strong anterior bristles; promargin with three well separated teeth and retromargin with six denticles close to each other. Labium and sternum light yellow. Legs light yellowish brown. Leg measurements: leg I 6.65 (1.66, 0.61, 1.89, 1.65, 0.84), II 5.36 (1.42, 0.47, 1.43, 1.29, 0.75), III 4.56 (1.22, 0.49, 1.01, 1.20, 0.64), IV 7.38 (2.00, 0.58, 1.75, 2.08, 0.97). Leg formula: 4123. Femur I with two dorsal spines and four prolateral spines, femur II with one dorsal spine and two prolateral spines, femora III–IV with one dorsal spine; tibia I with seven proventral spines and eight retroventral spines, tibia II with seven pairs of ventral spines; metatarsi I–II with four pairs of ventral spines. Abdomen oval, dorsum black, anterior half with a narrow dorsal scutum, posterior half with several black transversal stripes; venter light grey, with black longitudinal stripes.

Palp (Figs 5A–D, 6A–C). Femur distally with an apophysis on ventral side and a retrolateral concavity. RTA broad, with sharp apex, base with a triangular process dorsally. Embolus slender, needle-like, slightly curved. Tegular apophysis sclerotized and thumb-shaped, situated at the apex of the bulb, separate from the embolus base.

Female (Fig. 4C–D). Total length 3.27–4.29 (n = 5). One paratype: body 4.29 long; carapace 1.72 long, 1.44 wide; abdomen 2.57 long, 1.62 wide. Carapace yellowish brown. Eye diameters: AME 0.11, ALE 0.10, PME 0.09, PLE 0.10. Eye interdistances: AME–AME 0.04, AME–ALE 0.01, PME–PME 0.11, PME–PLE 0.05, ALE–PLE 0.11. MOA 0.26 long, front 0.24 wide, back 0.31 wide. Clypeus 0.13 high. Leg measurements: I 6.63 (1.71, 0.65, 1.91, 1.56, 0.80); II 5.48 (1.44, 0.60, 1.42, 1.28, 0.74); III 4.67 (1.24, 0.56, 0.97, 1.20, 0.70); IV 7.29 (1.90, 0.63, 1.76, 2.00, 1.00). Leg formula: 4123. Femur I with two dorsal spines and four prolateral spines, femur II with one dorsal spine and three prolateral spines, femora III–IV with one dorsal spine; tibia I with eight pairs of ventral spines, tibia II with eight proventral spines and seven retroventral spines; metatarsus I with four pairs of ventral spines, metatarsus II with




Figure 5. *Otacilia yangmingensis* sp. n. **A** left male palp, ventral view **B** same, dorsal view **C** same, prolateral view **D** same, retrolateral view **E** epigyne, ventral view **F** same, cleared by potassium hydroxide, ventral view **G** vulva, dorsal view.




Figure 6. *Otacilia yangmingensis* sp. n. **A** left male palp, ventral view **B** same, dorsal view **C** same, retro-lateral view **D** epigyne, ventral view **E** vulva, dorsal view. Scale bars equal for **A** and **B**, equal for **D** and **E**.

four proventral spines and three retroventral spines. Abdomen light grey, anterior half without dorsal scutum, posterior half dark with several indistinct chevron-like black stripes dorsally. Other characters as in male.

Epigyne (Figs 5E–F, 6D): median plate narrow, with parallel lateral margin; copulatory openings situated centrally, covered with mating plugs (Fig. 5E), connected with a pair of shallow concavities anteriorly, and the concavities have distinct anterior and inner lateral margins. Vulva (Figs 5G, 6E): copulatory ducts thick, posteriorly swollen, connected to a pair of large, transparent long ovoid bursae; spermathecae located posteriorly and small, close to each other; bursae and spermathecae connected by slender, slightly curved connecting tubes.

Distribution. Known only from the type localities, Hunan, China (Fig. 13).

***Otacilia curvata* sp. n.**

<http://zoobank.org/EFCA1B66-8035-41A4-BB6F-634B4F16BB01>

Figs 7–9

Type material. Holotype ♂, China, *Hunan Province*: Shuangpai County, Mt. Yangming, around the Forest Park Service (26°03'36.698"N, 111°56'12.707"E), 539 m a.s.l., 24 September 2015, Chi Jin leg. Paratypes: 5♀3♂, same data as for holotype; 2♀4♂, Shuangpai County, Mt. Yangming, Wanshou Temple (26°06'27.490"N, 111°55'19.186"E), 1375 m a.s.l., 26 September 2015, Chi Jin leg.; 2♀1♂, Shuangpai County, Mt. Yangming, Hongjun Pavilion (26°04'34.924"N, 111°56'19.223"E), 1324 m a.s.l., 27 September 2015, Xiangbo Guo and Jingchao He leg.

Etymology. The specific name is derived from the Latin “curvatus”, meaning curved and refers to the shape of the DTA of the male palp; adjective.

Diagnosis. The male can be distinguished from all other *longituba* group species, except *O. bifurcata* Dankittipakul & Singtripop, 2014, *O. loriot* Jäger & Wunderlich, 2012 and *O. namkhan* Jäger & Wunderlich, 2012, by having a long RTA and a long DTA and can be distinguished from them by the needle-like embolus (embolus of these three species claw-like, knife-shaped and semicircular respectively) (Figs 8A–D, 9A–C). The female of the new species can be easily distinguished from all of the other *longituba* group species, except *O. microstoma* Wang et al., 2015, by the copulatory ducts longitudinal and close together, and it can be distinguished from *O. microstoma* by the present of glandular appendages and sigmoid connecting tubes (glandular appendages absent and connecting tubes V-shaped in *O. microstoma*) (Figs 8E–G, 9D–E; Wang et al. 2015: figs 1D–E, 2F–G).

Description. Male (Fig. 7A–B). Total length 2.51–2.80 (n = 5). Holotype: body 2.67 long; carapace 1.37 long, 1.15 wide; abdomen 1.30 long, 0.96 wide. Carapace yellowish brown, with black marginal bands; middle with broad longitudinal black stripe, from ocular area to the posterior margin of carapace; fovea longitudinal, dark brown. Diameter of eyes: AME 0.08, ALE 0.09, PME 0.06, PLE 0.10. Eye interdistances: AME–AME 0.03, AME–ALE 0.01, PME–PME 0.08, PME–PLE 0.06, ALE–PLE 0.05. MOA 0.19 long, front 0.19 wide, back 0.20 wide. Clypeus 0.12 high. Chelicerae with two strong anterior bristles; promargin with three well-separated teeth and retromargin with five denticles close to each other. Labium and sternum dark yellow. Legs light yellowish brown; all femora with distal black annulus; patellae I–II all black, patellae III–IV absenting black patches; tibia I almost all black, tibiae II–IV with black proximal and distal annulus; metatarsus I distal half part black, metatarsi II–IV with black distal annulus. Measurements of legs: leg I 4.95 (1.31, 0.51, 1.42, 1.21, 0.50), II 3.90 (1.06, 0.40, 1.02, 0.91, 0.51), III 3.36 (0.86, 0.44, 0.68, 0.86, 0.52), IV 4.90 (1.29, 0.45, 1.09, 1.41, 0.66). Leg formula: 1423. Femora I–III lack dorsal spines, femur IV with one dorsal spine, femur I with three proateral spines; tibia I with six proventral spines and seven retroventral spines, tibia II with six proventral spines and five retroventral spines; metatarsus I with four pairs of ventral spines, metatarsus II with four proventral spines and three retroventral spines. Abdomen oval,



Figure 7. *Otacilia curvata* sp. n. **A** male habitus, dorsal view **B** same, ventral view **C** female habitus, dorsal view **D** same, ventral view.




Figure 8. *Otacilia curvata* sp. n. **A** left male palp, ventral view **B** same, dorsal view **C** same, prolateral view **D** same, retrolateral view **E** epigyne, ventral view **F** same, cleared by potassium hydroxide, ventral view **G** vulva, dorsal view.




Figure 9. *Otacilia curvata* sp. n. **A** left male palp, ventral view **B** same, dorsal view **C** same, retrolateral view **D** epigyne, ventral view **E** vulva, dorsal view **F** schematic course of internal duct system. Scale bars equal for **A** and **B**, equal for **D** and **E**.

dorsum black, anterior half with a narrow dorsal scutum, posterior half with several chevron-like black stripes; venter light grey.

Palp (Figs 8A–D, 9A–C). Femur distally with an inflated hump on ventral side. RTA thick in proximal part and abruptly tapering at half of its length. DTA with anterior and posterior margins parallel in proximal part from the dorsal view, then abruptly curved to the prolateral side of bulb, tapering and with an enlarged blunt apex. Embolus short, needle-like. Conductor membranous, close to and as long as the embolus. Tegular apophysis absent but with a tegular ridge.

Female (Fig. 7C–D). Total length 2.77–2.85 (n = 7). One paratype: body 2.77 long; carapace 1.41 long, 1.20 wide; abdomen 1.36 long, 1.01 wide. Eye diameters: AME 0.08, ALE 0.09, PME 0.07, PLE 0.10. Eye interdistances: AME–AME 0.04,

AME–ALE 0.01, PME–PME 0.07, PME–PLE 0.07, ALE–PLE 0.07. MOA 0.21 long, front 0.18 wide, back 0.19 wide. Clypeus 0.11 high. Leg measurements: I 5.11 (1.31, 0.54, 1.46, 1.29, 0.51); II 4.10 (1.10, 0.49, 1.03, 0.97, 0.51); III 3.46 (0.91, 0.43, 0.73, 0.86, 0.53); IV 5.08 (1.35, 0.50, 1.12, 1.41, 0.70). Leg formula: 1423. Femur II with one dorsal spines and two pro lateral spines, tarsus II with six proventral spines and five retroventral spines, other segments with the same spination as male. Abdomen light grey, anterior half lacks dorsal scutum. Other characters as in male.

Epigyne (Figs 8E–F, 9D): median plate absent; copulatory openings situated centrally, tiny. and trumpet-shaped. Vulva (Figs 8G, 9E–F): copulatory ducts longitudinal, connecting with a pair of large, transparent, long, ovoid bursae; spermathecae located posteriorly, small and ovoid, close to each other; bursae and spermathecae connected by strong, curved, sigmoid connecting tubes. Glandular appendages present, as long as the diameter of one spermatheca.

Distribution. Known only from the type localities, Hunan, China (Fig. 13).

***Otacilia submicrostoma* sp. n.**

<http://zoobank.org/64BE3E6B-B7E9-40EB-A1F0-39A01D85D844>

Figs 10–12

Type material. Holotype ♂, China, *Hunan Province*: Sangzhi County, Bamaoxi Town, Mt. Tianping, Watch Tower (29°47'11.854"N, 110°05'28.838"E), 1626 m a.s.l., 15 September 2015, Chi Jin leg. Paratypes: 11♀7♂, same data as for holotype; 19♀19♂, Sangzhi County, Bamaoxi Town, Mt. Tianping (29°46'07.921"N, 110°04'22.159"E), 1330 m a.s.l., 16 September 2015, Xiangbo Guo and Jingchao He leg.; 2♀6♂, Sangzhi County, Bamaoxi Town, Mt. Tianping (29°46'35.332"N, 110°05'54.474"E), 1520 m a.s.l., 17 September 2015, Chi Jin leg.

Etymology. The species is named for its similarity to *O. microstoma* Wang et al., 2015; adjectival.

Diagnosis. The male can be distinguished from all other *longituba* group species, except *O. mira* Fu, Zhang & Zhang, 2016, *O. mustela* Kamura, 2008 and *O. parva* Deeleman-Reinhold, 2001, by having only one tibial apophysis and no conductor and can be distinguished from them by the RTA base with a small triangular process (Figs 11A–D, 12A–C). The female of the new species can be easily distinguished from all of the other *longituba* group species by the long, S-shaped connecting peculiar tubes (Figs 11E–F, 12D).

Description. Male (Fig. 10A–B). Total length 2.65–2.99 (n = 33). Holotype: body 2.99 long; carapace 1.52 long, 1.29 wide; abdomen 1.47 long, 1.04 wide. Carapace yellowish brown, with black marginal bands; fovea longitudinal, brown. Eye diameters: AME 0.09, ALE 0.10, PME 0.09, PLE 0.10. Eye interdistances: AME–AME 0.04, AME–ALE 0.01, PME–PME 0.11, PME–PLE 0.05, ALE–PLE 0.08. MOA




Figure 10. *Otacilia submicrostoma* sp. n. **A** male habitus, dorsal view **B** same, ventral view **C** female habitus, dorsal view **D** same, ventral view.




Figure 11. *Otacilia submicrostoma* sp. n. **A** left male palp, ventral view **B** same, dorsal view **C** same, prolateral view **D** same, retrolateral view **E** epigyne, ventral view **F** same, cleared by potassium hydroxide, ventral view **G** vulva, dorsal view.

0.22 long, front 0.21 wide, back 0.30 wide. Clypeus 0.15 high. Chelicerae with two strong anterior bristles; promargin with three well-separated teeth and retromargin with seven denticles close to each other. Labium and sternum dark yellow. Legs light




Figure 12. *Otacilia submicrostoma* sp. n. **A** left male palp, ventral view **B** same, dorsal view **C** same, retrolateral view **D** epigyne, ventral view **E** vulva, dorsal view **F** schematic course of internal duct system. Scale bars equal for **A** and **B**, equal for **D** and **E**.

yellowish brown, all femora, patellae, tibiae, metatarsi distally with black annulus. Measurements of legs: leg I 5.45 (1.42, 0.54, 1.55, 1.30, 0.64), II 4.54 (1.20, 0.47, 1.18, 1.08, 0.61), III 3.96 (1.04, 0.44, 0.83, 1.03, 0.62), IV 5.72 (1.56, 0.50, 1.26, 1.57, 0.83). Leg formula: 4123. Femur I with two dorsal spines and four prolateral spines, femur II with one dorsal spine and one prolateral spine, femora III–IV with one dorsal spine; tibia I with six proventral spines and seven retroventral spines, tibia II with six pairs of ventral spines; metatarsus I with four pairs of ventral spines, metatarsus II with four proventral spines and three retroventral spines. Abdomen oval, dorsum black, anterior half with a narrow dorsal scutum, posterior half with several black transverse stripes; venter light grey, with black scattered patches.

Palp (Figs 11A–D, 12A–C). Femur distally with an apophysis on ventral side and a retrolateral concavity. RTA broad, with relatively sharp apex extending along the cym-




Figure 13. Distribution of new species the genus *Otacilia* from Hunan, China.

bium retrolaterally, basally with a small triangular process. Embolus slender, needle-like. Tegular apophysis and conductor absent.

Female (Fig. 10C–D). Total length 3.02–3.48 (n = 32). One paratype: body 3.48 long; carapace 1.60 long, 1.36 wide; abdomen 1.88 long, 1.25 wide. Carapace yellowish brown. Eye diameters: AME 0.09, ALE 0.09, PME 0.10, PLE 0.10. Eye interdistances: AME–AME 0.04, AME–ALE 0.01, PME–PME 0.10, PME–PLE 0.06, ALE–PLE 0.09. MOA 0.25 long, front 0.20 wide, back 0.29 wide. Clypeus 0.14 high. Leg measurements: I 5.71 (1.47, 0.59, 1.64, 1.40, 0.61); II 4.87 (1.28, 0.52, 1.20, 1.23, 0.64); III 4.11 (1.09, 0.47, 0.85, 1.04, 0.66); IV 5.98 (1.60, 0.53, 1.33, 1.64, 0.88). Leg formula: 4123. Femur I with two dorsal spines and four prolateral spines, femur II with one dorsal spine and two prolateral spines, femora III–IV with one dorsal spine; tibiae and metatarsi I and II with the same spination as male. Abdomen light grey, anterior half lacking dorsal scutum, posterior half dark with several indistinct chevron-like black stripes dorsally. Other characters as in male.

Epigyne (Figs 11E–F, 12D): median plate absent; copulatory openings situated centrally, tiny and pore-like. Vulva (Figs 11G, 12E–F): copulatory ducts thick and short, connected with a pair of large, transparent, long, ovoid bursae; spermathecae located posteriorly, large and ovoid, close to each other; bursae and spermathecae connected by strongly curved, S-shaped connecting tubes. Glandular appendages present, as long as one spermatheca's diameter.

Distribution. Known only from the type localities, Hunan, China (Fig. 13).

Acknowledgments

Thanks to Xiangbo Guo and Jingchao He for their assistance during the field work in Hunan Province, China. The English of the manuscript was kindly reviewed by Dr J. MacDermott. This work was supported by the National Natural Science Foundation of China (No. 31372154), and by the Program of Ministry of Science and Technology of the Republic of China (2015FY210300).

References

- Banks N (1892) A classification of North American spiders. *Canadian Entomologist* 24: 88–97. doi: 10.4039/Ent2488-4
- Dankittipakul P, Singtripop T (2014) New species and new records of the spider genus *Otacilia* Thorell, 1897 (Araneae, Corinnidae) from Southeast Asia. *Revue Suisse de Zoologie* 121(3): 383–394.
- Deeleman-Reinhold CL (2001) Forest spiders of South East Asia: with a revision of the sac and ground spiders (Araneae: Clubionidae, Corinnidae, Liocranidae, Gnaphosidae, Prodidomidae and Trochanterriidae). Brill, Leiden, 591 pp.
- Fu LN, Jin C, Zhang F (2014) Three new species of the genus *Otacilia* Thorell (Araneae: Phrurolithidae) from China. *Zootaxa* 3869(4): 483–492. doi: 10.11646/zootaxa.3869.4.10
- Fu LN, He JC, Zhang F (2015) Species of the genus *Otacilia* from Hainan Island, China (Araneae: Phrurolithidae). *Zoological Systematics* 40(4): 436–450.
- Fu LN, Zhang ZS, Zhang F (2016a) Description of two new *Otacilia* species from Anhui, China (Araneae: Phrurolithidae). *Acta Zoologica Academiae Scientiarum Hungaricae* 62(2): 133–142. doi: 10.17109/AZH.62.2.133.2016
- Fu LN, Zhang ZS, Zhang F (2016b) New *Otacilia* species from Southwest China (Araneae: Phrurolithidae). *Zootaxa* 4107(2): 197–221. doi: 10.11646/zootaxa.4107.2.4
- Jäger P, Wunderlich J (2012) Seven new species of the spider genus *Otacilia* (Araneae: Corinnidae) from China, Laos and Thailand. *Beiträge zur Araneologie* 7: 251–271.
- Kamura T (2005) Spiders of the genus *Otacilia* (Araneae: Corinnidae) from Japan. *Acta Arachnologica*, Tokyo 53: 87–92. doi: 10.2476/asjaa.53.87
- Thorell T (1897) Viaggio di Leonardo Fea in Birmania e regioni vicine. LXXIII. Secondo saggio sui Ragni birmani. I. Parallelodontes. Tubitelariae. *Annali del Museo Civico di Storia Naturali di Genova* 17: 161–267.
- Wang LY, Chen HM, Zhou KX, Zhang F, Zhang ZS (2015) Diversity of spiders in Fanjing Mountain Nature Reserve, Guizhou, China, I: Six new species of Phrurolithidae (Araneae). *Zootaxa* 4012(3): 447–464. doi: 10.11646/zootaxa.4012.3.2
- Wang LY, Zhang F, Zhang ZS (2012) Ant-like sac spiders from Jinyun Mountain Natural Reserve of Chongqing, China (Araneae: Corinnidae). *Zootaxa* 3431: 37–53.
- World Spider Catalog (2016) World Spider Catalog. Natural History Museum Bern. <http://wsc.nmbe.ch> [accessed 7 July 2016]
- Yaginuma T (1952) Two new species (*Phrurolithus* and *Ariamnes*) found in Japan. *Arachnological News* 21: 13–16.