

The complete mitochondrial genomes of five Agrilinae (Coleoptera, Buprestidae) species and phylogenetic implications

Zhonghua Wei¹

¹ *The Key Laboratory of Southwest China Wildlife Resources Conservation of the Ministry of Education, College of Life Sciences, China West Normal University, 637009, Nanchong, Sichuan Province, China*

Corresponding author: Zhonghua Wei (wzh1164@126.com)

Academic editor: Dmitry Telnov | Received 21 January 2022 | Accepted 18 March 2022 | Published 6 April 2022

<http://zoobank.org/F8957AFF-24AE-44E5-9577-43E3160A778B>

Citation: Wei Z (2022) The complete mitochondrial genomes of five Agrilinae (Coleoptera, Buprestidae) species and phylogenetic implications. *ZooKeys* 1092: 195–212. <https://doi.org/10.3897/zookeys.1092.80993>

Abstract

Five complete mitochondrial genomes of five species from the subfamily Agrilinae were sequenced and annotated, including *Coraebus diminutus* Gebhardt, 1928 (15,499 bp), *Coraebus cloueti* Théry, 1893 (15,514 bp), *Meliboeus sinae* Obenberger, 1935 (16,108 bp), *Agrilus sichuanus* Jendek, 2011 (16,521 bp), and *Sambus femoralis* Kerremans, 1892 (15,367 bp). These mitogenomes ranged from 15,367 to 16,521 bp in length and encoded 37 typical mitochondrial genes: 13 protein-coding genes (13 PCGs), 2 ribosomal RNA genes (2 rRNAs), 22 transfer RNA genes (22 tRNAs), and a control region (CR). Most of PCGs had typical ATN start codons and terminated with TAR or an incomplete stop codon T-. Among these five mitogenomes, Leu2, Ile, Phe, Ser2, Gly, Met, and Val were the seven most frequently encoded amino acids. Interestingly, in *A. sichuanus*, a 774 bp insertion was present at *trnW* and *trnC* junction, which is unusual in Buprestidae. Additionally, phylogenetic analyses were performed based on three kinds of nucleotide matrices (13 PCGs, 2 rRNAs, and 13 PCGs + 2 rRNAs) using Bayesian inference and maximum-likelihood methods. The results showed that the clade of Buprestidae was well separated from outgroups and all Agrilinae species formed to a single highly supported clade. The tribe Coraebini was polyphyletic, as the genus *Meliboeus* (Coraebini) clustered with the genus *Trachys* (Tracheini). The rRNA genes had important impact for the tree topology of Agrilinae. Compared to the tribes Tracheini and Agrilini, the tribe Coraebini is a younger group.

Keywords

Comparative analysis, mitogenome, phylogenetic analysis

Introduction

The superfamily Buprestoidea, which contains the families Buprestidae and Schizopodidae, differs from other groups of the Elateriformia by their serrate antennae, hypognathous head, transverse suture of metaventrite present, and two connate basal abdominal ventrites (Bellamy and Volkovitsh 2016). The buprestid beetles are a large group containing six subfamilies, 521 genera, and more than 15,000 species widely distributed in the world (Bellamy 2008; Kubáň et al. 2016). The adults exhibit a broad range of host utilization in leaves, flowers, and stems, whereas larvae are mostly internal feeders on roots and stems, or feed on foliage of woody or herbaceous plants (Bellamy and Volkovitsh 2016). Only adults of the Australian *Xyroscelis crocata* were reported to feed on the sap of the host plant *Macrozamia communis* (Bellamy 1997).

Although taxonomists have made important contributions to the buprestid classification of subfamilies and tribes based on several morphological characteristics (Cobos 1980, 1986; Tôyama 1987; Hołyński 1988, 1993, 2009; Bellamy 2003), the problems of the overall classification in Buprestoidea remain unsettled.

In the past two decades, molecular systematic approaches have been used to resolve unsettled classification and phylogenetic relationships in Insecta (Short and Fikáček 2013; Cline et al. 2014; Robertson et al. 2015; Kundrata et al. 2017; Gimmel et al. 2019; Lee et al. 2020). As to Buprestidae, Bernhard et al. (2005) first used molecular phylogenetic methods based on three mitochondrial markers (*nad1*, *12S*, and *16S*) and confirmed that the *Agrilus viridis* complex, which is widely distributed across Eurasia, is monophyletic. Pentinsaari et al. (2014) and Pellegrino et al. (2017) used mitochondrial markers to evaluate the diversity of *A. viridis* complex, their results suggest that different feeding forms of *A. viridis* represent distinct species. Subsequently, Evans et al. (2015) performed the first large-scale phylogenetic trees combining nuclear and mitochondrial data from 141 species to understand the higher-level relationships in Buprestidae. In that study, the monophyly of the family Schizopodidae and subfamilies Agrilinae, Julodinae, and Galbellinae were strongly supported, while the interrelationships of Chrysochroinae and Buprestinae remained uncertain. Hansen et al. (2016) used molecular systematic methods based on nuclear and mitochondrial data (*coi* and *ak*) to investigate the relationships within *Chrysobothris femorata* species group, and their results showed that some morphological species were not well separated. Kelnarova et al. (2019) provided a molecular phylogeny of *Agrilus* species from the Northern Hemisphere and their results suggest that DNA barcoding is a powerful species identification to *Agrilus*.

During this time, the mitogenome emerged as a valuable source for higher-level phylogenetic analyses, evolutionary strategies, and genetic diversity analyses (Saccone et al. 1999; Krzywinski et al. 2011; Cameron 2014; Qin et al. 2015; Song et al. 2019; Wang et al. 2019). Several buprestid mitogenomes have been sequenced and reported, such as the mitogenome of *Chrysochroa fulgidissima* (Schönherr, 1817) by Hong et al. (2009); the mitogenome of *Agrilus planipennis* Fairmaire, 1888 by Duan et al. (2017), who also performed phylogenetic analyses based on 13 PCGs of 45 mitogenomes of coleopterans;

Table 1. Information on the mitogenomes of Buprestidae and two outgroups used in this study.

No.	Taxa	Accession no.	Genome size (bp)	A%	A+T%	AT skew	GC skew	References
1	<i>Coraeus diminutus</i>	OK189521	15,499	38.34	68.42	0.12	-0.25	This study
2	<i>Coraeus cloueti</i>	OK189520	15,514	38.53	69.27	0.11	-0.25	This study
3	<i>Meliboeus sinae</i>	OK189522	16,108	40.18	72.42	0.11	-0.22	This study
4	<i>Sambus femoralis</i>	OK349489	15,367	40.98	73.23	0.12	-0.18	This study
5	<i>Agrilus sichuanus</i>	OK189519	16,521	40.19	71.73	0.12	-0.21	This study
6	<i>Agrilus planipennis</i>	KT363854	15,942	40.25	71.90	0.12	-0.24	Duan et al. 2017
7	<i>Agrilus mali</i>	MN894890	16,204	40.34	74.46	0.08	-0.18	Sun et al. 2020
8	<i>Coraeus cavifrons</i>	MK913589	15,686	38.94	69.79	0.12	-0.18	Cao and Wang 2019b
9	<i>Trachys auricollis</i>	MH638286	16,429	38.94	71.05	0.10	-0.20	Xiao et al. 2019
10	<i>Trachys troglodytiformis</i>	KX087357	16,316	41.03	74.62	0.10	-0.19	Unpublished
11	<i>Trachys variolaris</i>	MN178497	16,771	39.92	72.11	0.11	-0.21	Cao and Wang 2019a
12	<i>Melanophila acuminata</i>	MW287594	15,853	38.74	75.66	0.02	-0.25	Peng et al. 2021
13	<i>Anthaxia chinensis</i>	MW929326	15,881	40.12	73.61	0.09	-0.29	Chen et al. 2021
14	<i>Chrysobroa fulgidissima</i>	EU826485	15,592	40.31	69.92	0.15	-0.24	Hong et al. 2009
15	<i>Acmaeodera</i> sp.	FJ613420	16,217	38.11	68.41	0.11	-0.25	Sheffield et al. 2009
16	<i>Heterocerus parallelus</i> (outgroup)	KX087297	15,845	41.90	74.03	0.13	-0.24	Unpublished
17	<i>Dryops ernesti</i> (outgroup)	KX035147	15,672	39.04	72.98	0.07	-0.23	Unpublished

the mitogenome of *Trachys variolaris* Saunders, 1873 by Cao and Wang (2019a); and the mitogenome of *Coraeus cavifrons* Descarpentries & Villiers, 1967 by Cao and Wang (2019b). More detailed information of buprestid mitogenomes is presented in Table 1.

Currently, the subfamily Agrilinae contains four tribes (Agrilini, Coraebini, Aphanisticini, and Tracheini); however, the phylogenetic placement of several genera of this subfamily remains unstable. The genera in the tribes Coraebini and Agrilini were revised by Kubáň et al. (2000). In that study, the genus *Sambus* in the tribe Coraebini was transferred to Agrilini based on the female behavior of ovipositing on rather smooth surfaces of living plants. Later, Kubáň (2016) placed the genera *Sambus*, *Parasambus*, and *Pseudagrilus* in *incertae sedis*. In order to solve these problems, we contribute mitogenomic data of five species of buprestids, *Coraeus diminutus* Gebhardt, 1928, *Coraeus cloueti* Théry, 1893, *Meliboeus sinae* Obenberger, 1935, *Agrilus sichuanus* Jendek, 2011, and *Sambus femoralis* Kerremans, 1892, and perform a molecular phylogenetic analysis in this study. The phylogenetic trees of 15 species from nine genera belonging to four subfamilies of Buprestidae were constructed based on the newly sequenced and previously reported mitogenomes (Table 1).

Material and methods

Sampling and DNA extraction

Specimens of five species were collected using an entomological net. Among them, *C. diminutus*, *C. cloueti*, *M. sinae*, and *A. sichuanus* were collected in the Dayaoshan Mountains in Guangxi Zhuang Autonomous Region, and *S. femoralis* was collected at

Yingjiang County in Yunnan Province, China. Specimens were immediately preserved in 95% ethanol in the field after collected and then stored at $-24\text{ }^{\circ}\text{C}$ in the laboratory. The specimens were identified based on morphological characteristics under a Leica M205 FA stereomicroscope. Total DNA was extracted from muscle tissues using the Ezup Column Animal Genomic DNA Purification Kit (Shanghai, China) following the manufacturer's instructions.

Sequencing, sequence assembly, annotation, and heterogeneity

DNA sequencing and de novo assembly of each mitogenome were performed by Beijing Aowei Gene Technology Co. Ltd (Beijing, China). 22 tRNA genes were identified using the MITOS webserver, with the parameters of the Invertebrate Mito genetic code (Bernt et al. 2013). Their secondary structures were plotted manually from the MITOS predictions using Adobe Illustrator. Every sequence of tRNA genes was manually checked separately. The PCGs were identified as open reading frames corresponding to the 13 PCGs. The rRNAs and control regions were identified by the boundaries of the tRNA genes. The tRNA secondary structures were identified using tRNAscan-SE (Lowe and Chan 2016). Mitogenome maps (Suppl. material 1: Fig. S1) were produced using Organellar Genome DRAW (OGDRAW) (Greiner et al. 2019). The Base composition and relative synonymous codon usage values were determined using MEGA 6.0 (Kumar 2016). Strand asymmetry was calculated using the formulae $\text{AT-skew} = (A - T) / (A + T)$, and $\text{GC-skew} = (G - C) / (G + C)$ (Perna and Kocher 1995). In the control region (CR), tandem repeat elements were detected by Tandem Repeats Finder (Benson 1999). The heterogeneous analysis of the 13 PCGs and two rRNAs datasets were performed using AliGROOVE 1.06 (Kück et al. 2014), and the nucleotide diversity (Pi) and the ratio of Ka/Ks of PCGS were calculated with DnaSP v. 5 (Librado and Rozas 2009).

Phylogenetic analyses

Phylogenetic trees for *A. sichuanus*, *C. diminutus*, *C. cloueti*, *M. sinae*, *S. femoralis*, and 10 other buprestid species belonging to four subfamilies were reconstructed by three separate datasets (13 PCGs, 2 rRNAs, and 13 PCGs + 2 rRNAs) using different best-fit models (Table 4). The mitogenomes of *Heterocerus parallelus* (Heteroceridae) and *Dryops ernesti* (Dryopidae) were used as outgroups, as they are phylogenetically distant from Buprestidae in the suborder Polyphaga (Xiao et al. 2019). The phylogenetic analyses were performed using PhyloSuite v. 1.2.2 (Zhang et al. 2020). Nucleotide sequences of the 13 PCGs and 2 rRNAs of all 17 mitogenomes were aligned using ClustalW (Thompson et al. 1994) and trimmed using trimAl v. 1.2 (Capella-Gutiérrez et al. 2009). The best-fit model for three datasets was determined by ModelFinder based on Bayesian information criterion. The maximum-likelihood (ML) and Bayesian inference (BI) methods were used to reconstruct the phylogenetic trees by IQ-tree v. 1.6.8 (Guindon et al. 2010) and MrBayes v. 3.2.6 program respectively (Ronquist et al.

2012). Bayesian analyses were run with two independent chains spanning 2,000,000 generations, four Markov chains, sampling at every 100 generations, and a burn-in period of 0.25 for each chain. The phylogenetic trees were edited and visualized by Figtree v. 1.4.3.

Results and discussion

Genome organization and base composition

The complete mitogenomes of the buprestids *A. sichuanus*, *C. diminutus*, *C. cloueti*, *M. sinae*, and *S. femoralis* have the following GenBank accession numbers attributed to them: OK189519, OK189521, OK189520, OK189522, OK349489. The mitogenomes of these five species contained the 37 typical mitochondrial genes (13 PCGs, 22 tRNAs, and 2 rRNAs) and a control region (CR) (Table 2). The composition and arrangement of the mitochondrial genes in these species (Table 2) were highly similar as those in most other buprestid species (Duan et al. 2017; Cao and Wang 2019a, 2019b; Xiao et al. 2019; Chen et al. 2021; Peng et al. 2021).

Four of the 13 PCGs (*nad1*, *nad4L*, *nad4*, and *nad5*), eight tRNAs (*trnQ*, *trnV*, *trnL1*, *trnP*, *trnH*, *trnF*, *trnY*, and *trnC*), and two rRNAs (*rrnL* and *rrnS*) are encoded on the N-strand, whereas the other 23 genes (9 PCGs and 14 tRNAs) are encoded on the J-strand. The mitogenome sequence of these five buprestid species ranged in size from 15,367 to 16,521 bp.

The mean A + T nucleotide contents of five complete mitogenomes were similar: 68.42% in *C. diminutus*, 69.27% in *C. cloueti*, 72.42% in *M. sinae*, 71.73% in *A. sichuanus*, and 73.23% in *S. femoralis*. The entire mitogenomes had a higher A + T contents of 68.42–73.23% (66.05–72.50% for PCGs, 70.95–74.03% for tRNA genes, 75.20–77.33% for rRNA genes, and 74.17–78.38% for the CR) than G + C contents, which is consistent with the typical base of buprestid mitogenomes. The overall AT skews in these five complete mitogenomes were 0.12, 0.11, 0.11, 0.12, and 0.12, respectively. These five species showed a positive TA skew, suggesting that a slight AT bias which are similar to those observed in other buprestid species (Duan et al. 2017; Cao and Wang 2019a, 2019b; Xiao et al. 2019; Chen et al. 2021; Peng et al. 2021).

Protein-coding regions, codon usage, and nucleotide diversity

The total lengths of PCGs in these five buprestid species ranged from 11,090 to 11,158 bp, accounting for 67.54–72.17% of the entire mitogenomes. Similar to the other buprestid mitogenomes, *nad5* and *atp8* were found to be the largest (1708–1723 bp) and smallest (156–159 bp) genes, respectively. The majority of PCGs strictly started with an ATN (ATA/ATT/ATC/ATG) start codon, except for the *nad1* starting with TTG. All PCGs strictly terminated with TAR (TAG/TAA) or an incomplete stop codon T-. Similar to most previously sequenced members of Buprestidae, the AT skew

Table 2. The five newly annotated Buprestidae mitogenomes. The order of these five species in the table is as follows: *Agrilus sichuanus*, *Coraeus diminutus*, *Coraeus cloneti*, *Melibeus sinae*, and *Sambus femoralis*. – not determined.

Gene	Strand	Position From	To	Start codons	Stop condons	Anticodon	Intergenic nucleotides
<i>trnI</i>	J	1/1/1/1/1	65/63/63/64/65			GAT	-3/-3/-3/5-3
<i>trnQ</i>	N	63/61/61/70/63	131/129/129/138/131			AAG	-1/0/0/0/-1
<i>trnM</i>	J	131/129/129/138/131	199/196/196/205/196			CAA	0/0/0/0/0
<i>nad2</i>	J	200/197/197/206/197	1222/1219/1219/1231/1210	ATC/ATT/ATT/ATC/ATT	TAA/TAG/TAA/TAA/TAA		1/1/-2/0/-2
<i>trnW</i>	J	1224/1221/1218/1232/1209	1293/1286/1283/1303/1273			ACA	77/4/-8/-13/13/-8
<i>trnC</i>	N	2068/1279/1276/1296/1266	2130/1339/1336/1356/1326			GCA	0/2/2/0/0
<i>trnY</i>	N	2131/1342/1339/1357/1327	2195/1404/1401/1419/1387			GAA	9/1/1/1/1
<i>coxI</i>	J	2205/1406/1403/1421/1389	3735/2936/2933/2951/2919	-/-/-/-/-	TAA/TAA/TAA/TAA/TAA		0/0/0/0/0
<i>trnL2</i>	J	3736/2937/2934/2952/2920	3802/3003/3001/3016/2984	ATT/ATA/ATA/ATC/ATT	TAA/TAA/TAA/TAA/TAA	AAG	0/0/0/0/0
<i>cox2</i>	J	3803/3004/3002/3017/2985	4484/3670/3668/3698/3666			CAA	0/0/0/0/0
<i>trnK</i>	J	4485/3671/3669/3699/3667	4553/3740/3738/3768/3736			GAC	0/0/0/0/0
<i>trnD</i>	J	4554/3741/3739/3769/3737	4618/3803/3802/3830/3798				0/0/0/0/0
<i>atp8</i>	J	4619/3804/3803/3831/3799	4777/3962/3961/3989/3954	ATT/ATA/ATC/ATT/ATA	TAG/TAA/TAA/TAA/TAG		0/-7/-7/-7-7
<i>atp6</i>	J	4771/3956/3955/3983/3948	5445/4630/4629/4657/4622	ATG/ATG/ATG/ATG/ATG	TAA/TAA/TAA/TAA/TAA		-1/-1/-1/-1/-1
<i>cox3</i>	J	5445/4630/4629/4657/4622	6233/5416/5415/5443/5405	ATG/ATG/ATG/ATG/ATG	TAG/TAA/TAA/TAA/TAA		8/0/0/0/0
<i>trnG</i>	J	6242/5417/5416/5444/5406	6306/5477/5475/5509/5469	ATT/ATT/ATT/ATT/ATT	TAG/TAG/ATG/ATG/ATG	ACC	0/0/0/0/0
<i>nad3</i>	J	6307/5478/5477/5510/5470	6660/5831/5830/5863/5823			AGC	-2/-2/-2/-2/-2
<i>trnA</i>	J	6659/5830/5829/5862/5822	6721/5890/5889/5924/5884			ACG	0/-1/-1/-1/0
<i>trnR</i>	J	6722/5890/5889/5924/5885	6781/5952/5951/5988/5947			ACG	1/-1/-1/-1/1
<i>trnN</i>	J	6783/5952/5951/5988/5949	6849/6017/6016/6051/6013			GAA	0/0/0/0/0
<i>trnS1</i>	J	6850/6018/6017/6052/6014	6916/6075/6074/6117/6080			ACA	1/0/7/-1/0
<i>trnE</i>	J	6918/6076/6082/6117/6081	6982/6139/6143/6179/6143			AAC	-1/-4/-4/-1/-1
<i>trnF</i>	N	6982/6136/6140/6179/6143	7045/6198/6202/6240/6207	ATA/ATT/ATT/ATT/ATA	TAA/TAA/TAA/TAA/TAA	GAA	0/0/0/0/0
<i>nad5</i>	N	7046/6199/6203/6241/6208	8768/7915/7919/7960/7915			GAG	0/0/0/0/0
<i>trnH</i>	N	8769/7916/7920/7961/7916	8830/7977/7981/8026/7978				0/0/0/0/0
<i>trnH</i>	N	8831/7978/7982/8027/7979	10,166/9295/9299/9362/9308	ATG/ATG/ATG/ATG/ATG	TAA/TAA/TAA/TAA/TAA		0/0/0/0/0
<i>nad4</i>	N	10,160/9289/9293/9356/9302	10,444/9576/9580/9640/9589	ATG/ATG/ATG/ATG/ATA	TAA/TAA/TAA/TAA/TAA		0/0/0/0/0
<i>nad4L</i>	N	10,449/9580/9584/9643/9591	10,511/9642/9646/9704/9654				0/0/0/0/0
<i>trnT</i>	J	10,511/9642/9646/9704/9654	10,574/9704/9708/9769/9717			AGA	-1/-1/-1/-1/-1
<i>trnP</i>	J	10,576/9706/9710/9771/9719	11,079/10,185/10,189/10,259/10,192	ATT/ATA/ATA/ATG/ATT	TAA/TAA/TAA/TAA/TAA	AGG	1/1/1/1/1
<i>cyb</i>	J	11,079/10,185/10,189/10,259/10,192	12,224/11,327/11,331/11,401/11,334	ATG/ATG/ATG/ATG/ATG	TAA/TAG/ATG/ATG/ATG		-1/-1/-1/-1/-1
<i>trnS2</i>	J	12,233/11,326/11,330/11,400/11,333	12,298/11,391/11,395/11,465/11,400			ACA	8/-2/-2/-2/-2
<i>nad1</i>	N	12,316/11,411/11,415/11,485/11,415	13,266/12,361/12,365/12,432/12,365				17/9/9/19/14
<i>trnL1</i>	N	13,268/12,363/12,367/12,433/12,367	13,334/12,427/12,431/12,495/12,434	TTG/TTG/TTG/TTG/TTG	TAA/TAA/TAA/TAG/ATA	AAG	1/1/1/0/0
<i>trnL</i>	N	13,335/12,428/12,432/12,496/12,435	14,605/13,693/13,697/13,757/13,692				0/0/0/0/0
<i>trnV</i>	N	14,606/13,694/13,698/13,758/13,693	14,674/13,762/13,766/13,826/13,761			AAC	0/0/0/0/0
<i>trnS</i>	N	14,675/13,763/13,767/13,827/13,762	15,379/14,480/14,483/14,531/14,457				0/0/0/0/0
A + T rich region		15,380/14,481/14,484/14,532/14,458	16,521/15,499/15,516/16,108/15,367				0/0/0/0/0

Table 3. Summarized mitogenomic characteristics of the five buprestid species in this study.

Species	PCGs			rRNAs			tRNA			CR		
	Size (bp)	A+T content	AT skew	Size (bp)	A+T content	AT skew	Size (bp)	A+T content	AT skew	Size (bp)	A+T content	AT skew
<i>A. sichuanus</i>	11,158	70.08	-0.15	1976	75.96	-0.13	1444	74.03	-0.0009	1142	74.17	0.06
<i>C. diminutus</i>	11,093	66.05	-0.14	1984	75.20	-0.11	1477	70.95	0.03	1019	77.72	0.02
<i>C. cloueti</i>	11,093	67.09	-0.15	1983	75.39	-0.11	1414	71.22	0.019	1031	78.27	0.02
<i>M. sinæ</i>	11,135	70.70	-0.15	1967	77.33	-0.11	1435	72.13	0.007	1577	78.38	0.13
<i>S. femoralis</i>	11,090	72.50	-0.16	1954	75.69	-0.13	1430	73.85	0.03	910	75.82	0.18

Table 4. Best-fit models of three datasets used for phylogeny.

	ML method	BI method
13 PCGs	GTR+F+I+G4	GTR+F+I+G4
2 rRNAs	TVM+F+I+G4	GTR+F+I+G4
13 PCGs +2 rRNAs	GTR+F+I+G4	GTR+F+I+G4

(0.11–0.12) of these five PCGs (Table 3) were similar among the 15 buprestid species. Summaries of the numbers of amino acids in the annotated PCGs and relative synonymous codon usage are presented in Figs 1 and 2. Overall codon usage among the sequenced buprestid mitogenomes was found to be similar, with Leu2, Ile, Phe, Ser2, Gly, Met, and Val being the seven most frequently coded amino acids.

The nucleotide diversity (Pi) of the 13 PCGs among five species of Agrilinae is provided (Fig. 3), which ranged from 0.202 to 0.375. In these genes, *nad2* (Pi = 0.375) presented the highest variability, followed by *nad6* (Pi = 0.346), *nad4* (Pi = 0.300), and *nad5* (Pi = 0.290); *cox1* (Pi = 0.20) exhibited the lowest variability. The ratio of Ka/Ks (Fig. 4) for each gene of the 13 PCGs was calculated. The values of *nad4* and *nad4L* are distinctly higher than others, which suggests that the genes *nad4* and *nad4L* have a relatively higher evolutionary rate.




Figure 1. Numbers of different amino acids in the mitogenomes of the five buprestid species; the stop codon is not included. AS: *Agrilus sichuanus*, CC: *Coraeus cloueti*, CD: *Coraeus diminutus*, MS: *Melioboeus sinæ*, and SF: *Sambus femoralis*.




Figure 2. RSCU (relative synonymous codon usage) of the mitogenomes of the five buprestid species; the stop codons are not included.




Figure 3. Nucleotide diversity (Pi) of 13 PCGs among five newly sequenced Agrilinae mitogenomes.




Figure 4. The ratio of Ka/Ks of 13 PCGs among the 15 reported Buprestidae mitogenomes.

tRNA, rRNA genes, and heterogeneity

The length of *rrnL* genes ranged from 1258 bp (*S. femoralis*) to 1271 bp (*A. sichuanus*), whereas *rrnS* ranged from 696 bp (*S. femoralis*) to 718 bp (*C. diminutus*). The A + T content of the rRNA genes ranged from 75.20% (*C. diminutus*) to 77.33% (*M. sinae*) (Table 3). Compared with those in other sequenced buprestid mitogenom-




Figure 5. The predicted secondary structures of the tRNA-Ser in the mitogenomes of the five buprestid species.

es, the rRNA genes in these five newly sequenced buprestid mitogenomes are highly conserved (Hong et al. 2009; Duan et al. 2017; Cao and Wang 2019a, 2019b; Xiao et al. 2019; Sun et al. 2020; Chen et al. 2021; Peng et al. 2021). These rRNAs were located between the CR and *trnL1*, and separated by *trnV*. The total lengths of the 22 tRNA genes ranged from 1414 bp (*C. cloueti*) to 1444 bp (*C. diminutus*), whereas individual tRNA genes typically ranged in size from 58 to 70 bp, among which, eight tRNAs were encoded on the N-strand and the remaining 14 encoded on the J-strand. The secondary structures of tRNAs showed a standard clover-leaf structure (Suppl. material 1: Figs S2–S6), except for tRNA-Ser (Fig. 5) which lacks or has an unusual dihydrouridine arm, which forms a loop commonly found in other insects (Xiao et al. 2011; Park et al. 2012; Yu et al. 2016; Yan et al. 2017; Yu and Liang 2018; Li et al. 2019). In *A. sichuanus*, the longest intergenic nucleotide (774 bp) was located between *trnW* and *trnC*, which is an interesting and specific phenomenon in Buprestidae. The degree of heterogeneity of the 13 PCGs dataset was higher than that of the two rRNAs dataset (Suppl. material 1: Fig. S7). Additionally, the heterogeneity in sequence divergences was slightly stronger for *Coraebus* than for other buprestid genera (Suppl. material 1: Fig. S7).

Control region

The CR, also known as the A + T-rich region (Wolstenholme 1992), was the largest non-coding region and located between *trnI* and *rrnS*. The length of CR ranged from 910 bp (*S. femoralis*) to 1577 bp (*M. sinae*). The A + T content (74.17–78.38%) of the CR of these five species was found to be higher than that of the whole genome (68.42–73.23%), PCGs (66.05–72.50%), rRNAs (75.20–77.33%), and tRNAs (70.95–73.85%) (Table 3). Moreover, the compositional analysis revealed that the mitogenomes of the five buprestid species had a positive AT skew (0.02–0.18) in the CR. In these five species, only *C. cloueti* and *C. diminutus* had no tandem repeat element detected; however, those of *A. sichuanus* (20 and 40 bp), *M. sinae* (53 bp), and *S. femoralis* (265 bp) had different lengths.

Phylogenetic analyses

Both ML and BI trees using three datasets produced identical topologies (Figs 6–8), (Buprestinae + ((Chrysochroninae + Polycestinae) + Agrilinae)), in terms of subfamily-level relationship. The monophyly of Buprestidae is corroborated again, as all the buprestid species converged together as an independent clade, and two outgroup taxa obviously separated from the buprestid clade. The target species *C. diminutus*, *C. cloueti*, *Meliboeus sinae*, *Agrilus sichuanus*, and *Sambus femoralis*, as well as other species of Agrilinae, converged together as an independent clade. And the target species, *M. sinae*,




Figure 6. Phylogenetic relationships of 15 selected buprestid species using both BI and ML analyses based on 13 PCGs of mitogenomes. The numbers on the branches show posterior probability (BI tree), whereas the values under branches are bootstrap (ML tree).




Figure 7. Phylogenetic relationships of 15 selected buprestid species using both BI and ML analyses based on 2 rRNAs of mitogenomes. The numbers on the branches show posterior probability (BI tree), whereas the values under branches are bootstrap (ML tree).




Figure 8. Phylogenetic relationships of 15 selected buprestid species using both BI and ML analyses based on 13 PCGs + 2 rRNAs of mitogenomes. The numbers on the branches show posterior probability (BI tree), whereas the values under branches are bootstrap (ML tree).

was most closely related to the genus *Trachys* with high value support (Figs 6–8) which is inconsistent with the previous studies (Kubáň et al. 2000; Evans et al. 2015). The relationship of Agrilinae clades obtained from 2 rRNAs and 13 PCGs + 2 rRNAs datasets are identical but with different topology from the 13 PCGs dataset. In the topology generated from the 13 PCGs dataset, *S. femoralis* and *Agrilus* were clustered into a single branch with high support value (Fig. 6, ML: 77, BI: 1) whereas, in the topology generated from the 2 rRNAs and 13 PCGs + 2 rRNAs datasets, *S. femoralis* split from base of the Agrilinae clades (Figs 7, 8). Based on these results the position of the genus *Sambus* in the tribe Agrilini was not suitable and suspect. The different tree topologies suggested that the rRNA genes were extremely valuable for the phylogenetic analysis of Agrilinae. Coraebini is the most diverse tribe in Agrilinae, and 10 subtribes are defined (Kubáň et al. 2000). The genus *Meliboeus* (Meliboeina) and *Coraebus* (Coraebina) in different clades suggested that the tribe Coraebini was polyphyletic, which is consistent with the previous study of Evans et al. (2015). The samples used in this study might be too limited for a comprehensive phylogeny of Buprestidae which still needs a deep study in the future.

Conclusions

In this study, five mitogenomes (15,367–16,521 bp) were newly sequenced and annotated, including representatives from the tribes Coraebini and Agrilini in subfamily Agriinae. The mitogenomes of the genera *Sambus* and *Meliboeus* are reported for the first time. These five sequences showed a positive AT skew, and the amino acids Leu, Ile, Phe, Ser2, Gly, Met, and Val were most frequently used. The secondary structures of tRNA-Ser are absent the D-arm, which is similar to other orders of Insecta. The rRNA genes are valuable for phylogenetic analyses of Agrilinae as they could affect the tree topologies. The results show that Coaebini is polyphyletic, and the genus *Sambus* belongs to neither Coaebini nor Agrilini. However, more mitogenome samplings are needed to resolve the phylogeny of the Buprestidae in the future to better understand the phylogenetics of jewel beetles.

Acknowledgements

I am sincerely grateful to Lanrui Wang (Qingyang, Gansu, China) and Yingqi Liu (China Agricultural University, Beijing, China) for their guidance in using the software. I also thank Dr Hui-Feng Zhao (Langfang Normal University, Hebei, China) and Menglin Wang (China West Normal University, Sichuan, China) for revising the manuscript. This work was supported by the Doctoral Scientific Research Foundation of China West Normal University (20E054).

References

- Bellamy CL (1997) Phylogenetic relationships of *Xyroscelis* (Coleoptera: Buprestidae). *Invertebrate Systematics* 11(4): 569–574. <https://doi.org/10.1071/IT94026>
- Bellamy CL (2003) An illustrated summary of the higher classification of the superfamily Buprestoidea (Coleoptera). *Folia Heyrovskyana (Supplementum 10)*: 1–197.
- Bellamy CL (2008) A World Catalogue and Bibliography of the Jewel Beetles (Coleoptera: Buprestoidea). Volumes 1–4. Pensoft series faunistica No. 76–79, Sofia/Moscow, [8] + 2684 pp.
- Bellamy CL, Volkovitsh M (2016) 18 Buprestoidea Crowson, 1955. In: Beutel RG, Leschen RAB (Eds) *Handbook of Zoology, Arthropoda: Insecta, Volume 1: Morphology and Systematics (Archostemata, Adephaga, Myxophaga, Polyphaga partim)*, 2nd edn. Walter de Gruyter, Berlin/Boston, 543–552. <https://doi.org/10.1515/9783110373929-021>
- Benson G (1999) Tandem repeats finder: A program to analyze DNA sequences. *Nucleic Acids Research* 27(2): 573–580. <https://doi.org/10.1093/nar/27.2.573>
- Bernhard D, Fritzsche G, Glöckner P, Würst C (2005) Molecular insights into speciation in the *Agrilus viridis*-complex and the genus *Trachys* (Coleoptera: Buprestidae). *European Journal of Entomology* 102(4): 599–605. <https://doi.org/10.14411/eje.2005.083>
- Bernt M, Donath A, Juhling F, Externbrink F, Florentz C, Fritzsche G, Stadler PF (2013) MITOS: Improved de novo metazoan mitochondrial genome annotation. *Molecular Phylogenetics and Evolution* 69(2): 313–319. <https://doi.org/10.1016/j.ympev.2012.08.023>
- Cameron SL (2014) Insect mitochondrial genomics: Implications for evolution and phylogeny. *Annual Review of Entomology* 59(1): 95–117. <https://doi.org/10.1146/annurev-ento-011613-162007>
- Cao LM, Wang XY (2019a) The complete mitochondrial genome of the jewel beetle *Trachys variolaris* (Coleoptera: Buprestidae). *Mitochondrial DNA, Part B, Resources* 4(2): 3042–3043. <https://doi.org/10.1080/23802359.2019.1666053>
- Cao LM, Wang XY (2019b) The complete mitochondrial genome of the jewel beetle *Coraebus cavifrons* (Coleoptera: Buprestidae). *Mitochondrial DNA, Part B, Resources* 4(2): 2407–2408. <https://doi.org/10.1080/23802359.2019.1636730>
- Capella-Gutiérrez S, Silla-Martínez JM, Gabaldón T (2009) TrimAl: A tool for automated alignment trimming in large-scale phylogenetic analyses. *Bioinformatics (Oxford, England)* 25(15): 1972–1973. <https://doi.org/10.1093/bioinformatics/btp348>
- Chen B, Wei ZH, Shi AM (2021) The complete mitochondrial genome of the jewel beetle, *Anthaxia chinensis* (Coleoptera: Buprestidae). *Mitochondrial DNA, Part B, Resources* 6(10): 2962–2963. <https://doi.org/10.1080/23802359.2021.1973920>
- Cline AR, Smith TR, Miller K, Moulton M, Whiting M, Audisio P (2014) Molecular phylogeny of Nitidulidae: assessment of subfamilial and tribal classification and formalization of the family Cybocephalidae (Coleoptera: Cucujoidea). *Systematic Entomology* 39(4): 758–772. <https://doi.org/10.1111/syen.12084>
- Cobos A (1980) Ensayo sobre los géneros de la subfamilia Polycestinae (Coleoptera, Buprestidae) (Parte I). *EOS. Revista Española de Entomología* 54: 15–94.
- Cobos A (1986) *Fauna Iberica de Coleopteros Buprestidae*. Consejo Superior de Investigaciones Científicas, Madrid, 364 pp.

- Duan J, Quan GX, Mittapalli O, Cusson M, Krell PJ, Doucet D (2017) The complete mitogenome of the Emerald Ash Borer (EAB), *Agrilus planipennis* (Insecta: Coleoptera: Buprestidae). Mitochondrial DNA, Part B, Resources 2(1): 134–135. <https://doi.org/10.1080/23802359.2017.1292476>
- Evans AM, Mckenna DD, Bellamy CL, Farrell BD (2015) Large-scale molecular phylogeny of metallic wood-boring beetles (Coleoptera: Buprestoidea) provides new insights into relationships and reveals multiple evolutionary origins of the larval leaf-mining habit. Systematic Entomology 40(2): 385–400. <https://doi.org/10.1111/syen.12108>
- Gimmel ML, Bocakova M, Gunter NL, Leschenm RAB (2019) Comprehensive phylogeny of the Cleroidea (Coleoptera: Cucujiformia). Systematic Entomology 44(3): 527–558. <https://doi.org/10.1111/syen.12338>
- Greiner S, Lehwerk P, Bock R (2019) OrganellarGenomeDRAW (OGDRAW) version 1.3.1: Expanded toolkit for the graphical visualization of organellar genomes. Nucleic Acids Research 47(W1): W59–W64. <https://doi.org/10.1093/nar/gkz238>
- Guindon S, Dufayard J, Lefort V, Anisimova M, Hordijk W, Gascuel O (2010) New algorithms and methods to estimate maximum-likelihood phylogenies: Assessing the performance of PhyML 3.0. Systematic Biology 59(3): 307–321. <https://doi.org/10.1093/sysbio/syq010>
- Hansen JA, Moulton JK, Klingeman WE, Oliver JB, Windham MT, Trigiano RN, Reding ME (2016) Molecular systematics of the *Chrysobothris femorata* species group (Coleoptera: Buprestidae). Annals of the Entomological Society of America 108(5): 950–963. <https://doi.org/10.1093/aesa/sav080>
- Hołyński RB (1988) Remarks on the general classification of Buprestidae Leach as applied to Maoraxiina. Folia Entomologica Hungarica 49(1): 49–54.
- Hołyński RB (1993) A reassessment of the internal classification of the Buprestidae Leach (Coleoptera). Crystal. Series Zoologica (Göd) 1: 1–42.
- Hołyński RB (2009) Taxonomic structure of the subtribe Chrysochroina Cast. with review of the genus *Chrysochroa* Dej. Gondwana, Warszawa, 391 pp.
- Hong MY, Jeong HC, Kim MJ, Jeong HU, Lee SH, Kim I (2009) Complete mitogenome sequence of the jewel beetle, *Chrysochroa fulgidissima* (Coleoptera: Buprestidae). Mitochondrial DNA Mapping, Sequencing, and Analysis 20(2–3): 46–60. <https://doi.org/10.1080/19401730802644978>
- Kelnarova I, Jendek E, Grebennikov VV, Bocak L (2019) First molecular phylogeny of *Agrilus* (Coleoptera: Buprestidae), the largest genus on Earth, with DNA barcode database for forestry pest diagnostics. Bulletin of Entomological Research 109(2): 200–211. <https://doi.org/10.1017/S0007485318000330>
- Krzywinski J, Li C, Morris M, Conn JE, Lima JB, Pova MM, Wilkerson RC (2011) Analysis of the evolutionary forces shaping mitochondrial genomes of a Neotropical malaria vector complex. Molecular Phylogenetics and Evolution 58(3): 469–477. <https://doi.org/10.1016/j.ympev.2011.01.003>
- Kubáň V (2016) Tribe Agrilini, genera incertae sedis. In: Löbl I, Löbl D (Eds) Catalogue of Palaearctic Coleoptera. Volume 3, Scarabaeoidea, Scirtoidea, Dascilloidea, Buprestoidea, Byrrhoidea. Revised and updated edition; Leiden, Boston, 549 pp.

- Kubáň V, Majer K, Kolibáč J (2000) Classification of the tribe Coraebini Bedel, 1921 (Coleoptera, Buprestidae, Agrilinae). *Acta Musei Moraviae. Scientiae Biologicae* (Brno) 85: 185–287.
- Kubáň V, Volkovitsh MG, Kalashian MJ, Jendek E (2016) Family Buprestidae Leach, 1815. In: Löbl I, Löbl D (Eds) *Catalogue of Palaearctic Coleoptera. Volume 3, Scarabaeoidea, Scirtoidea, Dascilloidea, Buprestoidea, Byrrhoidea. Revised and Updated Edition.* Apollo Books, Stenstrup, 432–574.
- Kück P, Meid SA, Groß C, Wägele JW, Misof B (2014) AliGROOVE—visualization of heterogeneous sequence divergence within multiple sequence alignments and detection of inflated branch support. *Bioinformatics* (Oxford, England) 15: e294. <https://doi.org/10.1186/1471-2105-15-294>
- Kumar S, Stecher G, Tamura K (2016) MEGA7: Molecular Evolutionary Genetics Analysis version 7.0 for bigger datasets. *Molecular Biology and Evolution* 33(7): 1870–1874. <https://doi.org/10.1093/molbev/msw054>
- Kundrata R, Jäch MA, Bocak L (2017) Molecular phylogeny of the Byrrhoidea–Buprestoidea complex (Coleoptera, Elateriformia). *Zoologica Scripta* 46(2): 150–164. <https://doi.org/10.1111/zsc.12196>
- Lee MH, Lee S, Leschen RAB, Lee S (2020) Evolution of feeding habits of sap beetles (Coleoptera: Nitidulidae) and placement of Calonecrinae. *Systematic Entomology* 45(4): 911–923. <https://doi.org/10.1111/syen.12441>
- Li R, Shu XH, Li XD, Meng L, Li BP (2019) Comparative mitogenome analysis of three species and monophyletic inference of Catantopinae (Orthoptera: Acridoidea). *Genomics* 111(6): 1728–1735. <https://doi.org/10.1016/j.ygeno.2018.11.027>
- Librado P, Rozas J (2009) DnaSP v5: A software for comprehensive analysis of DNA polymorphism data. *Bioinformatics* (Oxford, England) 25(11): 1451–1452. <https://doi.org/10.1093/bioinformatics/btp187>
- Lowe TM, Chan PP (2016) tRNAscan-SE On-line: Integrating search and context for analysis of transfer RNA genes. *Nucleic Acids Research* 33(W1): W686–W689. <https://doi.org/10.1093/nar/gkw413>
- Park JS, Cho Y, Kim MJ, Nam SH, Kim I (2012) Description of complete mitochondrial genome of the black-veined white, *Aporia crataegi* (Lepidoptera: Papilionoidea), and comparison to papilionoid species. *Journal of Asia-Pacific Entomology* 15(3): 331–341. <https://doi.org/10.1016/j.aspen.2012.01.002>
- Pellegrino I, Curletti G, Liberatore F, Cucco M (2017) Cryptic diversity of the jewel beetles *Agrilus viridis* (Coleoptera: Buprestidae) hosted on hazelnut. *The European Zoological Journal* 84(1): 465–472. <https://doi.org/10.1080/24750263.2017.1362050>
- Peng XJ, Liu J, Wang Z, Zhan QZ (2021) The complete mitochondrial genome of the pyrophilous jewel beetle *Melanophila acuminata* (Coleoptera: Buprestidae). *Mitochondrial DNA. Part B, Resources* 6(3): 1059–1060. <https://doi.org/10.1080/23802359.2021.1899079>
- Pentinsaari M, Mutanen M, Kaila L (2014) Cryptic diversity and signs of mitochondrial introgression in the *Agrilus viridis* species complex (coleoptera: Buprestidae). *European Journal of Entomology* 111(4): 475–486. <https://doi.org/10.14411/eje.2014.072>
- Perna NT, Kocher TD (1995) Patterns of nucleotide composition at fourfold degenerate sites of animal mitochondrial genomes. *Journal of Molecular Evolution* 3(3): 353–358. <https://doi.org/10.1007/BF01215182>

- Qin J, Zhang YZ, Zhou X, Kong XB, Wei SJ, Ward RD, Zhang AB (2015) Mitochondrial phylogenomics and genetic relationships of closely related pine moth (Lasiocampidae: *Dendrolimus*) species in China, using whole mitochondrial genomes. *Genomics* 16(1): e428. <https://doi.org/10.1186/s12864-015-1566-5>
- Robertson JA, Ślipiński A, Moulton M, Shockley FW, Gorgi A, Lord NP, McKenna DD, Tomaszewska W, Forrester J, Miller KB, Whiting MF, McHugh JV (2015) Phylogeny and classification of Cucujoidea and the recognition of a new superfamily Coccinelloidea (Coleoptera: Cucujiformia). *Systematic Entomology* 40(4): 745–778. <https://doi.org/10.1111/syen.12138>
- Ronquist F, Teslenko M, Der Mark PV, Ayres DL, Darling AE, Höhna S, Huelsenbeck JP (2012) MrBayes 3.2: Efficient Bayesian phylogenetic inference and model choice across a large model space. *Systematic Biology* 61(3): 539–542. <https://doi.org/10.1093/sysbio/sys029>
- Saccone C, De Giorgi C, Gissi C, Pesole G, Reyes A (1999) Evolutionary genomics in Metazoa: The mitochondrial DNA as a model system. *Gene* 238(1): 195–209. [https://doi.org/10.1016/S0378-1119\(99\)00270-X](https://doi.org/10.1016/S0378-1119(99)00270-X)
- Sheffield NC, Song H, Cameron SL, Whiting MF (2009) Nonstationary evolution and compositional heterogeneity in beetle mitochondrial phylogenomics. *Systematic Biology* 58(4): 381–394. <https://doi.org/10.1093/sysbio/syp037>
- Short AEZ, Fikáček M (2013) Molecular phylogeny, evolution, and classification of the Hydrophilidae (Coleoptera). *Systematic Entomology* 38(4): 723–752. <https://doi.org/10.1111/syen.12024>
- Song F, Li H, Liu GH, Wang W, James P, Colwell DD, Tran A, Gong S, Cai WZ, Shao R (2019) Mitochondrial genome fragmentation unites the parasitic lice of eutherian mammals. *Systematic Biology* 68(3): 430–440. <https://doi.org/10.1093/sysbio/syy062>
- Sun HQ, Zhao WX, Lin RZ, Zhou ZF, Huai WX, Yao YX (2020) The conserved mitochondrial genome of the jewel beetle (Coleoptera: Buprestidae) and its phylogenetic implications for the suborder Polyphaga. *Genomics* 112(5): 3713–3721. <https://doi.org/10.1016/j.ygeno.2020.04.026>
- Thompson JD, Higgins DG, Gibson TJ (1994) CLUSTAL W: Improving the sensitivity of progressive multiple sequence alignment through sequence weighting, position-specific gap penalties and weight matrix choice. *Nucleic Acids Research* 22(22): 4673–4680. <https://doi.org/10.1093/nar/22.22.4673>
- Tôyama M (1987) The systematic positions of some buprestid genera (Coleoptera, Buprestidae). *Elytra* 15: 1–11.
- Wang WQ, Huang YX, Bartlett CR, Zhou FM, Meng H, Qin DZ (2019) Characterization of the complete mitochondrial genomes of two species of the genus *Aphaena* Guérin-Méneville (Hemiptera: Fulgoridae) and its phylogenetic implications. *International Journal of Biological Macromolecules* 141: 29–40. <https://doi.org/10.1016/j.ijbiomac.2019.08.222>
- Wolstenholme DR (1992) Animal mitochondrial DNA: Structure and evolution. *International Review of Cytology* 141: 173–216. [https://doi.org/10.1016/S0074-7696\(08\)62066-5](https://doi.org/10.1016/S0074-7696(08)62066-5)
- Xiao JH, Jia JG, Murphy RW, Huang DW (2011) Rapid evolution of the mitochondrial genome in chalcidoid wasps (Hymenoptera: Chalcidoidea) driven by parasitic lifestyles. *PLoS ONE* 6(11): e26645. <https://doi.org/10.1371/journal.pone.0026645>

- Xiao LF, Zhang SD, Long CB, Guo QY, Xu JS, Dai XH, Wang JG (2019) Complete mitogenome of a leaf-mining buprestid Beetle, *Trachys auricollis*, and its phylogenetic implications. *Genes* 10(12): e992. <https://doi.org/10.3390/genes10120992>
- Yan L, Zhang M, Gao Y, Pape T, Zhang D (2017) First mitogenome for the subfamily Miltoigramminae (Diptera: Sarcophagidae) and its phylogenetic implications. *European Journal of Entomology* 114: 422–429. <https://doi.org/10.14411/eje.2017.054>
- Yu F, Liang AP (2018) The complete mitochondrial genome of *Ugyops* sp. (Hemiptera: Delphacidae). *Journal of Insect Science* 18(3): e25. <https://doi.org/10.1093/jisesa/iey063>
- Yu P, Cheng X, Ma Y, Yu D, Zhang J (2016) The complete mitochondrial genome of *Brachythemis contaminata* (Odonata: Libellulidae). *Mitochondrial DNA A DNA Mapping Sequencing Analysis* 27: 2272–2273. <https://doi.org/10.3109/19401736.2014.984176>
- Zhang D, Gao F, Jakovlić I, Zou H, Zhang J, Li WX, Wang GT (2020) PhyloSuite: An integrated and scalable desktop platform for streamlined molecular sequence data management and evolutionary phylogenetics studies. *Molecular Ecology Resources* 20(1): 348–355. <https://doi.org/10.1111/1755-0998.13096>

Supplementary material I

Figures S1–S7

Authors: Zhonghua Wei

Data type: Images (pdf file)

Explanation note: Figure S1. The mitogenome maps of *Agrilus sichuanus*, *Coraebus cloueti*, *Coraebus diminutus*, *Meliboeus sinae*, and *Sambus femoralis*. Figure S2. The secondary cloverleaf structure for the tRNAs of *Agrilus sichuanus*. Figure S3. The secondary cloverleaf structure for the tRNAs of *Coraebus cloueti*. Figure S4. The secondary cloverleaf structure for the tRNAs of *Coraebus diminutus*. Figure S5. The secondary cloverleaf structure for the tRNAs of *Meliboeus sinae*. Figure S6. The secondary cloverleaf structure for the tRNAs of *Sambus femoralis*. Figure S7. Heterogeneous sequence divergence within datasets 13 PCGs and 2 rRNAs of Buprestidae species.

Copyright notice: This dataset is made available under the Open Database License (<http://opendatacommons.org/licenses/odbl/1.0/>). The Open Database License (ODbL) is a license agreement intended to allow users to freely share, modify, and use this Dataset while maintaining this same freedom for others, provided that the original source and author(s) are credited.

Link: <https://doi.org/10.3897/zookeys.1092.80993.suppl1>