RESEARCH ARTICLE



# Complete mitochondrial genomes of two catfishes (Siluriformes, Bagridae) and their phylogenetic implications

Renyi Zhang<sup>1\*</sup>, Lei Deng<sup>1\*</sup>, Xiaomei Lv<sup>1</sup>, Qian Tang<sup>1</sup>

School of Life Sciences, Guizhou Normal University, 116 Baoshan Road, Guiyang, Guizhou, 550001, China

Corresponding author: Renyi Zhang (zhangrenyi@gznu.edu.cn)

Academic editor: Tihomir Stefanov   Received 12 April 2022   Accepted 12 July 2022   Published	29 July 2022
https://zoobank.org/7A01B01B-92F7-469B-B2BE-81F1B9324D2F	

**Citation:** Zhang R, Deng L, Lv X, Tang Q (2022) Complete mitochondrial genomes of two catfishes (Siluriformes, Bagridae) and their phylogenetic implications. ZooKeys 1115: 103–116. https://doi.org/10.3897/zooKeys.1115.85249

#### Abstract

The mitochondrial genome (mitogenome) has been widely used as a molecular marker to investigate phylogenetic analysis and evolutionary history in fish. However, the study of mitogenomes is still scarce in the family Bagridae. In this study, the mitogenomes of *Tachysurus brachyrhabdion* and *T. gracilis* were sequenced, annotated, and analyzed. The mitogenomes were found to be 16,532 bp and 16,533 bp, respectively, and each contained 37 typical mitochondrial genes, which are 13 protein-coding genes (PCGs), 22 tRNA genes, two rRNA genes, and a control region. All PCGs begin with the codon ATG, except for the cytochrome c oxidase subunit 1 (*COI*) gene, while seven PCGs end with an incomplete termination codon. All tRNA genes can fold into their typical cloverleaf secondary structures, except for tRNA<sup>Ser(AGY)</sup>, which lacks the dihydrouracil arm. The Ka/Ks ratios for all PCGs are far lower than one. Phylogenetic analyses based on Bayesian inference (BI) and maximum likelihood (ML) showed that the two clades in Bagridae excluded *Rita rita*. The monophyly of *Tachysurus* supports previous research and the traditional classification that *Leiocassis, Pseudobagrus, Pelteobagrus,* and *Tachysurus* belong to one genus (*Tachysurus*). These findings provide a phylogenetic basis for future phylogenetic and taxonomic studies of Bagridae.

#### Keywords

bagrid catfish, mitogenome, phylogenetic analysis, Tachysurus brachyrhabdion, Tachysurus gracilis

<sup>\*</sup> These authors contributed equally to this work.

#### Introduction

The family Bagridae, commonly known as bagrid catfish, is widely distributed in Asia and Africa, with about 225 species in 19 genera (Fricke et al. 2022). It is one of the most diverse and complicated groups within Siluriformes. Regan (1911) first promoted bagrid fishes as a single-family and named Bagridae, including two subfamilies: Chrysichthyinae and Bagrinae. Based on osteological features, Jayaram (1968) divided Bagridae into five subfamilies: Ritinae, Chiysichthyinae, Bagrinae, Bagroidinae, and Auchenoglaninae. In 1992, Mo (1992) divided the traditional bagrid catfishes into three monophyletic groups: Caroteidae, Austroglanididae, and Bagridae; the latter Bagridae contained two subfamilies, Ritinae and Bagridae is monophyletic, with only *Rita* excluded (Sullivan et al. 2006; Vu et al. 2018). However, the phylogenetic relationships between different genera of Bagridae are not well understood, such as the monophyly or the validity of *Leiocassis, Pseudobagrus, Pelteobagrus*, and *Tachysurus* were classified into *Tachysurus* according to morphological analysis (Cheng et al. 2021; Shao et al. 2021).

Tachysurus brachyrhabdion Cheng, Ishihara & Zhang, 2008 was firstly named Pseudobagrus brachyrhabdion in 2008 (Cheng et al. 2008). It is mainly distributed in the Yuanjiang and Xiangjiang rivers, including the southwest of Hunan Province and the northeast of Guizhou Province in China. Tachysurus gracilis Li, Chen & Chan, 2005 was firstly named Pseudobagrus gracilis in 2005 (Li et al. 2005). It inhabits in the freshwater drainages of southern China. They are essential economic species of freshwater fishes in local areas.

Mitochondria are eukaryotic organelles that play essential roles in oxidative phosphorylation and other biochemical functions. Similar to other vertebrates, fish mitochondrial DNA (mtDNA) is a circular double-stranded molecule, and it is independent of the nuclear genome (Xiao and Zhang 2000). Fish mtDNA is generally small (15–18 kb), containing 13 protein-coding genes (PCGs), 22 transfer RNA (tRNA) genes, two ribosomal RNA (rRNA) genes, and a non-coding region (D-loop) (Brown 2008; Zhang and Wang 2018; Zhang et al. 2021). The mitochondrial genome has been used to study fish species identification, genome evolution, and phylogenetic studies because of the advantages of its small size, multiple copies, maternal inheritance, rapid evolution rate, and lack of introns (Boore 1999; Xiao and Zhang 2000; Zhang and Wang 2018; Zhang et al. 2021).

In this study, the mitochondrial genomes of two catfishes (*T. brachyrhabdion* and *T. gracilis*) were sequenced, assembled, and compared to reveal their evolutionary relationship. These mitochondrial genomes will provide a phylogenetic basis for future phylogenetic and taxonomic studies of Bagridae.

#### Materials and methods

#### Sample collection and DNA extraction

Specimens of *T. brachyrhabdion* and *T. gracilis* were collected from Jiangkou County (27°46'12"N, 108°46'56"E) and Liping County (26°17'51"N, 109°7'25"E), Guizhou,

China, respectively. The samples were preserved in 95% ethanol and stored at -20 °C until DNA extraction. Specimens were recognized as *T. brachyrhabdion* and *T. gracilis* by traditional morphology. The voucher specimens were deposited in the fish specimen room, School of Life Science, Guizhou Normal University under the voucher numbers GZNUSLS201909001~006 and GZNUSLS201907029~030 for *T. brachyrhabdion* and GZNUSLS202005279 for *T. gracilis*. Specimens GZNUSLS201907029 and GZ-NUSLS202005279 were destroyed for the molecular analysis. Total genomic DNA was extracted from muscle tissues using a standard high salt method (Sambrook et al. 1989). The integrity of the genomic DNA was evaluated via 1% agarose gel electrophoresis, and the concentration and purity of DNA were measured using an Epoch 2 Microplate Spectrophotometer (Bio Tek Instruments, Inc, Vermont, USA).

# PCR amplification and sequencing

The whole mitogenomes of *Tachysurus* species were amplified in overlapping PCR fragments using 13 primer pairs designed based on the mitogenome of *T. brevicaudatus* (GenBank accession number: NC\_021393) by Primer Premier 5.0 software (Lalitha 2000) (Suppl. material 3). PCR amplification were performed as described previously (Zhang et al. 2021). The PCR products were fractionated by electrophoresis through 1% agarose gel electrophoresis. The lengths of fragments were determined by comparison with the DL2000 DNA marker (TaKaRa, Japan). The PCR products were sent to Sangon Biotech. Co., Ltd. (Shanghai, China) for sequencing.

#### Sequence analysis and gene annotation

After sequencing, the sequence fragments were edited and assembled using the SeqMan software of DNAStar (DNASTAR Inc., Madison, WI, USA) to obtain the complete mitogenome sequences. Assembled mitogenome sequences were annotated using the MitoAnnotator on the MitoFish homepage (Sato et al. 2018). tRNA genes and their secondary structures were predicted with MITOS (Bernt et al. 2013) and tRNAscan-SE 2.0 (Chan et al. 2021). The base composition, codon usage, and relative synonymous codon usage (RSCU) values were calculated using MEGA 6.0 (Tamura et al. 2013). Strand asymmetry was calculated using the following formulae: AT-skew = (A-T)/(A+T) and GC-skew = (G-C)/(G+C) (Perna and Kocher 1995). The ratio of non-synonymous substitutions (Ka), synonymous substitutions (Ks), and evolutionary rates (Ka/Ks) of each PCG was calculated using DnaSP v. 6.0 (Rozas et al. 2017).

# Phylogenetic analysis

For phylogenetic analysis, sequences of 32 bagrid catfishes were downloaded from Gen-Bank. Additionally, *Cyprinus carpio* (NC\_001606.1), *Silurus asotus* (NC\_015806.1), *Liobagrus andersoni* (NC\_032035.1), and *L. styani* (NC\_034647.1) were used as outgroups. The species used in the analysis are listed in Suppl. material 4. The shared 13 concatenated PCGs were extracted and recombined to construct a matrix using PhyloSuite v.1.1.16 (Zhang et al. 2020). The 13 PCGs were aligned separately using MAFFT v.7.313 (Katoh and Standley 2013) and concatenated into a sequence matrix using PhyloSuite v.1.1.16 (Zhang et al. 2020). The optimal partitioning scheme and nucleotide sequence substitution model of each partition were estimated using PartitionFinder v.2.1.11 (Lanfear et al. 2017) with the Corrected Akaike information criterion (AICc) criteria and greedy algorithm. Bayesian inference (BI) analysis was performed using MrBayes v.3.2.6 (Ronquist et al. 2012) with the models determined by PartitionFinder (Lanfear et al. 2017). Two independent runs of four Markov Chain Monte Carlo (MCMC) chains (one cold chain and three heated chains) were performed for one million generations sampling every 100 generations. The first 25% of the generations was discarded as burnin, and the remaining trees were used to generate a majority rule consensus tree. Maximum likelihood (ML) analysis was carried out using IQ-TREE v.1.6.8 (Nguyen et al. 2015) with 10,000 bootstrap replicates using the ultrafast bootstrapping algorithm (Minh et al. 2013). The phylogenetic trees were visualized and edited using FigTree v1.4.2 (http://tree.bio.ed.ac.uk/software/figtree/).

#### Results

#### Genome structure, organization, and base composition

The entire mitogenome sequences of the two catfishes had lengths of 16,532 bp for *T. brachyrhabdion* and 16,533 bp for *T. gracilis* (GenBank accessions MW712739 and OM759888, respectively) (Fig. 1, Table 1). Both sequences contained 13 PCGs (*ND1-6, COI-III, ND4L, ATP6, ATP8*, and *Cytb*), 22 tRNA genes (one for each amino acid, two for Leucine and Serine), two rRNA genes (12S and 16S rRNA), and a non-coding region (D-loop) (Fig. 1, Table 1). Among these genes, one PCG and eight tRNA genes were encoded on the minority strand (L-strand), while another 28 genes were encoded on the majority strand (H-strand) (Fig. 1, Table 1).

The overall base composition for both species was very similar, 31.02% A, 27.05% T, 15.55% G, and 26.38% C for *T. brachyrhabdion* and 31.03% A, 27.14% T, 15.52% G, and 26.31% C for *T. gracilis* (Table 2). The third codon position of PCGs had the highest A+T content (65.75% for *T. brachyrhabdion* and 66.04% for *T. gracilis*), while the first codon position of PCGs had the lowest A+T content (49.00% for *T. brachyrhabdion* and 49.11% for *T. gracilis*) (Table 2). In addition, skew metrics of the mitogenomes showed positive AT-skew and negative GC-skew (Table 2), indicating that As and Cs were more abundant than Ts and Gs.

#### Protein-coding genes

The mitogenomes of *T. brachyrhabdion* and *T. gracilis* had one PCG (*ND6*) encoded on the L-strand and the remaining PCGs on the H-strand. Both mitogenomes had 11,405 bp coding for PCGs, accounting for 3627/3626 amino acids (Table 2, Suppl.



Figure 1. Circular map of the two catfishes mitochondrial genomes.

material 5). Both species had very similar codon usage, with the most commonly used amino acids being Leu (13.81%, 13.87%), Ser (9.98%, 10.04%), and Ile (8.88%, 8.80%) (Suppl. material 5). The RSCU values of the mitogenomes of the two species are summarized in Fig. 2. All PCGs start with ATG except for *COI*, which used GTG. For these protein-coding genes, the most common stop codon was TAA, although *ND1* used TAG and some used the incomplete stop codon T – or TA-.

To investigate the evolutionary patterns under different selective pressures among 13 PCGs in bagrid catfishes, the value of Ka/Ks was calculated for each PCG, respectively (Fig. 3). The gene (*ATP8*) exhibited the highest ratio of all the PCGs, whereas *COI* had the lowest ratio. However, the Ka/Ks ratios for all PCGs were far lower than one (Fig. 3).

#### Ribosomal, transfer RNA genes and control region

Both the *T. brachyrhabdion* and *T. gracilis* mitogenomes contained two rRNA genes: the large ribosomal RNA subunit (16S rRNA) and small ribosomal RNA subunit (12S rRNA). The 16S rRNA was located between tRNA<sup>Val</sup> and tRNA<sup>Leu (UUR)</sup>, and the 12S rRNA was located between tRNA<sup>Phe</sup> and tRNA<sup>Val</sup>. The 12S rRNA genes and the 16S rRNA genes of both mitogenomes were 953 bp and 1679 bp, respectively.

Twenty-one tRNA genes produced the typical cloverleaf secondary structure, while tRNA<sup>Ser (AGY)</sup> gene lacked the dihydrouracil (DHU) arm (Figs S1 and S2). The sizes of the tRNA genes ranged from 67 bp (tRNA<sup>Cys</sup> and tRNA<sup>Ser (AGY)</sup>) to 75 bp (tRNA<sup>Leu</sup> (<sup>UUR)</sup>) in both *T. brachyrhabdion* and *T. gracilis* (Table 1).

The putative control regions were located between *tRNA*<sup>*Pro*</sup> and *tRNA*<sup>*Phe*</sup> in the two bagrid catfishes. The control regions of *T. brachyrhabdion* and *T. gracilis* were 892 bp in size. The average A+T content of the CRs (62.33%–62.67%) was higher than that of the whole genomes, PCGs, rRNAs, or tRNAs (57.14%–58.24%) (Table2).

**Table 1.** Organization of mitochondrial genome of *T. brachyrhabdion* (TB) and *T. gracilis* (TG). H refers to the majority strand and L refers to the minority strand. Position numbers refer to positions on the majority strand.

Gene	Strand	Nucleotid	le number	Length	Intergenic	nucleotide	Anticodon	Start/Stop codons	
		ТВ	TG	(bp) <u>TB</u> TC		TG	_	Start	Stop
tRNA <sup>Phe</sup>	Н	1-70	1-70	70	0	0	GAA		
12S rRNA	Н	71-1023	71-1023	953	0	0			
$tRNA^{Val}$	Н	1024-1095	1024-1095	72	0	0	TAC		
16S rRNA	Н	1096-2774	1096-2774	1679	0	0			
tRNA <sup>Leu (UUR)</sup>	Н	2775-2849	2775-2849	75	0	0	TAA		
ND1	Н	2850-3824	2850-3824	975	2	2		ATG	TAG
tRNA <sup>11e</sup>	Н	3827-3898	3827-3898	72	-1	-1	GAT		
$tRNA^{Gln}$	L	3898-3968	3898-3968	71	-1	-1	TTG		
$tRNA^{Met}$	Н	3968-4037	3968-4037	70	0	0	CAT		
ND2	Н	4038-5082	4038-5082	1045	0	0		ATG	Т
$tRNA^{Trp}$	Н	5083-5153	5083-5153	71	2	2	TCA		
$tRNA^{Ala}$	L	5156-5224	5156-5224	69	1	1	TGC		
$tRNA^{Asm}$	L	5226-5298	5226-5298	73	32	32	GTT		
$tRNA^{C_{js}}$	L	5331-5397	5331-5397	67	0	0	GCA		
$tRNA^{Tyr}$	L	5398-5468	5398-5469	71/72	1	1	GTA		
COI	Н	5470-7020	5471-7021	1551	0	0		GTG	TAA
tRNA <sup>Ser (UCN)</sup>	L	7021-7091	7022-7092	71	4	4	TGA		
$tRNA^{Asp}$	Н	7096-7168	7097-7169	73	14	14	GTC		
COII	Н	7183-7873	7184-7874	691	0	0		ATG	Т
$tRNA^{Lys}$	Н	7874-7947	7875-7948	74	1	1	TTT		
ATPase 8	Н	7949-8116	7950-8117	168	-10	-10		ATG	TAA
ATPase 6	Н	8107-8789	8108-8790	683	0	0		ATG	TA
COIII	Н	8790-9573	8791-9574	784	0	0		ATG	Т
$tRNA^{Gh}$	Н	9574-9647	9575-9648	74	0	0	TCC		
ND3	Н	9648-9996	9649-9997	349	0	0		ATG	Т
$tRNA^{Arg}$	Н	9997-10067	9998-10068	71	0	0	TCG		
ND4L	Н	10068-10364	10069-10365	297	-7	-7		ATG	TAA
ND4	Н	10358-11738	10359-11739	1381	0	0		ATG	Т
$tRNA^{His}$	Н	11739-11808	11740-11809	70	0	0	GTG		
tRNA <sup>Ser (AGY)</sup>	Н	11809-11875	11810-11876	67	3	3	GCT		
tRNA <sup>Leu (CUN)</sup>	Н	11879-11951	11880-11952	73	0	0	TAG		
ND5	Н	11952-13778	11953-13779	1827	-4	-4		ATG	TAA
ND6	L	13775-14290	13776-14291	516	0	0		ATG	TAA
$tRNA^{Glu}$	L	14291-14359	14292-14360	69	2	2	TTC		
Cytb	Н	14362-15499	14363-15500	1138	0	0		ATG	Т
$tRNA^{Thr}$	Н	15500-15572	15501-15573	73	-2	-2	TGT		
$tRNA^{Pro}$	L	15571-15640	15572-15641	70	0	0	TGG		
D-loop	Н	15641-16532	15642–16533 16533	892	0	0			

#### Phylogenetic analysis

To determine the phylogenetic relationship between *T. brachyrhabdion* and *T. gracilis* in the Characidae, we selected the concatenated nucleotide sequences of the combined mitochondrial gene set (13 PCGs) from 34 bagrid catfishes. As shown in Fig. 4, the phylogenetic analysis of the two tree models (BI and ML) using the combined mitochondrial gene set well supported the tree topologies and yielded identical results. Phylogenetic analysis reveals bagrid catfishes could be separated into two clades excluding *Rita rita* (Hamilton, 1822) with strong support (Fig. 4). Clade I included *Hemibagrus, Sperata*, and *Mystus* (BS = 100%, PP = 100%). Clade II was only composed of *Tachysurus* (BS = 100%, PP = 100%). Several monophyletic clades of *Tachysurus*,



**Figure 2.** Relative synonymous codon usage (RSCU) in the mitogenomes of the *T. brachyrhabdion* **A** and *T. gracilis* **B**.

**Table 2.** Nucleotide composition of the mitochondrial genomes of *T. brachyrhabdion* (TB) and *T. gracilis* (TG).

	Length(bp)		A%		Τ%		G%		C%		A+T%		AT-skew		GC-skew	
	TB	TG	TB	TG	TB	TG	TB	TG	TB	TG	TB	TG	TB	TG	TB	TG
genome	16532	16533	31.02	31.03	27.05	27.14	15.55	15.52	26.38	26.31	58.07	58.17	0.0683	0.0669	-0.2585	-0.2580
PCGs	11405	11405	28.94	28.99	29.13	29.25	15.37	15.30	26.56	26.46	58.07	58.24	-0.0032	-0.0045	-0.2668	-0.2673
1 <sup>ss</sup> codon position	3806	3806	27.06	27.17	21.94	21.94	25.49	25.35	25.51	25.54	49.00	49.11	0.1046	0.1065	-0.0005	-0.0036
2 <sup>nd</sup> codon position	3806	3806	18.61	18.63	40.87	40.95	13.55	13.53	26.97	26.89	59.47	59.58	-0.3743	-0.3746	-0.3312	-0.3307
3 <sup>rd</sup> codon position	3806	3806	41.17	41.17	24.59	24.87	7.05	7.00	27.19	26.95	65.75	66.04	0.2522	0.2467	-0.5880	-0.5876
rRNA	2632	2632	34.77	34.65	22.38	22.34	19.64	19.72	23.21	23.29	57.14	56.99	0.2168	0.2160	-0.0833	-0.0830
tRNA	1566	1567	29.12	29.16	28.10	28.33	22.48	22.34	20.30	20.17	57.22	57.50	0.0179	0.0144	0.0507	0.0511
D-loop region	892	892	31.28	31.17	31.39	31.17	13.79	13.90	23.54	23.77	62.67	62.33	-0.0018	0.0000	-0.2613	-0.2619



Figure 3. Ka/Ks of each PCG from 33 bagrid catfishes mitogenomes (Rita rita was excluded).



**Figure 4.** Phylogenetic tree obtained from BI and ML analysis based on the 13 PCGs dataset. The numbers at the nodes separated by "/" indicate the posterior probability (BI) and bootstrap value (ML).

*Sperata, Mystus,* and *Liobagrus* were strongly supported (Fig. 4). Nevertheless, the genus *Hemibagrus* was paraphyletic (Fig. 4). In addition, our phylogenetic trees showed that *T. brachyrhabdion* and *T. gracilis* clustered together forming a group.

#### Discussion

Over the past two to three decades, mitochondrial genes and genomes have been frequently used in fish studies (Xiao and Zhang 2000; Brown 2008; Zhang and Wang 2018). In this study, the complete mitochondrial genomes of *T. brachyrhabdion* and *T. gracilis* were sequenced. The order and arrangement of the two mitogenomes were identical to that of other bagrid catfishes (Liang et al. 2014; Tian et al. 2016; Liu et al. 2019). The genomes had a total length of 16,532 bp and 16,533 bp, with A + T contents at 58.07% and 58.17%, respectively. These A+T biases were within the known range (56.37–59.80%) reported for mitochondrial genomes in closely related fishes (Tian et al. 2016; Liu et al. 2019).

Most of the PCGs of these two species had ATG as the start codon except *COI* that had GTG as the start codon. The *COI* gene usually uses GTG as the start codon in other fishes, such as *Lateolabrax, Sinocyclocheilus multipunctatus*, and *Microphysogobio elongatus* (Shan et al. 2016; Zhang and Wang 2018; Zhang et al. 2021). The PCGs of these two bagrid catfishes had an incomplete stop codon that was automatically filled by post-transcriptional polyadenylation (Ojala et al. 1981). This was a common character in metazoans mitogenomes (Shan et al. 2016; Jiang et al. 2019; Li et al. 2021). Twenty-one tRNA genes showed the typical cloverleaf secondary structure (Suppl. material 1, 2), while the tRNA-Ser<sup>AGY</sup> gene lacked the dihydrouracil (DHU) arm as noted in other fish species (Zhang and Wang 2018; Zhang et al. 2021). The low Ka/Ks value for each PCG indicated that they were all under strong purifying selection. The DNA barcoding gene *COI* had the lowest evolutionary rate, consistent with the results observed from other fish groups (Sun et al. 2021; Yu et al. 2021).

Phylogenetic analysis revealed Bagridae was supported as a monophyletic group with *Rita rita* excluded (Fig. 4), which is consistent with previous phylogenetic studies (Sullivan et al. 2006; Vu et al. 2018). The phylogenetic tree suggested the classification of *R. rita* should be further revised and perfected. Clade I was composed of two monophyletic groups (*Sperata* and *Mystus*) and one paraphyletic group (*Hemibagrus*) (Fig. 4), which is consistent with the previous study of Liu et al. (2019). The traditional genera *Leiocassis*, *Pseudobagrus*, *Pelteobagrus*, and *Tachysurus* were not each monophyletic (Hardman 2005; Ku et al. 2007; Liu et al. 2019), but instead were grouped into *Tachysurus* as found in recent research (Cheng et al. 2021; Shao et al. 2021). Our results supported that the species in *Leiocassis, Pseudobagrus*, *Pelteobagrus*, *nelteobagrus*, and *Tachysurus*. Thus, the current concept of *Tachysurus* us includes the traditional genera *Leiocassis*, *Pseudobagrus*, and *Tachysurus*. Due to limited sample size and molecular data, the phylogeny discussed in this research should be regarded as preliminary. The growing number of available mitogenomes will improve our understanding of the phylogeny and classification of bagrid catfish.

#### Conclusions

This study reported the complete mitochondrial genome sequences of two bagrid catfishes. The study showed that mitogenomes of Bagridae were conserved in structure, gene order, and nucleotide composition. Phylogenetic analysis confirmed that Bagridae is monophyletic group with *Rita rita* excluded and the traditional classification that *Leiocassis*, *Pseudobagrus*, *Pelteobagrus*, and *Tachysurus* belong to one genus.

#### Acknowledgements

This work was supported by the Joint Fund of the National Natural Science Foundation of China and the Karst Science Research Center of Guizhou Province (Grant No. U1812401), Guizhou Provincial Science and Technology Foundation (Qiankehejichu[2018]1113), Natural Science Foundation of Guizhou Educational Committee (QianjiaoheKY[2021]306) and the Undergraduate Research Training Program of Guizhou Normal University (DK2019A023).

#### References

- Bernt M, Donath A, Juhling F, Externbrink F, Florentz C, Fritzsch G, Putz J, Middendorf M, Stadler PF (2013) MITOS: Improved de novo metazoan mitochondrial genome annotation. Molecular Phylogenetics and Evolution 69(2): 313–319. https://doi.org/10.1016/j. ympev.2012.08.023
- Boore JL (1999) Animal mitochondrial genomes. Nucleic Acids Research 27(8): 1767–1780. https://doi.org/10.1093/nar/27.8.1767
- Brown KH (2008) Fish mitochondrial genomics: Sequence, inheritance and functional variation. Journal of Fish Biology 72(2): 355–374. https://doi.org/10.1111/j.1095-8649.2007.01690.x
- Chan PP, Lin BY, Mak AJ, Lowe TM (2021) tRNAscan-SE 2.0: Improved detection and functional classification of transfer RNA genes. Nucleic Acids Research 49(16): 9077–9096. https://doi.org/10.1093/nar/gkab688
- Cheng JL, Ishihara H, Zhang E (2008) *Pseudobagrus brachyrhabdion*, a new catfish (Teleostei: Bagridae) from the middle Yangtze River drainage, South China. Ichthyological Research 55(2): 112–123. https://doi.org/10.1007/s10228-007-0020-3
- Cheng JL, Shao WH, Lopez JA, Zhang E (2021) *Tachysurus lani*, a new catfish species (Teleostei: Bagridae) from the Pearl River basin, South China. Ichthyological Exploration of Freshwaters 30(4): 299–315. https://doi.org/10.23788/ief-1156
- Fricke R, Eschmeyer WN, Van der Laan R (2022) Eschmeyer's catalog of fishes: genera, species, references. http://researcharchive.calacademy.org/research/ichthyology/catalog/fishcatmain.asp [Electronic version accessed 8 Mar. 2022]
- Hardman M (2005) The phylogenetic relationships among non-diplomystid catfishes as inferred from mitochondrial cytochrome b sequences; the search for the ictalurid sister taxon (Otophysi: Siluriformes). Molecular Phylogenetics and Evolution 37(3): 700–720. https:// doi.org/10.1016/j.ympev.2005.04.029
- Jayaram KC (1968) Contributions to the study of bagrid fishes (Siluroidea: Bagridae). 3. A systematic account of the Japanese, Chinese, Malayan and Indonesian genera. Treubia 27(2–3): 287–386. https://doi.org/10.14203/treubia.v27i2-3.1560
- Jiang L, Peng L, Tang M, You Z, Zhang M, West A, Ruan Q, Chen W, Merila J (2019) Complete mitochondrial genome sequence of the Himalayan Griffon, *Gyps himalayensis* (Accipitriformes: Accipitridae): Sequence, structure, and phylogenetic analyses. Ecology and Evolution 91(15): 8813–8828. https://doi.org/10.1002/ece3.5433

- Katoh K, Standley DM (2013) MAFFT multiple sequence alignment software version 7: Improvements in performance and usability. Molecular Biology and Evolution 30(4): 772–780. https://doi.org/10.1093/molbev/mst010
- Ku X, Peng Z, Diogo R, He S (2007) MtDNA phylogeny provides evidence of generic polyphyleticism for East Asian bagrid catfishes. Hydrobiologia 579(1): 147–159. https://doi. org/10.1007/s10750-006-0401-z
- Lalitha S (2000) Primer Premier 5. Biotech Software & Internet Report 1(6): 270–272. https:// doi.org/10.1089/152791600459894
- Lanfear R, Frandsen PB, Wright AM, Senfeld T, Calcott B (2017) PartitionFinder 2: New methods for selecting partitioned models of evolution for molecular and morphological phylogenetic analyses. Molecular Biology and Evolution 34(3): 772–773. https://doi. org/10.1093/molbev/msw260
- Li J, Chen XL, Chan B (2005) A new species of *Pseudobagrus* (Teleostei: Siluriformes: Bagridae) from southern China. Zootaxa 1067(1): 49–57. https://doi.org/10.11646/zootaxa.1067.1.3
- Li R, Ying X, Deng W, Rong W, Li X (2021) Mitochondrial genomes of eight Scelimeninae species (Orthoptera) and their phylogenetic implications within Tetrigoidea. PeerJ 9: e10523. https://doi.org/10.7717/peerj.10523
- Liang HW, Meng Y, Li Z, Zhang Y, Zou GW (2014) Complete mitochondrial DNA genome of *Pseudobagrus brevicaudatus* (Siluriformes: Bagridae). Mitochondrial DNA 25(3): 169– 170. https://doi.org/10.3109/19401736.2013.792060
- Liu Y, Wu PD, Zhang DZ, Zhang HB, Tang BP, Liu QN, Dai LS (2019) Mitochondrial genome of the yellow catfish *Pelteobagrus fulvidraco* and insights into Bagridae phylogenetics. Genomics 111(6): 1258–1265. https://doi.org/10.1016/j.ygeno.2018.08.005
- Minh BQ, Nguyen MA, von Haeseler A (2013) Ultrafast approximation for phylogenetic bootstrap. Molecular Biology and Evolution 30(5): 1188–1195. https://doi.org/10.1093/ molbev/mst024
- Mo TE (1992) Anatomy, relationships and systematics of the Bagridae (Teleostei: Siluroidei) with a hypothesis of siluroid phylogeny. Copeia 1992(4): 1132–1134. https://doi. org/10.2307/1446657
- Nguyen LT, Schmidt HA, von Haeseler A, Minh BQ (2015) IQ-TREE: A fast and effective stochastic algorithm for estimating maximum-likelihood phylogenies. Molecular Biology and Evolution 32(1): 268–274. https://doi.org/10.1093/molbev/msu300
- Ojala D, Montoya J, Attardi G (1981) tRNA punctuation model of RNA processing in human mitochondria. Nature 290(5806): 470–474. https://doi.org/10.1038/290470a0
- Perna NT, Kocher TD (1995) Patterns of nucleotide composition at fourfold degenerate sites of animal mitochondrial genomes. Journal of Molecular Evolution 41(3): 353–358. https:// doi.org/10.1007/BF01215182
- Regan CT (1911) LXV. The classification of the Teleostean fishes of the order Ostariophysi. 2. Siluroidea. Annals & Magazine of Natural History 8(47): 553–577. https://doi. org/10.1080/00222931108693067
- Ronquist F, Teslenko M, van der Mark P, Ayres DL, Darling A, Hohna S, Larget B, Liu L, Suchard MA, Huelsenbeck JP (2012) MrBayes 3.2: Efficient Bayesian phylogenetic inference and model choice across a large model space. Systematic Biology 61(3): 539–542. https://doi.org/10.1093/sysbio/sys029

- Rozas J, Ferrer-Mata A, Sanchez-DelBarrio JC, Guirao-Rico S, Librado P, Ramos-Onsins SE, Sanchez-Gracia A (2017) DnaSP 6: DNA sequence polymorphism analysis of large data sets. Molecular Biology and Evolution 34(12): 3299–3302. https://doi.org/10.1093/molbev/msx248
- Sambrook JE, Maniatis TE, Fritsch EF (1989) Molecular cloning: a laboratory manual. Cold spring harbor laboratory press.
- Sato Y, Miya M, Fukunaga T, Sado T, Iwasaki W (2018) MitoFish and MiFish pipeline: A mitochondrial genome database of fish with an analysis pipeline for environmental DNA metabarcoding. Molecular Biology and Evolution 35(6): 1553–1555. https://doi.org/10.1093/ molbev/msy074
- Shan B, Song N, Han Z, Wang J, Gao T, Yokogawa K (2016) Complete mitochondrial genomes of three sea basses Lateolabrax (Perciformes, Lateolabracidae) species: Genome description and phylogenetic considerations. Biochemical Systematics and Ecology 67: 44–52. https://doi.org/10.1016/j.bse.2016.04.007
- Shao WH, Cheng JL, Zhang E (2021) Eight in one: Hidden diversity of the bagrid catfish *Tachysurus albomarginatus* s.l. (Rendhal, 1928) widespread in lowlands of south China. Frontiers in Genetics 12: 713793. https://doi.org/10.3389/fgene.2021.713793
- Sullivan JP, Lundberg JG, Hardman M (2006) A phylogenetic analysis of the major groups of catfishes (Teleostei: Siluriformes) using rag1 and rag2 nuclear gene sequences. Molecular Phylogenetics and Evolution 41(3): 636–662. https://doi.org/10.1016/j.ympev.2006.05.044
- Sun CH, Liu HY, Xu N, Zhang XL, Zhang Q, Han BP (2021) Mitochondrial genome structures and phylogenetic analyses of two tropical characidae fishes. Frontiers in Genetics 12: e627402. https://doi.org/10.3389/fgene.2021.627402
- Tamura K, Stecher G, Peterson D, Filipski A, Kumar S (2013) MEGA6: Molecular evolutionary genetics analysis version 6.0. Molecular Biology and Evolution 30(12): 2725–2729. https://doi.org/10.1093/molbev/mst197
- Tian H, Que Y, Zhao N, Chen F, Zhu B, Huang D, Chang J, Liao X (2016) The complete mitochondrial genome of the spotted longbarbel catfish, *Hemibagrus guttatus* (Siluriformes, Bagridae). Mitochondrial DNA A: DNA Mapping. Sequencing, and Analysis 27(1): 467– 468. https://doi.org/10.3109/19401736.2014.900670
- Vu Q, Thi OT, Lan PTT, Linh TT, Thuy BD (2018) Molecular phylogeny of catfishes (Teleostei: Siluriformes) inferred from mitochondrial markers-implications for lower Mekong River basin. European Journal of Advanced Research in Biological and Life Sciences 6(3): 1–12.
- Xiao WH, Zhang YP (2000) Genetics and evolution of mitochondrial DNA in fish. Shui Sheng Sheng Wu Hsueh Bao 24(4): 384–391. https://doi.org/1000-3207(2000)04-0384-08
- Yu P, Zhou L, Yang WT, Miao LJ, Li Z, Zhang XJ, Wang Y, Gui JF (2021) Comparative mitogenome analyses uncover mitogenome features and phylogenetic implications of the subfamily Cobitinae. BMC Genomics 22(1): 50. https://doi.org/10.1186/s12864-020-07360-w
- Zhang RY, Wang X (2018) Characterization and phylogenetic analysis of the complete mitogenome of a rare cavefish, *Sinocyclocheilus multipunctatus* (Cypriniformes: Cyprinidae). Genes & Genomics 40(10): 1033–1040. https://doi.org/10.1007/s13258-018-0711-3
- Zhang D, Gao F, Jakovlic I, Zou H, Zhang J, Li WX, Wang GT (2020) PhyloSuite: An integrated and scalable desktop platform for streamlined molecular sequence data management

and evolutionary phylogenetics studies. Molecular Ecology Resources 20(1): 348-355. https://doi.org/10.1111/1755-0998.13096

Zhang RY, Tang Q, Deng L (2021) The complete mitochondrial genome of *Microphysogobio* elongatus (Teleostei, Cyprinidae) and its phylogenetic implications. ZooKeys 1061: 57–73. https://doi.org/10.3897/zookeys.1061.70176

# Supplementary material I

#### Figure S1. Predicted tRNA structures of Tachysurus brachyrhabdion

Authors: Renyi Zhang, Lei Deng1, Xiaomei Lv, Qian Tang Data type: image

Copyright notice: This dataset is made available under the Open Database License (http://opendatacommons.org/licenses/odbl/1.0/). The Open Database License (ODbL) is a license agreement intended to allow users to freely share, modify, and use this Dataset while maintaining this same freedom for others, provided that the original source and author(s) are credited.

Link: https://doi.org/10.3897/zookeys.1115.85249.suppl1

# Supplementary material 2

# Figure S2. Predicted tRNA structures of Tachysurus gracilis

Authors: Renyi Zhang, Lei Deng1, Xiaomei Lv, Qian Tang Data type: image

Copyright notice: This dataset is made available under the Open Database License (http://opendatacommons.org/licenses/odbl/1.0/). The Open Database License (ODbL) is a license agreement intended to allow users to freely share, modify, and use this Dataset while maintaining this same freedom for others, provided that the original source and author(s) are credited.

Link: https://doi.org/10.3897/zookeys.1115.85249.suppl2

# Supplementary material 3

# Table S1. Primers used for PCR

Authors: Renyi Zhang, Lei Deng1, Xiaomei Lv, Qian Tang Data type:molecular data

Copyright notice: This dataset is made available under the Open Database License (http://opendatacommons.org/licenses/odbl/1.0/). The Open Database License (ODbL) is a license agreement intended to allow users to freely share, modify, and use this Dataset while maintaining this same freedom for others, provided that the original source and author(s) are credited.

Link: https://doi.org/10.3897/zookeys.1115.85249.suppl3

### Supplementary material 4

#### Tables S2. Species, GenBank accession number and length of mitogenomes used in this study

Authors: Renyi Zhang, Lei Deng1, Xiaomei Lv, Qian Tang

Data type: molecular data

Copyright notice: This dataset is made available under the Open Database License (http://opendatacommons.org/licenses/odbl/1.0/). The Open Database License (ODbL) is a license agreement intended to allow users to freely share, modify, and use this Dataset while maintaining this same freedom for others, provided that the original source and author(s) are credited.

Link: https://doi.org/10.3897/zookeys.1115.85249.suppl4

#### Supplementary material 5

# Table S3. Number of codons in *T. brachyrhabdion* (TB) and *T. gracilis* (TG) for mitochondrial PCGs

Authors: Renyi Zhang, Lei Deng1, Xiaomei Lv, Qian Tang

Data type: molecular data

Copyright notice: This dataset is made available under the Open Database License (http://opendatacommons.org/licenses/odbl/1.0/). The Open Database License (ODbL) is a license agreement intended to allow users to freely share, modify, and use this Dataset while maintaining this same freedom for others, provided that the original source and author(s) are credited.

Link: https://doi.org/10.3897/zookeys.1115.85249.suppl5