

A new species of *Sinopoda* from China, with first description of the male of *S. wuyiensis* Liu, 2021 (Araneae, Sparassidae)

Yang Zhong^{1,2,3,4}, Meng-Yun Zeng¹, Chao-Lan Gu¹,
Hui-Liang Yu^{3,4}, Jing-Yuan Yang^{3,4}

1 School of Nuclear Technology and Chemistry & Biology, Hubei University of Science and Technology, Xianning 437100, Hubei, China **2** Hubei Key Laboratory of Radiation Chemistry and Functional Materials, Hubei University of Science and Technology, Xianning 437100, Hubei, China **3** Shennongjia National Park Administration, Shennongjia 442421, Hubei, China **4** Hubei Province Key Laboratory of Conservation Biology for Shennongjia Golden Monkey, Shennongjia 442421, Hubei, China

Corresponding authors: Yang Zhong (hubeispider@aliyun.com), Jing-Yuan Yang (snjbhq@163.com)

Academic editor: Gergin Blagoev | Received 13 April 2022 | Accepted 18 July 2022 | Published 9 August 2022

<https://zoobank.org/83FFFD94-67F9-47CE-A7E8-5D1B05F3FEF9>

Citation: Zhong Y, Zeng M-Y, Gu C-L, Yu H-L, Yang J-Y (2022) A new species of *Sinopoda* from China, with first description of the male of *S. wuyiensis* Liu, 2021 (Araneae, Sparassidae). ZooKeys 1116: 121–132. <https://doi.org/10.3897/zookeys.1116.85303>

Abstract

One new species of the genus *Sinopoda* Jäger, 1999, *S. muyuensis* **sp. nov.** (♂, ♀), is described and figured from the Shennongjia Forestry District, Hubei Province, China. In addition, the male of *Sinopoda wuyiensis* Liu, 2021 is described for the first time from the Wuyishan National Nature Reserve, Fujian Province, China.

Keywords

Biodiversity, Fujian, Hubei, huntsman spiders, taxonomy

Introduction

The genus *Sinopoda* was established by Jäger (1999). As currently recognised, it comprises 137 species, of which 71 species were recorded from China, representing 51.8% of the global species [World Spider Catalog (WSC) 2022]. The genus has been reported from East Asia (85 species in China, Japan, and Korea), Southeast Asia (50 species in Brunei, Indonesia, Laos, Malaysia, Myanmar, Thailand, and Vietnam) and South Asia (a single species in India) (Zhang et al. 2021).

We have found two *Sinopoda* species present in collections from China in the past year. The discovery of one of these species, *Sinopoda muyuensis* sp. nov. from Hubei represents the first record of this species in Shennongjia Forestry District. Furthermore, specimens of *Sinopoda wuyiensis* Liu, 2021 from Fujian have allowed us to provide the first description of the male of the species.

Materials and methods

Specimens were examined and measured with an Olympus SZX7 stereomicroscope. Positions of tegular appendages are given according to clock positions, based on the left palp in ventral view. Male and female copulatory organs were examined and illustrated after dissection from the spider bodies; vulvae were cleared in a warm 10% potassium hydroxide (KOH) solution. All photographs were captured with a KUY NICE industrial digital camera (20.0 megapixels) mounted on an Olympus CX43 dissecting microscope, and assembled using Helicon Focus 3.10.3 image stacking software. Photographic images were then edited using Adobe Photoshop CC 2018. All measurements were obtained using an Olympus SZX7 stereomicroscope and are given in millimetres (mm).

Leg measurements are shown as: total length (femur, patella, tibia, metatarsus, tarsus). Number of macrosetae is listed for each segment in the following order: prolateral, dorsal, retrolateral, ventral (in femora and patellae ventral spines are absent and fourth digit is omitted in the setation formula). Abbreviations used in the text and figures are given below: **AB** = anterior bands, **ALE** = anterior lateral eye, **AME** = anterior median eye, **AW** = anterior width of carapace, **C** = conductor, **CH** = clypeus height, **dRTA** = dorsal branch of RTA, **E** = embolus, **EA** = embolic apophysis, **FD** = fertilization duct, **FE** = femur, **GA** = glandular appendage, **LL** = lateral lobes, **LS** = lobal septum, **MS** = membranous sac, **Mt** = metatarsus, **OL** = opisthosoma length, **OW** = opisthosoma width, **Pa** = patella, **PI** = posterior incision of LL, **PL** = carapace length, **PLE** = posterior lateral eyes, **PME** = posterior median eyes, **Pp** = palp or palpus, **PP** = posterior part of spermathecae, **PW** = carapace width, **RTA** = retrolateral tibial apophysis, **SP** = spermophor, **SS** = slit sensillum, **ST** = subtegulum, **T** = tegulum, **Ta** = tarsus, **Ti** = tibia. I, II, III, IV—legs I to IV, **TP** = tegular protrusion, **vRTA** = ventral branch of RTA, **HUST** = School of Nuclear Technology and Chemistry and Biology, Hubei University of Science and Technology, Xianning, Hubei, China.

Taxonomy

Family Sparassidae Bertkau, 1872

Subfamily Heteropodinae Thorell, 1873

Genus *Sinopoda* Jäger, 1999

Type species. *Sarotes forcipatus* Karsch, 1881

Diagnosis. See Jäger (1999), Liu et al. (2008), Zhang et al. (2015), and Grall and Jäger (2020).

Composition. Within the genus *Sinopoda*, five groups were established by Jäger and Ono (2000), Grall and Jäger (2020) and Zhang et al. (2021): *S. anguina*-group (*S. anguina* Liu, Li & Jäger, 2008, *S. bifurca* Grall & Jäger, 2020, *S. bispina* Grall & Jäger, 2020, *S. fornicata* Liu, Li & Jäger, 2008, *S. improcera* Zhong et al., 2019, *S. lata* Zhong et al., 2019, *S. longicymbialis* Grall & Jäger, 2020, *S. mamillata* Zhong, Cao & Liu, 2017, *S. nanphagu* Grall & Jäger, 2020, *S. phiset* Grall & Jäger, 2020, *S. rotunda* Grall & Jäger, 2020 and *S. tuber* Grall & Jäger, 2020); *S. chiangmaiensis*-group (*S. chiangmaiensis* Grall & Jäger, 2020, *S. lot* Grall & Jäger, 2020 and *S. phathai* Grall & Jäger, 2020); *S. globosa*-group (*S. globosa* Zhang, Zhang & Zhang, 2015, *S. longiducta* Zhang, Zhang & Zhang, 2015, *S. mi* Chen & Zhu, 2009, *S. ovata* Zhong et al., 2019, *S. triangula* Liu, Li & Jäger, 2008 and *S. yaanensis* Zhong et al., 2019); *S. tumefacta*-group (*S. crassa* Liu, Li & Jäger, 2008, *S. debiscens* Zhong et al., 2019, *S. erromena* Zhong et al., 2019, *S. tumefacta* Zhong et al., 2019, *S. yanlingensis* Zhong et al., 2019 and *S. yaojingensis* Liu, Li & Jäger, 2008); *S. okinawana*-group [*S. forcipata* (Karsch, 1881), *S. cochlearia* Zhang, Zhang & Zhang, 2015, *S. derivata* Jäger & Ono, 2002, *S. fasciculata* Jäger Gao & Fei, 2002, *S. guangyuanensis* Zhong et al., 2018, *S. hamata* (Fox, 1937), *S. koreana* (Paik, 1968), *S. okinawana* Jäger & Ono, 2000, *S. tanikawai* Jäger & Ono, 2000, *S. wangi* Song & Zhu, 1999]. Ninety-six other species have not yet been grouped.

***Sinopoda muyuensis* sp. nov.**

<https://zoobank.org/4BCEBD12-8C2A-44EA-9180-DDE1FB4ACD39>

Figs 1, 2, 4A, B, E–H, 5A, B, 6

Type material. *Holotype*: ♂ (HUST 0003), **CHINA: Hubei**: Shennongjia Forestry District, Muyu Town, Guanmenshan Scenic Area; 31.45°N, 110.40°E; alt. 1200 m; 10.XII.2021; Y. Zhong leg. *Paratypes* (HUST): 2♂, 3♀, same data as holotype.

Etymology. ‘Muyu’ refers to the type locality of this species, Muyu Town.

Diagnosis. The male of *Sinopoda muyuensis* sp. nov. resembles *S. angulata* Jäger, Gao & Fei, 2002 (Zhu et al. 2020: figs 1A–C, 2A–C) and *S. yichangensis* Zhu, Zhong & Yang, 2020 (Zhu et al. 2020: figs 4A–C, 5A–C; Gong and Zhong 2021: figs 2A–C) in having the embolus distally filiform, as long as the embolic apophysis, and RTA arising subdistally from tibia, but the new species can be separated from *S. angulata* by the posterior margins of the embolic apophysis being distinctly humped (smooth in




Figure 1. *Sinopoda muyuensis* sp. nov., holotype male **A–C** palp, left (**A** prolateral view **B** ventral view **C** retrolateral view). Abbreviations: C – conductor, dRTA – dorsal branch of RTA, E – embolus, EA – embolic apophysis, SP – spermophore, T – tegulum, vRTA – ventral branch of RTA. Red arrow: embolic apophysis projection. Scale bar: 0.5 mm.

S. angulata); from *S. yichangensis* by the tip of embolic apophysis with a pointed end (blunt in *S. yichangensis*). Females are similar to those of *S. angulata* (Zhu et al. 2020: fig. 2D, E) in having the epigyne with a lobal septum $\sim 1/4$ of the epigynal width, posterior part of the spermathecae considerably larger than the glandular projection, and the angle of the diverging internal duct system $\sim 80^\circ$, but distinguished by the vulva with its internal duct system not touching (touching along the median line in *S. angulata*); glandular appendages extending into the median half of the internal duct system (anterior half in *S. angulata*); ends of internal duct system nearly straight in dorsal view (bent at 180° in *S. angulata*) (Figs 1, 2).




Figure 2. *Sinopoda muyuensis* sp. nov., paratype female **A** epigyne **B** vulva (**A** ventral view **B** dorsal view). Abbreviations: AB – anterior bands, FD – fertilisation duct, GA – glandular appendage, LL – lateral lobes, LS – lobal septum, MS – membranous sac, PP – posterior part of spermathecae, SS – slit sensillum. Scale bar: 0.5 mm.

Description. Male. PL 5.3, PW 4.7, AW 2.5, OL 5.8, OW 3.9. Eyes: AME 0.19, ALE 0.31, PME 0.22, PLE 0.32, AME–AME 0.20, AME–ALE 0.16, PME–PME 0.28, PME–PLE 0.36, AME–PME 0.38, ALE–PLE 0.25, CH AME 0.19, CH ALE 0.26. Setation: Palp: 131, 101, 1100; Fe: I–III 323, IV 321; Pa: I–IV 101; Ti: I–IV 2326; Mt: I–II 1014, III 2026, IV 3036. Measurements of palp and legs: Palp 6.8 (2.4, 1.3, 1.3, –, 1.8), I 17.6 (4.4, 1.4, 5.5, 4.9, 1.4), II 19.1 (5.6, 1.7, 6.5, 3.9, 1.4), III 15.7 (4.9, 1.5, 4.8, 3.3, 1.2), IV 15.9 (5.3, 1.5, 4.8, 3.3, 1.0). Leg formula: II-I-IV-III. Cheliceral furrow with three anterior and four posterior teeth, and with ~ 36 denticles (Fig. 4A). Carapace yellowish brown dorsally, with yellow transversal stripe posteriorly, with shallow fovea and radial furrows. Chelicerae deep reddish brown. Sternum yellow with brown margin. Endites and labium deep yellowish brown, with margin deep brown. Legs yellowish brown, covered by short spines. Opisthosoma yellowish brown dorsally, with three pairs of dark patches laterally. Opisthosoma uniformly yellowish brown with some brown patches ventrally (Fig. 4E, F).

Palp as in Fig. 1. Cymbium distinctly longer than tibia. Embolus S-shaped, arising from tegulum at nearly the 6-o’ clock-position in ventral view. Conductor arising at 12- to 1-o’ clock-position from tegulum. RTA arising mesially to distally from tibia, with distinct brush of stiff setae. dRTA slender, finger-shaped, vRTA roughly rectangular in retrolateral view.

Female. PL 5.7, PW 4.8, AW 3.1, OL 5.9, OW 3.9. Eyes: AME 0.16, ALE 0.25, PME 0.18, PLE 0.25, AME–AME 0.27, AME–ALE 0.09, PME–PME 0.33, PME–PLE 0.35, AME–PME 0.32, ALE–PLE 0.28, CH AME 0.20, CH ALE 0.25. Setation: Palp: 131, 101, 2026, 1014; Fe: I–III 323, IV 321; Pa: I–IV 000; Ti: I–III 2026, IV 2024; Mt: I–II 1014, III 2026, IV 3036. Measurements of palp and legs: Palp 5.8 (1.9, 0.8, 1.3, –, 1.8), I 14.6 (4.5, 2.4, 4.1, 2.3, 1.3), II 16.9 (5.2, 2.4, 4.1, 3.8, 1.4), III 13.2 (4.4, 2.0, 3.0, 2.6, 1.2), IV 14.2 (4.7, 1.8, 3.4, 3.0, 1.3). Leg formula: II-I-IV-III. Cheliceral furrow with three anterior and four posterior teeth, and with ~ 45 denticles (Fig. 4B).

Copulatory organ as in Fig. 2. Epigynal field wider than long, with short anterior bands and one slit sensillum on each side close to the epigynal field. Lateral lobes fused, with some fusion bubbles along median line. Fertilisation ducts arising posterolaterally. Membranous sac between fertilisation ducts almost rectangular.

Colouration in ethanol as in males, but generally slightly darker, Opisthosoma brown dorsally (Fig. 4G, H).

Distribution. Known only from the type locality (Fig. 6).

Sinopoda wuyiensis Liu, 2021

Figs 3, 4C, D, I–L, 5C, 6

Sinopoda wuyiensis Liu, in Zhang et al. 2021: 20, fig. 9A–D (holotype female from Wuyishan National Nature Reserve, Fujian Province, deposited in College of Life Science, Hubei University LJ-202002-ZY, examined)




Figure 3. *Sinopoda wuyiensis* Liu, 2021 **A–C** palp, left, ventral view **D** epigyne **E** vulva (**A** prolateral view **B, D** ventral view **C** retrolateral view **E** dorsal view). Abbreviations: AB – anterior bands, C – conductor, dRTA – dorsal branch of RTA, E – embolus, EA – embolic apophysis, FD – fertilisation duct, GA – glandular appendage, LL – lateral lobes, LS – lobal septum, MS – membranous sac, PP – posterior part of spermathecae, SP – spermophore, T – tegulum, TP – regular protrusion, vRTA – ventral branch of RTA. Scale bars: 0.5 mm.




Figure 4. **A, B** cheliceral dentition of *Sinopoda muyuensis* sp. nov. **C, D** cheliceral dentition of *Sinopoda wuyiensis* Liu, 2021 **E–H** habitus of *Sinopoda muyuensis* sp. nov. **I–L** habitus of *Sinopoda wuyiensis* Liu, 2021 (**A, C, F, J** male, ventral view **E, I** male, dorsal view **B, D, H, L** female, ventral view **G, K** female, dorsal view). Scale bars: 0.5 mm (**A–D**); 5 mm (**E–L**).

Material examined. 3♂, 4♀ (HUST 0004), **CHINA: Fujian:** Wuyishan National Reserve, Guadun Village; 27.58°N, 117.48°E; 16.XI.2021; Y. Zhong leg.

Diagnosis. Males of *S. wuyiensis* can be distinguished from other *Sinopoda* species by the combination of the following characters: bases of the tegulum with a distinct sub-triangular protrusion, embolic apophysis reduced, distinctly narrower and shorter




Figure 5. **A, B** Photograph of living *Sinopoda muyuensis* sp. nov. **C** photograph of living *Sinopoda wuyiensis* Liu, 2021 **A** male **B, C** female.




Figure 6. Distribution records of *Sinopoda* species in China **1** *Sinopoda muyuensis* sp. nov. **2** *Sinopoda wuyiensis* Liu, 2021.

than the embolus, and the dRTA about twice as long as the vRTA in ventral view. Females of *S. wuyiensis* may be recognised by the following combination of characters: epigynal field almost fusiform, lateral lobes fused without visible seam, with their anterior and posterior margins almost parallel, and glandular appendages laterad, as wide as posterior part of spermathecae (Fig. 3).

Description. Male. PL 4.5, PW 4.3, AW 2.5, OL 5.3, OW 3.0. Eyes: AME 0.18, ALE 0.28, PME 0.26, PLE 0.31, AME–AME 0.24, AME–ALE 0.10, PME–PME 0.24, PME–PLE 0.28, AME–PME 0.33, ALE–PLE 0.23, CH AME 0.10, CH ALE 0.14. Setation: Palp: 131, 101, 1021; Fe: I–III 323, IV 321; Pa: I–IV 000; Ti: I–IV 2226; Mt: I 1014, II 2024, III–IV 3036. Measurements of palp and legs: Palp 6.4 (1.9, 1.2, 1.4, –, 1.9), I 18.8 (5.1, 2.2, 5.5, 4.7, 1.3), II 21.9 (6.0, 2.2, 6.2, 5.9, 1.6), III 16.8 (5.1, 2.0, 4.6, 3.6, 1.5), IV 18.3 (4.6, 2.1, 5.1, 4.9, 1.6). Leg formula: II–I–IV–III. Cheliceral furrow with three anterior and four posterior teeth, and with ~ 38 denticles (Fig. 4C). Carapace deep yellowish brown dorsally, laterally and posteriorly with dark brown U-shaped pattern, with shallow fovea and radial furrows. Chelicerae deep reddish brown. Sternum yellow with brown margin. Endites and labium deep yellow.

lowish brown, with margin deep brown. Legs deep yellowish brown, covered by short spines. Opisthosoma brown dorsally, median part with four brown dots, posteriorly dark brown. Opisthosoma uniformly greyish brown with some brown patches ventrally (Fig. 4I, J).

Palp as in Fig. 3A–C. Cymbium $\sim 2\times$ longer than tibia in ventral view. Embolus slightly curved, arising from tegulum at nearly the 6-o’ clock-position in ventral view. Conductor curving distally, arising at 12- to 1-o’clock-position from tegulum. Sper-mophore almost straight. RTA arising mesially to distally from tibia, with distinct brush of stiff setae. vRTA wider than dRTA, dRTA long, in ventral view proximal part wide and tip tapering.

Female. PL 5.4, PW 4.7, AW 3.0, OL 5.3, OW 3.6. Eyes: AME 0.21, ALE 0.33, PME 0.21, PLE 0.39, AME–AME 0.26, AME–ALE 0.07, PME–PME 0.33, PME–PLE 0.46, AME–PME 0.35, ALE–PLE 0.25, CH AME 0.15, CH ALE 0.20. Seta-tion: Palp: 131, 101, 2121 1014; Fe: I–III 323, IV 321; Pa: I–IV 000; Ti: I–III 2026, IV 2226; Mt: I–II 1014, III 2024, IV 3036. Measurements of palp and legs: Palp 5.8 (2.0, 0.6, 1.0, –, 2.2), I 15.7 (4.7, 1.7, 4.1, 3.6, 1.6), II 16.2 (4.8, 1.8, 4.5, 3.7, 1.4), III 13.9 (4.4, 1.9, 3.5, 3.0, 1.1), IV 15.1 (4.4, 1.9, 4.1, 3.6, 1.1). Leg formula: II-I-IV-III. Colouration in ethanol as in males, but opisthosoma distinctly darker dorsally and ventrally (Fig. 4K, L; see Zhang et al. (2021) for others described).

Distribution. Known only from the type locality (Fig. 6).

Acknowledgements

The manuscript benefitted greatly from comments by Gergin Blagoev, Galina N. Azarkina, Tamas Szuts, and İlhan Coşar. We thank Guofei Ma (Shennongjia National Park Administration), Jiangwei Zheng (HUST), Han Dong (HUST), and Wei Wang (HUST) for assistance in collecting specimens. This study was financially supported by the National Natural Sciences Foundation of China (32000303), Technology Innovation Project of Hubei Province (2019AC161): Biodiversity research, monitoring and demonstration in Shennongjia National Park, Hubei Province Key Laboratory of Conservation Biology for Shennongjia Golden Monkey Foundation (No. SNJKL2021003), and the program of Hubei University of Science and Technology (2021ZX12).

References

- Gong LJ, Zhong Y (2021) First description of the female of *Sinopoda yichangensis* Zhu, Zhong & Yang, 2020 (Araneae, Sparassidae). ZooKeys 1067: 93–100. <https://doi.org/10.3897/zookeys.1067.72419>
- Grall E, Jäger P (2020) Forty-seven new species of *Sinopoda* from Asia with a considerable extension of the distribution range to the south and description of a new species group

- (Sparassidae: Heteropodinae). Zootaxa 4797(1): 1–101. <https://doi.org/10.11646/zootaxa.4797.1.1>
- Jäger P (1999) *Sinopoda*, a new genus of Heteropodinae (Araneae, Sparassidae) from Asia. The Journal of Arachnology 27: 19–24.
- Jäger P, Ono H (2000) Sparassidae of Japan. I. New species of *olios*, *Heteropoda*, and *Sinopoda* with notes on some known species (Araneae: Sparassidae: Sparassinae and Heteropodinae). Acta Arachnologica 49: 41–60. <https://doi.org/10.2476/asjaa.49.41>
- Liu J, Li SQ, Jäger P (2008) New cave-dwelling huntsman spider species of the genus *Sinopoda* (Araneae: Sparassidae) from southern China. Zootaxa 1857(1): 1–20. <https://doi.org/10.11646/zootaxa.1857.1.1>
- World Spider Catalog WSC (2022) World Spider Catalog, version 23.0. Natural History Museum, Bern. <https://doi.org/10.24436/2> [accessed on 27 July 2022]
- Zhang BS, Zhang ZS, Zhang F (2015) Three new *Sinopoda* species (Araneae: Sparassidae) from southern China. Zootaxa 3974(1): 59–75. <https://doi.org/10.11646/zootaxa.3974.1.4>
- Zhang H, Zhong Y, Zhu Y, Agnarsson I, Liu J (2021) A molecular phylogeny of the Chinese *Sinopoda* spiders (Sparassidae, Heteropodinae): implications for taxonomy. PeerJ 9: e11775. [26 pp] <https://doi.org/10.7717/peerj.11775>
- Zhu Y, Zhong Y, Yang TB (2020) One new species of the genus *Sinopoda* from Hubei Province, with description of the male of *Sinopoda angulata* (Araneae, Sparassidae). Biodiversity Data Journal 8: e55377. <https://doi.org/10.3897/BDJ.8.e55377>