

# Two new species of *Anacaena* Thomson, 1859 (Coleoptera, Hydrophilidae) from Northern Luzon, Philippines

Enrico Gerard S. Sanchez<sup>1</sup>, Emmanuel D. Delocado<sup>1</sup>, Hendrik Freitag<sup>1</sup>

<sup>1</sup> *Ateneo Biodiversity Research Laboratory, Department of Biology, School of Science and Engineering, Ateneo de Manila University, Quezon City, 1108 Philippines*

Corresponding author: Enrico Gerard S. Sanchez ([enrico.sanchez@obf.ateneo.edu](mailto:enrico.sanchez@obf.ateneo.edu))

---

Academic editor: Mariano Michat | Received 24 April 2022 | Accepted 1 June 2022 | Published 12 July 2022

---

<http://zoobank.org/F9F1E9A2-CB92-49CB-8D4C-FB3FAC592F96>

---

**Citation:** Sanchez EGS, Delocado ED, Freitag H (2022) Two new species of *Anacaena* Thomson, 1859 (Coleoptera, Hydrophilidae) from Northern Luzon, Philippines. *ZooKeys* 1112: 11–25. <https://doi.org/10.3897/zookeys.1112.85752>

---

## Abstract

Two species of *Anacaena* Thomson, 1859, *A. angatubay* **sp. nov.** and *A. auxilium* **sp. nov.**, are described from Northern Luzon, Philippines. The new species can be distinguished through colour, body shape, surface puncturation and characteristic aedeagi. Descriptions are provided and complemented with habitus photographs and drawings of the aedeagi. Data on genus distribution in the Philippines are reviewed and an updated Philippine checklist is provided.

## Keywords

Biodiversity assessment, high altitude, species inventory, taxonomy

## Introduction

*Anacaena* Thomson, 1859 is the most speciose genus under the tribe Anacaenini Hansen, 1991 (Short and Fikáček 2011; Archangelsky et al. 2016). Like other Anacaenini representatives, *Anacaena* are small, typically ranging from 1.5 to 3.3 mm in length (Komarek et al. 2003; Archangelsky et al. 2016). Adults are often collected on side pools at riverbanks, specifically in leaf litter or bottom gravel (Komarek 2010; Komarek and Freitag 2014). While *Anacaena* are deemed to be generally aquatic, representatives have been documented also from entirely terrestrial habitats (Komarek 2011).

Compared to other representatives of Hydrophilidae Latreille, 1802, *Anacaena* can be distinguished by examination of its second maxillary palpomere, which is always slightly expanded relative to the third one. Additional distinguishing features include the dorsal face of the head lacking metallic lustre, metafemora with slight hydrofuge pubescence on the proximal portion and a prosternum which is not carinate (Komarek et al. 2003). Meanwhile, species identification is done through the comparison of aedeagi. Other features which can discriminate congeners include elytral puncturation, colour, metafemoral pubescence and infuscation and colour of the maxillary palpomeres (Komarek 2012; Komarek and Freitag 2014; Archangelsky et al. 2016).

The genus *Anacaena* was established in 1859 for *Anacaena globulus* (Paykull, 1798), which was initially described as “*Hydrophilus globulus*”, based on characters of the femur and spurs on the tibia (Thomson 1859). For the next 140 years, not much study on *Anacaena* was done until the works of Gentili (1993, 1996, 2002), who increased the species number to 29. In the last two decades, there was a five-fold increase in the inventory of *Anacaena* species attributed significantly to the revisions on representatives from the Oriental region by Komarek (2006, 2009, 2010, 2011, 2012, 2013) and Komarek and Freitag (2014). Numerous new species were described from museum collections accumulated through expeditions in the past 30 years (Komarek 2010, 2013).

The massive increase can be partly attributed also to the synonymisation of “*Enigmata* Hansen, 1999”, “*Gentilina* Hebauer, 2003”, “*Grodum* Hansen, 1999”, “*Hebauerina* Gentili, 2002”, “*Omnioops* Perkins & Short, 2004” and “*Paranacaena* Blackburn, 1888”, given molecular (Komarek and Beutel 2007) and morphological evidence (Komarek 2009). The synonymisations were summarised by Short and Fikáček (2011). Conversely, years later, some species previously identified as *Anacaena* were transferred to *Crenitulus* Winters, 1926 and *Paracymus* Thomson, 1867 (Komarek 2012; Fikáček and Vondráček 2014). Currently, there are at least 150 *Anacaena* species worldwide, with a key to species for China, Indonesia, Philippines and continental Southeast Asia (Komarek 2010, 2012, 2013; Komarek and Freitag 2014).

For species from New Guinea, grouping and clustering were proposed by Komarek (2009), based on patterns of elytral punctures, eye constriction and median lobe details. Another species group, *A. yunnanensis*-group, was established by sole morphological means for species found in China (Komarek 2012). However, subsequent studies did not observe these species groups (Komarek 2010, 2011, 2013).

The only study on Philippine *Anacaena* (Komarek and Freitag 2014) lists 15 species. The recorded distribution is limited to Greater Luzon, Greater Mindoro, Greater Palawan and Greater Mindanao. While Philippine endemism of Hydrophilidae is high at 54% (Freitag et al. 2016), it is remarkable that all *Anacaena* described from the Philippines are endemic (Komarek and Freitag 2014).

Recent expeditions uncovered an enormous richness of endemic insect species in high altitudes (Komarek and Freitag 2014, 2020). It is of paramount importance to continue surveying these high-elevation localities to elucidate entomofaunal diversity

further. In this contribution, two new species, *A. angatbuhay* sp. nov. and *A. auxilium* sp. nov., are described morphologically from high elevation (> 1000 m a.s.l.) localities in Ifugao, Ilocos Sur and Mountain Province in Northern Luzon.

## Materials and methods

### Taxon Sampling

Specimens examined came from preserved collections and recent field expeditions. Adults were collected from the field using light traps (see Freitag 2015; Delocado and Freitag 2021). Specimens were stored in vials with 95% ethanol and were placed in a freezer (−20 °C).

### Morphological analysis

Selected *Anacaena* specimens were dissected by separating the terminal abdominal part. The terminal parts of the abdomen, including the aedeagus, were mounted on a microscope slide with small amounts of lactic acid for about 24 hours to clear sclerotised tissue. The aedeagi were examined using an OLYMPUS CX21 compound microscope and were compared to type specimens of described Philippine *Anacaena*. Specimens were identified using the key for Philippine *Anacaena* (Komarek and Freitag 2014). Aedeagi and dissected portions were photographed using a DinoEye Eyepiece camera attachment (AnMo Electronics Corp., New Taipei City, Taiwan), where multiple images from various focal points were taken. These images were then stacked by utilising CombineZP (Hadley 2010). Vector image files were generated in Adobe Photoshop (Adobe, San Jose, CA, USA).

Habitus photographs were taken using a CANON EOS 6D with MP-E 65 mm f/2.8 Macro Photo lens (Canon, Tokyo, Japan). Multiple images were stacked using Helicon Focus 7.6.1 software (Helicon Soft, Kharkiv, Ukraine). Type specimens of the new species were deposited at the Philippine National Museum of Natural History, Manila Philippines (**PNM**), Ateneo de Manila University, Quezon City, Philippines (**AdMU**), and Museum für Naturkunde Berlin, Germany (**ZMB**). Holotype labels were quoted verbatim from labels with backslashes (\) indicating line break. Morphological terminology follows the respective Hydrophiloidea chapter of the Handbook of Zoology (Archangelsky et al. 2016).

The following abbreviations were used:

- EL** elytra length;
- EW** elytra width;
- PL** pronotum length;
- PW** pronotum width;
- TL** total length;
- TW** total width.

## Taxonomy

Family Hydrophilidae Latreille, 1802

Genus *Anacaena* Thomson, 1859

*Anacaena angatubay* sp. nov.

<http://zoobank.org/A237568F-6B56-4F69-A5F0-08D4D2311949>

Figs 1–2, 5


**Type locality.** PHILIPPINES • Luzon, Ifugao, Banaue, Sumigar Bridge; mountain creek, secondary forest; 16°59'37"N, 121°02'51"E; ca. 1700 m a.s.l.

**Type material.** *Holotype*: PHILIPPINES • ♂ (PNM: GS076), “PHIL.: Luzon, Ifugao, Banaue, \ Sumigar Bridge; mt. creek, sec. forest; \ 16°59'37"N; 121°02'51"E; ca. 1700 m a.s.l.; \ Nov. 1997, leg. Mey (455)L”; GS076, specimen and terminal parts of the abdomen, including genitalia, were glued separately on the entomological card. *Paratypes*: PHILIPPINES • 4♂♂ (ADMU: GS068, GS075; ZMB: GS069): same data as holotype.

**Description.** (Fig. 1). TL 2.4 mm (2.1–2.5 mm); TW 1.5 mm (1.3–1.5 mm); EL 1.7 mm (1.5–1.9 mm). Body form oval, moderately convex (Fig. 2); elytra about 3.2 times as long as pronotum (dorsal view).

**Head:** Clypeus dark brown to black, moderately large, anterior margin straight, with indistinct antero-lateral angle. Frons black; frontoclypeal suture visible. Labrum black. Puncturation regular, coarse, strongly impressed throughout; interstices as wide as diameter of one puncture; series of densely arranged punctures along inner margin of eyes absent. Ventral punctures irregular, coarse, shallowly impressed, interstices as wide as diameter of one puncture. Maxillary palpomeres dark yellow to brown; palpomeres 1–3 dark yellow; palpomere 2 moderately inflated, lateral margins black. Palpomere 4 widest toward mid-length wider proximally than distally; inner margin straight; outer margin distinctly convex; proximal half yellow; anterior 0.3 black, but apex yellow. Mentum with fine setae on lateral margins; anterior margin with distinct median incision. Labial palpi stout, not longer than lateral edge of mentum; palpomere 3 about twice as long as palpomere 2. Eyes not constricted anteriorly, dorsal and ventral portions of almost equal size. Antennae capitate, 9-segmented; scape triangular, broadest at base; pedicel oval; antennomere 1 (segment 3) elongate oval, pointed bluntly at apex; antennomeres 2–4 smallest in length and width, paler than adjacent segments, indistinct intermediate segments, slightly longer than antennomere 1, decreasing in size distally; antennomere 5–7 darkest of all antennomeres; antennomeres 5 and 6 of equal length, with numerous, thick, erect, dark brown setae. Segment 7 (first club segment) irregularly globular; segment 8 globular. Terminal segment slightly longer than segments 7+8, widest near mid-length, curved on lateral margins, asymmetrical apicad; setae distributed throughout, dark brown, long, thick, erect, but setae on apical one-third longer. Preocular patches absent.

**Thorax:** PW/PL = 2.93; PL/PW = 0.34. Pronotum dark brown on the disc, light brown anterior and laterad, with narrow light brown to yellow margins not wider than



**Figure 1–4.** Habitus of new *Anacaena* species **1, 2** *Anacaena angatbuhay* sp. nov. **3, 4** *Anacaena auxilium* sp. nov. in **(1, 3)** dorsal and **(2, 4)** lateral views. Scale bar: 1 mm.

diameter of eye. Pronotal punctures large, sparse, deeply impressed on the disc, shallowly impressed laterad; interstices as wide as diameter of 2–5 punctures. Anterior margin curving slightly inwards on both sides behind the eye, gradually curving outwards starting from lateral 0.15–0.2 on both ends. Lateral margins almost straight, with setae more numerous in the anterior portion. Postero-lateral angles slightly rounded, ca. 80–85°. Posterior margin thickly bordered, almost straight. Prosternum flat. EL/

EW = 1.2; EL/PL = 4.1; EW/PW = 1.1; TL/EW = 1.8. Elytra dark brown, darkest on disc, with very narrow yellow lateral margins; setae thin, very long, some setae one-fifth length of elytra, unevenly and sparsely distributed, but denser on the disc. Elytral punctures coarse, moderately impressed, arrangement denser on the disc, spacing as wide as punctures; rows of coarser punctures on lateral portion present, but not very distinct. Mesoventrite distinctly elevated medially, with protuberance subtly bulging.

**Legs:** Light brown, but tarsi amber-coloured. Procoxa with spine-like setae. Metafemur with minute pubescence on proximal one-fourth near outer margin, hair-line direction horizontal and recumbent. Tibia with long, thick, brown, spine-like setae along lateral margins, pointed distally; distal end of tibia with coupled pairs of setae distinctly longer than proximal setae, with exterior pair longer than interior pair; setae uneven in length, with terminal pair of setae at least 1.5 as long as adjacent setae; tibiae length ratio (protibia: mesotibia: metatibia) 1.0: 1.3: 2.0. Metatarsus size slightly longer than metatibia. Tarsi length ratio (protarsus: mesotarsus: metatarsus) 1.0: 2.0: 2.4.

**Aedeagus:** (Fig. 5) Parameres symmetrical, apices rounded; apical region widens laterally; inner margins slightly concaving until apical one-third, then forming a mesal gap, converging to base; outer margins slightly biconvex with pronounced lateral expansion towards posterior 0.4 then slightly narrowing; combined width of parameres approximately the same or slightly broader as phallobase; basal portion 5 times as wide as apical part. Median lobe with apex rounded, broadens starting in anterior 0.6, abruptly bulging prominently at posterior 0.25, then narrowing before converging at the base; base of median lobe about 4 times as wide as width of apex; basal apophyses extending about one-third into the phallobase, distinctly curving outwards, narrowing basad. Phallobase longer than parameres, longer than wide; median reinforcement or pigmented line absent.

**Differential diagnosis.** *A. angatbuhay* sp. nov. (Fig. 1) resembles *A. philippina* Komarek & Freitag, 2014 and *A. zamboangana* Komarek & Freitag, 2014 in terms of colour and size. However, the new species is different from these congeners in that its labial palpi are stout compared to the slender labial palpi of *A. philippina* and *A. zamboangana*. Additionally, the new species has spine-like setae on procoxa unlike *A. philippina* and *A. zamboangana*.

Moreover, in terms of the male genitalia, *A. angatbuhay* sp. nov. and *A. philippina* exhibit similarities in the apical portion of their median lobes. While the median lobe of *A. philippina* has straight margins, the median lobe of the new species has a distinctly pointed lateral expansion towards the middle, loosely resembling a diamond shape. Additionally, the phallobase of *A. angatbuhay* sp. nov. is considerably broader than that of *A. philippina*.

**Distribution.** The species is only known from the type locality (Fig. 7).

**Remarks.** No external sexual dimorphism is observed.

**Etymology.** The species epithet alludes to the Angat Buhay (literal translation: lifting lives) anti-poverty flagship programme of Her Excellency Maria Leonor “Leni” Gerona Robredo, the 14<sup>th</sup> Vice President of the Republic of the Philippines who concluded her term this year. It is in honour of the exemplary service of her office, with

emphasis on engaging with local communities, uplifting the marginalised and heeding the needs of the health care sector, especially in the time of the pandemic. The term is used as a noun in apposition.

***Anacaena auxilium* sp. nov.**

<http://zoobank.org/E3BC4B40-3228-4317-8213-CF2BE3A13748>


Figs 3–4, 6

**Type locality.** PHILIPPINES • Luzon, Mountain Province, Bauko; Besang mountain river, Chico River tributary; rural/agricultural area; boulders; 17°00'14"N, 120°53'49"E; ca. 1080 m a.s.l.

**Type material.** *Holotype*: PHILIPPINES • ♂ (PNM: GS066), “PHIL.: Luzon, Mountain Province, \ Bauko; Besang mt. river, Chico River tributary; \ rural/agricultural area; boulders, \ 17°00'14"N; 120°53'49"E; ca. 1080 m a.s.l.; \ Nov. 1997, leg. Mey (451)L”; GS066, specimen and terminal parts of the abdomen, including genitalia, were glued separately on the entomological card. *Paratypes*: PHILIPPINES • 1♂, 2♀♀ (ADMU: GS009, GS067; ZMB: GS010): same data as holotype. • 1♂, 2♀♀ (ADMU: GS034, GS035, GS038): Luzon, Ilocos Sur, Suyo Municipality, Tagudin-Cervantes-Sabangan Road, Besang Pass Area; secondary forest, mountain creek; 16°57'17"N; 120°38'57"E; ca. 1200 m a.s.l.; 15 Apr. 2019, leg. Freitag, Garces, Pangantihon “(448)L”.

**Description.** (Fig. 3). TL 2.8 mm (2.3–2.8 mm); TW 1.7 mm (1.4–1.7 mm); EL 1.8 mm (1.5–1.9 mm). Body form oval, moderately convex (Fig. 4); elytra about 3.4 times as long as pronotum (dorsal view), greatest width at anterior one-third.

**Head:** Clypeus black, moderately large, straight anteriorly, with distinct anterolateral angle. Frons black; frontoclypeal suture visible laterally. Labrum black. Punctuation regular, coarse, shallowly impressed; interstices as wide as diameter of one puncture mediad, approximately twice as wide as diameter of punctures laterad; series of densely arranged punctures along inner margin of eyes absent. Ventral punctures obsolete. Maxillary palpomeres yellow to brown; palpomeres 1–3 dark yellow; palpomere 2 moderately inflated. Palpomere 4 almost entirely infuscated, lighter towards palpomere 3, widest towards mid-length; inner margin slightly convex; outer margin distinctly convex. Mentum with fine setae on lateral margins; anterior margin with median incision. Labial palpi stout, not longer than lateral edge of mentum; palpomere 3 slightly longer than palpomere 2. Eyes not constricted anteriorly, dorsal and ventral portions of almost equal size. Antennae capitate, 9-segmented; scape parallel-sided; pedicel narrowest towards mid-length; antennomere 1 (segment 3) not distinctly elongated, broadest at base, narrows distally; antennomeres 2–4 smallest in length and width, lighter than adjacent segments, smaller than antennomere 1 (segment 3), increasing size distally; antennomere 5–7 darkest of all antennomeres; antennomeres 5 and 6 of equal length; setae along lateral margins black, long, thick, erect. Segment 7 irregularly shaped, asymmetrical; segment 8 globular. Terminal segment as long as seg-



**Figure 5–6.** Aedeagi of **5** *Anacaena angatbuhay* sp. nov. holotype (GS076) **6** *Anacaena auxilium* sp. nov. holotype (GS066) in dorsal view. Scale bar: 0.25 mm.

ments 7 and 8, widest at base, subparallel on lateral margins, asymmetrical apicad with apex present on distal side; setae distributed throughout, black, long, thick, erect, but setae on apical one-third longer. Preocular patches absent.

**Thorax:**  $PW/PL = 2.62$ ;  $PL/PW = 0.38$ . Pronotum largely black, with narrow light brown or yellow lateral margins not wider than diameter of eye. Puncturation of two series: larger, coarser series evenly spaced, with interstices as wide as diameter of one puncture; smaller, fine series near postero-medial margin, irregularly scattered, one-fourth as wide as larger series. Anterior margin curving inwards on both sides from antero-lateral angle, with deepest point on 0.2 laterad and curving outwards mediad. Lateral margins slightly curved, without setae. Postero-lateral angle ca.  $70\text{--}80^\circ$ . Posterior margin almost straight

with curvature laterad. Prosternum flat. EL/EW = 1.13; EL/PL = 4.13; EW/PW = 1.11; TL/EW = 1.69. Elytra black, with very narrow brown lateral margins and posterior area; setae absent. Shoulder regions not accentuated. Punctures coarse, strongly impressed especially near the disc, arrangement dense, spacing as wide as punctures near the disc, wider than punctures laterad; depressions or rows of coarser punctures on lateral portion absent. Mesoventrite distinctly elevated medially, with slightly pointed protuberance.


**Legs:** Dark brown, but tarsi lighter. Spine-like setae on procoxa present. Metafemur pubescent on proximal one-fourth and near outer margin, hairline direction horizontal and recumbent. Tibia with long, thick, black, spine-like setae along lateral margins, pointing distally; terminal pair of setae at least 1.5 times as long as adjacent setae; tibiae length ratio (protibia: mesotibia: metatibia) 1: 1.3: 1.5. Metatarsus size as long as metatibia or slightly shorter. Tarsi length ratio (protarsus: mesotarsus: metatarsus) 1.0: 1.4: 2.2.

**Aedeagus:** (Fig. 6) Parameres apical region blunt, broadly pointing medially, slightly asymmetrical; inner margins almost straight on anterior half, widening medially on posterior half, mesal gap tapering to a narrow opening about 0.25 times as long as parameres; outer margins convex, bulging at anterior 0.5; combined width of parameres approximately the same or slightly broader as diameter of phallobase; basal portion 4 times as wide as apical part. Median lobe with apex blunt, apical portion subparallel, broadening starting in anterior 0.25, broadest before basal apophyses; base of median lobe not distinctly connected with parameres, about 2 times as wide as the width of the apex of median lobe; basal apophyses barely extending beyond the anterior portion of phallobase, narrow and subparallel. Phallobase longer than parameres, longer than wide; median reinforcement or pigmented line absent.

**Differential diagnosis.** *A. auxilium* sp. nov. (Fig. 3) is remarkably larger than most Philippine congeners. In length, *A. auxilium* sp. nov. (2.3–2.8 mm) overlaps with *A. cordillera* Komarek & Freitag, 2014 (2.6–3.1 mm) and *A. levistriata* Komarek & Freitag, 2014 (2.1–2.7 mm). However, the new species has a smaller body width than *A. cordillera* (1.3 mm vs. > 1.5–1.8 mm) and a longer elytral length (1.5 mm vs. < 1.20–1.27 mm) than both congeners. Additionally, the new species has stout labial palpi which are not longer than the lateral edge of the mentum, while both *A. cordillera* and *A. levistriata* have labial palpi longer than the lateral edge of the mentum. Additionally, the labial palpi of *A. auxilium* sp. nov. are stout, while the labial palpi of *A. cordillera* are slender. Moreover, pre-ocular patches are absent in *A. auxilium* sp. nov., while these are distinct in *A. cordillera*. Remarkably, the new species has a bipunctuate elytra with large, coarse series and small, fine series, while *A. cordillera* only exhibits the latter and *A. levistriata* only exhibits the former.

While the parameres of the new species and *A. levistriata* look similar, *A. auxilium* sp. nov. can be differentiated in that its median lobe is distinctly longer (ca. 0.4×) than its parameres. Meanwhile, the apical portions of the median lobe and parameres of *A. cordillera* are clearly different from those of the new species.

**Distribution.** The species is documented from two nearby localities, namely Ilocos Sur and Mountain Province (Fig. 7).



**Figure 7.** Map showing the updated distribution of *Anacaena* in Northern Luzon, including Komarek & Freitag (2014) data.

**Remarks.** Specimens from a preserved collection during 1997 fieldwork showed no pertinent phenotypic plasticity to specimens collected in a field expedition 22 years later. No external sexual dimorphism is observed.

**Etymology.** The species epithet ‘auxilium’ is Latin for ‘help’. The new species is dedicated to the Mary Help of Christians Seminary system to which the first author expresses gratitude for constant inspiration. Additionally, ‘help’ alludes to the vulnerable state of freshwater biodiversity in the tropics.

### Updated checklist of the species of *Anacaena* in the Philippines

1. *Anacaena albay* Komarek & Freitag, 2014: Luzon Island (Albay)
2. *Anacaena amplocomata* Komarek & Freitag, 2014: Luzon Island (Mountain Province)
3. *Anacaena angatbuhay* sp. nov.: Luzon Island (Ifugao)
4. *Anacaena apo* Komarek & Freitag, 2014: Mindanao Island (Davao)
5. *Anacaena auxiliium* sp. nov.: Luzon Island (Ilocos Sur, Mountain Province)
6. *Anacaena balabag* Komarek & Freitag, 2014: Mindanao Island (Cotabato)
7. *Anacaena cordillera* Komarek & Freitag, 2014: Luzon Island (Ifugao, Mountain Province)
8. *Anacaena davao* Komarek & Freitag, 2014: Mindanao Island (Davao)

9. *Anacaena destructa* Komarek & Freitag, 2014: Leyte, Mindanao Island (Davao)
10. *Anacaena emergens* Komarek & Freitag, 2014: Tinaga Island, Mindoro, Palawan, Busuanga
11. *Anacaena hemisphaerica* Komarek & Freitag, 2014: Mindanao Island (Misamis Occidental)
12. *Anacaena levistriata* Komarek & Freitag, 2014: Luzon Island (Mountain Province, Rizal)
13. *Anacaena philippina* Komarek & Freitag, 2014: Luzon Island (Mountain Province, Benguet, Laguna), Mindoro, Leyte, Mindanao Island (Agusan del Sur)
14. *Anacaena princesa* Komarek & Freitag, 2014: Palawan
15. *Anacaena quezona* Komarek & Freitag, 2014: Luzon Island (Quezon Province)
16. *Anacaena sulcata* Komarek & Freitag, 2014: Luzon Island (Benguet, Mountain Province)
17. *Anacaena zamboangana* Komarek & Freitag, 2014: Mindanao Island (Zamboanga del Sur)

## Discussion

The total number of Philippine *Anacaena* increased to 17. All known species of Philippine *Anacaena* are endemic, affirming Komarek and Freitag (2014). Thirteen species are aquatic and four have unknown habitats. The documentation of *A. auxilium* sp. nov. from Ilocos Sur is the first *Anacaena* record from the province. Although *Anacaena* has been previously documented in Ifugao and Mountain Province in Northern Luzon, this study uncovered greater diversity in higher altitudes (Fig. 7), as projected in Freitag et al. (2016).

Previous studies have identified that higher altitudes foster higher insect diversity (Mengual et al. 2006; Komarek and Freitag 2014; Kaltenbach et al. 2020a). Although *A. levistriata* and *A. philippina* have been documented in areas ranging from lower (ca. 10–400 m a.s.l. in Mindoro and Rizal Province) to higher (ca. 1450 m a.s.l. in Mountain Province) altitudes, most high-elevation species have a distribution which is limited to their corresponding altitude. This pattern has been documented in the localities of Mountain Province (ca. 1450 m a.s.l.) and Ifugao (ca. 1700 m a.s.l. for *A. angatbuhay* sp. nov.) in Northern Luzon, Davao (ca. 1830 m a.s.l.) and Misamis (ca. 1500 m a.s.l.) in Mindanao Island. This trend in species-specific distribution to a limited altitudinal range has been documented in beetles (Maveety et al. 2011; Koutroumpa et al. 2013; Sheth et al. 2019).

External morphological characters are helpful for species identification in some Philippine *Anacaena*. In particular, body measurements (size, body width, elytra length), colour and features, such as the shape of the labial palpi, puncturation in the pronotum and elytra and presence of procoxal setae, can be used to discriminate congeners (Komarek 2009, 2011; Short and Fikáček 2011). More importantly, careful examination and comparison of the male genitalia is important in species identification

(Dufour 1848; Rudoy and Ribera 2016). This was demonstrated in how *A. angatbuhay* sp. nov. and *A. philippina* are similar in size and colour, but are differentiated after comparing the aedeagi. On the contrary, female genitalia cannot be used to determine species-level identification with certainty (Komarek 2013).

The discovery of rather inconspicuous species in megadiverse countries is crucial, especially amidst biodiversity decline (Bálint et al. 2011). In the past decades, insect decline has been happening rapidly and even proceeding four-fold compared to that of vertebrates (Dudgeon et al. 2006; Carrizo et al. 2017; Darwall et al. 2018). While terrestrial insects face high extinction rates, the speed at which aquatic insect decline happens is much faster (Sánchez-Bayo and Wyckhuys 2019). Moreover, the magnitude at which this occurs may be higher than current estimates because of the lack of baseline data for tropical aquatic insects (Wagner et al. 2021).

Following that Freitag et al. (2016) estimated that only one-third of Philippine aquatic and riparian coleopteran diversity has been described, massive efforts are being done by the Ateneo Biodiversity Research Laboratory to uncover the remaining undescribed beetle species (Komarek and Freitag 2014, 2020; Vidal et al. 2017; Sabordo et al. 2020; Delocado and Freitag 2021), as well as those from other insect orders (Garces et al. 2018, 2020; Kaltenbach et al. 2020a, 2020b; Pelingen and Freitag 2020; Pelingen et al. 2021). While the pandemic has hampered efforts in species discovery, especially in the Global South (Bang and Khadakkar 2020; Ramvilas et al. 2021), this contribution nonetheless demonstrates that species inventory can proceed by surveying existing preserved collections and exploring previously unsampled localities.

## Acknowledgements

Sincere thanks are due to the Municipality of Suyo (Ilocos Sur) for their prior informed consent on the aquatic arthropod sampling and the Bureau of Fisheries and Aquatic Resources (BFAR) for issuing the respective Gratuitous Permit. We also want to express our gratitude to Dr Wolfram Mey (ZMB) for loaning the specimens used in this study in the scope of the BIO-PHIL programme. We want to thank the members of the Ateneo Biodiversity Research Laboratory (Dr Jhoana Garces, Clister Pangantihon, Arthien Pelingen, Marc Sabordo) for their efforts in preserving and maintaining specimens examined in this study. We also thank Dr Martin Fikáček (National Sun Yat-sen University) for his continuous support on this project. The first author would like to express his sincerest gratitude to Caitlin Rodil and Alvin Dy for their technical assistance in crafting digital illustrations.

The study was partly funded by the School of Science and Engineering Industry 4.0 Research Fund (SI4–013) of the Ateneo de Manila University and the Biodiversity Teaching in a Philippine-Cambodian-German Network (BIO-PHIL). Training of the first author and examination of specimens in a remote lab-at-home set-up was made possible by the SC Johnson Environmental Leadership Development Fund. Publication assistance was provided by the Office of the Vice President for the Loyola Schools of the Ateneo de Manila University.

## References

- Archangelsky M, Beutel RG, Komarek A (2016) 12 Hydrophiloidea. In: Beutel RG, Leschen RAB (Eds) Handbook of Zoology, Coleoptera, Beetles, Volume 1: Morphology and Systematics (Archostemata, Adephaga, Myxophaga, Polyphaga partim). Walter de Gruyter, Berlin & New York, 231–272.
- Bálint M, Domisch S, Engelhardt CHM, Haase P, Lehrian S, Sauer J, Theissing K, Pauls SU, Nowak C (2011) Cryptic biodiversity loss linked to global climate change. *Nature Climate Change* 1: 313–318. <https://doi.org/10.1038/nclimate1191>
- Bang A, Khadakkar S (2020) Biodiversity conservation during a global crisis: Consequences and the way forward. *Proceedings of the National Academy of Sciences of the United States of America* 117(48): 29995–29999. <https://doi.org/10.1073/pnas.2021460117>
- Carrizo SF, Jähnig SC, Bremerich V, Freyhof J, Harrison I, He F, Langhans SD, Tockner K, Zarfl C, Darwall W (2017) Freshwater Megafauna: Flagships for Freshwater Biodiversity under Threat. *Bioscience* 67(10): 919–927. <https://doi.org/10.1093/biosci/bix099>
- Darwall W, Bremerich V, De Wever A, Dell AI, Freyhof J, Gessner MO, Grossart HP, Harrison I, Irvine K, Jähnig SC, Jeschke JM, Lee JJ, Lu C, Lewandowska AM, Monaghan MT, Nejtgaard JC, Patricio H, Schmidt-Kloiber A, Stuart SN, Thieme M, Tockner K, Turak E, Weyl O (2018) The *Alliance for Freshwater Life*: A global call to unite efforts for freshwater biodiversity science and conservation. *Aquatic Conservation* 28(4): 1015–1022. <https://doi.org/10.1002/aqc.2958>
- Delocado ED, Freitag H (2021) Two new species of *Byrrhinus* Motschulsky, 1858 (Coleoptera, Limnichidae, Limnichinae) from Negros, Philippines. *ZooKeys* 1070: 51–72. <https://doi.org/10.3897/zookeys.1070.70531>
- Dudgeon D, Arthington AH, Gessner MO, Kawabata ZI, Knowler DJ, Lévêque C, Naiman RJ, Prieur-Richard AH, Soto D, Stiassny MLJ, Sullivan CA (2006) Freshwater biodiversity: Importance, threats, status and conservation challenges. *Biological Reviews* 81(2): 163–182. <https://doi.org/10.1017/S1464793105006950>
- Dufour L (1848) Anatomie générale des dipteres. *Annales des Sciences Naturelles*: 244–264.
- Fikáček M, Vondráček D (2014) A review of *Pseudorygmodus* (Coleoptera: Hydrophilidae), with notes on the classification of the Anacaenini and on distribution of genera endemic to southern South America. *Acta Entomologica Musei Nationalis Pragae* 54(2): 479–514.
- Freitag H (2015) Clues to aquatic beetle research in Southeast Asia: A multitude of ecological niches, microhabitats and deduced field sampling techniques. *Entomology, Ornithology & Herpetology* 4(4): e1000166. [6 pp] <https://doi.org/10.4172/2161-0983.1000166>
- Freitag H, Jäch MA, Wewalka G (2016) Diversity of aquatic and riparian Coleoptera of the Philippines: Checklist, state of knowledge, priorities for future research and conservation. *Aquatic Insects* 37(3): 177–213. <https://doi.org/10.1080/01650424.2016.1210814>
- Garces JM, Bauernfeind E, Freitag H (2018) *Sparsorythus sescarorum*, new species from Mindoro, Philippines (Ephemeroptera, Tricorythidae). *ZooKeys* 795: 13–30. <https://doi.org/10.3897/zookeys.795.28412>
- Garces JM, Sartori M, Freitag H (2020) Integrative taxonomy of the genus *Dudgeodes* Sartori, 2008 (Insecta, Ephemeroptera, Teloganodidae) from the Philippines with description of

- new species and supplementary descriptions of Southeast Asian species. *ZooKeys* 910: 93–129. <https://doi.org/10.3897/zookeys.910.48659>
- Gentili E (1993) *Paranacaena* Blackburn, 1889: a valid genus (Coleoptera, Hydrophilidae). *Giornale Italiano di Entomologia* 6: 285–296.
- Gentili E (1996) Notes on the Anacaenini from Australia and Melanesia with descriptions of new species (Coleoptera, Hydrophilidae). *Giornale Italiano di Entomologia* 8: 177–189.
- Gentili E (2002) Descrizione di nuove specie del genere *Paranacaena* Blackburn, 1889 (Coleoptera, Hydrophilidae). *Giornale Italiana di Entomologia* 10: 77–97.
- Hadley A, Hadley Web (2010) [Internet]. CombineZP; c2010 [cited 2019 August 01]. <https://combinezp.software.informer.com/>
- Kaltenbach T, Garces J, Gattolliat J-L (2020a) A new genus of Baetidae (Insecta, Ephemeroptera) from Southeast Asia. *European Journal of Taxonomy* 612: 1–32. <https://doi.org/10.5852/ejt.2020.612>
- Kaltenbach T, Garces JM, Gattolliat JL (2020b) The success story of *Labiobaetis* Novikova & Kluge in the Philippines (Ephemeroptera, Baetidae), with description of 18 new species. *ZooKeys* 1002: 1–114. <https://doi.org/10.3897/zookeys.1002.58017>
- Komarek A (2006) Taxonomic revision of *Anacaena* Thomson, 1859 III. The Indian Subcontinent (Coleoptera: Hydrophilidae). *Koleopterologische Rundschau* 76: 283–314.
- Komarek A (2009) Taxonomic revision of *Anacaena* Thomson, 1859 V. New Guinea (Coleoptera: Hydrophilidae). *Koleopterologische Rundschau* 79: 197–254.
- Komarek A (2010) Taxonomic revision of *Anacaena* Thomson, 1859 VII. Indonesia except Papua (Coleoptera: Hydrophilidae). *Koleopterologische Rundschau* 80: 113–118.
- Komarek A (2011) Taxonomic revision of *Anacaena* Thomson, 1859 VIII. Taiwan (Coleoptera: Hydrophilidae). *Koleopterologische Rundschau* 81: 229–236.
- Komarek A (2012) Taxonomic revision of *Anacaena* Thomson, 1859 IX. The People's Republic of China (Coleoptera: Hydrophilidae). *Koleopterologische Rundschau* 82: 235–284.
- Komarek A (2013) A revision of *Anacaena* Thomson, 1859 X. Laos, Malaysia, Myanmar, Thailand, and Vietnam. *Entomologica Basiliensia et Collectionis Frey* 34: 137–177.
- Komarek A, Freitag H (2014) Revision of *Anacaena* Thomson, 1859 XI. Republic of the Philippines (Coleoptera: Hydrophilidae). *Koleopterologische Rundschau* 84: 235–276.
- Komarek A, Freitag H (2020) Taxonomic revision of *Agraphhydrus* Regimbart, 1903 V. Philippine species and their first DNA barcodes (Coleoptera: Hydrophilidae: Acidocerinae). *Koleopterologische Rundschau* 90: 201–242.
- Komarek A, Jäch MA, Ji L (2003) Hydrophilidae: I. Check list and key to Palearctic and Oriental genera of aquatic Hydrophilidae (Coleoptera). In: Jäch MA, Li J (Eds) *Water Beetles of China*, vol. 3. Zoologisch-Botanische Gesellschaft in Österreich Wiener Koleopterologenverein, Vienna, 383–395.
- Komarek A, Beutel RG (2007) Phylogenetic analysis of Anacaenini (Coleoptera: Hydrophilidae: Hydrophilinae) based on morphological characters of adults. *Systematic Entomology* 32(2): 205–226. <https://doi:10.1111/j.1365-3113.2006.00359.x>
- Koutroumpa FA, Rougon D, Bertheau C, Lieutier F, Roux-Morabito G (2013) Evolutionary relationships within European *Monochamus* (Coleoptera: Cerambycidae) highlight the

- role of altitude in species delineation. *Biological Journal of the Linnean Society* 109(2): 354–376. <https://doi.org/10.1111/bij.12042>
- Maveety SA, Browne RA, Erwin TL (2011) Carabidae diversity along an altitudinal gradient in a Peruvian cloud forest (Coleoptera). *ZooKeys* 147: 651–666. <https://doi.org/10.3897/zookeys.147.2047>
- Mengual X, Ståhls G, Vujić A, Marcos-García MÁ (2006) Integrative taxonomy of Iberian *Merodon* species (Diptera, Syrphidae). *Zootaxa* 1377: 1–26.
- Pelting AL, Freitag H (2020) Description of *Neoperla mindoroensis* sp. nov., the first record of a stonefly from Mindoro, Philippines (Plecoptera, Perlidae), and identification of its life stages using COI barcodes. *ZooKeys* 954: 47–63. <https://doi.org/10.3897/zookeys.954.53746>
- Pelting AL, Murányi D, Freitag H (2021) An additional new species and records of *Neoperla* Needham, 1905 (Plecoptera, Perlidae) from the Philippines. *Tijdschrift voor Entomologie* 164(1–3): 91–105. <https://doi.org/10.1163/22119434-bja10016>
- Ramvilas G, Dhyan S, Kumar B, Sinha N, Raghavan R, Selvaraj G, Divakar N, Anoop VK, Shalu K, Sinha A, Kulkarni A, Das S, Molur S (2021) Insights on COVID-19 impacts, challenges and opportunities for India's biodiversity research: From complexity to building adaptations. *Biological Conservation* 255: e109003. <https://doi.org/10.1016/j.biocon.2021.109003>
- Rudoy A, Ribera I (2016) The macroevolution of size and complexity in insect male genitalia. *PeerJ* 4: e1882. <https://doi.org/10.7717/peerj.1882>
- Sabordo MR, Delocado E, Freitag H (2020) Two new species of the genus *Ancyronyx* Erichson, 1847 from the island of Negros, Philippines (Insecta, Coleoptera, Elmidae). *Tijdschrift voor Entomologie* 163(1): 13–30. <https://doi.org/10.1163/22119434-20192087>
- Sánchez-Bayo F, Wyckhuys KAG (2019) Worldwide decline of the entomofauna: A review of its drivers. *Biological Conservation* 232: 8–27. <https://doi.org/10.1016/j.biocon.2019.01.020>
- Sheth SD, Padhye AD, Ghate HV (2019) Factors affecting aquatic beetle communities of Northern Western Ghats of India (Arthropoda: Insecta: Coleoptera). *Annales de Limnologie* 55: 1–13. <https://doi.org/10.1051/limn/2018030>
- Short AEZ, Fikáček M (2011) World catalogue of the Hydrophiloidea (Coleoptera): Additions and corrections II (2006–2010). *Acta Entomologica Musei Nationalis Pragae* 51(1): 83–122.
- Thomson CG (1859) *Skandinaviens Coleoptera, synoptiskt bearbetade*. Tryckt uti Lundberg-ska Boktryckeriet; 1859–68., Lund, 290 pp. <https://doi.org/10.5962/bhl.title.138677>
- Vidal AR, Go KCTS, Freitag H (2017) Hydraenidae (Insecta: Coleoptera) of Mindoro, Philippines. I: *Hydraena* Kugelann, 1794 of the Baroc River Basin, Roxas, Oriental Mindoro with description of three new species. *Aquatic Insects* 38(1–2): 1–20. <https://doi.org/10.1080/01650424.2017.1303517>
- Wagner DL, Grames EM, Forister ML, Berenbaum MR, Stopak D (2021) Insect decline in the Anthropocene: Death by a thousand cuts. *Proceedings of the National Academy of Sciences of the United States of America* 118(2): e2023989118. <https://doi.org/10.1073/pnas.2023989118>