

First record of the flat-skulled woolly bat *Kerivoula depressa* and the Indochinese woolly bat *K. dongduongana* (Chiroptera, Vespertilionidae) in China

Xiaoling Liang¹, Huixian Xie¹, Yunnan Li¹,
Zhenglanyi Huang¹, Song Li², Yi Wu¹, Wenhua Yu¹

1 Key Laboratory of Conservation and Application in Biodiversity of South China, School of Life Sciences, Guangzhou University, Guangzhou 510006, China **2** Kunming Natural History Museum of Zoology, Kunming Institute of Zoology, Chinese Academy of Sciences, Kunming 650223, China

Corresponding authors: Yi Wu (wuyi@gzhu.edu.cn); Wenhua Yu (wenhua_yu@gzhu.edu.cn)

Academic editor: W. Bogdanowicz | Received 30 April 2022 | Accepted 19 December 2022 | Published 22 February 2023

<https://zoobank.org/D6FFAFDE-5310-4BA1-AD68-8FE762189718>

Citation: Liang X, Xie H, Li Y, Huang Z, Li S, Wu Y, Yu W (2023) First record of the flat-skulled woolly bat *Kerivoula depressa* and the Indochinese woolly bat *K. dongduongana* (Chiroptera, Vespertilionidae) in China. ZooKeys 1149: 1–15. <https://doi.org/10.3897/zookeys.1149.85821>

Abstract

Recent studies have revealed that the *Kerivoula depressa* complex should be divided into two species, *K. depressa* distributed mainly in Myanmar, Vietnam, Laos and Cambodia, and *K. dongduongana* found only in the Annamite Mountains of Vietnam, Laos and Cambodia. In November 2018 and April 2019, 24 woolly bats were collected by two-band harp traps in Xishuangbanna, Yunnan, China. Based on morphological, morphometric, and phylogenetic (*COI*, *Cytb*, and *RAG2* gene sequences) analyses, these bats were identified as *K. depressa* and *K. dongduongana*, representing two new species records for the country. Including the new records, six *Kerivoula* species have been recorded in China, namely *K. depressa*, *K. dongduongana*, *K. furva*, *K. kachinensis*, *K. picta* and *K. titania*. To facilitate their identification and biological research in the future, we have provided an up-to-date key to all *Kerivoula* species occurring in China.

Keywords

COI, *Cytb*, *Kerivoula dongduongana*, morphology, morphometric analyses, new records, phylogenetic inferences, *RAG2*, Yunnan

Introduction

The genus *Kerivoula* (Gray, 1842) contains 27 species in the Indomalaya-Australasia and Afrotropic ecozones (Wilson and Mittermeier 2019). Among these, *Kerivoula hardwickii* (sensu stricto) (Horsfield 1824), the most widespread species of the genus has long been treated as a species complex, with up to six recognized subspecies (Ellerman and Morrison-Scott 1951; Hill 1965; Corbet and Hill 1992). Because of the lack of comparative material, Corbet and Hill (1992), Sinha (1999), and Simmons (2005) did not acknowledge any subspecies of *K. hardwickii*. Subsequently, Bates et al. (2007) divided the species into two morphological types, *K. hardwickii* (with domed skull) and *K. depressa* (Miller, 1906) (with flattened skull). However, molecular phylogenies indicate that the taxonomy and systematics of the *K. hardwickii* complex are still ambiguous because of the occurrence of multiple divergent lineages (Francis et al. 2007, 2010; Khan et al. 2010; Douangboubpha et al. 2016; Nguyen et al. 2016). Recent genetic analyses revealed that *K. depressa* can be divided into two distinct clades (Kuo et al. 2017). Tu et al. (2018) described *K. depressa* and *K. dongduongana* (Tu et al. 2018) based on phylogenetic analyses using *COI*, *Cytb*, and *RAG2* sequences from newly obtained specimens and those from previous studies across the Indo-China Peninsula and the Philippines (Hoofer et al. 2003; Stadelmann et al. 2004; Francis et al. 2010; Khan et al. 2010; Wu et al. 2012; Kruskop 2013; Kuo et al. 2014; Kuo et al. 2017). Currently, the *K. hardwickii* complex contains five species: *K. hardwickii* (sensu stricto), *K. kachinensis* (Bates et al., 2004), *K. furva* (Kuo et al., 2017), *K. depressa*, and *K. dongduongana* (Kuo et al. 2017; Tu et al. 2018).

In China, four species of woolly bats from the genus *Kerivoula* have been recorded, including *K. picta* (Pallas, 1767), *K. furva*, *K. kachinensis*, and *K. titania* (Bates et al., 2014) (Wu et al. 2012; Kuo et al. 2017; Tu et al. 2018; Yu et al. 2018; Wilson and Mittermeier 2019; Yu et al. 2022). In November 2018 and April 2019, a series of chiropteran surveys were conducted in the southwestern region of Yunnan Province, and 24 *Kerivoula* individuals were sampled. Based on morphology, morphometric analyses, and phylogenetic inferences using *COI*, *Cytb*, and *RAG2* sequences, they were identified as *K. depressa* and *K. dongduongana*, which represent two new records of *Kerivoula* in China. In this paper, we provided details about these findings, new distribution information and an up-to-date key to identify all *Kerivoula* species occurring in China.

Materials and methods

Specimen sampling and morphological measurements and analyses

In November 2018 and April 2019, 24 *Kerivoula* bats were collected using two-band harp traps during field surveys in Xishuangbanna Tropical Botanical Garden, Yunnan, China (21°57'17"N, 101°15'26"E and 21°30'58"N, 101°30'38"E). All field survey

and sample collection protocols complied with the current laws of Yunnan Province, China. We followed the guidelines of the American Society of Mammalogists (Sikes 2016) for the care and use of animals. All voucher specimens were determined to be adults based on the degree of epiphyseal-diaphyseal fusion (Brunet-Rossinni and Wilkinson 2009). The specimens were preserved in 75% ethanol and deposited at the School of Life Sciences, Guangzhou University, China.

External and skull measurements were taken with a digital caliper to the nearest 0.01 mm following Bates and Harrison (1997) and Bates et al. (2004). Body mass was measured with an electronic scale. Twenty-four adult specimens were examined using six external and eight craniodental measurements following Tu et al. (2018) and Yu et al. (2018), and further morphometric analyses were performed using 20 specimens (Table 1). We conducted a principal component analysis (PCA) and discriminant analysis of principal components (DAPC) of craniodental measurements using the R Core Team (2013) and R add-in packages: psych (Revelle 2013), ade4 (Dray et al. 2007), adegenet (Jombart 2008), FactoMineR (Le et al. 2008), and ggplot2 (Wickham 2016).

Phylogenetic inference

We followed the DNA extraction, amplification, and sequencing procedures according to Yu et al. (2022). Three gene sequences from all 24 voucher *Kerivoula* specimens were obtained (GenBank accession numbers: *COI*: OM716930–OM716952; *Cytb*: OM735691–OM735714; *RAG2*: OM735715–OM735736). These sequences were compared with 87 *COI*, 38 *Cytb*, and 28 *RAG2* sequences of the subfamily Kerivoulinae from NCBI nucleotide databases and with the three outgroup species, including *Myotis muricola* (Gray, 1846), *Harpiocephalus harpia* (Temminck, 1840), and *Murina cyclotis* (Dobson, 1872) (Hooper et al. 2003; Khan et al. 2010; Ruedi et al. 2012). Newly generated and downloaded sequences are detailed in Appendix 1. The final matrices of *COI*, *Cytb*, and *RAG2* contained 110 taxa and 734 bp, 62 taxa and 1220 bp, and 50 taxa and 1267 bp, respectively. We inferred phylogenetic relationships using both Bayesian and maximum likelihood (ML) approaches. Sequences were aligned with MUSCLE (Edgar 2004). Bayesian analyses were performed using MrBayes v.3.2.7 (Ronquist et al. 2012), and the best-fitting models of sequence evolution were selected by MrModeltest v.2.4 (Nylander 2004) using the Akaike information criterion (GTR+I+G for *COI*, HKY+I+G for *Cytb*, HKY+G for *RAG2*). Four independent Markov chains were run, and 10,000,000 Metropolis-coupled Markov Chain Monte Carlo generations with sampling every 1000 generations were set. The first 25% samples were discarded. ML analyses were performed in IQ-TREE (Minh et al. 2020) with the best model setting in ModelFinder (Kalyaanamoorthy et al. 2017) using Bayesian information criterion (TPM2+F+I+G4 for *COI* and *Cytb*, HKY+F+R2 for *RAG2*).

Table 1. External and craniodental measurements (mm) and body mass (g) of studied *Kerivoula* species, and variable loadings on principal components (PCs) and contribution of original variables in discriminant functions (DFs).

Characters	<i>Kerivoula depressa</i>			<i>Kerivoula dongluangana</i>		
	Yunnan, China (This study)	Vietnam and Cambodia Tu et al. 2018		Yunnan, China (This study)	Vietnam and Cambodia Tu et al. 2018	
	Mean ± SD (n) (Range)	Mean ± SD (n) (Range)		Mean ± SD (n) (Range)	Mean ± SD (n) (Range)	
MASS	3.53±0.49 (12) (3.00–4.50)	–		4.05±0.25 (12) (3.50–4.40)	4.50 (1)	–
HB	36.88±2.31 (12) (32.99–39.74)	–		38.43±2.84 (12) (34.40–42.75)	–	–
TL	39.13±1.35 (12) (36.68–41.49)	–		39.97±1.90 (12) (37.20–44.50)	38.00 (1)	–
E	11.55±0.91 (12) (9.96–13.15)	–		12.05±0.89 (12) (10.83–14.04)	–	–
HF	6.98±0.65 (12) (6.09–7.83)	–		7.57±0.48 (12) (7.00–8.43)	–	–
FA	30.75±1.08 (12) (28.82–32.29)	32.08±0.15 (4) (32.00–32.30)		34.02±0.94 (12) (32.56–35.86)	32.00±1.73 (3) (30.00–33.00)	–
TIB	15.95±0.44 (12) (15.38–16.61)	–		16.37±0.58 (11) (15.70–17.44)	18.00 (1)	–
GTL	13.75±0.17 (10) (13.57–14.08)	13.65±0.27 (5) (13.34–13.98)		14.28±0.30 (10) (13.71–14.59)	13.51±0.38 (7) (12.70–13.79)	0.98 –0.14 0.21 0.01
CCL	12.14±0.17 (10) (11.88–12.43)	12.51±0.23 (5) (12.17–12.75)		12.79±0.31 (10) (12.34–13.43)	12.43±0.36 (7) (11.68–12.76)	0.95 –0.24 0.41 0.13
M ³ –M ³	4.90±0.16 (10) (4.70–5.16)	5.09±0.16 (5) (4.84–5.24)		5.12±0.14 (10) (4.92–5.37)	4.99±0.17 (7) (4.70–5.15)	0.97 –0.02 0.04 0.01
ZB	7.95±0.18 (10) (7.71–8.35)	8.22±0.21 (5) (7.89–8.45)		8.29±0.35 (10) (7.63–8.67)	8.18±0.24 (7) (7.72–8.49)	0.98 –0.10 0.05 –
GBB	6.99±0.11 (10) (6.85–7.15)	7.12±0.05 (5) (7.06–7.19)		6.99±0.11 (10) (6.85–7.15)	7.14±0.20 (7) (6.73–7.31)	– – –
BH	4.91±0.10 (10) (4.74–5.06)	4.97±0.16 (5) (4.86–5.25)		4.47±0.29 (10) (3.98–4.97)	4.81±0.20 (7) (4.57–5.13)	0.60 0.80 0.29 0.75
C ¹ –M ³	5.11±0.09 (10) (4.94–5.22)	5.37±0.13 (5) (5.17–5.49)		5.23±0.14 (10) (5.00–5.50)	5.25±0.13 (7) (5.06–5.45)	0.97 0.07 – 0.04
ML	9.17±0.28 (10) (8.61–9.56)	9.52±0.21 (5) (9.17–9.75)		9.46±0.29 (10) (8.97–9.88)	9.42±0.26 (7) (9.00–9.78)	0.98 – 0.02 0.06
C ₁ –M ₃	5.47±0.14 (10) (5.26–5.65)	5.71±0.15 (5) (5.48–5.86)		5.40±0.22 (10) (4.92–5.71)	5.46±0.15 (7) (5.29–5.76)	0.96 –0.07 0.02 –

Results

Morphological examination

Kerivoula depressa is a moderate-sized species with a forearm (FA) length of 30.75 ± 1.08 mm. Ears are small and rounded, and the posterior margin of the pinnae has a deep, smoothly concave emargination just below the apex. Overall pelage color is buff brown to dark brown. The lower part of ventral hair is dark brown, whereas its tip is light brownish yellow. Dorsal fur is of black base but with dark brown tip (Fig. 1A–C). The domed skull is small, with the greatest length of 13.75 ± 0.17 mm. The mid-portion of the braincase exceeds the frontal region in height. Its lateral profile is flattened from the rostrum to the forehead. A sagittal crest is not evident, and the lambdoid crests are relatively weak. The dental formula is I 2/3, C 1/1, P 3/3, M 3/3. The second upper incisor (I^2) is about half of the first upper incisor (I^1) in height, and the latter is one half the height of the upper canine. The third upper premolar (P^4) is distinctly higher than the anterior two. The third upper molar (M^3) is degenerated. The crown area of the first and second lower molars is approximately equal and slightly larger than the last molar (M_3).

Kerivoula dongduongana, with FA of 34.02 ± 0.94 mm, is slightly larger than *K. depressa*. Its ears are rounded with a tiny, smooth depression near the tip. *Kerivoula dongduongana* is obviously yellower than *K. depressa* in pelage coloration (Fig. 1D–F).



Figure 1. Photographs of *K. depressa* (A–C voucher GZHU 19202, male) and *K. dongduongana* (D–F voucher GZHU 19198, female) representing their lateral view (A, D), ventral pelage (B, E) and dorsal pelage (C, F).




Figure 2. Skull morphology of *K. depressa* (A–D voucher GZHU 19222, female) and *K. dongduongana* (E–H voucher GZHU 19308, female). Scale bar: 10 mm.




Figure 3. Two-dimensional PCA and DAPC plots of *Kerivoula* species based on nine craniodental measurements **A** PCA plots for *K. depressa*, *K. dongduongana*, *K. furva*, *K. kachinensis*, *K. picta*, and *K. titania* showing projections of individual specimens and variable loadings on the first two principal components **B** projections of 46 specimens and variable loadings on two discriminant functions obtained from external and craniodental measurements.

The ventral pelage is golden brown, and the base and middle portions are medium brown with golden-brown tips. The dorsal pelage is dark brown with golden-brown tips. Craniodental features are similar to *K. depressa*, but *K. dongduongana* is characterized by a flatter and longer skull (Fig. 2C, G; Table 1).

Multivariate comparison analysis

PCA based on eight craniodental measurements revealed 96.2% of the total variance from the first two principal components (PCs) (86.9% and 9.3% for PC1 and PC2, respectively) in the scatter plot of the six morphological groups (Fig. 3A). For PC1,




Figure 4. Bayesian and ML trees from analysis of *COI*, *Cytb*, and *RAG2* sequences for the *K. hardwickii* complex. Values on the branches indicate posterior probabilities and bootstrap values, respectively. The terminals *K. picta*, *K. papillosa*, *K. titania*, *K. hardwickii* (sensu stricto), *K. kachinensis*, and *K. furva* each include multiple samples (see Appendix 1).

all measurements had positive loadings (Table 1), reflecting the skull size. Larger bats were characterized by higher PC1 scores; thus, specimens of *K. kachinensis* clustered to the right compared with those of other taxa (Fig. 3A). For PC2, all measurements had low loadings except for the braincase height (BH) (Table 1). Therefore, based on PC2, *K. picta* and *K. titania*, which had a larger BH, were clustered to the top of the plot, whereas *K. dongduongana* and *K. kachinensis* were assigned to the bottom (Fig. 3). For DAPC, we entered the first two PCs from the PCA results and obtained two discriminant functions (DFs) to distinguish among studied *Kerivoula* specimens. PCA and DAPC scatter plots showed that *K. depressa* and *K. dongduongana* specimens formed distinct and separated clusters (pale pink and green triangles in Fig. 3), although some scatter values overlapped with those from *K. furva*. Meanwhile, *K. picta*, *K. titania*, and *K. kachinensis* clustered into three distinguishable groups (Fig. 3).

Phylogenetic relationships of *Kerivoula*

Bayesian and ML trees using *COI*, *Cytb*, and *RAG2* matrices highly supported monophyly of the genus *Kerivoula* (Fig. 4) [posterior probabilities (PP)/bootstrap values (BS), 1/99 for *COI* and *Cytb*, 1/97 for *RAG2*] and revealed a similar well-supported topology. *Kerivoula picta* and *K. papillosa* occurred outside of the clade uniting all other examined species within *Kerivoula*. All inferences clustered our sequences with *K. depressa* (PP = 1 in *COI/Cytb*; BS: 99 for *COI*, 100 for *Cytb*) and *K. dongduongana* (PP = 1 in *COI/Cytb/RAG2*; BS: 99 for *COI*, 100 for *Cytb*, 97 for *RAG2*), thus verifying our morphological species identification results (Fig. 4). However, interspecific relationships of *K. depressa*, *K. dongduongana*, *K. kachinensis*, *K. furva*, and *K. hardwickii* (*sensu stricto*) remain ambiguous and could not be resolved herein.

Discussion

The major interspecific phylogenetic relationships of our analyses are comparable with those reported by Kuo et al. (2017) and Tu et al. (2018). Our studies similarly confirmed the monophyly of *K. depressa* and *K. dongduongana*. However, the topology of the phylogenetic tree based on the *RAG2* gene remains unresolved and needs further study. Finally, combining the results of external and craniodental examination and multivariate analyses, 24 specimens were determined as *K. depressa* and *K. dongduongana* (Table 1).

Our discovery of *K. depressa* and *K. dongduongana* in China indicates that six species of *Kerivoula* live in China. According to morphological analyses, *K. picta* is easily distinguished by its unique pelage color pattern and skull shape (Wilson and Mittermeier 2019), whereas *K. kachinensis* is the largest species with a distinctly flattened skull (Bates et al. 2004; Tu et al. 2018; Yu et al. 2022). As for the remaining four similar-sized species, *K. titania* has a distinctly longer tibia and higher braincase than the others (Kuo et al. 2017; Tu et al. 2018). In pelage coloration, *K. furva* has the darkest fur color, varying from black brown to black gray, whereas *K. depressa* and *K. dongduongana* are pale brown.

Among the four species, *K. dongduongana* has the shortest BH (Kuo et al. 2017; Tu et al. 2018). A key to the *Kerivoula* species occurring in China is provided in Appendix 2.

Until recently, *Kerivoula* species were considered forest-dependent (Wilson and Mittermeier 2019). They are known in the south of China across Yunnan to Taiwan, and from Hainan to Chongqing. It is worth noting that five of the “Chinese” *Kerivoula* species are found in the southwest region of Yunnan Province, which is often treated as a biodiversity hotspot near Myanmar, Laos, and Vietnam. Its unique terrain, vegetation, and environmental conditions, including a low latitude, warm and tropical forest, and humid micro-climate, appear suitable for inhabitation and colonization (Kruskop 2013; Wilson and Mittermeier 2019; Qian et al. 2020). The high diversity of woolly bats in tropical forest areas may indicate the origin in their diversification progress.

Based on a comparison of the recorded *Kerivoula* diversity from the bordering countries of Myanmar (five species), Laos (seven species), and Vietnam (eight species) (Wilson and Mittermeier 2019), we suggest that there is still a risk of underestimating the diversity of *Kerivoula* in China. More surveys should therefore be conducted, especially on the border/unexplored region and using effective sampling tools such as multi-bank harp traps.

Acknowledgements

The authors thank National Nature Reserve Administration of Xishuangbanna Dai Autonomous Prefecture for their guides and help during the field survey. We also thank Ruichang Quan from Xishuangbanna Tropical Botanical Garden, Chinese Academy of Sciences, for the assistance in the field survey. This study was financially supported by the National Sciences Foundation of China (31970394, 32192421), The Special Foundation for National Science and Technology Basic Research Program of China [2021FY100303].

References

- Bates PJJ, Harrison DL (1997) Bats of the Indian Subcontinent. Harrison Zoological Museum, Sevenoaks, Kent, 212–215.
- Bates PJJ, Struebig MJ, Rossiter SJ, Kingston T, Oo SSL, Mya KM (2004) A New Species of *Kerivoula* (Chiroptera: Vespertilionidae) from Myanmar (Burma). *Acta Chiropterologica* 6(2): 219–226. <https://doi.org/10.3161/001.006.0203>
- Bates PJJ, Struebig MJ, Hayes BD, Furey NM, Mya KM, Thong VD, Tien PD, Son NT, Harrison DL, Francis CM, Csorba G (2007) A new species of *Kerivoula* (Chiroptera: Vespertilionidae) from Southeast Asia. *Acta Chiropterologica* 9(2): 323–337. [https://doi.org/10.3161/1733-5329\(2007\)9\[323:ANSOKC\]2.0.CO;2](https://doi.org/10.3161/1733-5329(2007)9[323:ANSOKC]2.0.CO;2)
- Brunet-Rossinni AK, Wilkinson GS (2009) Methods for age estimation and the study of senescence in bats. In: Kunz TH, Parsons S (Eds) *Ecological & behavioral methods for the study of bats*. The Johns Hopkins University Press, Baltimore, 315–325.
- Corbet GB, Hill JE (1992) *The Mammals of Indomalayan Region: a Systematic Review*. Oxford University Press, Oxford, 488 pp.

- Douangboubpha B, Bumrungsri S, Satasook C, Wanna W, Soisook P, Bates PJJ (2016) Morphology, genetics and echolocation calls of the genus *Kerivoula* (Chiroptera: Vespertilionidae: Kerivoulinae) in Thailand. *Mammalia* 80(1): 21–47. <https://doi.org/10.1515/mammalia-2014-0004>
- Dray S, Dufour AB, Chessel D (2007) The ade4 Package– II. Two-table and K-table Methods. 7: 6–6.
- Edgar RC (2004) MUSCLE: Multiple sequence alignment with high accuracy and high throughput. *Nucleic Acids Research* 32(5): 1792–1797. <https://doi.org/10.1093/nar/gkh340>
- Ellerman JR, Morrison-Scott TC (1951) Check list of Palaearctic and Indian mammals 1758 to 1946. British Museum, Natural History, London, 810 pp.
- Francis CM, Kingston T, Zubaid A (2007) A new species of *Kerivoula* (Chiroptera: Vespertilionidae) from peninsular Malaysia. *Acta Chiropterologica* 9(1): 1–12. [https://doi.org/10.3161/1733-5329\(2007\)9\[1:ANSOKC\]2.0.CO;2](https://doi.org/10.3161/1733-5329(2007)9[1:ANSOKC]2.0.CO;2)
- Francis CM, Borisenko AV, Ivanova NV, Eger JL, Lim BK, Guillén-Servent A, Kruskop SV, Mackie I, Hebert PDN (2010) The Role of DNA Barcodes in Understanding and Conservation of Mammal Diversity in Southeast Asia. *PLoS ONE* 5: e12575. <https://doi.org/10.1371/journal.pone.0012575>
- Gray JE (1842) Descriptions of some new Genera and fifty unrecorded Species of Mammalia. *Annals & Magazine of Natural History* 10(65): 255–267. <https://doi.org/10.1080/03745484209445232>
- Hill JE (1965) Asiatic bats of the genera *Kerivoula* and *Phoniscus* (Vespertilionidae), with a note on *Kerivoula aerea* Tomes. *Mammalia* 29(4): 524–556. <https://doi.org/10.1515/mamm.1965.29.4.524>
- Hoofer SR, Reeder SA, Hansen EW, Van Den Bussche RA (2003) Molecular phylogenetics and taxonomic review of Noctilionoid and Vespertilionoid bats (Chiroptera: Yangochiroptera). *Journal of Mammalogy* 84(3): 809–821. <https://doi.org/10.1644/BWG-034>
- Horsfield T (1824) Zoological researches in Java, and the neighbouring islands. Printed for Kingsbury, Parbury and Allen, London. <https://doi.org/10.5962/bhl.title.44848>
- Jombart T (2008) adegenet: A R package for the multivariate analysis of genetic markers. *Bioinformatics* 24(11): 1403–1405. <https://doi.org/10.1093/bioinformatics/btn129>
- Kalyaanamoorthy S, Minh BQ, Wong TKF, von Haeseler A, Jermiin LS (2017) ModelFinder: Fast model selection for accurate phylogenetic estimates. *Nature Methods* 14(6): 587–589. <https://doi.org/10.1038/nmeth.4285>
- Khan FAA, Solari S, Swier VJ, Larsen PA, Abdullah MT, Baker RJ (2010) Systematics of Malaysian woolly bats (Vespertilionidae: *Kerivoula*) inferred from mitochondrial, nuclear, karyotypic, and morphological data. *Journal of Mammalogy* 91(5): 1058–1072. <https://doi.org/10.1644/09-MAMM-A-361.1>
- Kruskop SV (2013) Bats of Vietnam: Checklist and an Identification Manual (2nd edn.). *Tovarišchestvo nauchnykh izdanii KMK, Moscow*, 154–160.
- Kuo HC, Chen SF, Fang YP, Flanders J, Rossiter SJ (2014) Comparative range wide phylogeography of four endemic Taiwanese bat species. *Molecular Ecology* 23(14): 3566–3586. <https://doi.org/10.1111/mec.12838>
- Kuo HC, Soisook P, Ho YY, Csorba G, Wang CN, Rossiter SJ (2017) A Taxonomic Revision of the *Kerivoula hardwickii* Complex (Chiroptera: Vespertilionidae) with the Description

- of a New Species. *Acta Chiropterologica* 19(1): 19–19. <https://doi.org/10.3161/15081109ACC2017.19.1.002>
- Le S, Josse J, Husson F (2008) FactoMineR: An R Package for Multivariate Analysis. *Journal of Statistical Software* 25(1): 1–18. <https://doi.org/10.18637/jss.v025.i01>
- Minh BQ, Schmidt HA, Chernomor O, Schrempf D, Woodhams MD, von Haeseler A, Lanfear R (2020) IQ-TREE 2: New Models and Efficient Methods for Phylogenetic Inference in the Genomic Era. *Molecular Biology and Evolution* 37: 1530–1534. <https://doi.org/10.1093/molbev/msaa015>
- Nguyen ST, Motokawa M, Oshida T, Endo H (2016) A morphological analysis of the skull size and shape of Kerivoulinae (Chiroptera: Vespertilionidae) from Vietnam. *The Journal of Veterinary Medical Science* 78(2): 187–198. <https://doi.org/10.1292/jvms.15-0270>
- Nylander JAA (2004) MrModeltest v2. Program distributed by the author. Evolutionary Biology Centre, Uppsala University.
- Qian LS, Chen JH, Deng T, Sun H (2020) Plant diversity in Yunnan: Current status and future directions. *Plant Diversity* 42(4): 281–291. <https://doi.org/10.1016/j.pld.2020.07.006>
- R Core Team (2013) R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna. <http://www.R-project.org/>
- Revelle W (2013) Psych: Procedures for psychological, psychometric, and personality Research, 1. R Package Version 1.0–95.
- Ronquist F, Teslenko M, van der Mark P, Ayres DL, Darling A, Höhna S, Larget B, Liu L, Suchard MA, Huelsenbeck JP (2012) MrBayes 3.2: Efficient Bayesian phylogenetic inference and model choice across a large model space. *Systematic Biology* 61(3): 539–542. <https://doi.org/10.1093/sysbio/sys029>
- Ruedi M, Biswas J, Csorba G (2012) Bats from the wet: two new species of tube-nosed bats (Chiroptera: Vespertilionidae) from Meghalaya, India. *Revue Suisse de Zoologie* 119(1): 111–135. <https://doi.org/10.5962/bhl.part.150145>
- Sikes RS (2016) the Animal Care Use Committee of the American Society of Mammalogists Guidelines of the American Society of Mammalogists for the use of wild mammals in research and education. *Journal of Mammalogy* 97: 663–688. <https://doi.org/10.1093/jmammal/gyw078>
- Simmons NB (2005) Order Chiroptera. In: Wilson DE, Reeder DM (Eds) *Mammal Species of the World: a Taxonomic and Geographic Reference* (3rd ed.). Johns Hopkins University Press, London, 312–529.
- Sinha YR (1999) Records of the zoological survey of India: contribution to the knowledge of bats (Mammalia: Chiroptera) from North East Hills, India. *Zoological Survey of India, Calcutta*, 52 pp.
- Stadelmann B, Jacobs DS, Schoeman C, Ruedi M (2004) Phylogeny of African *Myotis* Bats (Chiroptera, Vespertilionidae) Inferred from Cytochrome b Sequences. *Acta Chiropterologica* 6(2): 177–192. <https://doi.org/10.3161/001.006.0201>
- Tu VT, Hassanin A, Furey NM, Son NT, Csorba G (2018) Four species in one: multigene analyses reveal phylogenetic patterns within Hardwicke's woolly bat, *Kerivoula hardwickii*-complex (Chiroptera, Vespertilionidae) in Asia. *Hystrix, the Italian Journal of Mammalogy* 29(1): 111–121. <https://doi.org/10.4404/hystrix-00017-2017>
- Wickham H (2016) ggplot2: Elegant Graphics for Data Analysis. Springer-Verlag New York. ISBN 978-3-319-24277-4. <https://ggplot2.tidyverse.org>

Wilson DE, Mittermeier RA (2019) Handbook of the Mammals of the World (Vol. 9). Bats. Lynx Edicions, Barcelona, 894–900.

Wu Y, Li YC, Lin LK, Harada M, Chen Z, Motokawa M (2012) New Records of *Kerivoula titania* (Chiroptera: Vespertilionidae) from Hainan Island and Taiwan. *Mammal Study* 37(1): 69–72. <https://doi.org/10.3106/041.037.0109>

Yu WH, Li F, Csorba G, Xu ZX, Wang XY, Guo WJ, Li YC, Wu Y (2018) A revision of *Kerivoula hardwickii* and occurrence of *K. furva* (Chiroptera: Vespertilionidae) in China. *Zootaxa* 4461(1): 45–45. <https://doi.org/10.11646/zootaxa.4461.1.2>

Yu W, Lin C, Huang Z, Liu S, Wang Q, Quan R, Li S, Wu Y (2022) Discovery of *Kerivoula kachinensis* and a validity of *K. titania* (Chiroptera: Vespertilionidae) in China. *Mammalia* 86(3): 303–308. <https://doi.org/10.1515/mammalia-2021-0102>

Appendix I

Table A1. Sample used in molecular analyses, with GenBank accession numbers for *COI*, *Cytb* and *RAG2* genes provided. Newly generated sequences in this study are shown in bold.

Taxon	Voucher	GenBank accession numbers			Location	Reference
		<i>COI</i>	<i>Cytb</i>	<i>RAG2</i>		
<i>Kerivoula depressa</i>	GZHU 18213	OM716942	OM735703	OM735730	Yunnan, China	This study
	GZHU 18246	–	OM735713	OM735736	Yunnan, China	This study
	GZHU 19082	OM716943	OM735707	OM735727	Yunnan, China	This study
	GZHU 19083	OM716948	OM735704	OM735729	Yunnan, China	This study
	GZHU 19122	OM716946	OM735712	OM735731	Yunnan, China	This study
	GZHU 19133	OM716947	OM735714	OM735734	Yunnan, China	This study
	GZHU 19173	OM716944	OM735708	OM735732	Yunnan, China	This study
	GZHU 19174	OM716949	OM735710	OM735725	Yunnan, China	This study
	GZHU 19202	OM716950	OM735709	OM735728	Yunnan, China	This study
	GZHU 19218	OM716952	OM735705	OM735735	Yunnan, China	This study
	GZHU 19222	OM716951	OM735706	OM735733	Yunnan, China	This study
	GZHU 19325	OM716945	OM735711	OM735726	Yunnan, China	This study
<i>Kerivoula dongduongana</i>	GZHU 19134	OM716941	OM735697	OM735715	Yunnan, China	This study
	GZHU 19150	OM716940	OM735696	OM735720	Yunnan, China	This study
	GZHU 19197	OM716936	OM735691	OM735717	Yunnan, China	This study
	GZHU 19198	OM716937	OM735692	OM735721	Yunnan, China	This study
	GZHU 19200	OM716938	OM735693	OM735722	Yunnan, China	This study
	GZHU 19201	OM716939	OM735694	OM735724	Yunnan, China	This study
	GZHU 19301	OM716930	OM735698	OM735718	Yunnan, China	This study
	GZHU 19302	OM716931	OM735699	OM735719	Yunnan, China	This study
	GZHU 19305	OM716932	OM735700	–	Yunnan, China	This study
	GZHU 19306	OM716933	OM735701	–	Yunnan, China	This study
	GZHU 19307	OM716935	OM735695	OM735723	Yunnan, China	This study
	GZHU 19308	OM716934	OM735702	OM735716	Yunnan, China	This study
<i>Murina cyclotis</i>	VN11-1199	KF772775	MH137367	MH137498	Viet Nam	Tu et al. 2018
<i>Harpioccephalus harpia</i>	VN11-1288	MH137300	MH137366	MH137497	Viet Nam	Tu et al. 2018
<i>Myotis muricola</i>	VN11-1186	MH137299	MH137365	MH137496	Viet Nam	Tu et al. 2018
<i>Kerivoula picta</i>	VN11-1565	MH137303	–	–	Cambodia	Tu et al. 2018
	VN11-1576	MH137304	MH137370	MH137501	Cambodia	Tu et al. 2018
	VN11-1577	MH137305	MH137371	MH137502	Cambodia	Tu et al. 2018
<i>Kerivoula papillosa</i>	20467	MH137301	MH137368	MH137499	Thailand	Tu et al. 2018
	21719	MH137302	MH137369	MH137500	Cambodia	Tu et al. 2018

Taxon	Voucher	GenBank accession numbers			Location	Reference
		<i>COI</i>	<i>Cytb</i>	<i>RAG2</i>		
<i>K. dongduongana</i>	ROM 110828	JF443952	–	–	Vietnam	Tu et al. 2018
	CPV10-295	MH137318	–	–	Cambodia	Tu et al. 2018
	CPV10-297	MH137319	MH137382	MH137510	Cambodia	Tu et al. 2018
<i>K. dongduongana</i>	ROM MAM 111278	HM540688	–	–	Viet Nam	Francis et al. 2010
	VN11-1158	MH137320	MH137383	MH137511	Vietnam	Tu et al. 2018
	VN11-1178	MH137321	MH137384	MH137512	Vietnam	Tu et al. 2018
	ROM MAM 111298	HM540687	–	–	Viet Nam	Francis et al. 2010
	ROM MAM 111277	HM540704	–	–	Viet Nam	Francis et al. 2010
	CPV10-292	MH137317	–	–	Cambodia	Tu et al. 2018
	ROM MAM 110605	HM540690	–	–	Laos	Francis et al. 2010
	ROM MAM 110604	HM540705	–	–	Laos	Francis et al. 2010
	23028	MH137315	–	–	Viet Nam	Tu et al. 2018
	23036	MH137316	MH137381	MH137509	Viet Nam	Tu et al. 2018
	PSUZYC-MM2011.48	KY034089	–	–	Thailand	Soisook et al. 2016
<i>Kerivoula titania</i>	PSUZYC-MM2011.15	KY034108	–	–	Thailand	Soisook et al. 2016
	VN11-0945	MH137361	–	–	Viet Nam	Tu et al. 2018
	VN11-0013	MH137349	–	–	Viet Nam	Tu et al. 2018
	23034	MH137340	MH137396	–	Viet Nam	Tu et al. 2018
	VN11-0010	MH137346	–	–	Viet Nam	Tu et al. 2018
	VN11-0025	MH137352	–	–	Viet Nam	Tu et al. 2018
	VN11-0026	MH137353	–	–	Viet Nam	Tu et al. 2018
	VN11-0027	MH137354	–	–	Viet Nam	Tu et al. 2018
	VN11-0030	MH137355	–	–	Viet Nam	Tu et al. 2018
	VN11-0031	MH137356	–	–	Viet Nam	Tu et al. 2018
	VN11-0944	MH137360	–	–	Viet Nam	Tu et al. 2018
<i>K. titania</i>	VN11-1832	MH137364	MH137402	–	Viet Nam	Tu et al. 2018
	CPV10-415	MH137343	–	–	Cambodia	Tu et al. 2018
	CPV10-363	MH137341	MH137398	MH137520	Cambodia	Tu et al. 2018
	CPV10-399	MH137342	MH137399	MH137521	Cambodia	Tu et al. 2018
	VN11-1188	MH137362	MH137401	MH137523	Viet Nam	Tu et al. 2018
	VN11-1193	MH137363	–	–	Viet Nam	Tu et al. 2018
	21942	MH137339	MH137395	MH137519	Viet Nam	Tu et al. 2018
	VN11-0002	MH137344	–	–	Viet Nam	Tu et al. 2018
	VN11-0009	MH137345	–	–	Viet Nam	Tu et al. 2018
	VN11-0011	MH137347	–	–	Viet Nam	Tu et al. 2018
	VN11-0012	MH137348	MH137400	MH137522	Viet Nam	Tu et al. 2018
	VN11-0014	MH137350	–	–	Viet Nam	Tu et al. 2018
	VN11-0018	MH137351	–	–	Viet Nam	Tu et al. 2018
	VN11-0035	MH137357	–	–	Viet Nam	Tu et al. 2018
	VN11-0040	MH137358	–	–	Viet Nam	Tu et al. 2018
	VN11-0044	MH137359	–	–	Viet Nam	Tu et al. 2018
	23041	–	MH137397	–	Vietnam	Tu et al. 2018
	21746	MH137306	MH137373	MH137503	Indonesia	Tu et al. 2018
<i>Kerivoula hardwickii</i> s. str.	VN11-1623	MH137309	MH137376	MH137505	Cambodia	Tu et al. 2018
	VN11-1593	MH137307	MH137374	MH137504	Cambodia	Tu et al. 2018
	VN11-1622	MH137308	MH137375	–	Cambodia	Tu et al. 2018
	17348	–	MH137372	–	Indonesia	Tu et al. 2018
<i>K. depressa</i>	ROM MAM 118429	HM540670	–	–	Laos	Francis et al. 2010
	AGS980322-65	HM540664	–	–	Laos	Francis et al. 2010
	ROM MAM 118186	HM540671	–	–	Laos	Francis et al. 2010
	ROM MAM 118026	HM540672	–	–	Laos	Francis et al. 2010
	ROM MAM 107721	HM540669	–	–	Viet Nam	Francis et al. 2010
	VN11-1554	MH137313	MH137379	MH137506	Cambodia	Tu et al. 2018
	VN11-1835	MH137314	MH137380	–	Vietnam	Tu et al. 2018
	ROM MAM 117982	HM540673	–	–	Laos	Francis et al. 2010
	ROM MAM 110602	HM540667	–	–	Laos	Francis et al. 2010
	ROM MAM 117971	HM540668	–	–	Laos	Francis et al. 2010
	ROM MAM 110585	HM540674	–	–	Laos	Francis et al. 2010
	CPV10-291	MH137310	MH137377	MH137507	Cambodia	Tu et al. 2018

Taxon	Voucher	GenBank accession numbers			Location	Reference
		COI	Cytb	RAG2		
<i>K. depressa</i>	CPV10-293	MH137311	–	–	Cambodia	Tu et al. 2018
	CPV10-409	MH137312	MH137378	MH137508	Cambodia	Tu et al. 2018
<i>Kerivoula kachinensis</i>	MAM 107718	HM540736	–	–	Viet Nam	Francis et al. 2010
	VN11-1831	MH137338	MH137394	–	Viet Nam	Tu et al. 2018
	VN11-0940	MH137337	MH137393	–	Viet Nam	Tu et al. 2018
	CPV10-416	MH137336	MH137392	MH137518	Cambodia	Tu et al. 2018
	ZMMU S-184667	GU684767	–	–	Viet Nam	Tu et al. 2018
	EDB 25747	HM540734	–	–	Laos	Francis et al. 2010
<i>Kerivoula furva</i>	16481	MH137335	MH137391	MH137513	Nepal	Tu et al. 2018
	VN11-0050	MH137330	–	–	Viet Nam	Tu et al. 2018
	VN11-0004	MH137324	MH137387	MH137515	Viet Nam	Tu et al. 2018
	VN11-0045	MH137325	–	–	Viet Nam	Tu et al. 2018
	VN11-0046	MH137326	–	–	Viet Nam	Tu et al. 2018
	VN11-0047	MH137327	–	–	Viet Nam	Tu et al. 2018
	VN11-0048	MH137328	–	–	Viet Nam	Tu et al. 2018
	VN11-0049	MH137329	–	–	Viet Nam	Tu et al. 2018
	VN11-0943	MH137333	MH137389	MH137516	Viet Nam	Tu et al. 2018
	VN11-0937	MH137331	MH137388	–	Viet Nam	Tu et al. 2018
	VN11-0942	MH137332	–	–	Viet Nam	Tu et al. 2018
	23024	MH137322	–	–	Viet Nam	Tu et al. 2018
	23025	MH137323	MH137385	MH137514	Viet Nam	Tu et al. 2018
	VN11-1361	MH137334	MH137390	MH137517	Viet Nam	Tu et al. 2018
	2005-670	–	MH137386	–	Nepal	Tu et al. 2018

Appendix 2

A key to the *Kerivoula* species occurring in China (both in English and Chinese)

- 1 Pelage color relatively bright. Fur orange. Wing membranes dark brown with red-dish brown markings / 体色鲜艳,体毛橙黄色;翼膜深棕色,具红褐色斑块
..... *Kerivoula picta* / 彩蝠
- Pelage color duller. Fur dark gray, brown, or blackish gray / 体色较暗淡,体毛暗灰色、棕色或黑灰色 2
- 2 Size larger, forearm length more than 40 mm. Greatest length of skull more than 16 mm, greatest breadth of braincase more than 8 mm / 体型较大,前臂长超过 40 mm; 颅全长大于 16 mm, 脑颅宽大于 8 mm
..... *Kerivoula kachinensis* / 克钦彩蝠
- Size smaller, forearm length less than 40 mm. Greatest length of skull less than 16 mm, greatest breadth of braincase less than 8 mm / 体型较小,前臂长不及 40 mm; 颅全长小于 16 mm, 脑颅宽小于 8 mm 3
- 3 Tibia length usually more than 18.5 mm. Greatest length of skull 14.5–16.0 mm, braincase height usually more than 5.3 mm / 胫骨长常大于 18.5 mm; 颅全长 14.5–16 mm, 颅高常大于 5.3 mm..... *Kerivoula titania* / 泰坦尼亚彩蝠
- Tibia length less than or equal to 18.5 mm. Greatest length of skull less than 14.5 mm, braincase height usually less than 5.3 mm / 胫骨长小于或等于 18.5 mm; 颅全长小于 14.5 mm, 颅高不及 5.3 mm..... 4

- 4 Greatest width across the outer edges of the third upper molars more than 5.3 mm, greatest length of mandible more than 9.8 mm. Dorsal pelage dark brown to blackish gray, broadly uniform in colour from bases to tips / 第三上臼齿宽大于5.3 mm, 下颌长大于9.8 mm; 背毛深棕色到黑灰色, 毛基到毛尖颜色大致相同 *Kerivoula furva* / 暗褐彩蝠
- Greatest width across the outer edges of the third upper molars less than 5.3 mm, greatest length of mandible less than 9.8 mm. Dorsal pelage buff brown to dark brown, its base and tip significantly different / 第三上臼齿宽小于5.3 mm, 下颌长小于9.8 mm; 背毛浅棕色到深棕色, 毛基与毛尖颜色明显不同 5
- 5 Braincase slightly flattened, braincase height more than 4.7 mm, more than 1/3 of greatest length of skull. Dorsal pelage buff brown to dark brown, ventral pelage dark brown / 脑颅略为扁平, 颅高大于4.7 mm, 超过颅全长1/3; 背毛浅棕色到深棕色, 腹毛深棕色 *Kerivoula depressa* / 平颅彩蝠
- Braincase flatter, braincase height usually less than 4.7 mm, less than 1/3 of greatest length of skull. Dorsal pelage dark brown to golden brown, ventral pelage golden brown / 脑颅更扁平, 颅高多小于4.7 mm, 不及颅全长1/3; 背毛深棕色到金棕色, 腹毛金棕色 *Kerivoula dongduongana* / 印支彩蝠