

An unusual species of *Pseudophanias* Raffray from West Tianmu Mountain, China (Coleoptera, Staphylinidae, Pselaphinae)

Zi-Wei Yin¹, Tie-Xiong Zhao²

¹ Laboratory of Systematic Entomology, College of Life Sciences, Shanghai Normal University, Shanghai 200234, China ² 44 East Genta Road, Zhuji, Shaoxing, Zhejiang 311800, China

Corresponding author: Zi-Wei Yin (pselaphinae@gmail.com)

Academic editor: Adam Brunke | Received 23 September 2022 | Accepted 25 October 2022 | Published 10 November 2022

<https://zoobank.org/B19E8A19-D8A0-4CA3-8620-D8532B447D17>

Citation: Yin Z-W, Zhao T-X (2022) An unusual species of *Pseudophanias* Raffray from West Tianmu Mountain, China (Coleoptera, Staphylinidae, Pselaphinae). ZooKeys 1129: 37–44. <https://doi.org/10.3897/zookeys.1129.95245>

Abstract

The genus *Pseudophanias* Raffray includes 19 species distributed in the Oriental Region. Here, the first species from continental China, *P. tianmuensis* Yin & Zhao **sp. nov.**, is described and illustrated from West Tianmu Mountain. This species can be readily separated from all congeners by the coarse vestiture, along with its male features. A previous key to *Pseudophanias* species of East and South Asia is modified to accommodate the new species.

Keywords

East China, new species, *Pseudophanias*, taxonomy, Zhejiang

Introduction

With greatly reduced maxillary palpi and often a C-shaped aedeagus, the genus *Pseudophanias* Raffray is an easily recognizable group among the Oriental genera of the ant-loving beetle tribe Tmesiphorini (Staphylinidae: Pselaphinae). After removing two species from former *Chandleriella* Hlaváč (now a junior synonym of *Pseudophanias*; Hlaváč 2000; Yin 2019), the genus currently comprises 19 species distributed in the

Oriental Region (Inoue et al. 2020; Inoue and Nomura 2020, 2021). Members are usually taken from the forest leaf-litter layer, in decomposing wood, and sometimes also with ants or termites. One species from Nepal was collected in a cave and possesses an exceptionally elongate body and appendages that are typical for a cavernicolous beetle (Yin et al. 2015).

In 2006 an unidentified pselaphine collected by our friend Xiao-Bin Song from a decomposing log on West Tianmu Mountain was made available for study. At first look, the beetle hardly recalled a *Pseudophanias* because of its unusual coarsely punctate body surface, subquadrate elytra, and cylindrical antennal clubs which are unknown among all previously described members of the genus. In the following 12 years, two more specimens were collected in the same area; one was sifted from leaf litter, and the other was, again, found in dead wood with termites. This species is confirmed to belong to *Pseudophanias* and is formally described here.

Materials and methods

The type material of the new species is deposited in the Insect Collection of Shanghai Normal University (SNUC), Shanghai, China. The label data of the material are quoted verbatim.

Dissected parts were mounted in Euparal on plastic slides pinned with the specimen. The habitus image of the beetle was photographed using a Canon 5D Mark III camera with Canon MP-E 65 mm f/2.8 1–5× macro lens, with a Canon MT-24EX Macro Twin Lite Flash as the light source. Images of the morphological details were produced using a Canon G9 camera mounted to an Olympus CX31 microscope under reflected or transmitted light. Zerene Stacker (v. 1.04) was used for image stacking. All images were modified and grouped into plates using Adobe Photoshop CC 2020.

Measurements were taken as follows: total body length was measured from the anterior margin of the clypeus to the apex of the abdomen; head length was measured from the anterior margin of the clypeus to the head base, excluding the occipital constriction; head width was measured across the eyes; the length of the pronotum was measured along the midline, the width of the pronotum equals pronotum's maximum width; the length of the elytra was measured along the suture; the width of the elytra was measured as the maximum width across both elytra; the length of the abdomen is the length of the dorsally exposed part of the abdomen along its midline, the width is abdomen's maximum width. Since one male paratype is completely disarticulated, only the holotype and the female paratype were measured. Abdominal tergites and sternites are numbered following Chandler (2001), in Arabic (starting from the first visible segment) and Roman (reflecting true morphological position) numerals, e.g., tergite 1 (IV), or sternite 1 (III). Paired appendages in the description of the new species are treated as singular.

Taxonomy

Pseudophanias tianmuensis Yin & Zhao, sp. nov.

<https://zoobank.org/4476E74C-644F-4667-8AF5-45C75D21638A>

Figs 1–3

Chinese common name: 天目隐须蚁甲

Type material. (3 exx.). **Holotype:** CHINA: ♂, ‘China: Zhejiang, Linan District, West Tianmu Mountain, 1100 m, 23.viii.2014, *Cryptomeria* leaf, sift, Tie-Xiong Zhao leg., 浙江省临安市西天目山(此路不通)’ (in SNUC). **Paratypes:** CHINA: 1 ♂ [completely disarticulated specimen], China: Zhejiang, Linan District, West Tianmu Mountain, 1150 m, 2.vii.2006, decomposing log, sift, Xiao-Bin Song leg., 浙江省临安市西天目山(停车场); 1 ♀, ‘China: Zhejiang, Linan District, West Tianmu Mountain, 1180 m, 18.viii.2019, decomposing log, sift, Zi-Wei Yin leg., 浙江省临安市西天目山(消防道)’ (both in SNUC).

Diagnosis. Body length 2.8–2.9 mm; dorsal vestiture coarsely punctate and densely setose. Head subrounded at base, with small vertexal and frontal foveae; antennomeres moniliform. Elytra with discal and lateral longitudinal ridges, areas between ridges impressed. Tergite 1 (IV) longer than 2 (V) at middle but clearly shorter than 2 and 3 (VI) combined. **Male:** antennal clubs formed by apical four enlarged, cylindrical antennomeres. Aedeagus symmetric dorso-ventrally; median lobe C-shaped laterally; parameres thin and elongate, each with two long apical and one preapical seta.

Description. Male. Body (Fig. 1A) length 2.91 mm; color red-brown, tarsi and mouthparts lighter. Dorsal vestiture coarsely punctate and covered with short, dense setae.

Head (Fig. 2A) subrounded at base, longer than wide, length 0.63 mm, width 0.49 mm; vertex weakly convex, with two small, asetose vertexal foveae (dorsal tentorial pits); antennal tubercles indistinct; frons with single small fovea, protruding anteriorly to form rostrum; clypeus sharply descending, its surface coarse, anterior margin rounded and slightly carinate; lacking ocular-mandibular carina. Venter with two small, well-separated gular foveae (posterior tentorial pits), lacking median carina/sulcus. Maxillary palpus (Fig. 2B) small in size, palpomere 1 minute, 2 pedunculate basally and broadened apically, 3 roundly trapezoidal, 4 broadly at approximately basal 1/3 and gradually narrowing apically. Compound eyes moderately prominent, reniform in lateral view, each composed of approximately 55 ommatidia. Antenna elongate, length 1.75 mm, distinct club (Fig. 2C) formed by greatly enlarged antennomeres 8–11 and approximately as long as antennomeres 1–7 combined (0.88 mm *vs* 0.87 mm); antennomere 1 large, subcylindrical, 2–7 moniliform, of similar shape, 8, 9 and 10 each much larger than 7, subquadrate, 11 longest, slightly narrower than 10 and as long as 9 and 10 combined.

Pronotum (Fig. 2A) approximately as long as wide, length 0.66 mm, width 0.63 mm, widest anterior middle; sides from widest point greatly convergent anteriorly




Figure 1. Habitus of *Pseudophanias tianmuensis* sp. nov. **A** male **B** female. Scale bars: 1.0 mm.

and subparallel posteriorly; disc distinctly convex to form hump at middle, lacking carina or sulcus; with small asetose median and lateral antebasal foveae; lacking basolateral foveae. Prosternum with anterior part much shorter than coxal part at middle, with small, asetose lateral procoxal foveae; hypomeron lacking ridge that demarcating it from pronotum; margin of coxal cavity non-carinate.

Elytra much wider than long, length 0.96 mm, width 1.24 mm, truncate at bases; each elytron (Fig. 2D) with two moderately large, asetose basal foveae; with one discal and one lateral ridge, discal stria formed by broad impressed area between ridges; humeral region lacking subhumeral fovea nor marginal stria; epipleural area broad; posterolateral margin with broad notch. Metathoracic wings fully developed.

Mesoventrite (Fig. 2E) with rough surface, with pair of thick admesal carinae, lateral area fully demarcated from metaventrite, openings of median mesoventral



Figure 2. Morphology of *Pseudophanias tianmuensis* sp. nov., male **A** head and pronotum **B** maxillary palpus **C** antennal club **D** elytron **E** meso- and metaventrите **F** Mesotarsal claws. Scale bars: 0.5 mm (**A**, **D**, **E**); 0.1 mm (**B**, **F**); 0.3 mm (**C**).

foveae widely separated, foveae greatly extended internally, lateral mesoventral foveae unforked, extended internally and overlapped with median foveae; intercoxal process short. Metaventrите (Fig. 2E) coarsely punctate, greatly prominent medially and with



Figure 3. Morphology and habitat of *Pseudophanias tianmuensis* sp. nov. **A–C** aedeagus **A** dorsal **B** lateral **C** ventral **D** paramere, lateral **E** general environment of the collecting site **F** decomposing tree branches inhabited by the beetle. Scale bars: 0.2 mm (**A–C**); 0.1 mm (**D**).

small crest; with large, asetose lateral mesocoxal foveae; posterior margin roundly emarginate at middle.

Legs elongate; femora coarsely punctate; each tarsus (Fig. 2F) with one major and one reduced pretarsal claw.

Abdomen widest at lateral margins of tergite 1 (IV), length 0.86 mm, width 1.15 mm. Tergite 1 (IV) slightly less than twice as long as 2 (V), with broad, deep basal sulcus and two basolateral foveae at lateral ends of sulcus, 2 and 3 (VI) successively shorter, lacking basal sulcus nor fovea, 4 (VII), much longer than 3, narrowed posteriorly, with one pair of basolateral foveae; tergites 1–3 each with pair of quadrangular

and 4 with pair of triangular paratergites, 5 (VIII) broadly rounded, with one pair of small basolateral foveae, posterior margin roundly emarginate at middle. Sternite 2 (IV) longest, with densely setose basal sulcus and pair of basolateral foveae at lateral ends of sulcus, 3 (V) and 4 (VI) each short at middle, combined approximately as long as 5 (VII), 3–5 lacking sulcus nor fovea at base, 6 (VIII) transverse, posterior margin slightly protruding at middle, 7 (IX) composed of pair of membranous lamellae.

Aedeagus (Fig. 3A–C) 0.51 mm long, dorso-ventrally symmetric; median lobe C-shaped in lateral view and broadened in apical half in dorso-ventral view, with large basal capsule and oval foramen; parameres (Fig. 3D) each thin, elongate and curved ventrally in lateral view, with two apical and one preapical macroseta and four short setae near base.

Female. Similar to male in external morphology (Fig. 1B). Antenna slightly shorter, club less enlarged (0.77 mm for antennomeres 1–7 vs. 0.82 mm for antennomeres 8–11); each compound eye composed of approximately 43 ommatidia; elytra slightly shorter; metathoracic wings also well developed; metaventricle lacking crest at middle. Measurements (as for male): body length 2.84 mm; length/width of head 0.62/0.49 mm, pronotum 0.63/0.65 mm, elytra 0.91/1.22 mm; abdomen 0.83/1.13 mm; length of antenna 1.59 mm.

Comparative notes. This species is unique among all known members of the genus in the coarsely punctate body surface, and a rather different type of the male antennal modifications.

Bionomics. The male holotype was collected by sifting *Cryptomeria* (Cupressaceae) leaves, while the other two individuals were found inside decomposing wood with termites. Although indicated by a set of morphological traits such as the compact body segments and moniliform antennomeres, it is still inconclusive as to whether the beetle has an obligate association with social insects, pending further observations in the field or at the laboratory.

Distribution. East China: Zhejiang.

Etymology. The specific epithet refers to West Tianmu Mountain, the type locality of this species.

Remarks. To accommodate the new species, the key to East and South Asian species of *Pseudophanias* by Inoue et al. (2020) is modified as follow:

- 0 Whole body surface roughly punctate and with dense, short setae; antennal club formed by enlarged, subcylindrical antennomeres 8–11; elytra roundly quadrate, broadly truncate at bases.....
- ...***Pseudophanias tianmuensis* Yin & Zhao, sp. nov. (East China: Zhejiang)**
- Body surface usually finely punctate (head with coarse punctuation in *P. excavatus* Inoue, Nomura & Yin) and with moderately long pubescence; antennal club formed by antennomere 11 alone, or by variously modified antennomeres 5–11, 8–11, or 9–11; elytra roundly triangular, and more strongly constricted at bases..... **1**

Acknowledgements

We thank Xiao-Bing Song (宋晓彬) for providing the first male specimen and associated collecting data 16 years ago. Donald S. Chandler (University of New Hampshire, Durham, USA) and Shûhei Nomura (National Museum of Nature and Science, Tsukuba, Japan) read the draft manuscript and provided helpful comments. This study was supported by the National Natural Science Foundation of China (no. 31872965) granted to ZWY.

References

- Chandler DS (2001) Biology, morphology, and systematics of the ant-like litter beetles of Australia (Coleoptera: Staphylinidae: Pselaphinae). *Memoirs on Entomology International* 15: [i–x,] 1–560.
- Hlaváč P (2000) *Chandleriella*, new genus of Tmesiphorini from Borneo (Coleoptera: Staphylinidae: Pselaphinae). *Entomological Problems* 31(1): 91–93.
- Inoue S, Nomura S (2020) A new species of the genus *Pseudophanias* Raffray, 1890 from Thailand (Coleoptera: Staphylinidae: Pselaphinae). *Japanese Journal of Systematic Entomology* 26(2): 336–339.
- Inoue S, Nomura S (2021) Two new species of *Pseudophanias* Raffray from Myanmar (Coleoptera: Staphylinidae: Pselaphinae). *Zootaxa* 4996(3): 591–599. <https://doi.org/10.11646/zootaxa.4996.3.11>
- Inoue S, Nomura S, Yin Z-W (2020) Three new species of *Pseudophanias* Raffray from Japan and Taiwan Island, and synonymy of *Chandleriella* Hlaváč with *Pseudophanias* (Coleoptera, Staphylinidae, Pselaphinae). *ZooKeys* 987: 135–156. <https://doi.org/10.3897/zookeys.987.53648>
- Yin Z-W (2019) First record of the genus *Chandleriella* Hlaváč (Coleoptera: Staphylinidae: Pselaphinae) from China, with description of a second species. *Zootaxa* 4571(3): 432–438. <https://doi.org/10.11646/zootaxa.4571.3.11>
- Yin Z-W, Coulon G, Bekchiev R (2015) A new species of *Pseudophanias* Raffray from a cave in central Nepal (Coleoptera: Staphylinidae: Pselaphinae). *Zootaxa* 4048(3): 446–450. <https://doi.org/10.11646/zootaxa.4048.3.10>