

Revision of the family Haliplidae (Insecta, Coleoptera) in Japan

Masakazu Hayashi¹, Tomofumi Iwata², Hiroyuki Yoshitomi³¹ Hoshizaki Green Foundation, Sono, Izumo, 691-0076, Japan² Toyama Science Museum, 1-8-31 Nishinakano-machi, Toyama, 939-8084, Japan³ Entomological Laboratory, Faculty of Agriculture, Ehime University, Tarumi 3-5-7, Matsuyama, 790-8566, JapanCorresponding authors: Masakazu Hayashi (hgf-haya@green-f.or.jp); Hiroyuki Yoshitomi (hymushi@agr.ehime-u.ac.jp)

Abstract

The Japanese members of Haliplidae were reviewed and 13 species in two genera are recognized. A new species, *Haliplus morii* **sp. nov.** is described from Honshu; it is similar to *Haliplus japonicus* Sharp, 1873, but belongs to a different subgenus. *Haliplus diruptus* J. Balfour-Browne, 1946, **syn. nov.** is treated as a junior synonym of *Haliplus kotoshonis* Kano & Kamiya, 1931. The records of *Haliplus davidi* Vondel, 1991 from Japan are regarded as misidentifications of *H. kotoshonis*. *Haliplus basinotatus latiusculus* Nakane, 1985, **syn. nov.** is treated as a junior synonym of *H. basinotatus*. *Haliplus angustifrons* Régimbart, 1892 known from south and southeast Asia, is newly recorded from Japan.

Key words: Aquatic beetles, Dytiscoidea, east Asia, Oriental Region, Palaearctic Region

Introduction

The Japanese Haliplidae includes 11 species in two genera (Nakajima et al. 2020). Two species of the genus *Peltodytes* Régimbart, 1879 are recorded (Kamiya 1936) but the genus *Haliplus* Latreille, 1802 is one of the most diverse of the family. Among the Japanese *Haliplus* that have been reviewed by Nakane (1963a, 1963b, 1985) and Satô (1984, 1985), there are disagreements in the taxonomic assignments of some species. Since then, van Vondel (1991, 1995, 1998, 2003a, 2003b, 2019) and van Vondel et al. (2006) have extensively examined the Old World species, and the elucidation of Asian species has progressed. However, for Japanese species, unresolved problems remain due to confusion between species whose type specimens may have disappeared (Mori 2007). As a result of re-examination of Japanese Haliplidae, 13 species in two genera, including one new species and one newly recorded species, were found, and are described and illustrated. Additionally, a review was conducted on Japanese Haliplidae, and their distributions, larval stages, and ecological notes have been added.

Materials and methods

Specimens studied herein are deposited at the following institutions and collections:

EUMJ	Ehime University, Matsuyama, Japan [Hiroyuki Yoshitomi];
HIPC	Hiroaki Iketake Private Collection, Nagoya, Japan;



Academic editor: Christopher Majka

Received: 26 December 2022

Accepted: 30 May 2023

Published: 3 July 2023

ZooBank: <https://zoobank.org/28659D39-3708-4035-84BF-A95AC91C59FF>

Citation: Hayashi M, Iwata T, Yoshitomi H (2023) Revision of the family Haliplidae (Insecta, Coleoptera) in Japan. ZooKeys 1168: 267–294. <https://doi.org/10.3897/zookeys.1168.99302>

Copyright: © Masakazu Hayashi et al. This is an open access article distributed under terms of the Creative Commons Attribution License ([Attribution 4.0 International – CC BY 4.0](https://creativecommons.org/licenses/by/4.0/)).

HOWP	Hoshizaki Institute for Wildlife Protection, Izumo, Japan [Masakazu Hayashi];
IIM	Ishikawa Insect Museum, Hakusan, Japan [Kohei Watanabe];
KAPC	Koji Arai Private Collection, Ranzan, Japan;
KPMNH	Kanagawa Prefectural Museum of Natural History, Odawara, Japan [Kyohei Watanabe];
KWPC	Kohei Watanabe Private Collection, Hakusan, Japan;
RUMF	Ryukyu University Museum Fujukan, Nishihara, Japan [Takeshi Sasaki];
TFPC	Takuya Fukuzawa Private Collection, Takao, Japan;
TIPC	Tomofumi Iwata Private Collection, Toyama, Japan;
TPM	Tochigi Prefectural Museum, Utsunomiya, Japan [Takashi Kurihara].

Minute structures on body surface of the new species were photographed under a scanning electron microscope (JCM-6000 Neoscope, JEOL Ltd., Tokyo, Japan). Male genitalia of the new species were photographed under a light microscope, Nikon Eclipse E600 with the digital camera (Digital Camera DS-L2, Nikon Ltd., Tokyo, Japan). The digital photographs were made by focus stacking, using a digital image processing software, Adobe Photoshop CS4 for Macintosh.

Stereoscopic (Leica S8 APO, Leica Microsystems Ltd., Wetzlar, Germany) and digital (HiROX KH-1300, HiROX Co. Ltd., Tokyo, Japan) microscopes were used to observe each part of the specimens except for the new species. Photographs were taken of the whole body, head, prosternal process, and male genitalia (penis, right and left paramere). Approximately ten images of the whole body were taken using a microscope camera (Digital Camera Unit DS-Fi1, Nikon Ltd., Tokyo, Japan) attached to a stereoscopic microscope, and focus stacking used to combine images using the image processing software CombineZM. The head and male genitalia were removed with tweezers after softening the dried specimens. The male genitalia were then placed in a 10% KOH solution for several hours to dissolve the flesh, and dissected under a stereomicroscope (Leica S8 APO).

The measurement abbreviations used in the text are shown below. The average is given in parenthesis after the range.

BT	(body thickness): maximum thickness of body height;
CED	(compound eyes distance): minimum distance between eyes;
EL	(elytral length): maximum length of the elytron (measured on the line of suture);
EW	(elytral width): maximum width of the elytra (two elytra);
HW	(head width): maximum width of the head (including compound eyes);
PL	(pronotal length): maximum length of pronotum;
PW	(pronotal width): maximum width of pronotum;
TL	total length (PL+EL).

Dried specimens were observed from the dorsal (HW, CED, PL, PW, EL, EW) and lateral (BT) views, and the maximum length at each measurement site was measured. For observation and measurement, a stereomicroscope (Olympus SZ40, Olympus Corporation, Tokyo, Japan) was used for PL, PW, EL, EW, and BT, and a stereomicroscope (Leica S8 APO) was used for HW and CED.



Figure 1. Map showing the island names included in this paper.

The terminology of body characters is followed by van Vondel (1991, 1992, 2019) and van Vondel et al. (2006). Fig. 1 shows the island names of Japan included in this paper.

Taxonomic accounts

Genus *Peltodytes* Régimbart, 1879

Peltodytes intermedius (Sharp, 1873)

Figs 2, 18A

Japanese name: Kogashira-mizumushi

Cnemidotus intermedius Sharp, 1873: 55.

Peltodytes intermedius: Kamiya 1936: 39; 1951: 122; Satô 1985: 181; Nakane 1963a: 55; 1987: 27; Matsui 1992: 2; Vondel 1995: 124; 2003a: 33; Takahashi 1998: 6; Dejima 2007: 83; 2010: 43; Tomisawa 2012: 42; Yoshitomi 2014: 25; Matsuo and Fukagawa 2016: 50; Mitamura et al. 2017: 136; Hayashi and Kadowaki 2019: 25; Nakajima et al. 2020: 19; Nakamine and Nakamine 2021: 2.

Material examined. 1 ex., Ehime Prefecture: Hakata-jima, 14–15. VIII.1997, H. Nakanishi leg. (EUMJ); 1 ex., Ehime Prefecture: Iwagi-jima, 16.VIII.1997, H. Nakanishi leg. (EUMJ); 1 ex., Ehime Prefecture: Ômi-shima, 13.VIII.1997, H. Nakanishi leg. (EUMJ); 3 exs., Ehime Prefecture: Ôshima, 23.V.1998, H. Nakanishi leg. (EUMJ); 1 ex., Ehime Prefecture: Ôshima, 13.VIII.1997, H. Nakanishi leg.




Figure 2. *Peltodytes intermedius* **A** habitus **B** head **C** prosternal process **D** penis **E** left paramere **F** right paramere. Scale bars: 1.0 mm (**A**); 0.5 mm (**B**); 0.25 mm (**C**); 0.1 mm (**D–F**).

(EUMJ); 1 ex., Kagoshima Prefecture: Chaban, Yoron-to, 10.VIII.1958, S. Ueno & Y. Morimoto leg. (EUMJ). Other specimens are listed in Suppl. material 1.

Measurements ($n = 10$). TL 3.38–4.82 (4.48) mm; HW 0.69–0.78 (0.73) mm; CED 0.19–0.23 (0.21) mm; PL 0.62–0.71 (0.67) mm; PW 1.45–1.64 (1.52) mm; EL 2.34–2.67 (2.45) mm; EW 1.86–2.16 (1.98) mm; BT 1.41–1.65 (1.53) mm; HW/CED 3.22–3.83 (3.49); PW/PL 2.18–2.35 (2.27); EL/EW 1.17–1.27 (1.22).

Biology. This species typically inhabits stagnant water environments such as ponds, paddies, and swamps (Nakajima et al. 2020). The larvae primarily feed on Zygnemataceae algae. The adults were collected using a sweep net in shallow water or captured by a light trap. A pupal chamber was observed within the mud in rearing condition (Hayashi 2015).

Immature stages. The larva was described by Fukuda et al. (1959) and illustrated in color by Mitamura et al. (2017) and Nakajima et al. (2020).

DNA barcodes. The sequences of COI (Cox1) gene of Japanese specimens are deposited in DNA Data Bank of Japan (DDBJ) (Hayashi and Ooi 2022): Aomori Prefecture ([LC633200](#)), Shimane Prefecture ([LC633201–LC633205](#)).

Distribution. JAPAN: Hokkaido, Honshu, Shikoku, Kyushu, Sado, Izu-shoto, Noto-jima of Nanao-wan in Ishikawa, Oki, Awaji-shima, Shodo-shima, Te-jima

in Kagawa, Geiyo-shoto in Seto-naikai (new records) (Iwagi-jima, Ômishima, Hakata-jima, Ôshima), Tsushima, Iki, Hirado-jima in Nagasaki, Goto-retto, Takashima of Imari-wan in Nagasaki, Tobase-jima of Yatsushiro-kai in Kumamoto, Amakusa-shoto in Kumamoto, Nansei shoto (Tanegashima, Yoron-jima: new record); Korea, China, Taiwan, Far East of Russia.

***Peltodytes sinensis* (Hope, 1845)**

Figs 3, 18B

Japanese name: Shina-kogashira-mizumushi

Haliplus sinensis Hope, 1845: 15.

Peltodytes sinensis: Kamiya 1936: 40; Satô 1985: 181; Nakane 1987: 27; Vondel 1995: 124; 2003a: 33; Aoyagi 2011a: 99; 2011b: 107; Matsuo and Fukagawa 2016: 51; Mitamura et al. 2017: 137; Aoyagi 2019: 34; Nakajima et al. 2020: 18; Aoyagi 2020: 87; Uchida et al. 2021: 183.

Material examined. 6 exs., Kagoshima Prefecture: Ohtsukan, Okinoerabu-jima, 3.VIII.1958, S. Ueno & Y. Morimoto leg. (EUMJ); 1 ex., Kagoshima Prefecture:




Figure 3. *Peltodytes sinensis* **A** habitus **B** head **C** prosternal process **D** penis **E** left paramere **F** right paramere. Scale bars: 1.0 mm (**A**); 0.5 mm (**B**); 0.25 mm (**C**); 0.1 mm (**D–F**).

Uchida, near, Taniyama, Okinoerabu-jima, 4.VIII.1958, S. Ueno & Y. Morimoto leg. (EUMJ). Other specimens are listed in Suppl. material 1.

Measurements ($n = 10$). TL 3.06–5.21 (4.76) mm; HW 0.75–0.82 (0.78) mm; CED 0.20–0.25 (0.22) mm; PL 0.63–0.80 (0.71) mm; PW 1.46–1.68 (1.58) mm; EL 2.43–2.76 (2.58) mm; EW 1.91–2.30 (2.12) mm; BT 1.46–1.76 (1.62) mm; HW/CED 3.26–3.73 (3.49); PW/PL 2.01–2.40 (2.21); EL/EW 1.17–1.28 (1.21).

Biology. This species usually lives in still, shallow waters such as ponds and paddies, and the adults were collected by sweep netting in shallow water and using light traps (Nakajima et al. 2020).

Immature stages. Unknown.

Distribution. JAPAN: Tsushima, Nansei shoto (Nakano-shima and Takara-jima of Tokara-retto, Amami-Ōshima, Kakeroma-jima, Tokuno-shima, Okinoerabu-jima (new record), Yoron-jima, Okinawa-jima, Yagaji-shima in Okinawa, Miyagi-jima in Okinawa, Iheya-jima, Kume-jima, Miyako-jima, Ikema-jima in Okinawa, Irabu-jima in Okinawa, Ishigaki-jima, Iriomote-jima, Yonaguni-jima); Korea, China, Taiwan, SE Asia.

Genus *Haliplus* Latreille, 1802

Subgenus *Nipponiplus* Vondel, 2019

Haliplus (*Nipponiplus*) *japonicus* Sharp, 1873

Figs 4, 18C

Japanese name: Kubiboso-kogashira-mizumushi

Haliplus japonicus Sharp, 1873: 55. Kamiya 1936: 44; Satō 1985: 181; Nakane 1963a: 55; 1985: 62; 1987: 29; Matsui 1992: 2; Vondel 1995: 122; Tomisawa 2012: 42; Matsuo and Fukagawa 2016: 51; Mitamura et al. 2017: 138; Hayashi and Kadowaki 2019: 25; Nakajima et al. 2020: 20.

Haliplus (*Haliplus*) *japonicus*: Satō, 1984: 1; Vondel 2003a: 31; Vondel et al. 2006: 250.

Haliplus (*Nipponiplus*) *japonicus*: Vondel 2019: 22.

Haliplus brevior Nakane, 1963a: 55. Nakane 1985: 62; 1987: 29. [synonymized with *H. minutus* by Satō (1984) but with *H. japonicus* by van Vondel et al. (2006)]

Material examined. Specimens examined in this study are listed in Suppl. material 1.

Measurements ($n = 10$). TL 2.69–4.08 (3.79) mm; HW 0.62–0.69 (0.67) mm; CED 0.29–0.32 (0.30) mm; PL 0.55–0.66 (0.62) mm; PW 1.05–1.20 (1.12) mm; EL 1.99–2.20 (2.10) mm; EW 1.50–1.71 (1.60) mm; BT 1.12–1.28 (1.19) mm; HW/CED 2.12–2.35 (2.24); PW/PL 1.65–2.04 (1.76); EL/EW 1.25–1.34 (1.32).

Biology. This species usually lives in fresh waters with abundant aquatic plants such as ponds, paddies, and streams (Nakajima et al. 2020). The larvae eat Zygnemataceae algae (Hayashi 2015). The adults were collected by sweep netting in shallow waters and are rarely attracted by light traps (Hayashi 2015). The pupation in mud with a pupal chamber was accomplished by laboratory rearing (Hayashi 2015).




Figure 4. *Haliplus japonicus* **A** habitus **B** head **C** prosternal process **D** penis **E** left paramere **F** right paramere. Scale bars: 1.0 mm (**A**); 0.5 mm (**B**); 0.25 mm (**C**); 0.1 mm (**D–F**).

Immature stages. The color photographs were provided by Mitamura et al. (2017) and Nakajima et al. (2020).

Distribution. JAPAN: Hokkaido, Honshu, Shikoku, Kyushu, Noto-jima, Oki, Go-to-retto, Taka-shima, Amakusa-shoto; China, Far East Russia.

***Haliplus (Nipponiplus) regimbarti* Zaitzev, 1908**

Figs 5, 18D

Japanese name: Taiwan-kogashira-mizumushi

Haliplus brevis Wehncke, 1880: 7. [nec Stephens 1828]

Haliplus regimbarti Zaitzev, 1908: 122. [replacement name for *Haliplus brevis* Wehncke, 1880]

Haliplus regimbarti: Vondel 1995: 122; Nakajima et al. 2020: 21.

Haliplus (Haliplus) regimbarti: Vondel et al. 2006: 257; Iwata 2016: 100.

Haliplus sauteri Zimmermann, 1924: 130. [synonymized by van Vondel 1995]

Material examined. Specimens examined in this study are listed in Suppl. material 1.




Figure 5. *Haliphus regimbarti* **A** habitus **B** head **C** prosternal process **D** penis **E** left paramere **F** right paramere. Scale bars: 1.0 mm **A**; 0.5 mm (**B**); 0.25 mm (**C**); 0.1 mm (**D–F**).

Measurements ($n = 4$). TL 2.87–4.44 (4.02) mm; HW 0.68–0.74 (0.72) mm; CED 0.31–0.34 (0.32) mm; PL 0.65–0.71 (0.68) mm; PW 1.23–1.35 (1.30) mm; EL 2.22–2.35 (2.31) mm; EW 1.76–1.90 (1.85) mm; BT 1.26–1.39 (1.35) mm; HW/CED 2.18–2.34 (2.23); PW/PL 1.84–1.98 (1.92); EL/EW 1.22–1.26 (1.25).

Biology. This species was collected from a small pond with submerged water plants (Iwata 2016) and a narrow stream with flourishing water plants flowing beside paddy fields (Nakajima et al. 2020).

Immature stages. Unknown.

Distribution. JAPAN: Nansei shoto (Yonaguni-jima); Taiwan, south China.

Subgenus *Haliphus* Latreille, 1802

Haliphus (*Haliphus*) *kamiyai* Nakane, 1963

Figs 6, 18D

Japanese name: Kamiya-kogashira-mizumushi

Haliphus kamiyai Nakane, 1963b: 25. Nakane 1963a: 55; 1985: 62; 1987: 29; Mitamura et al. 2017: 141; Nakajima et al. 2020: 20.

Haliphus (*Haliphus*) *kamiyai*: Vondel et al. 2006: 252.

Material examined. Specimens examined in this study are listed in Suppl. material 1.




Figure 6. *Haliplus kamiyai* **A** habitus **B** head **C** prosternal process **D** penis **E** left paramere **F** right paramere. Scale bars: 1.0 mm (**A**); 0.5 mm (**B**); 0.25 mm (**C**); 0.1 mm (**D–F**).

Measurements ($n = 10$). TL 2.64–4.24 (3.84) mm; HW 0.61–0.71 (0.66) mm; CED 0.34–0.36 (0.35) mm; PL 0.60–0.67 (0.64) mm; PW 1.06–1.25 (1.16) mm; EL 1.85–2.28 (2.08) mm; EW 1.46–1.71 (1.60) mm; BT 1.16–1.33 (1.24) mm; HW/CED 1.81–2.09 (1.89); PW/PL 1.76–1.90 (1.79); EL/EW 1.26–1.33 (1.29).

Biology. This species lives in still water with abundant aquatic plants such as, ponds, paddies, and streams (Nakajima et al. 2020). The larvae feed on Zygnemataceae algae (Watanabe and Yamasaki 2020).

Immature stages. Color photographs were provided by Watanabe and Yamasaki (2020).

Distribution. JAPAN: Honshu (Tohoku and Kanto Regions).

***Haliplus (Haliplus) morii* sp. nov.**

<https://zoobank.org/71E771A3-91A0-4F75-885E-4D1789CBCD8F>

Figs 7–10, 18D

Japanese name: Mizonashi-kogashira-mizumushi

Type series. **Holotype** (Fig. 7) [EUMJ] 1 male Ochifushi, Yuza-machi, Yamagata Pref., Japan, 21.II.1993, M. Takahashi leg. **Paratypes** [EUMJ, HOWP] 11 exs. (male and female) Same data as holotype.

Description. TL 3.0–3.3 mm, EW 1.5–1.6 mm. Body oval, tapering backwards, widest before the middle (Fig. 7).



Figure 7. *Haliphus morii* sp. nov. habitus. Scale bar: 1.0 mm.

Head. Black, strongly and densely punctured, labrum dark yellow. Distance between eyes 1.7–2.0× width of one eye. Antennae yellow. Palpi yellow.

Pronotum. Yellow to yellowish-brown. Plicae lacking (Fig. 8C). Coarsely punctate near base. Lateral sides margined, nearly straight.

Elytra. Yellow to yellow-brown, dark interrupted lines on primary puncture rows, darkened along suture, sometime indistinct marks connecting primary puncture rows. Completely margined. Primary puncture rows moderately strong, dense in first rows, ~ 36 or 37 punctures in first row. Secondary punctures moderately strong and dense along suture, moderately strong and sparse on intervals. All punctures darkened.

Ventral side. Yellow-brown, legs yellow-brown, slightly darkened towards coxae, elytral epipleura yellow-brown with strong punctures, reaching to sixth sternite. Prosternal process narrowed near coxae, grooved along each side, anterior edge weakly margined, moderately strongly punctured. Metasternal process flat or even slightly bulbous with a row of strong punctures on each side, else moderately punctured, emarginate in apical margin (Fig. 8F). Metacoxal plates reaching to fifth sternite, moderately strongly punctured, near suture weakly punctured, row of setae on posterior edge (Fig. 9C, D). Third and fourth sternites not fused with well indicated depression and around ridge. Fifth and sixth sternite with sparse transverse puncture row, last sternite weakly punctured in apical part. No setiferous striole on dorsal face of hind tibia, longer tibial spur of hind legs 2/3 length of first tarsal segment (Fig. 8D).




Figure 8. *Haliphus morii* sp. nov. SEM **A** habitus **B** lateral side **C** apical half of body **D** right hind leg **E** labium **F** prosternal process.

Male genitalia. Penis carved in apical and basal part; slender and round in apex (Fig. 10A). Right paramere triangular, apex with long setae and lacking small segment (Fig. 10C).

Similar species. This species may be confused with *H. japonicus*, but the latter has plicae on the pronotum.

Biology. The type locality is dominated by rice paddy fields in the plains.

Distribution. JAPAN (Tohoku Region).

Etymology. The species name is dedicated to Mr. Masato Mori, who first noticed the existence of this species.




Figure 9. *Haliphus morii* sp. nov. SEM A left pro leg B prosternal process C metacoxa D base of abdominal segments.



Figure 10. *Haliphus morii* sp. nov. male genitalia A penis B left paramere C right paramere. Scale bar: 0.1 mm.

***Haliplus (Haliplus) simplex* Clark, 1863**

Figs 11, 18E

Japanese name: Chibi-kogashira-mizumushi

Haliplus simplex Clark, 1863: 419. Vondel 1995: 123; Matsuo and Fukagawa 2016: 51; Mitamura et al. 2017: 142; Nakajima et al. 2020: 21.

Haliplus minutus Takizawa, 1931: 140. Kamiya 1936: 45; Satô 1985: 181; Nakane 1985: 62. [synonymized by van Vondel et al. (2006)]

Haliplus (Haliplus) minutus: Satô 1984: 1; Vondel 2003a: 31; Vondel et al. 2006: 265.

Material examined. Specimens examined in this study are listed in Suppl. material 1.

Measurements ($n = 10$). TL 2.49–3.75 (3.54) mm; HW 0.62–0.66 (0.64) mm; CED 0.32–0.35 (0.33) mm; PL 0.57–0.62 (0.59) mm; PW 1.04–1.14 (1.10) mm; EL 1.86–1.99 (1.92) mm; EW 1.50–1.61 (1.55) mm; BT 1.14–1.19 (1.15) mm; HW/CED 1.86–2.02 (1.92); PW/PL 1.75–1.94 (1.84); EL/EW 1.21–1.28 (1.24).

Biology. This species typically inhabits stagnant water environments such as ponds, paddies, and swamps, and the adults were collected by sweep netting in shallow water (Nakajima et al. 2020).

Immature stages. Unknown.

Distribution. JAPAN: Hokkaido, Honshu (Tohoku Region), Tsushima; Korea, China, Far East of Russia.




Figure 11. *Haliplus simplex* **A** habitus **B** head **C** prosternal process **D** penis **E** left paramere **F** right paramere. Scale bars: 1.0 mm (**A**); 0.5 mm (**B**); 0.25 mm (**C**); 0.1 mm (**D–F**).

Subgenus *Liaphlus* Guignot, 1928

***Haliplus (Liaphlus) angustifrons* Régimbart, 1892**

Figs 12, 18E

Japanese name: Usucha-kogashira-mizumushi

Haliplus angustifrons Régimbart, 1892: 112. van Vondel 1993: 292; Sheth et al. 2016: 361.

Haliplus kotoshonis: Vondel, 1991: 113 [misidentification]; Aoyagi 2014: 192 [misidentification].

Material examined. 2 exs., Kagoshima Prefecture: Amagi, Amagi-cho, Ôshima-gun, Tokuno-shima, 1.XI.2010, H. Iketake leg. (HIPC); 27 exs., Okinawa Prefecture: Ohgimi, Ohgimi-son [Okinawa-jima], 19.III.2014, R. Okano leg. (EUMJ); 1 ex., Okinawa Prefecture: Nakama, Onna-son [Okinawa-jima], 30.X.2011, H. Iketake leg. (HIPC); 1 ex., Okinawa Prefecture: Onna-dam, Sokei, Ginoza-son [Okinawa-jima], 30.X.2011, H. Iketake leg. (HIPC); 6 exs., Okinawa Prefecture: Nuuha, Ohgimi-son [Okinawa-jima], 8.I.1989, Y. Abe & T. Abe leg. (KPMNH); 1 ex., Okinawa Prefecture: Gima, Kumejima-cho, 19.V.2007, Y. Kamite leg. (EUMJ); 1




Figure 12. *Haliplus angustifrons* **A** habitus **B** head **C** prosternal process **D** penis **E** left paramere **F** right paramere. Scale bars: 1.0 mm (**A**); 0.5 mm (**B**); 0.25 mm (**C**); 0.1 mm (**D–F**).

ex., Okinawa Prefecture: Kanegusuku, Kumejima-cho, 20.V.2007, Y. Kamite leg. (EUMJ); 6 exs., Okinawa Prefecture: Uehara, Taketomi-cho (24.40°N, 123.79°E) [Iriomote-jima], 14.III.2017, T. Iwata leg. (TIPC); 1 ex., Okinawa Prefecture: Ishigaki-jima, 5.VI.1970, T. Hozumi leg. (EUMJ).

Measurements ($n = 10$). TL 3.66–5.36 (4.98) mm; HW 0.80–0.87 (0.84) mm; CED 0.27–0.30 (0.28) mm; PL 0.76–0.82 (0.79) mm; PW 1.64–1.76 (1.68) mm; EL 2.67–2.86 (2.76) mm; EW 2.04–2.25 (2.14) mm; BT 1.48–1.69 (1.59) mm; HW/CED 2.85–3.07 (2.95); PW/PL 2.07–2.17 (2.12); EL/EW 1.27–1.32 (1.29).

Biology. The above specimens were collected from small ponds.

Immature stages. Unknown.

Discussion. *Haliplus angustifrons* is widely distributed from south Asia to southeast Asia (van Vondel 1993; Sheth et al. 2016), but there are no records in east Asia. The pattern of dorsal marks, the shape of the prosternal process, and male genitalia are in agreement with the redescription of *H. angustifrons* (van Vondel 1993). The figures of “*H. kotoshonis*” given by van Vondel (1991, 1993) are not of true *H. kotoshonis*, and at least the records of Japanese specimens correspond to *H. angustifrons*.

Distribution. JAPAN (new records): Nansei shoto (Tokuno-shima, Okinawa-jima, Iheya-jima, Kume-jima, Ishigaki-jima, Iriomote-jima); Pakistan, India, Sri Lanka, Nepal, Myanmar, Laos, Vietnam.

***Haliplus (Liaphlus) basinotatus* Zimmermann, 1924**

Figs 13, 18F

Japanese name: Kurohoshi-kogashira-mizumushi

Haliplus basinotatus Zimmermann, 1924: 137. Nakane 1963a: 55; 1987: 30; Vondel 1995: 119; Matsuo and Fukagawa 2016: 52; Mitamura et al. 2017: 139; Hayashi and Kadowaki 2019: 25; Nakajima et al. 2020: 22; Watanabe 2021: 16.

Haliplus (Liaphlus) basinotatus: Vondel 2003a: 31.

Haliplus basinotatus latiusculus Nakane, 1985: 63. Syn. nov.

Material examined. Specimens examined in this study are listed in Suppl. material 1.

Measurements ($n = 10$). TL 3.30–5.34 (4.88) mm; HW 0.78–0.88 (0.83) mm; CED 0.30–0.38 (0.35) mm; PL 0.71–0.75 (0.74) mm; PW 1.43–1.59 (1.52) mm; EL 2.54–2.86 (2.66) mm; EW 1.95–2.17 (2.06) mm; BT 1.60–1.74 (1.65) mm; HW/CED 2.27–2.62 (2.35); PW/PL 1.95–2.14 (2.05); EL/EW 1.24–1.32 (1.27).

Biology. The larvae eat Characeae algae, and the adults were collected by sweep on shallow water (Nakajima et al. 2020).

Immature stages. A color illustration is given by Mitamura et al. (2017).

DNA barcodes. The COI (Cox1) gene sequence of a Japanese specimen are deposited in DDBJ (Hayashi and Ooi 2022): Shimane Prefecture ([LC633206](#)).

Discussion. Nakane (1985) described a subspecies for Japanese population, but we could not find any differences between the Asian (van Vondel 1995: fig. 53) and Japanese populations. In this paper we treat this subspecies as a junior synonym of nominotypical subspecies.

Distribution. JAPAN: Hokkaido, Honshu, Shikoku, Kyushu, Oki, Tsushima; Korea, China, Far East Russia.




Figure 13. *Haliplus basinotatus* **A** habitus **B** head **C** prosternal process **D** penis **E** left paramere **F** right paramere. Scale bars: 1.0 mm (**A**); 0.5 mm (**B**); 0.25 mm (**C**); 0.1 mm (**D–F**).

***Haliplus (Liaphlus) kotoshonis* Kano & Kamiya, 1931**

Figs 14, 18F

Japanese name: Koto-kogashira-mizumushi

Haliplus kotoshonis Kano & Kamiya, 1931: 2. Satô 1985: 181, pls 33–37; Nakane 1985: 63; 1987: 30; Nakajima et al. 2020: 21.

Haliplus (Liaphlus) kotoshonis: Satô 1984: 2.

Haliplus diruptus J. Balfour-Browne, 1946: 436. Vondel 1995: 120, figs 71–79. Syn. nov.

Haliplus davidi Vondel, 1991: 92. [synonymized with *Haliplus diruptus* by van Vondel (2017)]

Material examined. 1 ex., Kagoshima Prefecture: Takara-jima, Tokara, 2.VII.1960, M. Satô leg. (EUMJ); 2 exs., Kagoshima Prefecture: Takara-jima, Tokara, 20.VII.1961, Y. Hama leg. (EUMJ); 7 exs., Kagoshima Prefecture: Takara-jima, Tokara, 2.VI.1962, M. Satô leg. (EUMJ); 2 exs., Okinawa Prefecture:




Figure 14. *Haliphus kotoshonis* **A** habitus **B** head **C** prosternal process **D** penis **E** left paramere **F** right paramere. Scale bars: 1.0 mm (**A**); 0.5 mm (**B**); 0.25 mm (**C**); 0.1 mm (**D–F**).

Maesato, Ishigaki-jima, 4.IX.1975, T. Takahashi leg. (EUMJ); 1 ex., Okinawa Prefecture: Mt. Urabu-dake, Yonaguni-jima, 1.IV.1990, Y. Uchida leg. (EUMJ); 1 ex., Okinawa Prefecture: Shirahama, Iriomote-jima, 30.VI.2001, T. Nakamura leg. (TPM); 3 exs., Okinawa Prefecture: Okinawa-jima, Ryukyu, 26.VI.1995, T. Takara leg. (RUMF).

Measurements ($n = 10$). TL 3.39–4.94 (4.47) mm; HW 0.74–0.87 (0.80) mm; CED 0.31–0.38 (0.33) mm; PL 0.65–0.79 (0.73) mm; PW 1.29–1.53 (1.42) mm; EL 2.30–2.62 (2.42) mm; EW 1.76–2.01 (1.88) mm; BT 1.39–1.70 (1.50) mm; HW/CED 2.30–2.56 (2.38); PW/PL 1.87–2.04 (1.95); EL/EW 1.23–1.32 (1.30).

Biology. The above specimens were collected from small ponds.

Immature stages. Unknown.

Discussion. This species was described based on one specimen from “Kotosho” (Orchid Island), Taiwan (Kano and Kamiya 1931). The type specimen is likely to have been lost (Mita et al. 2015), but the black marks on the dorsum are unique and the original description and figures are recognizable. Nakane (1985, 1987, 1990) has repeatedly pointed out that *Haliphus diruptus* J. Balfour-Browne, 1946 should be a junior synonym of the species, but a formal treatment has not been done. Judging from our investigation of the specimens and the description of van Vondel (1995), *H. diruptus* is treated as a junior syn-

onym of this species in this paper. In addition, *Haliplus davidi* Vondel, 1991 was recorded from Japan (van Vondel 1998, 2003a), but was later treated as a junior synonym of *H. diruptus* (van Vondel 2017); therefore, the record of *H. davidi* from Japan is not *H. diruptus* but *H. kotoshonis*.

This study revealed that specimens of *Haliplus angustifrons* Régimbart, 1892 were mixed in with the specimens identified as “*H. kotoshonis*” from Japan. In a typical individual, the dorsal marks can easily distinguish between the two species. In particular, this species has a black mark on head, but the *H. angustifrons* does not have such a mark.

Distribution. JAPAN: Nansei shoto (Takara-jima of Tokara-retto, Okinawa-jima, Ishigaki-jima, Iriomote-jima, Yonaguni-jima); Taiwan, China, Vietnam, Myanmar.

Haliplus (Liaphlus) eximius Clark, 1863

Figs 15, 19A

Japanese name: Kiuro-kogashira-mizumushi

Haliplus eximius Clark, 1863: 418. Kamiya 1936: 42; Satô 1985: 181; Nakane 1985: 63; 1987: 29; Matsui 1992: 2; Vondel 1995: 121; Dejima 2007: 84; Matsuo and Fukagawa 2016: 52; Mitamura et al. 2017: 141; Hayashi and Kadowaki 2019: 25; Nakajima et al. 2020: 23; Watanabe and Ohba 2022: 34.

Haliplus (Liaphlus) eximius: Satô 1984: 2; Vondel 1991: 97; 1993: 299; 2003a: 31.




Figure 15. *Haliplus eximius* **A** habitus **B** head **C** prosternal process **D** penis **E** left paramere **F** right paramere. Scale bars: 1.0 mm (**A**); 0.5 mm (**B**); 0.25 mm (**C**); 0.1 mm (**D–F**).

Haliphus hiogoensis Kano & Kamiya, 1931: 1. Kamiya 1936: 44. [synonymized by Satô 1984]

Material examined. 1 ex., Ehime Prefecture: Ôshima, 13.VIII.1997, H. Nakanishi leg. (EUMJ). Other specimens examined in this study are listed in Suppl. material 1.

Measurements ($n = 10$). TL 3.26–4.71 (4.38) mm; HW 0.70–0.78 (0.75) mm; CED 0.27–0.33 (0.30) mm; PL 0.69–0.75 (0.73) mm; PW 1.33–1.54 (1.46) mm; EL 2.10–2.53 (2.36) mm; EW 1.66–1.94 (1.85) mm; BT 1.31–1.51 (1.44) mm; HW/CED 2.23–2.70 (2.49); PW/PL 1.89–2.08 (2.03); EL/EW 1.24–1.32 (1.27).

Biology. This species typically inhabits ponds, and the adults were collected by sweep nets over shallow water (Nakajima et al. 2020). The larvae eat Characeae algae (Yamazaki et al. 2020).

Immature stages. The color photograph was shown by Kitano and Sano (2011).

DNA barcodes. The COI (Cox1) gene sequence of one Japanese specimen were deposited in DDBJ (Hayashi and Ooi 2022): Shimane Prefecture ([LC633207](#)).

Distribution. JAPAN: central to west Honshu, Shikoku, Kyushu, Oki, Hiroshima in Kagawa, Geiyo-shoto (Ôshima: new record), Tsushima, Iki, Hirado-jima, Goto-retto, Amakusa-shoto, Nansei shoto (Yonaguni-jima); Korea, China, SE Asia.

Haliphus (Liaphlus) ovalis Sharp, 1884

Figs 16, 19B

Japanese name: Hime-kogashira-mizumushi

Haliphus ovalis Sharp, 1884: 440. Kamiya 1936: 46; Satô 1985: 181; Nakane 1963a: 55; 1985: 63; 1987: 30; Matsuo and Fukagawa 2016: 52; Hayashi and Kadowaki 2019: 25; Nakajima et al. 2020: 23.

Haliphus (Liaphlus) ovalis: Satô 1984: 3; Vondel 1991: 125; 2003a: 32.

Material examined. 3 exs., Kagoshima Prefecture: Amagi, Amagi-cho, Ôshima-gun, Tokuno-shima, 1.XI.2010, H. Iketake leg. (HIPC); 1 ex., Niigata Prefecture: Ohura, Aikawa-machi, Sadoga-shima, 27–30. V. 1989, Y. Abe & T. Abe leg. (KPMNH); 11 exs., Kagoshima Prefecture: Ontsukan, Okinoerabu-jima, 3.VIII.1958, S. Ueno & Y. Morimoto leg. (EUMJ); 3 exs., Kagoshima Prefecture: Minzuki-ike, Okinoerabu-jima, 16.VIII.1958, S. Ueno & Y. Morimoto leg. (EUMJ); 12 exs., Kagoshima Prefecture: Okinoerabu-jima, 4.VIII.1958, S. Ueno & Y. Morimoto leg. (EUMJ).

Measurements ($n = 10$). TL 3.80–5.97 (5.47) mm; HW 0.90–0.99 (0.95) mm; CED 0.37–0.43 (0.40) mm; PL 0.74–0.85 (0.80) mm; PW 1.57–1.75 (1.66) mm; EL 2.81–3.18 (3.06) mm; EW 2.10–2.38 (2.22) mm; BT 1.67–1.95 (1.79) mm; HW/CED 2.29–2.50 (2.44); PW/PL 1.99–2.16 (2.13); EL/EW 1.34–1.42 (1.39).

Biology. This species typically inhabits ponds, and the adults were collected by sweep nets over shallow water and a light trap (Nakajima et al. 2020).

Immature stages. The larva was illustrated by Satô and Yoshitomi (2018).

Discussion. *Haliphus chinensis* Falkenström, 1932 distributed in mainland China is closely similar to this species. van Vondel (1991) states that the distinguishing points of both species, *H. ovalis* and *H. chinensis*, are the distance between the eyes and the morphology of the apical curve of penis (smooth in *H. chinensis* and flexed in *H. ovalis*). But some individuals from Ryukyu




Figure 16. *Haliplus ovalis* **A** habitus **B** head **C** prosternal process **D** penis **E** left paramere **F** right paramere. Scale bars: 1.0 mm (**A**); 0.5 mm (**B**); 0.25 mm (**C**); 0.1 mm (**D–F**).

(Okinoerabu-jima and Tokunoshima), Shikoku (Ehime Prefecture), and Honshu (Niigata Prefecture) have a smooth apex of the penis. Further detailed comparisons of both species are necessary.

DNA barcodes. The sequences of COI (Cox1) gene of Japanese specimens are deposited in DDBJ (Hayashi and Ooi 2022): Shimane Prefecture ([LC633208–LC633212](#)).

Distribution. JAPAN: Hokkaido, Honshu, Shikoku, Kyushu, Sado (new record), Oki, Goto-retto, Nansei shoto (Tokuno-shima, Okinoerabu-jima: new record).

Haliplus (Liaphlus) sharpi Wehncke, 1880

Figs 17, 19C

Japanese name: Madara-kogashira-mizumushi

Haliplus sharpi Wehncke, 1880: 74. Satô 1985: 181; Nakane 1985: 63; 1987: 30; Vondel 1995: 123; Matsuo and Fukagawa 2016: 53; Mitamura et al. 2017: 140; Hayashi and Kadowaki 2019: 25; Nakajima et al. 2020: 24; Nakamine and Nakamine 2021: 2; Imasaka et al. 2021: 71; Watanabe and Ohba 2022: 34.

Haliplus (Liaphlus) sharpi: Satô 1984: 3; Vondel 1991: 129; 1993: 313; 2003a: 32.

Haliplus simplex: Kamiya 1936: 48. [misidentification]




Figure 17. *Haliphus sharpi* **A** habitus **B** head **C** prosternal process **D** penis **E** left paramere **F** right paramere. Scale bars: 1.0 mm (**A**); 0.5 mm (**B**); 0.25 mm (**C**); 0.1 mm (**D–F**).

Haliphus tsukushiensis Yoshimura, 1932: 102. Nakane 1963a: 55. [synonymized by Satô 1984]

Haliphus holmeni Vondel, 1991: 109. [synonymized by van Vondel 2017]

Material examined. 10 exs., Niigata Prefecture: Ohura, Aikawa-machi, Sado-ga-shima, 27–30. V.1989, Y. Abe & T. Abe leg. (KPMNH).

Measurements ($n = 10$). TL 3.22–4.81 (4.48) mm; HW 0.71–0.81 (0.75) mm; CED 0.28–0.35 (0.31) mm; PL 0.70–0.80 (0.75) mm; PW 1.35–1.51 (1.44) mm; EL 2.30–2.52 (2.39) mm; EW 1.75–1.95 (1.87) mm; BT 1.40–1.56 (1.49) mm; HW/CED 2.25–2.69 (2.41); PW/PL 1.86–2.04 (1.95); EL/EW 1.23–1.33 (1.27).

Biology. This species typically inhabits stagnant water environments such as ponds, paddies, and swamp (Nakanishi 2012; Watanabe and Hidaka 2013). The larvae feed on Characeae algae (Nakanishi 2012). The pupation occurred in the pupal chamber in laboratory rearing experiments (Nakanishi 2012).

Immature stages. The color photographs were provided by Nakanishi (2012) and Mitamura et al. (2017).

Distribution. JAPAN: Hokkaido, Honshu, Shikoku, Kyushu, Sado (new record), Oki, Tsushima, Iki, Azuchi-ôshima in Nagasaki, Hirado-jima, Goto-retto, Koshikishima-retto, Nansei shoto (Tanegashima); Korea, China, Taiwan.




Figure 18. Distribution maps based on specimens examined **A** *Peltodytes intermedius* **B** *Peltodytes sinensis* **C** *Haliplus japonicus* **D** *Haliplus kamiyai* (circle), *Haliplus morii* (star), *Haliplus regimbarti* (square) **E** *Haliplus simplex* (black circle), *Haliplus angustifrons* (white circle) **F** *Haliplus basinotatus* (black circle), *Haliplus kotoshonis* (white circle).




Figure 19. Distribution maps based on specimens examined **A** *Haliplus eximius* **B** *Haliplus ovalis* **C** *Haliplus sharpi*.

Key to the species of *Haliplus* from Japan

Revised from Nakajima et al. 2020.

- 1 Pronotum with plicae (a pair of short grooves on basal-middle part) **2**
- Pronotum without plicae (Fig. 8C) **5**
- 2 Prosternal process without margin; front margin shorter than basal margin (Fig. 11C). Outline of elytra triangular (Fig. 11A). Penis robust in lateral view, apex with spines (Fig. 11D) ***H. simplex***
- Prosternal process with margin; front margin as long as basal margin. Outline of elytra oval. Penis without apical spines in lateral view **3**
- 3 Lateral sides of prosternal process slightly narrowing on base (Fig. 6C). Elytral marks developed, cruciform pattern on center (Fig. 6A). Penis curved nearly to apex and extremely narrowed (Fig. 6D) ***H. kamiyai***
- Lateral sides of prosternal process slightly widened on base. Elytral marks not developed, small black spots on disc. Penis gently curved from middle to apex **4**

- 4 Lateral surfaces of prosternal process with indistinct and narrow margins (Fig. 4C). Elytra gray (Fig. 6A). Penis slender at apex (Fig. 4D). Right paramere rounded at apex (Fig. 4F)..... ***H. japonicus***
- Lateral surfaces of prosternal process with distinct and wide margins (Fig. 5C). Elytra yellowish colored (Fig. 5A). Penis robust at apex (Fig. 5E). Right paramere acute at apex (Fig. 5F)..... ***H. regimbarti***
- 5 Head, pronotum, and elytra without black mark (Fig. 15A, B). Penis entirely slender, apex acute (Fig. 15D) ***H. eximius***
- Head, pronotum, and elytra with some black marks or entirely black. Penis entirely robust, apex rounded **6**
- 6 Head entirely black (Fig. 7). Black marks on elytra indistinct (Fig. 7)..... ***H. morii* sp. nov.**
- Head partly black or without black mark. Black marks on elytra distinct **7**
- 7 Sutural black band usually reaches main 1st interval on elytral base **8**
- Sutural black band barely reached main 1st interval on elytral base **9**
- 8 Head with black mark (Fig. 17B). Size of eye moderate (Fig. 17B). Black band on elytral base clear in outline (Fig. 17A). Central surface of prosternal process flat in apical half (Fig. 17C) ***H. sharpi***
- Head without black mark (Fig. 12B). Size of eye large (Fig. 12B). Black band on elytral base unclear in outline (Fig. 12A). Central surface of prosternal process slightly depressed on apical half (Fig. 12C) ***H. angustifrons***
- 9 Black mark on head widened at base (Fig. 16B). Black marks on elytra arranged in diamond shape centrally; two black marks on 4th interval, and one black mark on tip of 6th interval (Fig. 16A)..... ***H. ovalis***
- Black mark on head not widened at base. Black marks on elytra are aligned in straight line centrally; single black mark on 4th and 8th intervals **10**
- 10 Black marks on central elytra connected by thin lines; most individuals with black transverse band on elytral base (Fig. 13A). Prosternal process slightly narrowed near procoxa (Fig. 13C) ***H. basinotatus***
- Black mark on central elytra connected by black bands; most individuals without black transverse band on elytral base (Fig. 14A). Prosternal process deeply narrowed near procoxa (Fig. 14C) ***H. kotoshonis***

Discussion

In this paper we recognize 13 species belonging to two genera from Japan. Mori (2007) stated that several unrecorded species of the genus *Haliphus* were known from Nansei shoto and other areas of Japan. These unrecorded species are considered to be *H. regimbarti* (recorded by Iwata 2016), *H. angustifrons* (recorded in this paper), and *H. morii* sp. nov. (described in this paper).

Acknowledgements

We would like to express our sincere gratitude to those who have cooperated greatly in the study of materials: Koji Arai, Takuya Fukuzawa, Hiroaki Iketake, Takashi Kurihara, Masato Mori, Jun Nakajima, Takeshi Sasaki, Daiki Uchida, Kohei Watanabe and Kyohei Watanabe.

Additional information

Conflict of interest

The authors have declared that no competing interests exist.

Ethical statement

No ethical statement was reported.

Funding

No funding was reported.

Author contributions

All authors contributed to this work.

Author ORCIDs

Masakazu Hayashi  <https://orcid.org/0000-0003-3708-9005>

Tomofumi Iwata  <https://orcid.org/0000-0002-7738-3199>

Hiroyuki Yoshitomi  <https://orcid.org/0000-0001-8416-9249>

Data availability

All of the data that support the findings of this study are available in the main text or Supplementary Information.

References

- Aoyagi M (2011a) A list of aquatic insects in Ikema Island, Okinawa, Japan. *Ryukyu no konchu* 35: 95–100. [In Japanese]
- Aoyagi M (2011b) A list of aquatic insects in some islands around Okinawa main Island, Okinawa, Japan. *Ryukyu no konchu* 35: 101–110. [In Japanese]
- Aoyagi M (2014) A new records of aquatic beetles from Iheya Island, Okinawa, Japan. *Ryukyu no konchu* 38: 192–196. [In Japanese]
- Aoyagi M (2019) A list of aquatic insects in Miyagi Island, Okinawa, Japan. *Ryukyu no konchu* 43: 31–35. [In Japanese]
- Aoyagi M (2020) A list of aquatic bugs and aquatic beetles in Irabu Island, Okinawa, Japan. *Ryukyu no konchu* 4: 85–87. [In Japanese]
- Clark H (1863) Descriptions of new East-Asiatic of Haliplidae and Hydroporidae. *The Transactions of the Entomological Society of London* 3: 417–428. <https://doi.org/10.1111/j.1365-2311.1862.tb01288.x>
- Dejima T (2007) A list of Hydradephaga (Coleoptera, Caraboidea) in Kagawa Prefecture. *Kagawa Seibutsu* 34: 83–96. [In Japanese]
- Dejima T (2010) A list of insects collected in Te-jima Island, Kagawa, Japan. *Heriguro* 31: 39–49. [In Japanese]
- Fukuda A, Kurosa K, Hayashi N (1959) Coleoptera. In: Kawada A (Ed.) *Illustrated Insect Larvae of Japan*. Hokuryukan, Osaka, 392–545. [760 pp.] [In Japanese]
- Hayashi M (2015) Aquatic insects of Sanin Region, west Japan I: Coleoptera (part 1). *Special Bulletin of the Hoshizaki Green Foundation* 15: 1–98. [In Japanese]
- Hayashi M, Kadowaki H (2019) Species number survey of Insecta of Oki Islands, Shimane Prefecture, Japan. *Special Bulletin of the Hoshizaki Green Foundation* 23: 1–76. [In Japanese]

- Hayashi M, Ooi K (2022) DNA barcodes of aquatic Adephaga (Coleoptera) in Shimane Prefecture, Japan. Special Bulletin of the Hoshizaki Green Foundation 30: 89–92. [In Japanese]
- Hope FW (1845) On the entomology of China, with descriptions of the new species sent to England by Dr. Cantor from Chusan and Canton. The Transactions of the Entomological Society of London 4(1): 4–17. <https://doi.org/10.1111/j.1365-2311.1845.tb01326.x>
- Imasaka S, Hosoya T, Kokubu K, Ito R, Arima K (2021) A list of Coleoptera collected in Koshiki Islands, Kagoshima, Japan, part 3 (July 2020). Korasana 96: 21–98. [In Japanese]
- Iwata T (2016) A new record of *Haliphus regimbarti* (Coleoptera: Haliplidae) from Japan. Japanese Journal of Systematic Entomology 22: 99–100.
- Kamiya K (1936) Family Gyrinidae Family Haliplidae. Fauna Nipponica 10: 1–55.
- Kamiya K (1951) A list of aquatic Coleoptera collected in Sado, Niigata, Japan. Bulletin of Essa kontyu dokokai 5(3/4): 122–124. [In Japanese]
- Kano T, Kamiya K (1931) Two new species of Haliplidae from Japan. Transactions of the Kansai Entomological Society 2: 1–4. [1 pl.]
- Kitano T, Sano S (2011) Haliplidae larvae eat algae. In: Karube H (Ed.) Suisei Kontyu Daihyakka. Kanagawa Prefectural Museum, Odawara, 48–49. [In Japanese]
- Matsui E (1992) A list of aquatic Coleoptera from Amakusa Islands, Kumamoto Pref., Japan. Bulletin of Kumamoto Kontyu dokokai 38(1): 1–20. [In Japanese]
- Matsuo T, Fukagawa G (2016) Notes on Haliplidae (Coleoptera) in Nagasaki Prefecture, Kyusyu, Japan. Nagasaki-ken Seibutsu Gakkaishi 78: 49–57. [In Japanese]
- Mita T, Kojima H, Okajima S (2015) Name-bearing types of Coleoptera (Insecta) in the Museum of Agriculture, Tokyo University of Agriculture. Journal of Agricultural Science, Tokyo University of Agriculture 60: 50–56. [In Japanese]
- Mitamura T, Hirasawa K, Yoshii S (2017) The Handbook of Japanese Aquatic Insects (Vol. 1): Coleoptera. Bun-ichi Sogo Shuppan, Tokyo, 352 pp. [In Japanese]
- Mori M (2007) Haliplidae. In: Nakane T, Obayashi K, Nomura S, Kurosawa Y (Eds) Iconographia Insectorum Japonicorum Color Naturali Edita (new edition, Vol. 2). Hokuryukan, Tokyo, 61–62. [pl. 28.] [In Japanese]
- Nakajima J, Hayashi M, Ishida K, Kitano T, Yoshitomi H (2020) Aquatic Coleoptera and Hemiptera of Japan. Bun-ichi Sogo Shuppan, Tokyo, 352 pp. [In Japanese]
- Nakamine A, Nakamine K (2021) Records of aquatic insects collected on Tanegashima Island in 2020. Bulletin of the Kagoshima Prefectural Museum 40: 1–8. [In Japanese]
- Nakane T (1963a) Haliplidae. In: Nakane T, Obayashi K, Nomura S, Kurosawa Y (Eds) Iconographia Insectorum Japonicorum Color naturali edita (Vol. 2). Hokuryukan, Tokyo, 55 pp. [pl. 28.] [In Japanese]
- Nakane T (1963b) New or little-known Coleoptera from Japan and its adjacent regions. XVI–XXII. Fragmenta Coleopterologica 4–12: 18–48.
- Nakane T (1985) A revisional study on Japanese species of the genus *Haliphus* Latreille (Coleoptera, Haliplidae). Kita-Kyushu no Konchu 32: 61–67. [pl. 6–7.] [In Japanese]
- Nakane T (1987) The beetles of Japan (new series) 80. Insects & Nature 22(11): 26–30. [In Japanese]
- Nakane T (1990) The beetles of Japan (new series) 91. Insects & Nature 25(10): 22–23. [In Japanese]
- Nakanishi K (2012) Biological notes on *Haliphus sharpi* Wehncke, 1880 (Coleoptera: Haliplidae). Elytra. New Series 2: 107–108.
- Régimbart M (1892) Insectes du Bengale Occidental. Annales de la Société Entomologiques de Belgique 36: 112–121.

- Satô M (1984) Taxonomic notes on the aquatic Coleoptera of Japan, I. Coleopterists' News 65: 1–4. [In Japanese]
- Satô M (1985) Haliplidae. In: Ueno S, Kurosawa Y, Satô M (Eds) The Coleoptera of Japan in Color (Vol. II). Hoikusha, Osaka, 180–181. [pl. 33.] [In Japanese]
- Satô M, Yoshitomi H (2018) Coleoptera. In: Kawai T, Tanida K (Eds) Aquatic insects of Japan: Manual with Keys and Illustrations (2nd edn.). Tokai University Press, Hadano, 707–790. [1730 pp.] [In Japanese]
- Sharp D (1873) The water beetles of Japan. The Transactions of the Entomological Society of London 1873: 45–67. <https://doi.org/10.1111/j.1365-2311.1873.tb00636.x>
- Sharp D (1884) XXI. The water-beetles of Japan. The Transactions of the Entomological Society of London 32: 439–464. <https://doi.org/10.1111/j.1365-2311.1884.tb01616.x>
- Sheth SD, Ghatge NV, Vondel BJ (2016) Illustrated redescription of *Haliphus* (*Liaphlus*) *arrowi* Guignot, 1936 (Coleoptera: Haliplidae) from the Western Ghats, India, and notes on the closely related *H. angustifrons* Régimbart, 1892. Zootaxa 4127(2): 355–364. <https://doi.org/10.11646/zootaxa.4127.2.7>
- Stephens JE (1828) Illustrations of British Entomology. Mandibulata (Vol. 2). Baldwin & Cradock, London, 200 pp. <https://doi.org/10.5962/bhl.title.50040>
- Takahashi T (1998) A list of Coleoptera collected in Awaji Island, Hyogo, Japan. Parnassius 47: 1–10. [In Japanese]
- Takizawa M (1931) The Haliplidae of Japan. Insecta Matsumurana 5(3): 137–143.
- Tomisawa A (2012) A list of aquatic insects collected in Noto Island, Ishikawa, Japan. Tokkuribachi 80: 41–43. [In Japanese]
- Uchida D, Furuhashi R, Kitano T, Sano S, Watanabe K (2021) Distributional records of aquatic Coleoptera from Kakeroma-jima Island, the Amami Islands, the Ryukyus, Japan. Elytra. New Series 11: 183–185.
- van Vondel BJ (1991) Revision of the Palaearctic species of *Haliphus* subgenus *Liaphlus* Guignot (Coleoptera: Haliplidae). Tijdschrift voor Entomologie 134: 75–144.
- van Vondel BJ (1992) Revision of the Palaearctic and Oriental species of *Peltodytes* Régimbart (Coleoptera: Haliplidae). Tijdschrift voor Entomologie 135: 275–297.
- van Vondel BJ (1993) Revision of the *Liaphlus* species of the Oriental Region excluding China (Coleoptera: Haliplidae). Tijdschrift voor Entomologie 136: 289–316.
- van Vondel BJ (1995) Haliplidae: Review of the Haliplidae of China. In: Jäch MA, Ji L (Eds) Water Beetles of China 1: 111–154.
- van Vondel BJ (1998) Haliplidae: Additional notes on the Haliplidae of China and neighboring countries (Coleoptera). In: Jäch MA, Ji L (Eds) Water Beetles of China 2: 131–136.
- van Vondel BJ (2003a) Haliplidae. In: Löbl I, Smetana S (Eds) Catalogue of Palaearctic Coleoptera (Vol. 1). Apollo Books, Vester Skerninge, 30–33.
- van Vondel BJ (2003b) Haliplidae: II. Additional notes on Chinese Haliplidae, with description of a new species. In: Jäch MA, Ji L (Eds) Water Beetles of China 3: 289–294.
- van Vondel BJ (2017) Five new synonymies in *Haliphus* subgen. *Liaphlus* Guignot, based on the variability of the left paramere (Coleoptera: Haliplidae). Koleopterologische Rundschau 87: 31–35.
- van Vondel BJ (2019) Features of the metacoxal air-storage space as additional characters for reconstructing the phylogeny of Haliplidae (Coleoptera). Tijdschrift voor Entomologie 162(1): 13–32. <https://doi.org/10.1163/22119434-20192081>
- van Vondel BJ, Holmen M, Petrov PN (2006) Review of the Palaearctic and Oriental species of the subgenus *Haliphus* s.str. (Coleoptera: Haliplidae: Haliphus) with descriptions of three new species. Tijdschrift voor Entomologie 149(2): 227–273. <https://doi.org/10.1163/22119434-900000203>

- Watanabe R (2021) Record of *Haliphus basinotatus* Zimmermann from Hokkaido, Japan. Sayabane, New Series 41: 16–17. [In Japanese]
- Watanabe K, Hidaka K (2013) Seasonal occurrence of adult *Haliphus sharpi* Wehncke in paddy fields of southwestern Shikoku, Japan. Hozen Seitai-gaku Kenkyu 18: 101–105. [In Japanese with English abstract]
- Watanabe R, Ohba S (2022) New records of four species of aquatic insects (Halipidae, Dytiscidae, Nepidae) from Tsushima Island, Japan. Nagasaki-ken Seibutsu Gakkaishi 91: 34–35. [In Japanese]
- Watanabe R, Yamasaki S (2020) Discovery of larvae of *Haliphus kamiyai* Nakane, 1963 (Coleoptera, Halipidae) and implications for its life cycle. Elytra. New Series 10(2): 365–371.
- Wehncke E (1880) Neue *Haliphus*. Stettiner Entomologische Zeitung 41: 72–75.
- Yamazaki S, Hirasawa K, Sano S (2020) Knowledge of larval and pupal period of *Haliphus eximius*. Sayabane, New Series 40: 17–18. [In Japanese]
- Yoshimura S (1932) On a new species of *Haliphus* from Japan. Mushi 5: 102–103.
- Yoshitomi H (2014) Water beetle fauna of Izu Islands. Sayabane, New Series 16: 26–31. [In Japanese]
- Zaitzev P (1908) Berichtigungen und Zusätze zu den Halipidae, Dytiscidae und Gyrinidae in den neuesten Katalogen der Coleopteren. Russkoe Entomologisches Obozrenie 7: 114–124.
- Zimmermann A (1924) Die Halipliden der Welt. Entomologische Blätter (für Biologie und Systematik der Käfer) 20(1): 1–16. [(2): 65–80; (3): 129–144; (4): 193–213.]

Supplementary material 1

The list of specimens examined in this study

Authors: Masakazu Hayashi, Tomofumi Iwata, Hiroyuki Yoshitomi

Data type: table (.xlsx)

Copyright notice: This dataset is made available under the Open Database License (<http://opendatacommons.org/licenses/odbl/1.0/>). The Open Database License (ODbL) is a license agreement intended to allow users to freely share, modify, and use this Dataset while maintaining this same freedom for others, provided that the original source and author(s) are credited.

Link: <https://doi.org/10.3897/zookeys.1168.99302.suppl1>