

A survey of *Pireneitega* from Tajikistan (Agelenidae, Coelotinae)

Xiaoqing Zhang¹, Yuri M. Marusik^{2,3,4}

1 College of Life Science, Shenyang Normal University, Shenyang, Liaoning 110034, China **2** Institute for Biological Problems of the North RAS, Portovaya Str. 18, Magadan 685000, Russia **3** Department of Zoology & Entomology, University of the Free State, Bloemfontein 9300, South Africa **4** Zoological Museum, University of Turku, FI-20014 Turku, Finland

Corresponding author: Yuri M. Marusik (yurmar@mail.ru)

Academic editor: S. Li | Received 13 September 2016 | Accepted 26 October 2016 | Published 23 November 2016

<http://zoobank.org/59A928AF-4609-484D-BF3D-6D59AC314BD6>

Citation: Zhang X, Marusik YM (2016) A survey of *Pireneitega* from Tajikistan (Agelenidae, Coelotinae). ZooKeys 635: 89–107. doi: 10.3897/zookeys.635.10487

Abstract

Five new species of *Pireneitega* species from Tajikistan are described: *P. zonsteini* **sp. n.** (♂♀), *P. muratovi* **sp. n.** (♀), *P. tyurai* **sp. n.** (♀), *P. ramitensis* **sp. n.** (♀) and *P. kovblyuki* **sp. n.** (♂). *Pireneitega major* (Kroneberg, 1875) is redescribed for the first time based on the lectotype designated here. DNA barcodes for the five new species are documented for future use and as proof of molecular differences between these species.

Keywords

Aranei, central Asia, description, new species, *Paracoelotes*, redescription, spider, taxonomy

Introduction

Coelotinae is the largest subfamily of Agelenidae with more than 650 species distributed in the Holarctic and southeast Asia (World Spider Catalog 2016). *Pireneitega* Kishida, 1955 with 30 species distributed across the Palaearctic (World Spider Catalog 2016, Zhang et al. 2016) is one of the most species-rich genera of the subfamily. It is relatively well studied in comparison to other species-rich (and non-monophyletic) genera *Coelotes* Blackwall, 1841 and *Draconarius* Ovtchinnikov, 1999. The species

of *Pireneitega* found in Caucasus and Xinjiang were recently revised (Kovblyuk et al. 2013; Zhang et al. 2016) but the genus remains poorly studied in Central Asia. Of three species known from central Asia (Mikhailov 2013: *P. birulai* (Ermolajev, 1927) (currently considered a junior synonym of *P. luctuosa* (L. Koch, 1878)), *P. fedotovi* (Charitonov, 1946) and *P. major* (Kroneberg, 1875)), *P. fedotovi* is known only from the original description and *P. major* only from two very short descriptions supplied with sketchy figures. A short trip by the junior author to Tajikistan revealed five new morphospecies of *Pireneitega*, each separated by distinct genetic gaps. The goal of this paper is to provide descriptions of the new species (including records of their molecular markers) and a redescription of *P. major* whose type locality lies in northern Tajikistan.

Material and methods

Specimens were examined and measured with a Leica M205C stereomicroscope. Images were captured with an Olympus C7070 wide zoom digital camera (7.1 megapixels) mounted on an Olympus SZX12 dissecting microscope. Epigynes and male palps were examined after dissection. Epigynes were cleared by boiling it in 10% KOH solution before taking photos of the dorsal view. All measurements are given in millimeters. *Pireneitega major* was photographed and drawn using an MBS-9 stereomicroscope with Pro-MicroScancamera. Leg measurements are given as: total length (femur, patella + tibia, metatarsus, tarsus).

Terminology used for copulatory organ characters in the text and figure legends follows Wang (2002) with some modifications.

Abbreviations used in the text and figure legends are:

A	epigynal atrium;
ALE	anterior lateral eye;
AME	anterior median eye;
AME-ALE	distance between AME and ALE;
AME-AME	distance between AME and AME;
ALE-PLE	distance between ALE and PLE;
CD	copulatory ducts;
CF	cymbial furrow;
CO	conductor;
d	dorsal;
E	embolus;
EB	embolic base;
ET	epigynal teeth;
FD	fertilization ducts;
Fe	femur;

H	epigynal hood;
MA	median apophysis;
Mt	metatarsus;
p	prolateral;
PA	patellar apophysis;
Pa	patella;
PLE	posterior lateral eye;
PME	posterior median eye;
PME-PLE	distance between PME and PLE;
PME-PME	distance between PME and PME;
R	receptacle;
r	retrolateral;
RTA	retrolateral tibial apophysis;
ST	subtegulum;
T	tegulum;
Ta	tarsus;
Ti	tibia;
v	ventral;
VTA	ventral tibial apophysis.

DNA barcodes were obtained for future use: a partial fragment of the mitochondrial gene cytochrome oxidase subunit I (COI) was amplified and sequenced for five new species using Primers LCO1490-oono (5'-CWACAAAYCATARRGATATTGG-3') (Folmer et al. 1994; Miller et al. 2010) and HCO2198-zz (5'-TAAACTTCCAGGT-GACCAAAAAATCA-3') (Folmer et al. 1994; Zhao & Li 2016). For additional information on extraction, amplification, and sequencing procedures, see Zhao et al. (2013). All sequences were blasted in GenBank; accession numbers are provided in Table 1.

Holotypes and some paratypes will be deposited in the Zoological Museum of the Moscow State University (ZMMU). Most paratypes are deposited in the Institute of Zoology, Chinese Academy of Sciences (IZCAS) in Beijing, China.

Table 1. Voucher specimen information.

Species	GenBank accession number	Sequence length	Collection localities
<i>P. zonsteini</i> sp. n.	KY024475	642bp	Env. of Dushanbe, Hissar Mt. Ridge 48 th km of Varzob Hwy
<i>P. muratovi</i> sp. n.	KY024477	642bp	Env. of Dushanbe Hissar, Mt. Ridge 20 th km of Varzob Hwy Gusgarf Vill.
<i>P. tyurai</i> sp. n.	KY024478	642bp	Khatlon Area Khovaling Distr., Obimazar River
<i>P. ramitensis</i> sp. n.	KY024476	642bp	Khatlon Area Hissar Mt. Range Ramit Reserve
<i>P. kovblyuki</i> sp. n.	KY024474	642bp	Tajikstan: Khatlon Area Dangara Distr Sanglogh

Taxonomy

Genus *Pireneitega* Kishida, 1955

Pireneitega Kishida, 1955: 11. Type species *Amaurobius roscidus* L. Koch, 1868 (= *P. segestriformis* (Dufour, 1820)) from Germany.

Paracoelotes Brignoli, 1982: 347. Type species *Coelotes armeniacus* Brignoli, 1978 from Turkey.

Note. *Pireneitega* was long considered a *nomen nudum* (Yaginuma, in Brignoli 1983: p. 468). Kishida (1955), in a general survey of Agelenidae, considered *Pireneitega* to have been described by himself in 1928, although he had no publications that year. The genus "*Pireneitega* Kishida, 1928 [Genotype: *roscida* (Koch, 1868)]" was considered among the tribe Tegenariini Kishida, 1928 (Kishida 1955: p. 11). Although eye pattern was mentioned in the key to the genera of "Tegenariini", Kishida (1955) did not provide a formal description of the genus. Brignoli (1982) described *Paracoelotes* (type species *Coelotes armeniacus* Brignoli, 1978) from Turkey. Subsequently, Wang and Jäger (2007) revalidated *Pireneitega* with *Paracoelotes* as a junior synonym.

Diagnosis. The chelicerae in most species of *Pireneitega* (including the type species) have 3 promarginal and 3 retromarginal teeth; other coelotines have either 2 or 4 retromarginal teeth (Zhang et al. 2016). The females can be distinguished by the widely separated epigynal teeth, the large atrium with subparallel margins, and the broad copulatory ducts (Fig. 2A–B); other coelotines usually have a small atrium and copulatory ducts. The males can be distinguished by the absence of a dorsal "apophysis" on the conductor, the small RTA, and the distinct median apophysis (Fig. 1A–C); other coelotines usually have a broad dorsal apophysis on the conductor and a reduced or indistinct median apophysis.

Composition. Thirty species of *Pireneitega* are known from Spain to Sakhalin (World Spider Catalog 2016; Mikhailov 2013). One species, *P. major*, was known from Tajikistan before the current study.

Pireneitega zonsteini sp. n.

<http://zoobank.org/1AF265B6-AAB0-4974-A8A7-94A906F8FBBF>

Figs 1–2, 8

Type material. Holotype ♂ (ZMMU): Tajikistan, environs of Dushanbe, Hissar Mt. Range, 48th km of Varzob Hwy, S exposed slope with *Juglans* litter & under stones, 38°55'31"N, 68°48'18"E, 1530 m, 7.05.2015 (Y.M. Marusik, M. Saidov). Paratypes: 1♂1♀ (IZCAS), same data as holotype.

Etymology. The species is named after Sergei Zonstein (University of Tel-Aviv, Israel) a partner of the junior author in the expedition to Tajikistan; noun (name) in genitive case.




Figure 1. Male palp of *Pireneitega zonsteini* sp. n., holotype. **A** Prolateral **B** Ventral **C** Retrolateral. Scale bar 0.1 mm.

Diagnosis. The male can be distinguished from all other *Pireneitega* species except *P. involuta* (Wang et al., 1990) by having a broad conductor and thick patellar apophysis. From *P. involuta* it is distinguished by the blunt tip of the patellar apophysis (*vs* a tapering tip in *P. involuta*) (Fig. 1; Wang et al. 1990: figs 13–15). The female can be distinguished from all other *Pireneitega* species except *P. fedotovi* by having a nearly trapezoidal atrium, long copulatory ducts, and short receptacles. From *P. fedotovi* it can be distinguished by its short epigynal teeth, about 0.5 times as long as length of the atrium (*vs* long epigynal teeth in *P. fedotovi*, about as long as the length of the atrium) (Fig. 2; Charitonov 1946: fig. 4).

Description. Male (holotype): Total length 8.90. Carapace 4.40 long, 3.50 wide. Abdomen 4.50 long, 2.80 wide. Eye sizes and interdistances: AME 0.15, ALE 0.20, PME 0.15, PLE 0.20; AME-AME 0.07, AME-ALE 0.06, PME-PME 0.15, PME-PLE 0.18. Leg measurements: I: 12.95 (3.50, 4.30, 3.15, 2.00); II: 12.25 (3.25, 4.00, 3.00, 2.00); III: 10.40 (3.15, 3.00, 3.25, 1.00); IV: 16.00 (4.50, 5.00, 4.25, 2.25). Carapace greenish, the radial grooves indistinct, with black lateral margins. Abdomen blackish, with yellow herringbone pattern.




Figure 2. *Pireneitega zonsteini* sp. n., female paratype and male holotype. **A** Epigyne, ventral **B** Vulva, dorsal **C** Male habitus, dorsal **D** Female habitus, dorsal **E** Female habitus, ventral. Scale bars equal for **D, E**.

Spinination in male

	Fe	Pt	Ti	Mt	Ta
I	3d 2p 1r	–	3-3v	3-3v	–
II	3d 1p 1r	–	2p 3-3v	2p 3-3v	–
III	3d 2p 2r	1p 1r	1d 2p 2r 3-3v	2p 2r 3-3v	–
IV	3d 2p 1r	1p 1r	2p 2r 3-3v	2p 2r 3-3v	–

Palp as in Fig. 1: patellar apophysis long, more than half length of tibia; tibia short, about 1/4 length of tarsus; VTA subequal to the tibial length, without pointed tip, extending beyond the tibia; RTA short, about 1/6 length of VTA; cymbial furrow long, more than half length of cymbium; conductor broad and with two spiraling loops; median apophysis broad and nearly triangular; embolus with broad base originating proximally on base of tegulum.

Female (paratype): Total length 10.0. Carapace 4.75 long, 3.65 wide. Abdomen 5.25 long, 3.45 wide. Eye sizes and interdistances: AME 0.20, ALE 0.25, PME 0.21, PLE 0.26; AME-AME 0.08, AME-ALE 0.05, PME-PME 0.17, PME-PL 0.20. Leg measurements: I: 12.50 (3.75, 4.25, 2.75, 1.75); II: 11.75 (3.50, 4.00, 2.75, 1.50); III: 10.60 (3.00, 3.50, 2.60, 1.50); IV: 15.00 (4.25, 4.75, 4.00, 2.00). Carapace yellow. Abdomen black, with yellow spots and herringbone pattern.

Epigyne as in Fig. 2A–B: epigynal teeth narrow and relatively short (shorter than width of atrium); septum short with weakly sclerotized tip, about 0.3 times as long as wide; atrium with well delimited posterior margin, about 1.3 times longer than wide, about 4 times longer than septum, subequal to width of septum; copulatory opening hidden by anterior margin of atrium; receptacles long, about 2 times longer than wide, separated by 2.5 times their diameters; copulatory ducts with 3 parts, the basal part running from receptacle posteriorly (*Bd*), median part running anteriorly (*Md*), and terminal part (*Td*) running posteriorly and leading to copulatory opening; median part as wide as terminal and 2 times longer than basal part; median part 1.5 times longer than receptacle; median parts touching each other; hoods indistinct.

Spinination in female

	Fe	Pt	Ti	Mt	Ta
I	3d 2p 1r	–	3-3v	3-3v	–
II	3d 1p 1r	–	1p 3-3v	1p 3-3v	–
III	3d 1p 2r	1p 1r	2p 2r 3-3v	2p 2r 3-3v	–
IV	3d 1p 1r	1p 1r	2p 2r 3-3v	1p 2r 3-3v	–

Distribution. Known only from the type locality (Fig. 8).

***Pireneitega muratovi* sp. n.**

<http://zoobank.org/A01FC654-273B-4E50-A278-052B957FBA4B>

Figs 3, 8

Type material. Holotype ♀ (ZMMU): Tajikistan: env of Dushanbe, Hissar Mt. Ridge, 20th km of Varzob Hwy, Gusgarf [Gushharf] Vill., N exposed slope with *Acer* litter & cliffs, 38°44'22"N, 68°47'33"E, 1750 m, 8.05.2015, Y. M. Marusik. Paratype: 1 ♀ (IZCAS), same data as holotype.

Etymology. The species is named after Tajikistan zoologist Rustam Muratov (Dushanbe, Tajikistan) who was very helpful in organizing the expedition to Tajikistan; noun (name) in genitive case.

Diagnosis. The female can be distinguished from all other *Pireneitega* species except *P. fedotovi*, *P. luniformis* (Zhu & Wang, 1994), and *P. major* by having narrow epigynal teeth and an elongate oval atrium. It can be distinguished from *P. fedotovi* by the pointed tip of septum (*vs* blunt tip in *P. fedotovi*), from *P. luniformis* by the elongate oval receptacles (*vs* spiralled in *P. luniformis*), and from *P. major* by its short epigynal teeth, *ca.* 0.8 times as long as length of the atrium (*vs* long epigynal teeth in *P. major*, about as long as the length of the atrium) (Figs 3, 7; Charitonov 1946: fig. 4; Zhu and Wang 1994: figs 5–6).

Description. Male: unknown.

Female (holotype): Total length 9.94. Carapace 4.49 long, 3.05 wide. Abdomen 5.45 long, 2.90 wide. Eye sizes and interdistances: AME 0.18, ALE 0.23, PME 0.24, PLE 0.30; AME-AME 0.10, AME-ALE 0.05, PME-PME 0.15, PME-PL 0.10. Leg measurements: I: 11.25 (3.25, 4.00, 2.50, 1.50); II: 10.30 (3.00, 3.50, 2.50, 1.30); III: 9.70 (2.75, 3.00, 2.65, 1.30); IV: 13.75 (3.75, 4.25, 4.00, 1.75). Carapace yellow, the radial grooves indistinct. Abdomen whitish-yellow, with green herringbone pattern.

Epigyne as in Fig. 3A–B: epigynal teeth narrow, their length equal to width of the narrowest part of the atrium; septum with well delimited tip, *ca.* 0.5 times as long as wide; copulatory opening distinct; atrium with well delimited posterior margin, about 1.4 times longer than wide, *ca.* 2 times longer than and 0.7 times as wide as septum; receptacles long, about 2.5 times as long as wide, bases of receptacles separated by 2 diameters; copulatory ducts with 3 parts, median part as long as receptacles, and anterior part slightly wider than receptacles; hoods indistinct.

Spination

	Fe	Pt	Ti	Mt	Ta
I	3d 2p 1r	–	3-3v	1p 3-3v	–
II	3d 3p 2r	–	2p 3-3v	3-3v	–
III	3d 3p 2r	1p 1r	2p 2r 3-3v	5p 4r 3-3v	1p 1r
IV	3d 1p 1r	1d 1p 1r	1d 2p 2r 3-3v	5p 5r 3-3v	2p 1r

Distribution. Known only from the type locality (Fig. 8).




Figure 3. *Pireneitega muratovi* sp. n., female holotype. **A** Epigyne, ventral **B** Vulva, dorsal **C** Habitus, dorsal **D** Habitus, dorsal **E** Habitus, ventral view. Scale bars equal for **C**, **D**, **E**.

***Pireneitega tyurai* sp. n.**

<http://zoobank.org/B14F37A9-6A33-446F-80AF-2C65472362D3>

Figs 4, 8

Type material. Holotype ♀ (ZMMU): Tajikstan: Khatlon Area, Khovaling Distr., Obimazar River, Sultan-Mazar, clay cliffs, 38°28'19"N, 70°04'01"E, 1854 m, 27.04.2015 (Y.M. Marusik). Paratypes: 4♀ (IZCAS), same data as holotype.

Etymology. The species is named after Sergei V. Tyura (Magadan, Russia) a friend of the junior author; noun (name) in genitive case.

Diagnosis. The female can be distinguished from all other *Pireneitega* species except *P. tianchiensis* (Wang et al., 1990) by having short receptacles and copulatory ducts. It can be distinguished from *P. tianchiensis* by the broad and long epigynal teeth, about 0.85 times as long as atrium (*vs* short and narrow epigynal teeth in *P. tianchiensis*, about 0.5 times as long as atrium) (Fig. 4A–B; Wang et al. 1990: figs 84–85).

Description. Male: unknown.

Female (holotype): Total length 5.15. Carapace 2.15 long, 1.75 wide. Abdomen 3.00 long, 2.00 wide. Eye sizes and interdistances: AME 0.10, ALE 0.13, PME 0.15, PLE 0.15; AME-AME 0.05, AME-ALE 0.10, PME-PME 0.02, PME-PLE 0.04. Leg measurements: I: 6.20 (1.90, 2.25, 1.25, 0.80); II: 5.10 (1.60, 1.75, 1.00, 0.75); III: 4.80 (1.50, 1.60, 1.00, 0.70); IV: 7.05 (2.05, 2.50, 1.50, 1.00). Carapace yellow, with black lateral margins. Abdomen blackish, with yellow herringbone pattern.

Epigyne as in Fig. 4A–B: epigynal teeth long (nearly as long as atrium); septum with weakly sclerotized tip, about 0.5 times as long as wide; atrium with weakly sclerotized posterior margin, about 0.7 times as long as wide, about 1.8 times longer than and 0.7 times as wide as septum; copulatory opening hidden; receptacles large, *ca.* 2 times longer than wide; copulatory ducts with two parts, terminal parts (Tp) not touching each other, about 0.5 length of receptacles, basal parts (Bp) shorter than width of receptacle; hoods indistinct.

Spination

	Fe	Pt	Ti	Mt	Ta
I	3d 2p	–	3-3v	3-3v	–
II	3d 1p 1r	1p	2p 3-3v	1p 3-3v	–
III	3d 1p 1r	1p 1r	2p 2r 3-3v	5p 4r 3-3v	2p 1r
IV	2d 1p 1r	1p 1r	2p 2r 3-3v	5p 4r 3-3v	2p 1r

Distribution. Known only from the type locality (Fig. 8).

***Pireneitega ramitensis* sp. n.**

<http://zoobank.org/C74C6BAE-DE7C-4A95-A4A2-5E5BFC45C341>

Figs 5, 8

Type material. Holotype ♀ (ZMMU): Tajikstan: Khatlon Area, Hissar Mt. Range, Ramit Reserve, 38°44'36"N, 69°18'30"E, 1324 m, 1.05.2015 (Y.M. Marusik). Paratypes: 4♀ (IZCAS), 2♀ (ZMMU), same data as holotype.




Figure 4. *Pireneitega tyurai* sp. n., female holotype. **A** Epigyne, ventral **B** Vulva, dorsal **C** Habitus, dorsal **D** Habitus, dorsal **E** Habitus, ventral. Scale bars equal for **C, D, E**.




Figure 5. *Pireneitega ramitensis* sp. n., female holotype. **A** Epigyne, ventral **B** Vulva, dorsal **C** Habitus, dorsal **D** Habitus, ventral **E** Habitus, ventral. Scale bars equal for **C**, **D**, **E**.

Etymology. The specific name is an adjective and refers to the type locality; adjective.

Diagnosis. The female can be distinguished from all other *Pireneitega* species except *P. muratovi* sp. n., *P. fedotovi*, *P. luniformis* and *P. major*, by having an elongate oval atrium, narrow epigynal teeth, and long copulatory ducts. It can be distinguished from *P. muratovi* sp. n. and *P. luniformis* by the narrow tip of the copulatory ducts (*vs* round tip in *P. muratovi* sp. n. and *P. luniformis*) and from *P. fedotovi* and *P. major* by the bent epigynal teeth (*vs* straight epigynal teeth in *P. fedotovi* and *P. major*) (Figs 3, 5, 7; Charitonov 1946: fig. 4; Zhu & Wang 1994: figs 5–6).

Description. Male: unknown.

Female (holotype): Total length 12.00. Carapace 4.50 long, 3.55 wide. Abdomen 7.50 long, 4.75 wide. Eye sizes and interdistances: AME 0.20, ALE 0.23, PME 0.25, PLE 0.20; AME-AME 0.10, AME-ALE 0.20, PME-PME 0.10, PME-PL 0.23. Leg measurements: I: 14.05 (4.00, 4.75, 3.45, 1.85); II: 13.40 (3.90, 4.50, 3.25, 1.75); III: 13.00 (3.75, 4.25, 3.25, 1.75); IV: 16.55 (4.75, 5.40, 4.40, 2.00). Carapace yellowish, with brown lateral margins. Abdomen pale-yellow, with brown spots.

Epigyne as in Fig. 5A–B: epigynal teeth pale, hyaline, long and thin, about 0.9 times as long as receptacles; septum with weakly sclerotized tip, *ca.* 0.5 times as long as wide, nearly triangular; copulatory ducts distinct; atrium about 1.4 times longer than wide, with well delimited posterior margin, *ca.* 2.8 times longer than and about as wide as septum; receptacles large, about 3 times longer than wide; receptacle bases separated by about 2 diameters; copulatory ducts with 3 parts, basal part about 2/3 of receptacle length, median part as long as receptacle, terminal part somewhat shorter than median part; hoods distinct.

Spination

	Fe	Pt	Ti	Mt	Ta
I	3d 2p 1r	–	1p 3-3v	1p 3-3v	–
II	3d 2p 2r	–	2p 3-3v	2p 3-3v	–
III	3d 3p 2r	1p 1r	2p 2r 3-3v	5p 4r 3-3v	2p 2r
IV	3d 2p 1r	1p 1r	2p 2r 3-3v	5p 4r 3-3v	2p 2r

Distribution. Known only from the type locality (Fig. 8).

***Pireneitega kovblyuki* sp. n.**

<http://zoobank.org/25787234-B768-4EB3-B6B2-781E025AB5D4>

Figs 6, 8

Type material. Holotype ♂ (ZMMU): Tajikistan, Khatlon Area, Dangara Distr., Sanglogh (=Sanglok) Mt. Range above Shar-Shar Pass, 38°17'56"N, 69°13'36"E, 1700–2060 m, 29.04.2015, (Y.M. Marusik). Paratypes: 3♂ (IZCAS), 2♂ (ZMMU), same data as holotype.

Etymology. The specific name is a patronym in honour of the well known arachnologist and friend of the junior author Mykola M. Kovblyuk (Simferopol, Ukraine); noun (name) in genitive case.




Figure 6. Male palp of *Pireneitega koblyuki* sp. n., holotype. **A** Prolateral **B** Ventral **C** Retrolateral. Scale bar 0.1 mm.

Diagnosis. The male can be distinguished from all other *Pireneitega* species except *P. tianchiensis* by having a hook-shaped conductor, and narrow cymbium. It can be distinguished from *P. tianchiensis* by the short cymbial furrow, *ca.* 1/10 length of cymbium (*vs* long cymbial furrow in *P. tianchiensis*, about 0.5 length of cymbium) (Fig. 6; Wang et al. 1990: figs 81–83).

Description. Male (holotype): Total length 7.90. Carapace 4.00 long, 3.00 wide. Abdomen 3.90 long, 2.65 wide. Eye sizes and interdistances: AME 0.15, ALE 0.20, PME 0.18, PLE 0.19; AME-AME 0.08, AME-ALE 0.07, PME-PME 0.13, PME-PLE 0.15. Leg measurements: I: 10.90 (3.25, 4.05, 2.00, 1.60); II: 9.85 (3.00, 3.50, 2.00,

1.35); III: 8.60 (2.75, 2.50, 2.10, 1.25); IV: 12.55 (3.70, 3.75, 3.50, 1.60). Carapace yellow, the radial grooves indistinct. Abdomen pale, with yellow herringbone pattern.

Palp as in Fig. 6A–C: patellar apophysis absent; tibia long, *ca.* 0.5 length of cymbium; VTA short and wide, about 1/3 length of tibia, without pointed tip; RTA short, about 1/5 length of VTA, poorly visible; cymbium long, its tip as long as or longer than genital bulb; conductor short, with hook-shaped, partially looped tip, tip located distally from tegulum; median apophysis broad and nearly triangular; embolus with broad, nearly tongue-shaped base, beginning at 6:30 o'clock position.

Spination

	Fe	Pt	Ti	Mt	Ta
I	3d 2p 1r	–	3-3v	3-3v	–
II	3d 3p 1r	1p	2p 3-3v	3p 3-3v	–
III	3d 2p 2r	1d1p 1r	1d 2p 2r 3-3v	5p 5r 3-3v	1p 1r
IV	3d 1p 1r	1p 1r	2p 2r 3-3v	5p 5r 3-3v	2p 1r

Female: Unknown.

Distribution. Known only from the type locality (Fig. 8).

Pireneitega major (Kroneberg, 1875)

Figs 7–8

Coelotes major Kroneberg, 1875: 15, pl. 1, fig. 6 (♀); Charitonov, 1946: 20, fig. 5 (♀).

Paracoelotes major: Ovtchinnikov, 1988: 142 (transferred from *Coelotes*).

Misidentifications:

Coelotes major: Schenkel 1936: 284, fig. 97 (♀); Hu & Wu 1989: 180, fig. 150.1–2 (♀).

Paracoelotes major: Song et al. 1999: 389, fig. 229Q–R (♀).

Material examined. Lectotype ♀ (ZMMU) with label «Ta 3845 1♀ ZMMU [Зоомузеті МГУ]» «Lectotypus» 2/VI; Аучи дагана [Auchi dagana] *Coelotes major* Kroneberg, 1875», ca 39°35'N, 69°05'E. Paralectotype: 1♀ (ZMMU) with 2 labels «Ta1059, 1, *Coelotes major*» «Туркестанская Учёная Экспедиция Императорскаго Общества Любителей Естествознания. Федченко [Turkestan Scientific Expedition of the Emperor's Society of Devotees of Natural Sciences. Fedchenko]» and «*Coelotes major* n. sp. Ta, No.1059, Кокангское ханство, Федченко [Kokand Khanate, Fedchenko]».

Comments. The figures of *P. major* presented by Schenkel (1936), Hu and Wu (1989), and Song et al. (1999; copied from Hu and Wu 1989) are of a species other than *P. major*, the identity of which is currently unknown. All records of this unknown species are from Xinjiang, China.

Diagnosis. This species is easily distinguished from other species of *Pireneitega* found in Tajikistan by its larger size (carapace length >6 mm *vs* <4.75) and having 5 spines on tarsus IV (*vs* other species with 0–4). The epigyne of *P. major* is most similar




Figure 7. Epigyne of *Pireneitega major*, lectotype. **A** Ventral **B** Dorsal.




Figure 8. Localities of *Pireneitega* species from Tajikistan. **1** *P. zonsteini* sp. n. **2** *P. muratovi* sp. n. **3** *P. tyurai* sp. n. **4** *P. ramitensis* sp. n. **5** *P. kovblyuki* sp. n. **6** *P. major*.

to that of *P. muratovi* sp. n. and *P. ramitensis* sp. n. It can be distinguished from *P. muratovi* sp. n. by its shorter receptacles with length/width ratio of 2.3 (*vs* 2.6 in *P. muratovi*), shape of copulatory ducts, and shorter teeth (cf. Figs 3A–B and 7A–B). *Pireneitega major* can be separated from *P. ramitensis* sp. n by its wider epigynal atrium and shorter, wider receptacles as well as by its shorter and wider copulatory ducts (cf. Figs 5A–B and 7A–B).

Description. Male: unknown.

Female: Lectotype. Total length 16.7. Carapace 7.0 long, 5.0 wide, fovea 1.25 long. Leg measurements: I: 19.75 (5.5, 2.5, 4.6, 4.65, 2.5); II: 18.6 (5.1, 2.5, 4.0, 4.5, 2.5); III: 17.2 (4.75, 2.2, 3.55, 4.6, 2.1); IV: 21.85 (5.75, 2.3, 5.0, 6.25, 2.55).

Spination

	Fe	Pt	Ti	Mt	Ta
I	3d 2p 2r	–	3-3v	3-3v 1vm	–
II	3d 3p 2r	–	2p 3-3v	1p 3-3v	–
III	3d 3p 2r	1p 1(0)r	2p 2r 3-3v	5p 4r 3-3v	2p 1-1v
IV	3d 2p 1r	1p 1r	2p 2r 3-3v	5p 4r 3-3v	2p 3r

Paralectotype ♀. Total length: 11.0. Carapace 6.0 long, 4.0 wide. Epigyne 0.51 wide.

Epigyne as in Fig. 7: epigynal teeth pale, hyaline, long and thin; septum with weakly sclerotized tip, about 0.4 times as long as wide, subtriangular; atrium as long as wide; receptacles large, about 2.5 times longer than wide; receptacle bases separated by *ca.* 2 diameters; copulatory ducts with 2 parts, basal part as long as receptacle, terminal part somewhat shorter receptacle.

Comments. Known from the type series females only. Exact locality is known for the lectotype only: Auchi lies on the northern macroslope of the Turkestan Mt Range (Fig. 8).

Acknowledgements

We thank Sergei Zonstein (Tel-Aviv, Israel; a partner of the junior author in the expedition to Tajikistan), Rustam Muratov and Murod Saidov (Dushanbe, Tajikistan) for field assistance and helping to organize the expedition; Kirill G. Mikhailov (Moscow, Russia) for allowing us to study the type material of *P. major*; Seppo Koponen (Turku, Finland) for providing museum facilities in Turku; and Zhe Zhao (Beijing, China) for helping to provide the molecular data. English of the final draft was kindly checked by Robert Bennett (Victoria, British Columbia, Canada). This study was supported by the National Natural Sciences Foundation of China to Guo Zheng (NSFC-31172121, 31372224, 31672315).

References

Brignoli PM (1982) On a few spiders from China (Araneae). Bulletin of the British Arachnological Society 5: 344–351.

- Brignoli PM (1983) A catalogue of the Araneae described between 1940 and 1981. Manchester University Press 468: 1–755.
- Charitonov DE (1946) New forms of spiders of the USSR. *Izvestiya Yestestvenno-Nauchnogo Instituta pri Molotovskom Gosudarstvennom Universitete imeni M. Gor'kogo* 12: 19–32. [In Russian]
- Folmer O, Black M, Hoeh W, Lutz R, Vrijenhoek R (1994) DNA primers for amplification of mitochondrial cytochrome c oxidase subunit I from diverse metazoan invertebrates. *Molecular Marine Biology and Biotechnology* 3(5): 294–299.
- Hu JL, Wu WG (1989) Spider from agricultural regions of Xinjiang Uygur Autonomous Region, China. Shandong University Publishing House, Jinan, 435 pp.
- Kishida K (1955) A synopsis of spider family Agelenidae. *Acta Arachnologica (Tokyo)* 14: 1–13. doi: 10.2476/asjaa.14.1
- Kovbyluk MM, Kastrygina ZA, Marusik YM, Ponomarev AV (2013) The spider genus *Pireneitega* Kishida, 1955 in the Caucasus (Aranei: Agelenidae: Coelotinae). *Arthropoda Selecta* 22: 59–73.
- Kroneberg A (1875) Araneae. In: Fedtschenko AP (Ed.) *Puteshestvie v Tourkestan*. *Izvestiya Imperatorskogo Obshchestva Lyubitelei Yestestvoznaniya* 19: 1–58. [In Russian]
- Mikhailov KG (2013) The spiders (Arachnida: Aranei) of Russia and adjacent countries: a non-annotated checklist. *Arthropoda Selecta*, Supplement 3: 1–262.
- Miller JA, Carmichael A, Ramirez MJ, Spagna JC, Haddad CR, Řezáč M, Johannesen J, Král J, Wang XP, Griswold CE (2010) Phylogeny of entelegyne spiders: affinities of the family Penestomidae (new rank), generic phylogeny of Eresidae, and asymmetric rates of change in spinning organ evolution (Araneae, Araneioidea, Entelegynae). *Molecular Phylogenetics and Evolution* 55: 786–804. doi: 10.1016/j.ympev.2010.02.021
- Ovtchinnikov SV (1988) [Materials on spider fauna of the superfamily Amaurobioidea of Kirghizia]. *Entomologiceskie issledovanija v Kirgizii* 19: 139–152. [In Russian]
- Schenkel E (1936) Schwedisch-chinesische wissenschaftliche Expedition nach den norwestlichen Provinzen Chinas. *Arkiv för Zoologi* 29(A1): 1–314.
- Song DX, Zhu MS, Chen J (1999) *The Spiders of China*. Hebei University of Science and Technology Publishing House, Shijiazhuang, 640 pp.
- Wang JF, Yin CM, Peng XJ, Xie LP (1990) New species of the spiders of the genus *Coelotes* from China (Araneae: Agelenidae). *Spiders in China: One Hundred New and Newly Recorded Species of the Families Araneidae and Agelenidae*. Hunan Normal University Press, 172–253.
- Wang XP (2002) A generic-level revision of the spider subfamily Coelotinae (Araneae, Amaurobioidea). *Bulletin of the American Museum of Natural History* 269: 1–150. doi: 10.1206/0003-0090(2002)269<0001:AGLROT>2.0.CO;2
- Wang XP, Jäger P (2007) A revision of some spiders of the subfamily Coelotinae F. O. Pickard-Cambridge 1898 from China: transfers, synonymies, and new species (Arachnida, Araneae, Amaurobioidea). *Senckenbergiana Biologica* 87: 23–49.
- World Spider Catalog (2016) World Spider Catalog. Natural History Museum Bern. <http://wsc.nmbe.ch> [accessed on October 10, 2016]

- Zhang XQ, Zhao Z, Zheng G, Li S (2016) Nine new species of the spider genus *Pireneitega* Kishida, 1955 (Agelenidae, Coelotinae) from Xinjiang, China. *ZooKeys* 601: 49–74. doi: 10.3897/zookeys.601.7893
- Zhao Z, Li S (2016) *Papiliocoelotes* gen. n., a new genus of Coelotinae (Araneae, Agelenidae) spiders from the Wuling Mountains, China. *ZooKeys* 585: 33–50. doi: 10.3897/zookeys.585.8007
- Zhao Z, Su TJ, Chesters D, Wang SD, Ho SYW, Zhu CD, Chen XL, Zhang CT (2013) The mitochondrial genome of *Elodia flavipalpis* Aldrich (Diptera: Tachinidae) and the evolutionary timescale of tachinid flies. *PLoS ONE* 8: e61814.
- Zhu CD, Wang JF (1994) Seven new species of the genus *Coelotes* from China (Araneae: Agelenidae). *Acta Zootaxonomica Sinica* 19: 37–45.