

# A new species of *Urophora* Robineau-Desvoidiy, 1830 (Diptera, Tephritidae) from Iran

Saeed Mohamadzade Namin<sup>1,†</sup>, Jamasb Nozari<sup>2,‡</sup>

**1** Department of Plant Protection, Faculty of Agriculture, Varamin-Pishva branch, Islamic Azad University, Varamin, Iran **2** Department of Plant Protection, Faculty of Agriculture, University of Tehran, Karaj, Iran

† [urn:lsid:zoobank.org:author:9063C413-5D65-44B7-9B94-E17684660E96](http://urn:lsid:zoobank.org:author:9063C413-5D65-44B7-9B94-E17684660E96)

‡ [urn:lsid:zoobank.org:author:D70429BD-0695-4A8A-8793-D84E45FF8C28](http://urn:lsid:zoobank.org:author:D70429BD-0695-4A8A-8793-D84E45FF8C28)

Corresponding author: Saeed Mohamadzade Namin ([mohamadzade@iauvaramin.ac.ir](mailto:mohamadzade@iauvaramin.ac.ir))

---

Academic editor: R. Meier | Received 13 August 2011 | Accepted 3 November 2011 | Published 8 December 2011

---

[urn:lsid:zoobank.org:pub:F30D708A-9B83-44F0-8DAE-F8163650EE71](http://urn:lsid:zoobank.org:pub:F30D708A-9B83-44F0-8DAE-F8163650EE71)

---

**Citation:** Mohamadzade Namin S, Nozari J (2011) A new species of *Urophora* Robineau-Desvoidiy, 1830 (Diptera, Tephritidae) from Iran. ZooKeys 152: 63–70. doi: 10.3897/zookeys.152.1911

---

## Abstract

*Urophora merzi* **sp. n.** reared from flower heads of *Centaurea behen* Linnaeus is described from Iran. It is similar to *U. campestris*, *U. sachalinensis*, *U. stylata*, *U. tsoii* and *U. vera* in wing pattern with 3 well developed crossbands and indistinct subbasal crossband, differing in aculeus tip with two pairs of diminished preapical steps and different host plants.

## Keywords

Tephritidae, *Urophora*, new species, Iran

## Introduction

The genus *Urophora* Robineau-Desvoidiy, 1830 with about 60 species is one of the largest genera of the family Tephritidae in the Palearctic Region (Norrbom et al. 1999). All species of known biology are associated with asteraceous plants and induce galls in their flower heads and, rarely, stems (White and Korneyev 1989). Some *Urophora* species are agents for biological control of astraceous weeds; *U. affinis* (Frauenfeld), *U.*

*cardui* (Linnaeus), *U. quadrifasciata* (Meigen), *U. sirunaseva* (Hering), *U. solstitialis* (Linnaeus) and *U. stylata* (Fabricius) successfully introduced to the Nearctic Region for biocontrol of weeds (Peschen and Harris 1975; Turner et al. 1994; Turner 1996 a, b; Wheeler and Stoops 1996).

While studying the tephritid flies fauna in Iran in 2008–2011 seasons, we collected and reared a previously undescribed species that infests the flower heads of *Centaurea behen* L. (Asteraceae). The new species is described and figured below.

## Material and methods

Materials were collected by standard sweeping net and rearing from flower heads of *Centaurea behen*. Morphological terminology follows White et al. (1999). The material examined minuten-pinned on side and deposited in collections of the following institutions:

- JAZM** Jalal Afashar Zoological Museum, College of Agriculture, University of Tehran, Karaj, Iran.  
**MHNG** Museum d'histoire naturelle, Genève, Switzerland.  
**SIZK** I. I. Schmalhausen Institute of Zoology, National Academy of Sciences of Ukraine, Kiev, Ukraine.  
**ZISP** Zoological Institute, Russian Academy of Sciences, St. Petersburg, Russia.

The following morphometric characters with their abbreviations are used: Body length (BL); wing length (WL); aculeus length (AL).

## Results

### Key to Western Palearctic species of *Urophora* with the *stylata*-like wing pattern (3 distinct crossbands, of them, apical and subapical fused)

(Corresponding to the couplet 94 in Korneyev and White 1999)

- |   |   |  |
|---|---|--|
| 1 | Apex of aculeus with two pairs of indistinct steps (Fig. 6); associated with <i>Centaurea behen</i> .....   | <b><i>U. merzi</i> sp. n.</b>                      |
| – | Apex of aculeus with 1–2 pairs of prominent primary steps. ....   | <b>2</b>   |
| 2 | Apex of aculeus with 1 pair of prominent primary steps .....  | <b>3</b>   |
| – | Apex of aculeus with 2 pairs of prominent, sharp steps .....  | <b>(see couplet 97 in Korneyev and White 1999)</b> |
| 3 | Apex of first flagellomere slightly pointed. Aculeus width between primary steps almost equal to distance from primary steps to apex (see Korneyev and White 1996: Fig. 21). Larvae in <i>Serratula</i> flower head galls. Armenia..... | <b><i>U. vera</i> Korneyev &amp; White</b>         |

- First flagellomere apically rounded. Aculeus apex between primary steps almost twice as wide as its length from primary steps level to tip (see Korneyev and White 1996: Fig. 17). Larvae in *Cirsium* flower head galls. Whole Europe and Western Asia to West Siberia and western China. ... *U. stylata* Fabricius

***Urophora merzi* Mohamadzade Namin, sp. n.**

urn:lsid:zoobank.org:act:2A468C69-CE7C-4C69-BE53-06168D739915

[http://species-id.net/wiki/Urophora\\_merzi](http://species-id.net/wiki/Urophora_merzi)

Figs 1–15


**Type material.** Holotype (female): **Iran:** Mazandaran Province, Haraz road, 10 km north east Abali, 35°50'N; 51°58'E, h 2360m, swept from flower heads of *Centaurea behen*, 20 May, 2011, S. Mohamadzade Namin leg. (JAZM).

Paratypes: 1 ♀, same collection data as in holotype, reared from flower heads of *Centaurea behen* Linnaeus, collected 13 September, 2008 & emerged 22 September 2008; 1 ♂, 1 ♀, Alburz Province, Chaloos road, Nesa, 36°04'N; 51°19'E, h 2200m, 22 June 2009, swept from *Centaurea behen*; 15 ♂, 18 ♀, same collection data as in holotype, 20 May, 2011, S. Mohamadzade Namin leg. (JAZM; some paratypes are deposited also in MHNG, SIZK, ZISP and first author's personal collection).

**Description. Head:** Yellow, except ocellar triangle, occiput and slender part of arista black. Length: height: width ratio = 1: 1: 1.25. Frons brown; face whitish yellow; Antenna yellow, scape with blackish setulae at dorso-apical margin; first flagellomere light yellow, 1.6 times as long as wide and distinctly rounded antro-ventrally; arista bare. Compound eye about as high as long. Gena 1.1 times as high as length of first flagellomere. Proboscis capitate with black setae. Two frontal and one orbital setae present. Postocellar, postocular, vertical and genal setae black and acuminate. Frons with black setulae around frontal setae (Fig. 4).


**Thorax:** General color black; mesonotal scutum densely covered with gray microtrichia and black setulae. Notopleura shining black. Pleuron black; only stripe in anterior half of anepisternum and postpronotal lobe yellow. Scutellum yellow; slightly convex, corners of scutellum black. Subscutellum and mediotergite black. All setae on thorax black and acuminate. Scutellum with 4 equal black setae; basal setae placed in yellow area. Halter yellow.

**Wing:** Hyaline with 3 well developed dark brown crossbands. Subbasal band reduced and only present as darkening near apex of cell bm and rarely bcu. Discal crossband complete, crossing wing from pterostigma through R-M crossvein into posterior margin. Preapical crossband complete, reaching posterior margin. Apical band well developed. In females, preapical and apical crossbands in 56.2% of type material fused in cell  $r_1$  (Fig. 2), in 31.2% fused in  $r_1$  and  $r_{2+3}$  (Figs 1, 11) and in 12.5% fused in  $r_1$ ,  $r_{2+3}$  and anterior half of  $r_{4+5}$  cells (Fig. 3). In males, preapical and apical crossbands in



**Figures 1–3.** *Urophora merzi* sp. n., 1 wing pattern of the holotype 2–3 variation of wing pattern in paratypes.

33.3% of specimens fused in cell  $r_1$ , in 50% fused in  $r_1$  and  $r_{2+3}$  and in 16.6% fused in  $r_1$ ,  $r_{2+3}$  and anterior half of  $r_{4+5}$  cells. In one male of type series discal and preapical crossbands narrowly joined in  $r_1$  cell and in one female and one male discal and preapical crossbands narrowly connected at posterior margin of wing. Pterostigma yellowish.



**Figures 4–8.** *Urophora merzi* sp. n., **4** head in profile **5** aculeus **6** aculeus tip **7** male terminalia **8** epandrium.


Distance between crossveins about 1.4 as long as dm-cu crossvein.  $R_{4+5}$  with 1 setula ventrally at node.

**Legs:** Completely yellow; fore femur in 60% of females and 55% of males with black stripe in dorsal side. All setae and setulae blackish (Figs 9, 10). Fore femur with two dorsal and one ventral rows of setae.

**Abdomen:** General color black, sparsely microtrichose, subshining with black setulae. Posterior margin of abdominal tergites, especially tergites 5–6 with long black setae. Oviscape 1.25 times as long as preabdomen, shining black with black hairs. Aculeus narrow, 11 times as long as wide, apically rounded, apex with two pairs of indistinct steps, as in Figs 5, 6, 13, 14. Tergite 5 of males as long as two preceding tergites with long setae in posterior margin. Epandrium as in Figs 8, 12 and glans as in Figs 7, 15.

**Measurements:** Male: BL = 3.5–4 mm (average 3.8), WL = 3.5–4.5 mm (average 3.9); female: BL = 4.5–6 mm (average 5.3), WL = 4–4.9 mm (average 4.3), AL = 1.5–2 mm (average 1.9) (n = 5).

**Etymology.** The species is named in honour of Dr Bernhard Merz, an outstanding Swiss dipterist, in recognition of his invaluable contribution into study of the order Diptera, especially family Tephritidae.



**Figures 9–15.** *Urophora merzi* sp. n., **9** ♂, total view, left **10** ♀ (Holotype), total view, left **11** wing pattern (Holotype) **12** aculeus **13** aculeus tip **14** epandrium **15** male terminalia.

**Discussion.** The new species is similar to *U. campestris* Ito (Japan), *U. sachalinensis* (Shiraki) (Russia and Japan), *U. stylata* Fabricius (Worldwide), *U. tsoii* Korneyev and White (Russia) and *U. vera* Korneyev and White (Armenia), sharing similar wing pattern (3 well developed crossbands and indistinct subbasal crossband, with apical and preapical crossbands fused along anterior margin of wing), yellow femora and antenna and black



notopleura, differing in the shape of aculeus apex. Apex of aculeus in *U. sachalinensis*, *U. stylata* and *U. vera* has one pair of steps. *U. campestris* and *U. tsoii* (both occurring in the Far East of the Palearctic Region) possess two pairs of distinct steps, whereas the aculeus tip in *U. merzi* sp. n. has two pairs of smoothed, almost indistinct steps. Also the new species is similar to *U. jaculata* Rondani (Italy and Greece), sharing similar aculeus apex and host plants of the genus *Centaurea*, differing in the subbasal crossband strongly reduced to a darkening near bm cell (distinct and reaching  $R_1$  in *U. jaculata*).

All the compared species are associated with different host plants: *U. campestris*, *U. sachalinensis* and *U. stylata* are associated with *Cirsium* spp., *Carduus* spp. and *Galactites tomentosa*; *U. tsoii* and *U. vera* with *Serratula* spp. and *U. jaculata* with *Centaurea solstitialis* (Korneyev and White 1999, 2000) whereas *U. merzi* sp. n. is associated with *Centaurea behen*.

## Acknowledgments

We wish to express our gratitude to Islamic Azad University Varamin-Pishva branch (Grant title: Faunistic study of fruit flies of family Tephritidae in West Azerbaijan and Mazandaran Provinces) for the support of this work. We thank Dr Valery A. Korneyev (Schmalhausen Institute of Zoology, NAS of Ukraine, Kiev, Ukraine) for reading early versions of this manuscript and suggesting many useful remarks. We are grateful to Mr. Masoumi (University of Tehran, Faculty of Horticulture) for identification of the host plant.

## References

- Korneyev VA, White IM (1996) Fruit flies of the genus *Urophora* R.-D. (Diptera, Tephritidae) of Eastern Palearctics. II. Review of species of the subgenus *Urophora* s. str. Communication 3. Entomological Review 76(4): 499–513.
- Korneyev VA, White IM (1999) Tephritids of the genus *Urophora* R.-D. (Diptera, Tephritidae) of East Palearctic: III. Key to palaeartic species. Entomological Review 78(3): 464–482.
- Korneyev VA, White IM (2000) Tephritids of the genus *Urophora* R.-D. (Diptera, Tephritidae) of East Palearctic: IV. Conclusion. Entomological Review 79(1): 239–253.
- Norrbom AL, Carroll LE, Thompson FC, White IM, Freidberg A (1999) Systematic Database of Names. In: Thompson FC (Ed) Fruit Fly Expert Identification System and Systematic Information Database. Myia, 65–299.
- Peschken DP, Harris P (1975) Host specificity and biology of *Urophora cardui* (Diptera: Tephritidae). A biocontrol agent for Canada thistle (*Cirsium arvense*). Canadian Entomologist 107: 1101–1110. doi: 10.4039/Ent1071101-10
- Turner CE, Sobhian R, Joley DB, Coombs EM, Piper GL, Maddox DM (1994) Establishment of *Urophora sirunaseva* (Hering) (Diptera: Tephritidae) for biological control of yellow star-thistle in the western United States. Pan-Pacific Entomologist 70: 206–211.

- Turner CE (1996a) Tephritidae in the biological control of weeds. In: McPherson BA, Steck GJ (Eds) Fruit fly pests: A world assessment of their biology and management. St. Lucie Press, Delray Beach, 157–164.
- Turner CE (1996b) Tephritid flies in the biological control of yellow starthistle. In: McPherson BA, Steck GJ (Eds) Fruit fly pests: A world assessment of their biology and management. St. Lucie Press, Delray Beach, 171–176.
- Wheeler AG, Stoops CA (1996) Establishment of *Urophora affinis* on spotted knapweed in Pennsylvania, with new eastern U.S. records of *U. quadrifasciata* (Diptera: Tephritidae). Proceedings of the Entomological Society of Washington 98: 93–99.
- White IM, Korneyev VA (1989) A revision of the western Palearctic species of *Urophora* Robineau-Desvoidy (Diptera: Tephritidae). Systematic Entomology 19(3): 327–374. doi: 10.1111/j.1365-3113.1989.tb00289.x
- White IM, Headrick DH, Norrbom AL, Carroll LE (1999) Glossary. In: Aluja M, Norrbom AL (Eds) Fruit Flies (Tephritidae): Phylogeny and Evolution of Behaviour. CRC Press, Boca Raton, London, New York, Washington DC, 881–924. doi: 10.1201/9781420074468.sec8